

# Nutrient digestibility and digestive tract development of the Kampong-crossbreed chicken fed *Salvinia molesta* leaf meal

*by* Binti Ma'rifah

---

**Submission date:** 04-Aug-2021 12:58PM (UTC+0700)

**Submission ID:** 1627592697

**File name:** Turnitin-Jurnal\_LRRD\_Binti\_Marifah.pdf (81.4K)

**Word count:** 1644

**Character count:** 8492

# Nutrient digestibility and digestive tract development of the Kampong-crossbreed chicken fed *Salvinia molesta* leaf meal

Binti Ma'rifah and Sugiharto Sugiharto

Department of Animal Science, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Central Java Indonesia (50275) [marifahbinti2908@gmail.com](mailto:marifahbinti2908@gmail.com)

## Abstract

The objective of the present study was to investigate the effect of dietary inclusion of *Salvinia molesta* on the nutrient digestibility and digestive tract development of the Kampong-crossbreed chicken. Four experimental diets with different levels of *Salvinia molesta* leaf meal i.e., 0, 6, 12 and 18% were formulated. The diets were offered to 100 of the 3-weeks-old Kampong-crossbreed chickens with the average body weight of  $218 \pm 0.54$  g. The birds were slaughtered at 10 weeks of age. Results showed that dietary inclusion of *Salvinia molesta* increased the feed and protein digestibility of chickens. The relative weights of proventriculus, gizzard, intestine, duodenum and cecum were lower in the treated birds than in control. The same condition was observed for the relative lengths of proventriculus, intestine, duodenum, ileum and cecum. The duodenal villi was higher in the birds fed *Salvinia molesta* than in control. Dietary inclusion of *Salvinia molesta* improved growth performance and feed conversion ratio of chickens. The treatments also lowered feed cost per gain of Kampong-crossbreed chickens. In conclusion, *Salvinia molesta* as a feed ingredient improved nutrient digestibility, duodenal villi and growth performance of the Kampong-crossbreed chickens.

Keywords: crossbreed chicken, gastrointestinal tract, performance

## Introduction

Kampong-crossbreed chicken has recently attracted huge interest from the consumers due to its unique taste and low cholesterol content in meat as compared to broiler meat (Nugraha et al 2016). To response the continuously increasing demand, farmers are now endeavoring to maximize the growth performance while reducing the cost of feed. The development and morphology of the intestine are the key factors for the growth and wellbeing of chickens (Yamauchi and Tarachai 2000; Sugiharto 2016). In general, the digestive tract of chickens begins to grow shortly after hatching and continue thereafter. A number of factors have been reported to affect the development of the chicken's digestive tract, one of which is the feed (Röhe 2013). Protein is generally considered as a vital nutrient for the normal growth and development of the gastrointestinal tract of chickens (Laudadio et al 2012). To meet the protein requirement, soybean meal has commonly been included in the chicken ration. Due to its high price, it is therefore necessary to find alternative other than soybean meal as a source of protein for chickens. One of the local cheap stuff that may be exploited as an alternative to soybean meal for the chicken is *Salvinia molesta*.

### *Salvinia molesta*

Is a free-floating aquatic plant that is abundantly grown in swamp, rivers and rice fields. According to McFarland et al (2004), the yearly production of *Salvinia molesta* can reach 45.6 to 109.5 t/ha. *Salvinia molesta* may be used as a feed ingredient for chickens as it contains of 12.9% crude protein (Leterme et al 2009). *Salvinia molesta* is also rich in essential minerals and amino acids needed for the growth of chickens. Nonetheless, *Salvinia molesta* has a high content of crude fiber (around 16.8%) (Adrizal 2002), which may limit the use of such stuff in the chicken rations. Apart from the latter concern, dietary fiber has been suggested to stimulate the growth and development of the digestive tract (Yu and Chiou 1996; Sumiati and Sumirat 2003) as well as improve the structure and morphology of the intestine of chickens (Oliveiera et al 2008). For these reasons, feeding *Salvinia molesta* was expected not only to substitute the protein-rich feedstuffs such as soybean meal, but also to improve the development of digestive tract of chickens and thus performances of the chickens. The objective of the present study was

to investigate the effect of dietary inclusion of *Salvinia molesta* on the nutrient digestibility and digestive tract development of the Kampong-crossbreed chicken.

means and standard deviation.

## Results and discussion

Data on the nutrient digestibility of the Kampong-crossbreed chicken fed *Salvinia molesta* leaf meal are presented in Table 3. Our present finding showed that dietary inclusion of *Salvinia molesta* increased the feed and protein digestibility of chickens. It has commonly been known that well-balanced essential amino acids is coupled with the improved nutrient digestibility in poultry. Previously, Beski et al (2015) pointed out that vegetable (plant)-based protein contain unbalanced essential amino acids. Conversely, aquatic ferns such as *Salvinia molesta* contain well-balanced essential amino acids (Leterme et al 2009). For this reason, the dietary incorporation of *Salvinia molesta* may therefore improve the balanced amino acids and thereby increased the feed and protein digestibility in the Kampong-crossbreed chickens. In addition to the amino acid balance, the increased content of fiber in the *Salvinia molesta* treated groups may also increase the digestibility of chicks due to the improvement in intestinal morphology, i.e., increased duodenal villi height, of chicks in the present study. The latter inference was supported by de Vries (2015) suggesting that moderate levels of fiber may improve the gastrointestinal development and function and thus nutrient digestibility and growth performance of chickens.

Table 3. Nutrient digestibility of Kampong-crossbreed chickens

Items	<i>Salvinia molesta</i>			
	0%	6%	12%	18%
Feed digestibility (%)	14.11 <sup>c</sup>	34.65 <sup>b</sup>	37.56 <sup>b</sup>	52.99 <sup>a</sup>
Protein digestibility (%)	82.06 <sup>c</sup>	83.73 <sup>b</sup>	83.95 <sup>b</sup>	86.01 <sup>a</sup>

<sup>a,b</sup> Means in a row with different superscripts are significantly different ( $p < 0.05$ )

Data on the relative weights of digestive tract and the height of duodenal villi are presented in Table 4. The relative weights of proventriculus, gizzard, intestine, duodenum and cecum were lower in the birds fed *Salvinia molesta* meal than in control. The same condition was observed for the relative lengths of proventriculus, intestine, duodenum, ileum and cecum. Since the data of the gastrointestinal size were presented as relative to the final body weight, the lower measurements of the gastrointestinal tract of the treated chickens seemed to be associated with the much greater final body weight of the treated chickens than that of the control (Table 5). Note that feeding *Salvinia molesta* meal increased the absolute weight of the small intestine of birds in the present study (the absolute weights of small intestine of the chickens fed *Salvinia molesta* meal of 0, 6, 12 and 18% were 18.51, 24.53, 27.10 and 25.51 g, respectively).

In this present study, the duodenal villi was higher in the birds fed *Salvinia molesta* than in control. The improved morphology of villi was most likely attributed to the higher fiber content of *Salvinia molesta* than of control, as de Vries (2015) and Alvarez et al (2007) reported that villi tended to increase with the increased content of fiber in the diets. The improved morphology of villi eventually results in expanded surface area for the nutrient absorption which is beneficial for the chickens (Lenhardt and Mozes 2003; Sarikhan et al 2010).

Table 4. Relative measurement of the gastrointestinal tract of Kampong-crossbreed chickens

Items	Salvinia molesta			
	0%	6%	12%	18%
<b>Relative weights</b>				
Proventriculus (%)	3.63 <sup>a</sup>	2.80 <sup>a</sup>	2.83 <sup>bc</sup>	2.68 <sup>c</sup>
Intestine (%)	3.63 <sup>a</sup>	2.80 <sup>b</sup>	2.83 <sup>b</sup>	2.68 <sup>b</sup>
Gizzard (%)	4.04 <sup>a</sup>	3.22 <sup>a</sup>	3.06 <sup>b</sup>	3.13 <sup>ab</sup>
Duodenum (%)	1.15 <sup>A</sup>	0.69 <sup>B</sup>	0.73 <sup>B</sup>	0.70 <sup>B</sup>
Jejunum (%)	1.27	1.14	1.16	1.18
Ileum (%)	1.21	0.97	0.95	0.80
Cecum (%)	1.01 <sup>A</sup>	0.66 <sup>B</sup>	0.58 <sup>B</sup>	0.55 <sup>B</sup>
Large intestine (%)	0.38	0.30	0.30	0.31
Pancreas (%)	0.38	0.40	0.37	0.37
<b>Relative lengths</b>				
Proventriculus (cm/100gBW)	0.89 <sup>A</sup>	0.60 <sup>B</sup>	0.50 <sup>B</sup>	0.56 <sup>B</sup>
Intestine (cm/100 g BW)	25.3 <sup>A</sup>	17.5 <sup>B</sup>	13.4 <sup>B</sup>	15.2 <sup>B</sup>
Duodenum (cm/100 g BW)	6.17	7.38	9.64	6.17
Jejunum (cm/100 g BW)	9.62 <sup>A</sup>	6.76 <sup>B</sup>	5.10 <sup>B</sup>	6.17 <sup>B</sup>
Ileum (cm/100 g BW)	5.16 <sup>A</sup>	3.78 <sup>B</sup>	3.04 <sup>B</sup>	3.30 <sup>B</sup>
Cecum (cm/100 g BW)	1.70	3.17	0.96	1.09
<b>Duodenal villi</b>				
Diameter (µm)	259	248	287	272
Height (µm)	52.4 <sup>B</sup>	47.9 <sup>B</sup>	57.1 <sup>B</sup>	78.4 <sup>A</sup>

<sup>A,B</sup> Means in a row with different superscripts are significantly different ( $p < 0.01$ )

<sup>a,b,e</sup> Means in a row with different superscripts are significantly different ( $p < 0.05$ )

Taken together, the increased height of intestinal villi and the reduced relative size of gastrointestinal tract (which may reduce the energy need for maintenances and thus save the energy for growth) eventually result in improved growth performance of the chickens. Indeed, the growth performance and FCR were better in the chickens fed *Salvinia molesta* meal than in control. Moreover, feeding *Salvinia molesta* meal resulted in lower feed cost per gain (Table 5). In the present study, dietary inclusion of *Salvinia molesta* reduced the use of soybean meal, which had much higher price compared to *Salvinia molesta*, in the rations of Kampong-crossbreed chicken. This consequently lowered the price of feed per kilogramme. Overall, the improved growth performance of chickens (with feeding *Salvinia molesta*) and the lowered feed price per kilogramme may simultaneously lowered the feed cost per gain of Kampong-crossbreed chickens in the current study.

Table 5. Final body weight and feed cost of Kampong-crossbreed chickens

Items	Salvinia molesta			
	0%	6%	12%	18%
Final body weight (g/bird)	511 <sup>B</sup>	876 <sup>A</sup>	960 <sup>A</sup>	952 <sup>A</sup>
Feed cost (IDR/kg feed)	17,977	16,261	15,520	14,816
Feed cost/gain (IDR/kg body weight)	35,152 <sup>a</sup>	18,565 <sup>b</sup>	16,171 <sup>b</sup>	15,569 <sup>b</sup>

<sup>A,B</sup> Means in a row with different superscripts are significantly different ( $p < 0.01$ )

<sup>a,b</sup> Means in a row with different superscripts are significantly different ( $p < 0.05$ )

IDR: Indonesian rupiah (Indonesian currency)

## Conclusion

- The use of *Salvinia molesta* as a feed ingredient improved nutrient digestibility, the duodenal villi and growth performance of the chickens.

# Nutrient digestibility and digestive tract development of the Kampong-crossbreed chicken fed *Salvinia molesta* leaf meal

## ORIGINALITY REPORT

5%

SIMILARITY INDEX

%

INTERNET SOURCES

5%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 F. Karadas, V. Pirgozliev \*, T. Acamovic, M.R. Bedford. " The effects of dietary phytase activity on the concentration of Coenzyme Q in the liver of young turkeys and broilers ", *British Poultry Abstracts*, 2005  
Publication 2%
- 2 Y Che Man. "Effect of rosemary and sage extracts on frying performance of refined, bleached and deodorized (RBD) palm olein during deep-fat frying", *Food Chemistry*, 2000  
Publication 1%
- 3 Eero Puolanne, Riitta Kivikari. "Determination of the buffering capacity of postrigor meat", *Meat Science*, 2000  
Publication 1%
- 4 Kissow, Hannelouise, Bolette Hartmann, Jens Juul Holst, Niels-Erik Viby, L rke Schmidt Hansen, Mette Marie Rosenkilde, Kristine Juul Hare, and Steen Seier Poulsen. "Glucagon-like peptide-1 (GLP-1) receptor agonism or DPP-4

inhibition does not accelerate neoplasia in carcinogen treated mice", Regulatory Peptides, 2012.

Publication

---

5

Jessie Fischer Moinuddin, Helen Wing-Tsit Lee. " Alimentary, blood and other changes due to feeding MnSO , MgSO and Na SO ", American Journal of Physiology-Legacy Content, 1960

Publication

---

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On