

# Seasonal and interannual coastal wind variability off the central Maluku Islands revealed by satellite oceanography

*by Anindya Wirasatriya*

---

**Submission date:** 19-Sep-2022 11:58PM (UTC+0700)

**Submission ID:** 1903717092

**File name:** he\_cenral\_Maluku\_Islands\_revealed\_by\_satellite\_oceanography.pdf (1,005.49K)

**Word count:** 4874

**Character count:** 26110

# Seasonal and interannual coastal wind variability off the central Maluku Islands revealed by satellite oceanography

Setiawan R.Y.<sup>1</sup>, Iskandar I.<sup>2\*</sup>, Wirasatriya A.<sup>3,4</sup>, Dwi Susanto R.<sup>5</sup>, Siswanto E.<sup>6</sup>, Pranowo W.S.<sup>7,8</sup>, Setiawati M.D.<sup>9</sup> and Mardiansyah W.<sup>2</sup>

- <sup>1</sup>Department of Fisheries, Faculty of Agriculture, Universitas Gadjah Mada, Yogyakarta, Indonesia
- <sup>2</sup>Department of Physics, Faculty of Mathematics and Natural Sciences, University of Sriwijaya, Indralaya, OI, South Sumatera, Indonesia
- <sup>3</sup>Department of Oceanography, Faculty of Fisheries and Marine Science, Diponegoro University, Semarang, Indonesia
- <sup>4</sup>Center for Coastal Rehabilitation and Disaster Mitigation Studies, Diponegoro University, Semarang, Indonesia
- <sup>5</sup>Department of Atmospheric and Oceanic Science, University of Maryland, College Park, MD, USA
- <sup>6</sup>Earth Surface System Research Center (ESS), Research Institute for Global Change (RIGC), Japan Agency for Marine-Earth Science and Technology (JAMSTEC), Yokohama, Japan
- <sup>7</sup>Marine Research Center, Ministry of Marine Affairs and Fisheries of the Republic of Indonesia, Jakarta, Indonesia
- <sup>8</sup>Department of Hydrography, Indonesia Naval Postgraduate School (STTAL), Jakarta, Indonesia
- <sup>9</sup>Research Center for Oceanography, National Research and Innovation Agency (BRIN), Jakarta, Indonesia

Received: 15/11/2021, Accepted: 04/01/2022, Available online: 05/01/2022  
 \*to whom all correspondence should be addressed: e-mail: iskhaq@mipa.unsri.ac.id

<https://doi.org/10.30955/gnj.004177>

## Graphical abstract




Figure. (Upper-left panel) Maps of the Maluku Islands (blue line), and (lower-left panel) of topography and bathymetry (m) of the central Maluku Islands (red box). MS, HS, and SS denote the Maluku, Halmahera, and Seram Seas. (Right panel) time series of wind stress, SST, chlorophyll-a, and Ekman transport over regions of 124°E-125.5°E, 3°S-4°S (Box A) and 126°E-127.5°E, 1°S-2.5°S (Box B).

## Abstract

The Maluku Islands (henceforth MI) are situated in the northeastern Indonesia. Ocean region off the central MI is

pivotal as it provides a course for the Indonesian Throughflow (ITF) through the Lifamatola passage. However, sea surface dynamics off the central MI is unknown until recently due to inadequate measurements. The current fact motivates the present study to decipher the coastal wind variability off the central MI and its effect on the sea surface by analysing long-term datasets (2007-2019) of satellite-derived sea surface wind, sea surface temperature (SST), and surface chlorophyll-a concentration. Possible influence of extreme climate events of the 2015 El Niño-Southern Oscillation (ENSO) and the 2019 Indian Ocean Dipole (IOD) on all oceanographic parameters was also examined. Results show that the prevailing southeasterly winds over the central MI induce SST cooling and phytoplankton bloom. Correlation analysis revealed that the ENSO and IOD play significant roles in defining spatial distribution of the coastal wind, SST, and phytoplankton bloom in the research area. In addition, the anomaly analysis exhibits distinct oceanographic features during the climate extreme events of 2015 and 2019. Collectively, results of the present research highlight the importance of coastal wind variability and extreme events in shaping the ocean surface characteristics and perhaps regional fisheries production.

**Keywords:** ENSO, IOD, Maluku Islands, sea surface temperature, winds

## 1. Introduction

Sea surface winds play a pivotal role in regulating ocean surface dynamics in the Indonesian Seas. Typically, the northwesterly (southeasterly) winds exert downwelling

(upwelling) in many coastal areas in the Indonesian Seas, as indicated by high (low) sea surface temperature (SST) (Kurniawati *et al.*, 2021; Muskananfola *et al.*, 2021; Setiawan *et al.*, 2019;2020; Siswanto *et al.*, 2020; Wirasatriya *et al.*, 2020a; 2021). Wind variability in the region is principally affected by the Australian Indonesian Monsoon (AIM) system, which generates the prevailing of northwesterly winds from December to February (Austral summer) and the southeasterly winds from June to September (Austral winter) (Gordon, 2005). The northwesterly winds carry moist air and cause heavy rainfall over the Indonesian Maritime Continent, while the southeasterly winds transport dry air and induce dry season over the region (Gordon, 2005). Sea surface condition of the Indonesian Seas is also modulated by the El Niño-Southern Oscillation (ENSO) and the Indian Ocean Dipole (IOD), i.e. the sea surface experiences fierce wind forcing during the El Niño/positive IOD year and weak wind during the La Niña/negative IOD year (Iskandar *et al.*, 2017; Kurniawati *et al.*, 2021; Setiawan *et al.*, 2019; Siswanto *et al.*, 2020; Susanto and Marra, 2005; Susanto *et al.*, 2006; Wirasatriya *et al.*, 2020b). These two climate modes may have exerted an essential role in controlling health of the marine environment and fish resource distribution in the Indonesian Seas. Hence, examining the variability of the AIM winds and their effect on the sea surface is essential for marine and coastal resource management of the Indonesian fisheries management areas.

The MI is situated in the northeastern Indonesia and comprises many small islands. Of particular interest here is the ocean region off the central MI provides a fishing hotspot for commercial fishes (Hutubessy *et al.*, 2014) as well as a pathway for the ITF via the Lifamatola passage (black arrow in Figure 1 right panel) (e.g., Sprintall *et al.*, 2014). The central MI waters are characterized by complex bathymetry and connected to the Maluku and Halmahera Seas in the north and the Banda Sea in the south. Furthermore, the region of interest is located in the Coral Triangle, one of the most diverse places due to its high species richness of coral, demersal fish, and sea turtle (DeVantier *et al.*, 2004; Veron *et al.*, 2006). This biodiverse region has delivered services through seafood resources and tourism for the people and the nation. Unfortunately, physical forcing like monsoon wind that controls sea surface conditions off the central MI is unknown due to lack of long-term *in situ* measurements. Variation of this wind may have played a substantial role in sustaining life of the marine realm in the ocean region off the central MI.

In addition to seasonal forcing, the MI also experiences interannual modulation by the Indo-Pacific climate modes, namely ENSO and IOD. The ENSO is an interannual climate anomaly in the tropical Pacific Ocean (e.g., McPhaden *et al.*, 2006). The El Niño event of 2015 has been suggested as one of the robust El Niño events in the record (Hu and Fedorov, 2017; Palmeiro *et al.*, 2017). It was characterized by considerable positive SST anomaly that dominated the tropical Pacific Ocean (L'Heureux *et al.*, 2017). Levine and McPhaden (2016) suggested the

multiple westerly wind bursts were liable for the advection of water mass from the Western Pacific Warm Pool into the central Pacific, which in turn enhanced the development of a strong El Niño like in 2015.

Meanwhile, the IOD is a climate mode associated with zonal SST gradient (east-west) in the tropical Indian Ocean (Saji *et al.*, 1999). Positive IOD event that occurred during boreal fall 2019 was categorized as an extreme event (Doi *et al.*, 2020; Ratna *et al.*, 2021). The 2019 positive IOD event's primary forcing was thought to be the existence of enhanced sea level pressure over Australia and declined sea level pressure over the South China Sea since May 2019 (Lu and Ren, 2020). This inter-hemispheric gradient of sea level pressure invoked the northward cross-equator flow over the Indonesian Seas and accelerated wind speed over the southeastern tropical Indian Ocean, leading to the evolution of positive IOD events via Kelvin waves forcing (Delman *et al.*, 2016; Lu and Ren, 2020). We hypothesize the two extreme climate events of 2015 and 2019 have profound effects on the Indonesian Seas' oceanic condition. Given the importance of the ocean region off the central MI, disentangling coastal wind variability and the detected effect of both extreme events on the sea surface will contribute to the successful prediction of future climate extreme events in the region.

The development of robust remote sensing technology has enabled us to investigate long-term surface ocean dynamics. Earlier investigations have shown the capability of the Moderate Resolution Imaging Spectroradiometer (MODIS) data in revealing long-term variability of chlorophyll-a and SST regionally and globally (e.g., Mao *et al.*, 2017; Muhammad *et al.*, 2016; Setiawan *et al.*, 2019). In the present work, we analyse 12 years (2007-2019) of satellite-derived sea surface wind, SST, and surface chlorophyll-a to decipher seasonal and interannual variations of the sea surface condition off the central MI. Moreover, we elucidate the effect of the 2015 El Niño and the 2019 positive IOD events on the ocean parameters. Results of this research can be used to manage better marine and fisheries resources in the central MI region.

## 2. Materials and methods

### 2.1. Materials

We employed KNMI Global Wind Level-3 data of spatial resolution of  $0.125^{\circ} \times 0.125^{\circ}$  of the Advanced Scatterometer (ASCAT) to unveil surface wind variation off the central MI. The analysis period of this study goes from January 2007 to December 2019. The ASCAT onboard the Meteorological Operational (MetOp) satellite can acquire wind data (speed and direction) in the coastal area and open ocean with an accuracy of  $\sim 2 \text{ m s}^{-1}$  (Verhoef and Stoffelen, 2013). The ASCAT wind data is available at the Copernicus Marine Environment Monitoring Service ([https://resources.marine.copernicus.eu/product-detail/WIND\\_GLO\\_WIND\\_L3\\_REP\\_OBSERVATIONS\\_012\\_005/SERVICES](https://resources.marine.copernicus.eu/product-detail/WIND_GLO_WIND_L3_REP_OBSERVATIONS_012_005/SERVICES)).

In order to decipher the effect of wind on the ocean surface off the central MI, Level-3 data of 11- $\mu\text{m}$  SST and chlorophyll-a retrieved from the Moderate Resolution

Imaging Spectroradiometer (MODIS) onboard the Aqua satellite were examined. The spatial resolution of the MODIS SST and chlorophyll-a data is  $0.04^\circ \times 0.04^\circ$  (Esaías et al, 1998), and both data have been documented to accurately record SST and chlorophyll concentration (e.g., Ghanea et al., 2016). The MODIS SST and chlorophyll-a data were downloaded from the Physical Oceanography Distributed Active Archive Center of NASA (<https://podaac-tools.jpl.nasa.gov/drive/files/allData/modis/L3/aqua>) and were analysed for the period of January 2007 to December 2019.

The Oceanic Niño Index (ONI) and the Dipole Mode Index (DMI) were used to determine ENSO events (El Niño and La Niña) and IOD events, respectively. Specifically, we used the Niño-3.4 SST anomaly index. The ONI data is generated by the National Ocean and Atmospheric Administration Climate Prediction Center and is accessible at <https://www.cpc.ncep.noaa.gov/data/indices/oni.ascii.txt>. The DMI data is generated by the Japanese Agency for Marine-Earth Science Technology (JAMSTEC) and is attainable at <https://stateoftheocean.osmc.noaa.gov/surind/dmi.php>.

## 2.2. Methods

The wind stress ( $\tau$ ) on the ocean surface and the Ekman Mass Transport (EMT) were calculated by employing the equation of Hsieh and Boer (1992),

$$\tau = \rho_a C_d U_{10}^2 \quad (1)$$

$$EMT_x = \frac{(\delta\tau_x + f\tau_y)}{\rho_w(f^2 + \delta^2)} \quad (2a)$$

$$EMT_y = \frac{(\delta\tau_y - f\tau_x)}{\rho_w(f^2 + \delta^2)} \quad (2b)$$

where  $\rho_a$  is the density of air ( $1.25 \text{ kg m}^{-3}$ ),  $\rho_w$  is the density of seawater ( $1.025 \times 10^3 \text{ kg m}^{-3}$ ),  $C_d$  is the drag coefficient,  $U_{10}$  is the wind speed of 10 m above sea level,  $\delta$  is frictional dumping parameter ( $480^{-1} \text{ days}$ ), and  $f$  is the Coriolis parameter.  $EMT_x$  and  $EMT_y$  express zonal and meridional direction of the EMT, respectively.


All satellite data were averaged into monthly means. The data were then computed into monthly climatological mean by following the equation of Wirasatriya et al. (2017),

$$\bar{X}(x,y) = \frac{1}{n} \sum_{i=1}^n x_i(x,y,t) \quad (3)$$

where  $\bar{X}(x,y)$  is monthly average or monthly climatology value at location  $(x,y)$ ,  $x_i(x,y,t)$  is the  $i$ -th value of the data at  $(x,y)$  and time  $t$ ,  $n$  is the number of data in a one-month period and the number of monthly data in one period of climatology (2007-2019). Note that hollow pixels were excluded in the calculation.


The monthly climatology time series for all oceanographic parameters were constructed by averaging all oceanographic parameters over the region of  $124^\circ\text{E}$ - $125.5^\circ\text{E}$ ,  $3^\circ\text{S}$ - $4^\circ\text{S}$  (Box A), and  $126^\circ\text{E}$ - $127.5^\circ\text{E}$ ,  $1^\circ\text{S}$ - $2.5^\circ\text{S}$  (Box B)

(Box B) to understand better the variations of ocean wind, SST, and surface chlorophyll-a in the region off the central MI. This time series is displayed in Figure 1b.



**Figure 1.** Maps of the Maluku Islands (blue line) and of topography and bathymetry (m) of the central Maluku Islands (red box). MS, HS, and SS denote the Maluku, Halmahera, and Seram Seas.

To examine the effects of the 2015 El Niño and the 2019 positive IOD, we performed monthly anomaly analysis to all oceanographic parameters. Furthermore, correlation analysis between oceanographic parameters and the ONI and DMI was computed to highlight the effect of extreme climate events on the ocean region off the central MI.




**Figure 2.** Time series of (a) wind stress, (b) SST, (c) chlorophyll-a, and (d) Ekman transport over regions of  $124^\circ\text{E}$ - $125.5^\circ\text{E}$ ,  $3^\circ\text{S}$ - $4^\circ\text{S}$  (Box A) and  $126^\circ\text{E}$ - $127.5^\circ\text{E}$ ,  $1^\circ\text{S}$ - $2.5^\circ\text{S}$  (Box B).


## 3. Results

Analyses of monthly climatology of sea surface wind stress over the central MI from December to February, and from June to September are demonstrated in Figure 2a, 3a, and 4a. It is seen that the northeasterly winds prevail over the investigated area from December to February with a magnitude of less than  $0.4 \times 10^{-1} \text{ N m}^{-2}$ . Meanwhile, the

southeasterly wind stresses prevail from June to September with a magnitude of more than  $0.4 \times 10^{-1} \text{ N m}^{-2}$ . Our results show the strongest wind occurred in July and August (Figure 2a).



**Figure 3.** Monthly climatology maps of (a) surface wind stress, (b) SST, (c) surface chlorophyll-a, and (d) EMT during the Austral summer.



**Figure 4.** Monthly climatology maps of (a) surface wind stress, (b) SST, (c) surface chlorophyll-a, and (d) EMT during the Austral winter.

SST off the central MI shows relatively higher SST ( $> 29^\circ\text{C}$ ) from December to February compared to that during the period of June-September (Figure 2b, 3b, and 4b). Indeed, the evolution of SST cooling is prominent during the latter time, with the coldest SST ( $< 28^\circ\text{C}$ ) takes place in August (Figure 2b).

Figure 2c, 3c, and 4c exhibit monthly climatology of surface chlorophyll-a concentration off the central MI during the Austral summer and Austral winter. There is a phytoplankton bloom occurrence off the central MI that

evolved during the Austral winter. We found a chlorophyll-a maximum of  $0.38 \text{ mg m}^{-3}$  in August, in contrast, a low phytoplankton biomass occupies the studied area during the Austral summer as indicated by chlorophyll-a concentration less than  $0.2 \text{ mg m}^{-3}$  (Figure 2c).


In line with the wind stress, strong EMT ( $> 5 \text{ m}^2 \text{ s}^{-1}$ ) observed in the ocean region off the central MI from June to September (Figure 2d, 3d, and 4d). On the other hand, a relatively weak EMT ( $< 4 \text{ m}^2 \text{ s}^{-1}$ ) persisted from December to February. The predominant direction of EMT during June-September (December to February) period is southwestward (northeastward).

#### 4. Discussion


The present research demonstrates that the coastal wind variability off the central MI is dynamically affected by the AIM winds. Climatological maps of surface wind from June to September show intense southeasterly wind stresses ( $> 0.6 \times 10^{-1} \text{ N m}^{-2}$ ) persisted in the area of interest. Simultaneously, there is an elevation in surface chlorophyll-a concentration, while the SST significantly decreases. We postulate coastal upwelling occurred in the region results in a phytoplankton bloom event (Figure 2c, 4b, and 4c). An upwelled of nutrient-rich water from the deeper depth to the surface may invoke phytoplankton bloom events off the central MI from June to September. Our inference is corroborated by the EMT values that denote higher values ( $> 5.5 \text{ m}^2 \text{ s}^{-1}$ ) during the period compared to December-February (Figure 2d, 3d, and 4d). Moreover, our results reveal the interaction between surface winds and central MI topography is apparent here, i.e. the existence of coastal mountains and small straits (indicated by black arrows in Figure 4a) shaped wind regime in the research site. The location and direction of enhanced wind stress and EMT are well fit with the position of significant SST cooling and phytoplankton bloom (Figure 4). Indeed, coastal mountains in the region cause celerity of the southeasterly wind stress via orography mechanism. This observation may imply that the surface ocean productivity and fish resource distribution in the region are likely determined by the variability of the coastal wind jet. We notice that orographic alteration of surface wind is commonplace in seas off the central MI and possibly in other regional seas within the Indonesian archipelago.

The northeasterly wind stresses seem unable to force coastal upwelling off the central MI from December to February. Hence, a phytoplankton bloom event in the investigated area is absent. It is evident here that the phytoplankton bloom diminished when the coastal wind stresses weakened ( $< 0.4 \times 10^{-1} \text{ N m}^{-2}$ ) (Figure 2c and 4a-c). The deceleration of wind forcing in this area also causes SST to increase ( $> 28^\circ\text{C}$ ). Besides the faint wind, we propose solar insolation maxima may promote high SST in the site. Furthermore, there is a high chlorophyll-a spot observed in the southeastern Halmahera Sea. According to Setiawan *et al.* (2020), the chlorophyll-a maxima there are induced by continental runoff, not coastal upwelling.

The present research also examines the ENSO and IOD's effect on the coastal wind variation off the central MI. We focus on the period of June-September due to the robustness of coastal wind stresses during this time. We found a remarkable positive correlation between the ENSO and coastal wind stresses from June to September (Figure 5a). The correlation between the ENSO and EMT yields a similar result (Figure 5d), demonstrating the ENSO strongly modulates the occurrence of phytoplankton bloom off the central MI through Ekman upwelling. This premise is corroborated by a pronounced negative correlation between the ENSO and SST from June to September (Figure 5b-c). Good correlations among oceanographic parameters here suggest the dominant influence of ENSO in governing surface ocean productivity off the central MI during the Austral winter.




**Figure 5.** Correlation between ONI and monthly (a) surface wind stress, (b) SST, (c) surface chlorophyll-a, (d) EMT during the Austral winter.




**Figure 6.** Correlation between DMI and monthly (a) surface wind stress, (b) SST, (c) surface chlorophyll-a, (d) EMT during the Austral winter.

The correlation between IOD and oceanographic parameters from June to September is depicted in Figure

6. Note that the IOD event indicates seasonal phase-lock: develops in boreal late spring/early summer, peaks in late summer/fall, and terminates in early winter. For this reason we performed the correlation analysis during the period of July – September when the IOD reached its peak. Likewise, the effect of IOD on the coastal wind, SST, phytoplankton biomass off the central MI is vigorous. Positive correlations between the IOD and surface wind stresses, phytoplankton bloom, and EMT (Figure 6) are noticeable during the Austral winter. Seemingly, coastal wind stresses are favourable for generating phytoplankton bloom and SST cooling when the IOD operates.



**Figure 7.** Anomaly maps of (a) surface wind stress, (b) SST, (c) surface chlorophyll-a, and (d) EMT during the Austral winter of 2015.



**Figure 8.** Anomaly maps of (a) surface wind stress, (b) SST, (c) surface chlorophyll-a, and (d) EMT during the Austral winter of 2019.

Moreover, the present research has evaluated the consequence of extreme climate events, particularly the 2015 El Niño and the 2019 positive IOD, on the coastal wind intensity from June to September. When an extreme El Niño event took place in 2015, positive anomalies are conspicuous in the surface wind stresses (Figure 7a), surface chlorophyll-a concentration (Figure 7c), and EMT (Figure 7d). This finding suggests that wind stresses'

strength and direction may have determined the spatial distribution of sea surface productivity off the central MI during this time frame, implying such anomaly may evoke a consequence on the regional seafood stock via a decline or increase in fisheries production.

During the 2019 positive IOD event (June-September), sea surface off the central MI showed significant positive anomalies in surface wind stress, surface chlorophyll-a concentration, EMT, and a negative anomaly in SST (Figure 8). The centre of positive chlorophyll-a anomaly is located congruently with the positions of wind stress, SST, and EMT anomalies (Figure 8). Indeed, the orography effect is visible during this event, i.e. the positive chlorophyll-a anomaly was generated by energetic wind stresses via mixed layer disturbances, leading to phytoplankton bloom and cold SST. Collectively, we conclude that the effects of the 2015 and 2019 extreme events on the sea surface off the central MI are robust. Furthermore, our results emphasized the significance of large-scale climate events in governing surface dynamics of the regional sea.

## 5. Conclusion

Our research has demonstrated that satellite-derived sea surface wind stress, SST, and surface chlorophyll-a concentration are applicable in revealing sea surface dynamics off the central MI. We discovered that southeasterly winds play a critical role in modifying sea surface condition of the study area, i.e. by lowering SST and generating phytoplankton bloom. The SST cooling and phytoplankton bloom occur in response to enhanced surface wind stress and EMT. On the other hand, the SST cooling and phytoplankton bloom deteriorated when wind stress and EMT weakened. The correlation analysis revealed that the ENSO and IOD affect all oceanographic parameters as denoted by positive correlations in wind stress, EMT, chlorophyll-a, and negative correlation in SST. In accordance with the correlation analysis, anomaly analysis suggests positive anomalies in wind stress, EMT, chlorophyll-a concentration, and negative anomaly in SST during the extreme events of 2015 and 2019. Overall, results of this research highlight the importance of coastal wind and extreme climate events in shaping ocean conditions and perhaps regional fisheries production.

## Funding

This work was supported by the Ministry of Education, Culture, Research and Technology, Republic of Indonesia through Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) Number: 150/E4.1/AK.04.PT/2021.

## Declaration of Competing Interest

The authors declare that there is no conflict of interest regarding the publication of this paper.

## Acknowledgement

It is supported by the Ministry of Education, Culture, Research and Technology, Republic of Indonesia through Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) Number: 150/E4.1/AK.04.PT/2021. RYS also acknowledges the Ministry of

Education, Culture, Research and Technology, Republic of Indonesia and the University of Gadjah Mada for the research grants (6/E1/KP.PTNBH/2021 and 2193/UN1/DITLIT/DITLIT/PT/2021). AW thanks to the University of Diponegoro for the research grant (185-98/UN7.6.1/PP/2021). The authors thank the E.U. Copernicus Marine Service Information and the Physical Oceanography Distributed Active Archive Center. The authors thank to Venny Sri and Wahdiny Napon for the massive supports. The authors also wish to thank the anonymous reviewers for their careful reading of the manuscript and constructive suggestions for improvement.

## References

- Delman A.S., Sprintall J.J., McClean L. and Talley L.D. (2016). Anomalous Java cooling at the initiation of positive Indian ocean dipole events. *Journal of Geophysical Research: Oceans*, **121**, 5805–5824. <https://doi.org/10.1002/2016JC011635>.
- DeVantier L., Alcalá A. and Wilkinson C. (2004). The Sulu-Sulawesi Sea: environmental and socioeconomic status, future prognosis and ameliorative policy options. *Ambio*, **33**(1–2), 88–97. <https://doi.org/10.1579/0044-7447-33.1.88>.
- Esaias W.E., Abbott M.R., Barton I., Brown O.B., Campbell J.W., Carder K.L., Clark D.K., et al. (1998). An overview of MODIS capabilities for ocean science observations. *IEEE Transactions on Geoscience and Remote Sensing*, **36**(4), 1250–1265. <https://doi.org/10.1109/36.701076>.
- Ghanea M., Moradi M., Kabiri K. and Mehdinia, A. (2016). Investigation and validation of MODIS SST in the Northern Persian Gulf. *Advances in Space Research*, **57**(1), 127–136. <https://doi.org/10.1016/j.asr.2015.10.040>.
- Gordon A.L. (2005). Oceanography of the Indonesian Seas and Their Throughflow. *Oceanography*, **18**(4), 14–27. <https://doi.org/10.5670/oceanog.2005.01>.
- Group T.W. (1988). The WAM model—A third generation ocean wave prediction model. *Journal of Physical Oceanography*, **18**, 1775–1810. [https://doi.org/10.1175/1520-0485\(1988\)018%3C1775:TWMTO%3E2.0.CO;2](https://doi.org/10.1175/1520-0485(1988)018%3C1775:TWMTO%3E2.0.CO;2).
- Hsieh W.W. and Boer G.J. (1992). Global climate change and ocean upwelling. *Fisheries Oceanography*, **1**(4), 333–338. <https://doi.org/10.1111/j.1365-2419.1992.tb00005.x>.
- Hu S. and Fedorov A.V. (2017). The extreme El Niño of 2015–2016 and the end of global warming hiatus. *Geophysical Research Letters*, **44**, 3816–3824. <https://doi.org/10.1002/2017GL072908>.
- Iskandar I., Sari Q.W., Setiabudidaya D., Yustian I. and Monger B. (2017). The distribution and variability of chlorophyll-a bloom in the southeastern tropical Indian Ocean using empirical orthogonal function analysis. *Biodiversitas*, **18**(4), 1546–1555. <https://doi.org/10.13057/biodiv/d180432>.
- Kurniawati N., Sari Q.W., Setiawan R.Y., Siswanto E., Fauziyah, Setiabudidaya D. and Iskandar I. (2021). Surface chlorophyll-a variations along the southern coast of Java during two contrasting Indian Ocean Dipole events: 2015 and 2016. *Journal of Sustainability Science and Management*, **16**(3), 116–127.
- Levine A.F.Z. and McPhaden M.J. (2016). How the July 2014 easterly wind burst gave the 2015–2016 El Niño a head start. *Geophysical Research Letters*, **43**, 6503–6510. <https://doi.org/10.1002/2016GL069204>.
- L'Heureux M.L., Takahashi K., Watkins A.B., Barnston A.G., Becker E.J. and Di Liberto T.E. (2016). Observing and predicting the 2015–16 El Niño. *Bulletin of the American Meteorological Society*, **98**(7), 1363–1382. <https://doi.org/10.1175/BAMS-D-16-0009.1>.

- Lu B. and Ren, H-L. (2020). what caused the extreme indian ocean dipole event in 2019? *Geophysical Research Letters*, **47**(11). <https://doi.org/10.1029/2020GL087768>.
- Muskananfolo M.R. and Jumsar W.A, (2021). Spatio-temporal distribution of chlorophyll-a concentration, sea surface temperature and wind speed using aqua-modis satellite imagery over the Savu Sea, Indonesia. *Remote Sensing Applications: Society and Environment*, **22**, 1–9. <https://doi.org/10.1016/j.rsase.2021.100483>.
- Palmeiro F.M., Iza M., Barriopedro D., Calvo N. and García-Herrera R. (2017). The complex behavior of El Niño winter 2015–2016. *Geophysical Research Letters*, **44**, 2902–2910.
- Satria F., Setyadi B. and Utama A. (2012). Species composition and distribution of deep-sea fishes of Sangihe and Talaud, North Sulawesi waters (in Indonesian). *BAWAL*, **4**(3), 149–159.
- Setiawan R.Y., Setyobudi E., Wirasatriya A., Muttaqin A.S. and Maslukah L. (2019). The influence of seasonal and interannual variability on surface Chlorophyll-a off the Western Lesser Sunda islands. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing (JSTARS)*, **12**(11), 4191–4197. <https://doi.org/10.1109/JSTARS.2019.2948385>.
- Setiawan R.Y., Wirasatriya A., Hernawan U., Leung S. and Iskandar I. (2020). Spatio-temporal variability of surface chlorophyll-a in the Halmahera Sea and its relation to ENSO and the Indian Ocean Dipole. *International Journal of Remote Sensing*, **41**(1), 284–299. <https://doi.org/10.1080/01431161.2019.1641244>.
- Siswanto E., Horii T., Iskandar I., Gaol J.L., Setiawan R.Y. and Susanto R.D. (2020). Impacts of climate changes on the phytoplankton biomass of the Indonesian Maritime Continent. *Journal of Marine Systems*, **212**, 1–15. <https://doi.org/10.1016/j.jmarsys.2020.103451>.
- Sprintall J., Gordon A.L., Koch-Larrouy A., Lee T., Potemra J.T., Pujiana K. and Wijffels S.E., (2014). The Indonesian seas and their role in the coupled Ocean-climate system. *Nature Geoscience*, **7**(7), 487–492. <https://doi.org/10.1038/ngeo2188>.
- Susanto R.D. and Marra J. (2005). Effect of the 1997/98 El Nino on Chlorophyll-a Variability along the Southern Coasts of Java and Sumatra. *Oceanography*, **4**(18), 124–127. <https://doi.org/10.5670/oceanog.2005.13>.
- Susanto R.D., Moore T.S. and Marra J. (2006). Ocean color variability in the Indonesian Seas during the SeaWiFS era. *G-cubed*, **7**, 1–16. <https://doi.org/10.1029/2005GC001009>.
- Verhoef A. and Stoffelen A. (2013). ASCAT Coastal Winds Validation Report, v1.5, May Technical Note SAF/OSI/CDOP/KNMI/TEC/RP/176, [http://projects.knmi.nl/scatterometer/publications/pdf/ASCAT\\_validation\\_coa.pdf](http://projects.knmi.nl/scatterometer/publications/pdf/ASCAT_validation_coa.pdf).
- Veron J., Devantier L.M., Turak E., Green A.L., Kininmonth S., Stafford-Smith M. and Peterson N. (2009). Delineating the Coral Triangle. *Galaxea, Journal of the Coral Reef Studies*, **11**, 91–100. <https://doi.org/10.3755/galaxea.11.91>
- Wirasatriya A., Setiawan J.D., Sugianto D.N., Rosyadi I.A., Haryadi H., Winarso G., Setiawan R.Y. and Susanto R.D. (2020b). Ekman dynamics variability along the southern coast of Java revealed by satellite data. *International Journal of Remote Sensing*, **41**(21), 8475–8496. <https://doi.org/10.1080/01431161.2020.1797215>.
- Wirasatriya A., Setiawan R.Y. and Subardjo P. (2017). The effect of ENSO on the variability of chlorophyll-a and sea surface temperature in the Maluku Sea. *IEEE Journal of Selected Topics in Applied Earth Observations and Remote Sensing (JSTARS)*, **10**(12), 5513–5518.
- Wirasatriya A., Sugianto D.N., Maslukah L., Ahkam M.F., Wulandari S.Y. and Helmi M. (2020a). Carbon dioxide flux in the Java Sea estimated from satellite measurements. *Remote Sensing Applications: Society and Environment*, **41**(21), 8475–8496. <https://doi.org/10.1016/j.rsase.2020.100376>.
- Wirasatriya A., Susanto R.D., Kunarso K., Jalil A.R., Ramdani F. and Puryajati A.D., 2021. Northwest monsoon upwelling within the Indonesian seas. *International Journal of Remote Sensing*, **42**(14), 5437–5458. <https://doi.org/10.1080/01431161.2021.1918790>.



# Seasonal and interannual coastal wind variability off the central Maluku Islands revealed by satellite oceanography

## ORIGINALITY REPORT

14%

SIMILARITY INDEX

%

INTERNET SOURCES

14%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

- 1 Febryanto Simanjuntak, Tang-Huang Lin. "Monsoon Effects on Chlorophyll-a, Sea Surface Temperature, and Ekman Dynamics Variability along the Southern Coast of Lesser Sunda Islands and Its Relation to ENSO and IOD Based on Satellite Observations", Remote Sensing, 2022  
Publication 2%
- 2 Kurnianingsih, Anindya Wirasatriya, Lutfan Lazuardi, Naoyuki Kubota, Nawi Ng. "IOD and ENSO-Related Time Series Variability and Forecasting of Dengue and Malaria Incidence in Indonesia", 2020 International Symposium on Community-centric Systems (CcS), 2020  
Publication 1%
- 3 Suci Amalia, Stevin Carolius Angga, Elvina Dhiaul Iftitah, Dias Septiana et al. "Immobilization of trypsin onto porous methacrylate-based monolith for flow-through protein digestion and its potential 1%

application to chiral separation using liquid chromatography", Heliyon, 2021

Publication

---

4

Anne-Carlijn Alderkamp, Véronique Garçon, Hein J.W. de Baar, Kevin R. Arrigo. "Short-term photoacclimation effects on photoinhibition of phytoplankton in the Drake Passage (Southern Ocean)", Deep Sea Research Part I: Oceanographic Research Papers, 2011

Publication

---

1 %

5

Hussen Seid Endris, Christopher Lennard, Bruce Hewitson, Alessandro Dosio, Grigory Nikulin, Guleid A. Artan. "Future changes in rainfall associated with ENSO, IOD and changes in the mean state over Eastern Africa", Climate Dynamics, 2018

Publication

---

1 %

6

Jessica Blunden, Derek S. Arndt. "State of the Climate in 2015", Bulletin of the American Meteorological Society, 2016

Publication

---

1 %

7

Esteban Abellán, Shayne McGregor, Matthew H. England, Agus Santoso. "Distinctive role of ocean advection anomalies in the development of the extreme 2015–16 El Niño", Climate Dynamics, 2017

Publication

---

<1 %

8

Shibghatullah Muhammady, Yudhi Kurniawan, Andrivo Rusydi, Yudi Darma. "Electronic and magnetic properties of new half-metallic ferromagnetic rutile  $Ti_{1-x}V_xNiO_2$  ( $x = y = 6.25\%$ ): A first-principles study", Journal of Physics and Chemistry of Solids, 2018

Publication

---

&lt;1 %

9

Wahyono -, Moh. Edi Wibowo, Ahmad Ashari, Muhammad Pajar Kharisma Putra. "Improvement of Deep Learning-based Human Detection using Dynamic Thresholding for Intelligent Surveillance System", International Journal of Advanced Computer Science and Applications, 2021

Publication

---

&lt;1 %

10

Mahyar Mohtadi. "Reconstructing the thermal structure of the upper ocean: Insights from planktic foraminifera shell chemistry and alkenones in modern sediments of the tropical eastern Indian Ocean", Paleoceanography, 09/10/2011

Publication

---

&lt;1 %

11

Bo Lu, Hong - Li Ren. "What Caused the Extreme Indian Ocean Dipole Event in 2019?", Geophysical Research Letters, 2020

Publication

---

&lt;1 %

12

ROBINSON MUGO. "Habitat characteristics of skipjack tuna (*Katsuwonus pelamis*) in the

&lt;1 %

western North Pacific: a remote sensing perspective : Skipjack tuna habitat from RS & GIS in western NP", Fisheries Oceanography, 08/04/2010

Publication

---

13

Siti Herlinda, Mimma Gustianingtyas, Suwandi Suwandi, Radix Suharjo, Jelly Milinia Puspita Sari, Ragil Putri Lestari. "Endophytic fungi confirmed as entomopathogens of the new invasive pest, the fall armyworm, *Spodoptera frugiperda* (JE Smith) (Lepidoptera: Noctuidae), infesting maize in South Sumatra, Indonesia", Egyptian Journal of Biological Pest Control, 2021

<1 %

Publication

---

14

Guang Yang, Xia Zhao, Yuanlong Li, Lin Liu, Fan Wang, Weidong Yu. "Chlorophyll variability induced by mesoscale eddies in the southeastern tropical Indian Ocean", Journal of Marine Systems, 2019

<1 %

Publication

---

15

Qinsheng Wei, Mingzhu Fu, Junchuan Sun, Qingzhen Yao, Baodong Wang, Xuehai Liu, Zhigang Yu. "Seasonal Physical Fronts and Associated Biogeochemical - Ecological Effects off the Jiangsu Shoal in the Western Yellow Sea, China", Journal of Geophysical Research: Oceans, 2020

<1 %

Publication

---

16

Omstedt, A.. "Modeling the variations of salinity and temperature in the large Gulfs of the Baltic Sea", *Continental Shelf Research*, 200302/03

Publication

<1 %

17

Susanto, R. Dwi, Amy Ffield, Arnold L. Gordon, and T. Rameyo Adi. "Variability of Indonesian throughflow within Makassar Strait, 2004–2009", *Journal of Geophysical Research*, 2012.

Publication

<1 %

18

Fatemeh Fallah, Dariush Mansoury. "Coastal upwelling by wind-driven forcing in the Caspian Sea: A numerical analysis", *Oceanologia*, 2022

Publication

<1 %

19

Purwanto Purwanto, Ike Sari Astuti, Fatchur Rohman, Kresno Sastro Bangun Utomo, Yulius Eka Aldianto. "Assessment of the dynamics of urban surface temperatures and air pollution related to COVID-19 in a densely populated City environment in East Java", *Ecological Informatics*, 2022

Publication

<1 %

20

Zhengyao Lu, Zhengyu Liu, Jiang Zhu, Kim M. Cobb. "A Review of Paleo El Niño-Southern Oscillation", *Atmosphere*, 2018

Publication

<1 %

21 Karina von Schuckmann, Pierre-Yves Le Traon, Neville Smith, Ananda Pascual et al. "Copernicus Marine Service Ocean State Report", Journal of Operational Oceanography, 2018  
Publication

---

22 Daniela I.V. Domeisen, Chaim I. Garfinkel, Amy H. Butler. "The Teleconnection of El Niño Southern Oscillation to the Stratosphere", Reviews of Geophysics, 2019  
Publication

---

23 Eric J. Webb, Brian I. Magi. "The Ensemble Oceanic Niño Index", International Journal of Climatology, 2022  
Publication

---

24 Nikita Pusparini, Budi Prasetyo, Ambariyanto, Ita Widowati. "The Thermocline Layer and Chlorophyll-a Concentration Variability during Southeast Monsoon in the Banda Sea", IOP Conference Series: Earth and Environmental Science, 2017  
Publication

---

25 Romero, Oscar E., Mahyar Mohtadi, Peer Helmke, and Dierk Hebbeln. "High interglacial diatom paleoproductivity in the westernmost Indo-Pacific Warm Pool during the past 130,000 years", Paleoceanography, 2012.  
Publication

---

26

Shoichiro Kido, Tomoki Tozuka. "Salinity Variability Associated with the Positive Indian Ocean Dipole and Its Impact on the Upper Ocean Temperature", *Journal of Climate*, 2017

Publication

---

<1 %

27

Yilong Lyu, Yuanlong Li, Xiaohui Tang, Fan Wang, Jianing Wang. "Contrasting Intraseasonal Variations of the Equatorial Pacific Ocean Between the 1997-1998 and 2015-2016 El Niño Events", *Geophysical Research Letters*, 2018

Publication

---

<1 %

28

Andrew S. Delman, Julie L. McClean, Janet Sprintall, Lynne D. Talley, Frank O. Bryan. "Process - Specific Contributions to Anomalous Java Mixed Layer Cooling During Positive IOD Events", *Journal of Geophysical Research: Oceans*, 2018

Publication

---

<1 %

29

Jorge Vazquez-Cuervo, Jose Gomez-Valdes, Marouan Bouali, Luis Miranda, Tom Van der Stocken, Wenqing Tang, Chelle Gentemann. "Using Saildrones to Validate Satellite-Derived Sea Surface Salinity and Sea Surface Temperature along the California/Baja Coast", *Remote Sensing*, 2019

Publication

---

<1 %

30

Laura Crews, Craig M. Lee, Luc Rainville, Jim Thomson. "Direct Observations of the Role of Lateral Advection of Sea Ice Meltwater in the Onset of Autumn Freeze Up", *Journal of Geophysical Research: Oceans*, 2022

Publication

&lt;1 %

31

Liu, Xin, Jing Wang, Xuhua Cheng, and Yan Du. "Abnormal upwelling and chlorophyll-*a* concentration off South Vietnam in summer 2007", *Journal of Geophysical Research*, 2012.

Publication

&lt;1 %

32

Michelle L. L'Heureux, Ken Takahashi, Andrew B. Watkins, Anthony G. Barnston et al. "Observing and Predicting the 2015/16 El Niño", *Bulletin of the American Meteorological Society*, 2017

Publication

&lt;1 %

33

Nicolas C. Jourdain, Matthieu Lengaigne, Jérôme Vialard, Takeshi Izumo, Alexander Sen Gupta. "Further Insights on the Influence of the Indian Ocean Dipole on the Following Year's ENSO from Observations and CMIP5 Models", *Journal of Climate*, 2016

Publication

&lt;1 %

34

Richard P. Dunne, Barbara E. Brown, Niphon Phongsuwan, Lalita Putchim. "The Indian Ocean Dipole and El Niño Southern Oscillation as major drivers of coral cover on shallow

&lt;1 %



reefs in the Andaman Sea", Global Change  
Biology, 2021

Publication

---

35

S. Meisel, K.-C. Emeis, U. Struck, I. Kristen.  
"Nutrient regime and upwelling in the  
northern Benguela since the middle Holocene  
in a global context – a multi-proxy approach",  
Fossil Record, 2011

Publication

---

36

Wan, Z.. "Validation of the land-surface  
temperature products retrieved from Terra  
Moderate Resolution Imaging  
Spectroradiometer data", Remote Sensing of  
Environment, 200211

Publication

---

37

Karina von Schuckmann, Pierre-Yves Le Traon,  
Neville Smith, Ananda Pascual et al.  
"Copernicus Marine Service Ocean State  
Report, Issue 4", Journal of Operational  
Oceanography, 2020

Publication

---

<1 %

<1 %

<1 %

---

Exclude quotes On

Exclude matches Off

Exclude bibliography On