

Study of Growth And N, P Content Of Microalgae *Chlorella vulgaris* Cultivated In Different Culture Media And Light Intensity

by Ita Widowati

Submission date: 28-Jul-2022 10:21AM (UTC+0700)

Submission ID: 1876044186

File name: C-8-Dedi.pdf (365.48K)

Word count: 3215

Character count: 16185

STUDY OF GROWTH AND N, P CONTENT OF MICROALGAE *Chlorella vulgaris* CULTIVATED IN DIFFERENT CULTURE MEDIA AND LIGHT INTENSITY

Article history

Received

4 December 2015

Received in revised form

12 February 2016

Accepted

26 February 2016


Dedi Edwin Satriaji*, Muhammad Zainuri, Ita Widowati

*Corresponding author

dediedwinsatriaji@gmail.com

Department of Marine Science, Diponegoro University,
Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275, Indonesia

Graphical abstract



Abstract

Different light intensity and different media types were used to test their effects on algal growth and biochemical composition of cell content to optimize the growth of algal biomass production. Analysis of the results includes growth calculation of *Chlorella vulgaris* obtained from 6 d observations for 12 h a day (06:00 am to 06:00 pm) with 3 h breaks in between observation. Laboratory analysis was done at the end of day 5th to determine the content of N, and P. Density of *C. vulgaris* were influenced by both light intensity and media. Differences media type and light intensity as separated parameters and their interactions significantly ($p < 0.05$) affected on the growth and nitrogen and phosphorus content of *C. vulgaris*. The highest density of 2.310×10^4 cells \cdot mL $^{-1}$ was on day 4 with walne media resulting in specific growth rate (μ) of 0.43 per day with a light intensity of 5 000 lux, and the lowest density of 725×10^4 cells \cdot mL $^{-1}$ was in day 4 with the NPK media and specific growth rate (μ) 0.25 per day with a light intensity of 4 000 lux. The highest nitrogen and phosphorus content (16.12 mg and 28.19 mg) was obtained from walne medium with a light intensity of 5000 lux, and the lowest (3.43 mg and 2.17 mg) was obtained from NPK with a light intensity of 4 000 lux.

Keywords: *Chlorella vulgaris* Beyerinck [Beijerinck], density, light intensity, media, specific growth rate

© 2016 Penerbit UTM Press. All rights reserved

1.0 INTRODUCTION

One of the challenges that Indonesian people face nowadays was the shortage of food sources as feed to support culture fisheries. Microalgae have a great potential for various applications including the production of compounds for food, feed and aquaculture [1]. *Chlorella vulgaris* Beyerinck [Beijerinck], is a natural resource that can be easily grown in a variety of water conditions. *Chlorella vulgaris* has a high potential value because it contains adequate nutrients in the form of proteins, amino acids, vitamins, and minerals. Considering the richness of the nutrients contained in *Chlorella vulgaris* and its highly extensive application on the food and cosmetic industries, it is necessary to realize

the cultivation of *C. vulgaris* by manipulating the medium of life and light intensity.

The culture media has a direct influence on cellular growth, as well as on the biomass composition of microalgae [2]. In this context, microalgae, like other microorganisms, requires sources of carbon, nitrogen, phosphorus and other micronutrients [3], and the nutrients that make up the medium must be dissolved (available) for biochemical uses by the microalgae. Microalgal cell growth rates and lipid content were affected by environmental parameters such as temperature, light intensity and frequency, gas composition, and nutrient level in the culture system. Light quantity and quality determine the amount of energy available to photosynthetic organisms to conduct their metabolic activities. Photosynthesis process is divided into stages of light reaction and a

dark reaction, each of which produce oxygen and carbohydrates. The reaction will be greatly influenced by the intensity of the light, the wavelength of light and nutrients available in the media [4]. Light intensity is also a major factor that determines the formation of carotenoid pigment, if the amount of light available is sufficient to inhibit the synthesis of chlorophyll. Chlorophyll and carotenoids will be synthesized in a balanced manner in the chloroplast. At the moment, the balance is changing as a result of the increase in carotenoids, the plastids structure will change and as a result will degrade chlorophyll but not carotenoids [5]. Studies show that light is important factors which affects the growth of autotrophic microalgae that study also showed *Chlorella* could grow with the intensity range 3 000 lux to 11 000 lux [6] (1 lux = 1 lm m²). The present study focuses on the growth and content of nitrogen (N), phosphor (P) microalgae *C. vulgaris*, to identify which treatment relevant and efficient of media and light intensity

2.0 EXPERIMENTAL

2.1 *Chlorella vulgaris* Culture

The microalgae *Chlorella vulgaris* was obtained from BBPAP (Brackish Water Aquaculture Research and Development Center) Jepara, Central Java. *Chlorella vulgaris* was cultivated with different culture media and light intensity. Experimental design used was a randomized block with nine culture bottles containing 5 L of sea water with three treatments and three replications. The treatments are Walne Pro-analysed, formulations, and NPK. Each treatment were treated with light densities of 3 000 lux, 4 000 lux, and 5 000 lux. Light intensity was measured with a Lux Meter.

2.2 Algal Growth Medium

The medium that was used in the cultivation of the microalgae was Walne, formulations, and NPK, compositions of walne medium is as follows Na₂EDTA 45 g L⁻¹, NaNO₃ 100 g L⁻¹, NaH₂PO₄·2H₂O 20 g L⁻¹, FeCl₃·6H₂O 1.3 g L⁻¹, MnCl₂·4H₂O 0.36 g L⁻¹, H₃BO₃ 33.60 g L⁻¹, Trace Metal Solution consisted of 1 ml from 100 ml solution CuSO₄·5H₂O 2.0 g L⁻¹, ZnCl₂ 2.1 g L⁻¹, CoCl₂·6H₂O 2.0 g L⁻¹, (NH₄)₆Mo₇O₂₄·4H₂O 0.9 g L⁻¹, Vitamins in 100 mL, B1 100 g L⁻¹, B12 5 g L⁻¹ [7].

2.3 Growth Evaluation

Cell concentration was determined by cell counting, 1 mL of algal suspension was collected via sampling tube. Cell density was directly counted with a Neubauer hemocytometer and an Olympus light microscope where it represents cell density mL⁻¹. Algal growth was evaluated daily in three replication. Specific growth rate was calculated using Eqs. (1), as follows [8]:

$$\mu = \frac{\ln[X_2 - X_1]}{t_2 - t_1} \quad (1)$$

Volumetric biomass productivity, was calculated using Eqs. (2), as follows [9]:

$$N = \frac{(\sum N_1 + \sum N_2)}{2} \times \frac{1}{1 \text{ mm} \times 0.2 \text{ mm} \times 0.1 \text{ mm}} \times \frac{1 \text{ mm}^3}{10^{-3} \text{ ml}} \quad (2)$$

where X₁ and X₂ are the biomass concentration on days t₁ and t₂, respectively, and N₁, N₂ is the number of observations.

Growth calculation and specific growth rate of *C. vulgaris* was obtained from 5 d observations for 12 h a day (06:00 am to 06:00 pm) with 3 h breaks in between observation times of 3 h intervals with a total of five observations per day. Laboratory analysis was done at the end of d 6 to determine the content of N, P.

2.4 Statistical Analysis

Cell densities were analyzed using polynomial orthogonal for obtaining the growth model of *C. vulgaris*. Prior to statistical analysis, normality and homogeneity of data were checked by using Shapiro-Wilk and Kolmogorov-Smirnov test, respectively. All data were statistically performed at a maximum significance level of 5 % by two-way analysis of variance (ANOVA) followed by tukey's honest significant difference (HSD) post hoc multicomparison test.

3.0 RESULTS AND DISCUSSION

From the polynomial orthogonal result, all treatment showed the same model of growth, even from different peak density.




Figure 1 The growth pattern of *C. vulgaris* at three light intensity (3 000 lux, 4 000 lux, 5 000 lux) and media (Walne, Formulation, NPK). Each value is the mean of three samples

Figure 1 shows the highest density from initial density $115 \times 10^4 \text{ cells} \cdot \text{mL}^{-1}$ to $(867 \text{ to } 2\,310) \times 10^4$

cells · mL⁻¹ with average 1 589 × 10⁴ cells mL⁻¹ with (μ) 0.43 per day was obtained in day 4 with the walne medium with a light intensity of 5 000 lux. The lowest density from initial density 112 × 10⁴ cells · mL⁻¹ to (416 to 725) × 10⁴ cells mL⁻¹ with average 570 × 10⁴ cells mL⁻¹ with (μ) 0.25 per day was obtained in day 4 with the medium NPK with a light intensity of 4 000 lux.

From the ANOVA result, light intensity and media seemed to have an effect on biomass production of *C. vulgaris*. The variation of light intensities and growth medium as separate parameters and their interactions significantly (p < 0.05) affected biomass of *C. vulgaris*. Figure 2 shows that light intensity and media affected biomass of *C. vulgaris*.




Figure 2 The biomass of *C. vulgaris* at three light intensity (3 000 lux, 4 000 lux, 5 000 lux) and media (Walne, Formulation, NPK). Each value is the mean of three samples

Figure 3 and Figure 4 shows the nitrogen and phosphorus content of *C. vulgaris*. Each light intensity and three different growth medium has maximum nitrogen and phosphorus content.




Figure 3 Phosphorus content of *C. vulgaris* at three light intensity (3 000 lux, 4 000 lux, 5 000 lux) and media (Walne, Formulation, NPK). Each value is the mean of three samples

The sample used for analysis of nitrogen and phosphorus as much as 500 mg of total dry weight of

the sample was obtained after harvest at the end of culture.




Figure 4 Nitrogen content of *C. vulgaris* at three light intensity (3 000 lux, 4 000 lux, 5 000 lux) and media (Walne, Formulation, NPK). Each value is the mean of three samples

This research also shows that nitrogen and phosphorus content of *C. vulgaris* cultivated in the 5 000 mL medium volume with the light intensity (3 000 lux, 4 000 lux, 5 000 lux) and three medium (Walne, Formulation and NPK) as separate parameters and their interactions significantly (p < 0.05) affected nitrogen and phosphorus content of *C. vulgaris*. Overall nitrogen and phosphorus content of all samples are shown in Table 2. The highest nitrogen and phosphorus content of 16.12 mg; 28.19 mg was in walne with a light intensity of 5 000 lux, and the lowest 3.43 mg and 2.17 mg was in NPK with a light intensity of 4 000 lux.

Table 1 shows the specific growth rates and maximum cell density, each light intensity and three different growth medium had maximum specific growth rate and density. The highest specific growth rates of 0.43 per day with optimal density 2 310 × 10⁴ cells · mL⁻¹ was obtained in day 4 walne medium with a light intensity of 5 000 lux, and the lowest specific growth rates of 0.25 per day with optimal density 725 × 10⁴ cells mL⁻¹ was obtained in day 4 on the NPK medium with a light intensity of 4 000 lux.

Light and medium are major factors that determine the growth and biochemical of *Chlorella vulgaris*. The result on three different culture media and light intensity shows the same graph model, even though that the optimal density was different. It was more likely due to the period of adaptation which was limited in the nutrient quantity. The optimal peak of microalga *C. vulgaris* cultivation could be varied due to the optimal quantity of nutrient, which was provided by the fertilizer and medium [10, 11]. During the mixing process of the fertilizer in the medium, the quantity of nutrient in the earlier process was less than the next period, which could be easier absorbed by the microalgae. Nutrient availability, on the other hand, has a significant impact on the growth and propagation of microalgae as well as on

their lipid and fatty acid composition. In general, when algal growth decreased and there was no requirement for the synthesis of any additional membrane compounds, the cells divert and deposit fatty acids [12-15]. Under nitrogen depletion condition, the growth rate was very low leading to higher lipid productivity [16].

Photosynthetic microalgae photosynthesize assimilate inorganic carbon for conversion into organic matter. Light is the source of energy which

drives this reaction and in this regard intensity. The growth characteristics depend on the light intensities [17]. The compounds that are produced in the light dependent phase (ATP, NADPH) are used in dark cycle to synthesize metabolic molecules essential for growth. Also, it has reported that, some enzymes of the pentose cycle of photosynthesis and CO₂ fixation are inactive during the illumination [18, 19].

Table 1 Maximum cell density and specific growth rate of *Chlorella vulgaris*

Light Intensity (Lux)	Walne Pro-analysed		Formulations		NPK		
	Maximum cell density	μ	Maximum cell density	μ	Maximum cell density	μ	
1	3 000	1729×10^4	0.30	1206×10^4	0.40	894×10^4	0.31
2	4 000	1419×10^4	0.37	1031×10^4	0.32	725×10^4	0.25
3	5 000	2310×10^4	0.43	919×10^4	0.30	856×10^4	0.26

Table 2 Nitrogen and phosphorus content of *Chlorella vulgaris*

Light Intensity (Lux)	Walne Pro-analysed		Formulations		NPK		
	Phosphorus (mg)	Nitrogen (mg)	Phosphorus (mg)	Nitrogen (mg)	Phosphorus (mg)	Nitrogen (mg)	
1	3 000	27.23	13.83	25.04	10.40	2.25	4.05
2	4 000	26.62	12.05	20.97	8.82	2.17	3.43
3	5 000	28.19	16.12	19.01	6.98	2.20	3.94

Media composition had significant effect on both the growth rate and the density of microalgae *C. vulgaris*. Microalgae were known to grow more abundantly in nutrient rich (eutrophic) waters leading frequently to algal blooms [20]. A higher light absorption rate would allow for the chloroplast to manufacture increased amounts of usable chemical energy which would result in an increase in algal growth (biomass).

Environmental parameters in this study include temperature, salinity, dissolved oxygen and pH were also observed. Overall, environmental parameters values are as follow: temperature was 27 °C to 29 °C, salinity was 30 ng L⁻¹ to 35 ng L⁻¹, dissolved oxygen was 5.4 mg L⁻¹ to 6.4 mg L⁻¹, and between pH 7 to pH 8. The observation of water quality during cultivation showed the optimal range for each variable water quality.

The pattern of biomass growth in this study showed that under low light intensity of 3 000 lux and 4 000 lux, *C. vulgaris* exhibited a slow growth and low biomass, but increased in light intensity of 5 000 lux in walne medium which resulted in highest biomass growth in this study.

4.0 CONCLUSION

This study has identified the effect of media type and light intensity as separate parameters and their interactions significantly ($p < 0.05$) on the growth of *Chlorella vulgaris* and nitrogen and phosphorus content. The highest density of $2\,310 \times 10^4$ cells mL⁻¹ was in day 4 with the walne medium and specific growth rate (μ) 0.43 per day with a light intensity of 5 000 lux, and the lowest density of 725×10^4 cells mL⁻¹ was in day 4 with the NPK medium and specific growth rate (μ) 0.25 per day with a light intensity of 4 000 lux. The highest nitrogen and phosphorus content (16.12 mg and 28.19 mg) was in walne with a light intensity of 5 000 lux, and the lowest (3.43 mg and 2.17 mg) was in NPK with a light intensity of 4 000 lux.

Acknowledgement

This research was funded by PNPB Universitas Diponegoro TA.2013 Number 023.04.2.189815/2014, 5 December 2013, for which the authors were gratefully acknowledge.

References

- [1] Brennan, L. and Owende, P. 2010. Biofuels from Microalgae – A Review of Technologies for Production, Processing, and Extractions of Biofuels and Co-products. *Renewable and Sustainable Energy Reviews*. 14: 557–577.
- [2] Richmond, A. 2004. Biological Principles of Mass Cultivation. In Richmond, A. (ed.). *Handbook of Microalgal Culture: Biotechnology and Applied Phycology*. London: Blackwell Publishing Ltd.
- [3] Kumar, A., S. Ergas, X. Yuan, A. Sahu, Q. Zhang, J. Dewulf, et al. 2010. Enhanced CO₂ Fixation and Biofuel Production via Microalgae: Recent Developments and Future Directions. *Trends Biotechnol.* 28: 371–380.
- [4] Zainuri, M. and Facta, M. 2006. Pengaruh Pengaturan Intensitas Cahaya yang Berbeda Terhadap Kelimpahan *Dunaliella* sp. dan Oksigen Terlarut dengan Simulator TRIAC dan Mikrokontroler AT89S52 [Effect of Different Light Intensity Settings to Abundance of *Dunaliella* sp. and Dissolved Oxygen with TRIAC and Microcontroller Simulator AT89S52]. *Jurnal Ilmu Kelautan*. 11(2): 67–71. [Bahasa Indonesia].
- [5] Zainuri, M and Kusumaningrum, H. P. 2014. Optimization and Stability of Total Pigments Production of Fusan from Protoplast Fusion of Microalgae *Dunaliella* and *Chlorella* *In Vivo*: Attempts on Production of Sustainable Aquaculture Natural Food. *International Journal of Marine and Aquatic Resource Conservation and Co-existence Research Article*. 1(1): 1–5.
- [6] Kitaya Y, Azuma H, and Kiyota M. 2005. Effects of Temperature, CO₂/O₂ Concentration and Light Intensity on Cellular Multiplication of Microalgae. *Euglena gracilis*. *Adv. Space. Re.* 35: 1584–1588.
- [7] Andersen, R. A. 2005. *Algal Culturing Technique*. UK: Elsevier Academic Press.
- [8] Das, P., W. Lei, S. S. Aziz, and J. P. Obbard. 2011. Enhanced Algae Growth in Both Phototrophic and Mixotrophic Culture under Blue Light. *Bioresour. Technol.* 102: 3883–3887.
- [9] Hadioetomo, R. S. 1993. *Mikrobiologi Dasar dalam Praktek Teknik dan Prosedur Dasar Laboratorium* [Basic Microbiology in Practice Laboratory Techniques and Procedures]. Jakarta: UI Press. [Bahasa Indonesia].
- [10] Borowitzka. 1992. *The Mass Culture of Dunaliella salina*. *Algal Biotechnology Laboratory*. Australia: Murdoch University.
- [11] Guedes, A. C., H. M. Amaro and F. X. Maleata. 2011. Microalga as Sources of Carotenoids. *Marine Drugs*. 9: 625–644.
- [12] Nigam, S., M. P. Rai, and R. Sharma. 2011. Effect of Nitrogen on Growth and Lipid Content of *Chlorella pyrenoidosa*. *Am J Biochem Biotechnol.* 7(3): 124–129.
- [13] Rodolfi, L., Zittelli, G. C., Bassi, N., Padovani, G., Biondi, N. and Bonini, G. 2009. Microalgae for Oil: Strain Selection, Induction of Lipid Synthesis and Outdoor Mass Cultivation in A Low-cost Photobioreactor. *Biotechnol Bioeng.* 102(1): 100–120.
- [14] Schuhmann, H., Lim, D. K. Y, Schenk, P. M. 2012. Perspectives on Metabolic Engineering for Increased Lipid Contents in Microalgae. *Biofuels*. 3(1): 71–86.
- [15] Thompson, G. A. 1996. Lipids and Membrane Function in Green Algae. *Biochim Biophys Acta*. 1302(1): 17–45.
- [16] Josephine, A., Niveditha, C., Radhika, A., et al. 2015. Analytical Evaluation of Different Carbon Sources and Growth Stimulators on the Biomass and Lipid Production of *Chlorella vulgaris* Implications for biofuels. *Biomass and Bioenergy*. 75: 170–179.
- [17] Khoeyi, Z. A., Seyfabadi, J. and Ramezanpour, Z. 2012. Effect of Light Intensity and Photoperiod on Biomass and Fatty Acid Composition of the Microalgae, *Chlorella vulgaris*. *Aquacult. Int.* 20: 41–49.
- [18] Bouterfas, R., Belkoura, M., Dauta A. 2006. The Effects of Irradiance and Photoperiod on The Growth Rate of Three Freshwater Green Algae Isolated from A Eutrophic Lake. *Limnetica*. 25: 647–656.
- [19] Ugwu CU, Aoyagi H, Uchiyama H. 2007. Influence of Irradiance, Dissolved Oxygen Concentration, and Temperature on The Growth of *Chlorella sorokiniana*. *Photosynthetica*. 45(2): 309–311.
- [20] Schenk, P. M., Thomas-Hall, S. R., Stephens, E. et al. 2008. Second Generation Biofuels: High-efficiency Microalgae for Biodiesel Production. *BioEnergy Res.* 1: 20–43.

Study of Growth And N, P Content Of Microalgae Chlorella vulgaris Cultivated In Different Culture Media And Light Intensity

ORIGINALITY REPORT

18%

SIMILARITY INDEX

%

INTERNET SOURCES

18%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Arruda Henrard Adriano, Martins da Rosa Gabriel, Moraes Luiza, Greque de Morais Michele, Alberto Vieira Costa Jorge. "The cultivation of microalgae Cyanobium sp. and Chlorella sp. in different culture media and stirring setting", African Journal of Microbiology Research, 2015
Publication 2%

- 2** Fiddy S. Prasetiya, Ikha Safitri, Ita Widowati, Bruno Cognie et al. "Does allelopathy affect co-culturing Haslea ostrearia with other microalgae relevant to aquaculture?", Journal of Applied Phycology, 2015
Publication 2%

- 3** Radia Bouterfas, M. Belkoura, A. Dauta. "The effects of irradiance and photoperiod on the growth rate of three freshwater green algae isolated from a eutrophic lake", Limnetica, 2006
Publication 2%

4

Shifa Helena, Muhammad Zainuri, Jusup Suprijanto. "Microalgae *Dunaliella salina* (Teodoresco, 1905) Growth Using the LED Light (Light Limiting Dioda) and Different Media", *Aquatic Procedia*, 2016

Publication

1 %

5

Nisha Phour Dhull, Raman Soni, Deepak Kumar Rahi, Sanjeev Kumar Soni. " Evaluation of autotrophic and mixotrophic regimen cells in various wastes water for its biochemical composition and biomass production ", *PeerJ*, 2014

Publication

1 %

6

C.M. Ho. "Autotrophic denitrification via a novel membrane-attached biofilm reactor", *Letters in Applied Microbiology*, 9/16/2001

Publication

1 %

7

Haiying Tang, Meng Chen, K.Y. Simon Ng, Steven O. Salley. "Continuous microalgae cultivation in a photobioreactor", *Biotechnology and Bioengineering*, 2012

Publication

1 %

8

"Algae for Biofuels and Energy", *Springer Nature*, 2013

Publication

1 %

9

Donghan Kang, Keugtae Kim, Youngho Jang, Heechun Moon, Dongjin Ju, Deokjin Jahng. "Nutrient removal and community structure

1 %

of wastewater-borne algal-bacterial consortia grown in raw wastewater with various wavelengths of light", International Biodeterioration & Biodegradation, 2018

Publication

10

Sekar Ashokkumar, Kuppusamy Manimaran, Keun Kim. "Cultivation and Identification of Microalgae (Diatom)", Wiley, 2015

Publication

11

F. Hadj-Romdhane, X. Zheng, P. Jaouen, J. Pruvost, D. Grizeau, J.P. Croué, P. Bourseau. "The culture of *Chlorella vulgaris* in a recycled supernatant: Effects on biomass production and medium quality", Bioresource Technology, 2013

Publication

12

Bui, V.P.. "Direct biooxidation of arenes to corresponding catechols with *E. coli* JM109 (pDTG602). Application to synthesis of combretastatins A-1 and B-1", Tetrahedron Letters, 20020408

Publication

13

Peer M. Schenk, Skye R. Thomas-Hall, Evan Stephens, Ute C. Marx, Jan H. Mussgnug, Clemens Posten, Olaf Kruse, Ben Hankamer. "Second Generation Biofuels: High-Efficiency Microalgae for Biodiesel Production", BioEnergy Research, 2008

Publication

1 %

1 %

1 %

1 %

14 Biofuel and Biorefinery Technologies, 2015. <1 %
Publication

15 Teresa Lopes Silva. "Oil Production Towards Biofuel from Autotrophic Microalgae Semicontinuous Cultivations Monitorized by Flow Cytometry", Applied Biochemistry and Biotechnology, 12/09/2008 <1 %
Publication

16 C. Scharff, N. Domurath, M. Wensch-Dorendorf, F.-G. Schröder. "Cultivation of microalgae in closed systems under artificial lighting", Acta Horticulturae, 2017 <1 %
Publication

17 Jeevan Raj Rajanren, Harun Mohamed Ismail, Padmesh Tirunelveli Narayanapillai. " Investigation on phototrophic growth of an indigenous for biodiesel production ", Environmental Progress & Sustainable Energy, 2015 <1 %
Publication

18 Minhas, Amritpreet K., Peter Hodgson, Colin J. Barrow, and Alok Adholeya. "A Review on the Assessment of Stress Conditions for Simultaneous Production of Microalgal Lipids and Carotenoids", Frontiers in Microbiology, 2016. <1 %
Publication

19

Kallio, P.T.. "Analysis of novel hemoglobins during microaerobic growth of HMP-negative *Escherichia coli*", *Enzyme and Microbial Technology*, 20070104

Publication

20

"Algal Biofuels", Springer Science and Business Media LLC, 2017

Publication

21

B. W. Kim, K. P. Chang, H. N. Chang. "Effect of light source on the microbiological desulfurization in a photobioreactor", *Bioprocess Engineering*, 1997

Publication

22

Khatoon, Helena, Norazira Abdu Rahman, Sanjoy Banerjee, Nazurah Harun, Siti Suhada Suleiman, Nur Hazwani Zakaria, Fathurrahman Lananan, Siti Hajar Abdul Hamid, and Azizah Endut. "Effects of different salinities and pH on the growth and proximate composition of *Nannochloropsis* sp. and *Tetraselmis* sp. isolated from South China Sea cultured under control and natural condition", *International Biodeterioration & Biodegradation*, 2014.

Publication

23

Nur Atikah Ahmad Latiffi, Radin Maya Saphira Radin Mohamed, Najeeha Mohd Apandi, Ramlah Mohd Tajuddin. "Preliminary

<1 %

<1 %

<1 %

<1 %

<1 %

Assessment of Growth Rates on Different Concentration of Microalgae *Scenedesmus* sp. in Industrial Meat Food Processing Wastewater", MATEC Web of Conferences, 2017

Publication

24

Suleiman Dauda, Mathias Ahii Chia, Sunday Paul Bako. "Toxicity of titanium dioxide nanoparticles to *Chlorella vulgaris* Beyerinck (Beijerinck) 1890 (Trebouxiophyceae, Chlorophyta) under changing nitrogen conditions", *Aquatic Toxicology*, 2017

<1 %

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Study of Growth And N, P Content Of Microalgae Chlorella vulgaris Cultivated In Different Culture Media And Light Intensity

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
