

KORESPONDENSI PAPER

JUDUL : Identification of Potential Locations for *Kappaphycus alvarezii* Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia

JURNAL : ILMU KELAUTAN: Indonesian Journal of Marine Sciences

No.	Activity	Date	Description	Page
1	Manuscript submission to IJMS	19 July 2022	Email from IJMS: Submission acknowledgement • Initial manuscript	2 3-26
2	Editor Decision.	27 July 2022	Email from IJMS: Editor Decision Revision required • Comments from reviewer A • Manuscripts comments from reviewer B: 47735RV	27 27 28-51
3	Revision #1 submission	16 August 2022	Email sent to IJMS: • Manuscript Revised 1: 47735RV 15082022 revised 1 • Rebuttal IJMS for Reviewer A and B	52 53-76 77-79
4	Revised Version Acknowledgement		Email from IJMS: • Revised Version Submitted	80
5	Editor Decision	30 August 2022	Email from IJMS: • Submission accepted	81
6	Proof Layout	8 September 2022	Email from IJMS: • Billing • Manuscripts 44735-15285-2-ED Proof Read	82 83-97
7	Payment and Revision #2 Submission	10 September 2022	Email sent to IJMS: Proof Read revised #2 • Manuscripts 44735-15285-2-ED Proof Read Revised #2 • Proof of Transfer Publication Fee	98 99-114 115
8	Published manuscript	September 2022	Manuscript Published on ILMU KELAUTAN: Indonesian Journal of Marine Science, Vol 27(3):	116-129



Sarjito Sarjito <sarjito@live.undip.ac.id>

[IK.IJMS] Submission Acknowledgement

1 message

Ambariyanto <ijms@live.undip.ac.id>

Tue, Jul 19, 2022 at 9:14 AM

Reply-To: sarjito - sarjito <sarjito@live.undip.ac.id>

To: sarjito - sarjito <sarjito@live.undip.ac.id>

sarjito - sarjito:

Thank you for submitting the manuscript, "Site selection of Kappaphycus alvarezii farming to optimize seaweed production using Geographic Information system with geospatial modeling at Spermonde Islands waters, South Sulawesi, Indonesia" to ILMU KELAUTAN: Indonesian Journal of Marine Sciences. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL: <https://ejournal.undip.ac.id/index.php/ijms/author/submission/47735>

Username: sarjito

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Ambariyanto

ILMU KELAUTAN: Indonesian Journal of Marine Sciences

INDONESIAN JOURNAL OF MARINE SCIENCE

<http://ejournal.undip.ac.id/index.php/ijms>

Site selection of *Kappaphycus alvarezii* farming to optimize seaweed production using Geographic Information system with geospatial modeling at Spermonde Islands waters, South Sulawesi, Indonesia

Sarjito Sarjito^{1*}, Hanny Ammaria¹, Muhammad Helmi^{2,3}, Slamet Budi Prayitno¹, Nurjannah Nurdin^{4,5}, Riza Yuliratno Setiawan⁶, Parichat Wetchayont⁷, Anindya Wirasatriya^{2,3}

¹Aquaculture Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro, Indonesia

Jl. Prof. Soedarto, SH., Tembalang, Semarang, Central Java (50275) Indonesia

²Oceanography Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro, Indonesia

³Center for Coastal Rehabilitation and Disaster Mitigation Studies (CoREM), Indonesia

⁴Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University, Indonesia

⁵Marine Science and Fisheries Faculty, Hasanuddin University, Indonesia

⁶Marine Science and Fisheries Faculty, Gadjah Mada University, Indonesia

⁷Department of Geography, Srinakharinwirot University, Thailand

e-mail: sarjito@live.undip.ac.id

Abstract

Indonesia is the largest producer of red algae in the world with one of the major *K. alvarezii* farming areas within the Indonesian seas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province. *K. alvarezii* production in Pangkep Regency increased year by year. However, this production still can be increased by expanding the utilized areas for seaweed farming. Proper site selection is crucial to guarantee the success of the seaweed farming development. We conducted a Geographic Information system (GIS) with geospatial modeling approach for the site selection of *K. alvarezii* cultivation at the Liukang Tuppabiring District, Spermonde Islands by combining various physical and chemical parameters obtained from in-situ data and hydrodynamics modeling. The parameters are Sea Surface Temperature, (SST), salinity pH, substrate type, current speed, wave height, DO, nitrate, phosphate, and Total Suspended Solid (TSS). We also include the accessibility and the restricted area of shipping lanes for determining the area suitability of *K. alvarezii* cultivation. This research demonstrates the cell-based geospatial modeling can be effectively used and found the suitable categorized area for *K. alvarezii* culture of 4,546.28 ha. The Potential productivity of wet-weight seaweed can reach 56,825 tons per cycle or 113,650 tons per year. This result indicates the GIS with geospatial modeling approach for developing the area of *K. alvarezii* culture which can be applied in the other area.

Keywords: Seaweed production, site selection, geospatial, hydrodynamics modeling

1.Introduction

Indonesia is the largest producer of red algae in the world (83% of global seaweed production) followed by the Philippines and several countries (Valderrama *et al.*, 2015; Kim *et al.*, 2017). Seaweed production in Indonesia comes not only from mariculture, and brackish water ponds but also from natural collections. However, the availability of natural seaweed can be threatened in line with their increasing demand so that seaweed cultivation is completely necessary (Pickering *et al.*, 2007). The increasing demand for seaweed was mostly driven by new markets in China, Eastern Europe, Brazil, and others (Bixler and Porse, 2011). Furthermore, Arbit *et al.* (2019) stated that 99.73% of Indonesia's seaweed production comes from cultivation. Therefore, seaweed farming is the best choice for fulfilling market demand. Moreover, one of the major *K. alvarezii* farming areas within the Indonesian seas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province (Jalil *et al.*, 2020).

Site selection is essential in aquaculture development which depends on the types of commodities and applied cultivation methods (Kaiser *et al.*, 2010; Junaidi *et al.*, 2018). For seaweed, the success of cultivation is intrinsically linked to physical and chemical parameters of seaweed farming areas such as turbidity, temperature, salinity, pH, DO, nitrate and phosphate concentration (Rusliani 2016, Jalil *et al.* 2020). Hydrodynamic aspects should also be considered in the site selection of *K. alvarezii* farming since the cultivation method uses the longline method which is prone to the strong current and wave. The existence of Indonesian Throughflow along the Makassar Strait (Susanto *et al.* 2016) and the occurrence of strong upwelling at the southern part of the South Sulawesi each summer (Setiawan and Kawamura 2011) may influence the hydrodynamics condition in the Spermonde Island. Besides, the existence of small islands among the water area makes the hydrodynamics condition at the Spermonde Islands becomes more complex. Spermonde Islands area also has dense shipping activities. Many shipping lanes have lied among these areas (Figure 1) which are should be avoided in the mariculture site selection. This shows the complex considerations for conducting site selection of seaweed farming at Spermonde Islands.

In dealing with the complex considerations as mentioned above, the use of the geospatial model which is part of Geographical Information Systems (GIS) becomes an effective solution for determining the proper mariculture site selection (Longdill *et al.* 2008; Micael *et al.* 2015). This approach, in general, is an analysis of layers that contains spatial data to evaluate and select a particular area based on its suitability criteria classified according to certain environmental aspects (Malczewski, 2006; Malczewski & Rinner, 2015). GIS utilization in mariculture development has been widely used in various types of commodities including seaweed (e.g., Radiarta *et al.*, 2011; Radiarta *et al.*, 2012; Sousa *et al.*, 2012; Nirmala *et al.*, 2014; Teniwut *et al.*, 2019) which were mainly used in-situ data and remote sensing data for the input of geospatial modeling. The application of GIS for site selection for aquaculture in the Spermonde Islands has conducted by Arifin *et al.* (2014). However, their analysis was only based on in-situ measurement which cannot represent the temporal change of ocean parameters. In the present study, we perform cell-based geospatial modeling for the determination of sustainable site selection of seaweed cultivation, which integrates various physical and chemical parameters obtained not only from in-situ data but also the hydrodynamic model. The combination of in-situ data and the hydrodynamic model becomes the new approach proposed in the present study. Considering the importance of hydrodynamics aspects for the site selection of seaweed culture which is not available in the remote sensing data, we built a 2D hydrodynamic model to represent the hydrodynamic condition in the study area. On the other hand, Indonesia as the maritime continent experiences four-season influenced by monsoon wind (Helmi *et al.*, 2018) which impacts the hydrodynamics conditions. Thus, in-situ and incidental measurements of current and wave cannot represent the hydrodynamics condition in the study area. This issue only can be resolved by performing the hydrodynamics model. Furthermore, we also simulate the potential productivity of *K. alvarezii* at the selected site. This research is expected to be a prototype of a geospatial model to explore the potential aquaculture area in small islands and can be scientific-based information for local governments, researchers, NGOs, and related stakeholders for planning and developing *K. alvarezii* cultivation in the study area.

2. Material and Method

2.1. Study Area

The study area is presented in Figure 1 which is located at the Spermonde islands waters precisely in the Liukang Tupabbiring District, Pangkep Regency, South Sulawesi. The Liukang Tupabbiring District area is 54.44 km² (4.89% of Pangkep Regency) and consists of 20 islands. The population of this district reached 29,680 inhabitants, generally as fishermen and sea farmers. The distribution of field station is located at 4°5'3.98"S - 4°59'51.137"S and 119°18'35.532"E - 119°26'19.849"E. There are 94 sampling points distributed among the island of Bontosua, Sanane, Panjenekang, Pankaiya, Langkadea, Balanglombo, Balangcaddi Island, and reef flat area.

2.2. Water Quality Data Collection

We conducted a field survey for collecting water quality data at 94 points which the depths are more than 2 meters. Survey site determination was using Garmin 76C GPS. The collection of water quality data are presented in Table 1. The in-situ measurements were Sea Surface Temperature, (SST), salinity, pH, and substrate type. For the chemical parameters such as DO, nitrate, phosphate, and Total Suspended Solid (TSS), the water samples were analyzed at Chemical Oceanography Laboratory, Marine Science Department, Hasanuddin University, Indonesia (May 2019). The dynamic and non-dynamic water parameter data were interpolated to obtain a spatial distribution pattern at the time of data retrieval. The data were compiled in a digital form of 5 x 5 meters cell size using spline and classified according to suitability class in Table 2.

2.3. Two dimensions Hydrodynamic Model

Two dimensions (2D) hydrodynamic model was used to simulate current and wave condition at the study area. The data for generating a hydrodynamic model are as follows:

1. High-resolution satellite imagery GeoEye-1 from ESRI online (<https://esriindonesia.co.id/arcgis-online>)
2. Bathymetry data from Indonesia Geospatial Agency (<http://batnas.big.go.id/>)
3. Surface wind and wave height data obtained from 6 hourly ERA-interim data with a grid interval of 0.125° from July 2018 to July 2019 (Dee *et al.* 2011)
4. Tidal elevation forecast from July 2018 to July 2019 (<https://www.dhigroup.com/download/mike-by-dhi-tools/coastandseetools/global-tide-model>)

The current model was constructed by continuity equations and momentum equations with the average of depth. On-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry, wind and tidal dynamics were used as an input for generating current model.

The initial process was making a mesh, and setting the boundary condition of the model using the mathematical continuity equation as follows:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = S$$

The horizontal momentum equation for component x is as follows (DHI, 2010) :

$$\frac{\partial u}{\partial t} + \frac{\partial u^2}{\partial x} + \frac{\partial uv}{\partial y} + \frac{\partial wu}{\partial z} = fv - g \frac{\partial h}{\partial x} - \frac{1}{r_0} \frac{\partial p_a}{\partial x} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial x} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{xx}}{\partial x} + \frac{\partial S_{xy}}{\partial y} \right) + F_u + \frac{\partial}{\partial z} \left(v_t \frac{\partial u}{\partial z} \right) + u_s S$$

While the horizontal momentum equation for component y is as follows (DHI, 2010):

$$\frac{\partial v}{\partial t} + \frac{\partial v^2}{\partial y} + \frac{\partial uv}{\partial x} + \frac{\partial wv}{\partial z} = -fu - g \frac{\partial h}{\partial y} - \frac{1}{r_0} \frac{\partial p_a}{\partial y} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial y} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{yx}}{\partial x} + \frac{\partial S_{yy}}{\partial y} \right) + F_v + \frac{\partial}{\partial z} \left(v_t \frac{\partial v}{\partial z} \right) + v_s S$$

where x, y, z = cartesian coordinates of $x, y,$ and z direction; t = time; u = water particle speed of direction x ; v = water particle speed of direction y ; w = water particle speed of direction z ; η = water surface elevation; d = sea depth; h = total depth of the sea; f = coriolis parameter ($f = 2\omega \sin\theta$); Ω = ratio of earth revolution; θ = latitude coordinates; g = acceleration of gravity; $S_{xx}, S_{xy}, S_{yy}, S_{yy}$ = component of radiation stress tensor; ν_t = vertical eddy viscosity; p_a = atmospheric pressure; ρ_0 = sea density; S = debits due to sources; u_s = particle velocity of air sources direction x ; and v_s = the particle speed of air sources direction y .

Wave modeling used directional decoupled parametric and fully spectral formulations based on wave eternity movements with the input data of on-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry data, wind speed and direction and wave data. Parameterization was created in the frequency domain by introducing the 0 and 1st moments of the wave spectrum as a variable (Azhar *et al.*, 2011). The arranger equation was the equilibrium wave force in cartesian coordinate formulated as follow (DHI, 2010):

$$\frac{\partial N}{\partial t} + \frac{\partial}{\partial \phi} C\phi N + \frac{\partial}{\partial \lambda} C\lambda N + \frac{\partial}{\partial \sigma} C\sigma N + \frac{\partial}{\partial \theta} C\theta N = \frac{S}{\sigma}$$

Source energy, S , shows the superposition of surface functions in various styles (Azhar *et al.*, 2011).

$$S = S_{in} + S_{nl} + S_{ds} + S_{bot} + S_{surf}$$

where $N_{x,\sigma,\theta,t}$ = power density, $\phi\lambda$ = coordinate spherical, C = speed of propagation of four-dimensional wave group, S = energy source, S_{in} = formation by wind, S_{nl} = energy transfer, non-linear, S_{ds} = wave energy due to white capping, S_{bot} = by bottom friction, S_{surf} = dissipation due to the depth-induced breaking.

We took the maximum value of the current speed and wave height in each grid that comes from 4 seasonal patterns characteristics, that is the west monsoon (December), transition 1 (May), east monsoon (July), and transition 2 seasons (October). The results of the 2D hydrodynamic model form the patterns of current speed and wave height every season were interpolated with a spline method with a cell size of 5 x 5 meters and analyzed using cell statistics to obtain the maximum value of current speed and wave height. The results of the distribution map were classified according to suitability class in table 2.

2.4. Determination of Area Suitability Criteria

The determination of the weighting and scoring was done to provide value in each criterion for supporting the feasibility of seaweed cultivation. The weight determination of each criterion was based on the literature research and later on was used to weight the variables by priority. The variable that is very influential for the growth of seaweed got a high score. The determination was based on the defined suitability matrix as shown in Table 2.

The total score multiplication and weight in table 2 were used to determine the class of water suitability of seaweed cultivation with quantitative analysis using the following equation :

$$Y = \sum a_i \cdot X_n$$

The range of suitability cultivation area of *K. Alvarezii* was adjusted to use the following equation :

$$I = \frac{(\sum a_i \cdot X_n) - (\sum a_i \cdot X_n)_{min}}{k}$$

where, Y = The final value, a_i = weight per parameter, and X_n = score of suitability level, I = interval of suitability class, k = number of suitability class.

Finally, the suitability interval was divided into 5 classes, 1) highly suitable (HS) is in the range of the value 42-50, 2) suitable category (S) is in the range of 33-41, C) moderately suitable (MS) range of 24-32, 4) less suitable (LS) are in the range of 15-23, and 5) not suitable (N) had the value range of < 15. Each parameter in table 2 was transformed in digital format using a spline interpolation method with a cell size of 5 x 5 meters resulting in an index map of water suitability.

2.5. Distance Suitability Analysis

Distance suitability analysis was also carried out as the other limiting factors out of the water parameters. Those were the distance to shipping lanes or sea lanes and the distance to a settlement which are arranged in a digital format with 5 x 5 -meter cell-sized using the buffer and euclidean distance method to consider the accessibility and disruption of marine traffic. These criteria are described in Table 3. The suitability of distance to the settlement was compiled using the data of inhabited island, namely Bontosua, Sanane, Panjenekang, Balanglombo, and Balangcaddi Islands, while the shipping lanes data used was inter-regional shipping routes. The results of the analysis were integrated with a water suitability index map.

2.6. Spatial Modelling

The spatial modelling in this study used cell-based modelling by integrating all oceanographic and water quality parameters based on the suitability classification according to table 2 with the following modelling algorithms:

$$WS = (T_i \times 0.5) + (pH_i \times 0.5) + (S_i \times 1) + (DO_i \times 0.5) + (N_i \times 1.5) + (F_i \times 1.5) + (B_i \times 0.5) + (HS_i \times 1.5) + (CS_i \times 1.5) + (TSS_i \times 0.5) + (SB_i \times 0.5)$$

with, WS = Water suitability, T_i = temperature index map, pH_i = pH index map S_i = salinity index map, DO_i = Dissolved Oxygen index map, N_i = Nitrate index map, F_i = phosphate index map, B_i = bathymetry index map, HS_i = significant wave height index map, CS_i = current speed index map, TSS_i = TSS index map, SB_i = seabed index map

The result of this modeling is continued by determining the classification of water suitability based on a class of suitability table 2. Then, integrating the results of water suitability above with the result of distance suitability analysis according to table 3 using cell statistics analysis.

2.7. Estimating Productivity

After obtaining the distribution of classes areas for all variable, the seaweed production can be estimated only by considering the area which is suitable or highly suitable. Longline method of seaweed cultivation unit was designed as 25 x 50 meter per unit, the distance between the rope was 1 meter and the planting distance of each tied seaweed was 25 cm (WWF, 2014). The initial weight of the seedlings was approximately 100 grams per bunch (Parakkasi *et al.*, 2020) with the wet weight crop was 500 grams per tie. The cultivation of seaweed with a floating rope/longline system produces an excellent product and can gain a weight up to four to five times from the initial weight of the seedlings (100 grams) in a period of cultivation time. The number of the allocated cultivation units was estimated to be 5 units/ha. The distance correction between units is 10 meters in width (twice longer than the average length of the fishing boat) for fishing traffic. This amount represents 62.5% of the maximum number of units i.e., 8 units/ha. To estimated production is the multiplication between the number of cultivation units (from the total suitable area) and once crop yields per unit with the following equations:

$$Y_e = N \times C_y$$

with, Ye = estimated yield per cycle (tons), N = number of cultivation unit, Cy = crop yield per cycle (tons). Briefly, the research method is summarized in the flowchart presented in Figure 2.

3. Result and Discussion

3.1. Distribution of Water Quality Parameters

Spatial distributions of the suitability of water quality parameters are shown in Figure 3. It can be seen that the water quality conditions based on the variable of SST, and pH, are highly suitable for seaweed cultivation in the study area. SST is the important limiting factor for marine organisms (Wirasatriya *et al.*, 2019). The range of measured temperature in the study area is very narrow i.e., only around 30°C. According to BSN (2005) the optimal temperature for seaweed culture is 26-32°C, so these temperatures are feasible for this activity. Whereas according to Andi *et al.* (2016), *K. alvarezii* grown well at the temperature range of 27°C–30°C. The high temperature of the water causes death on seaweed while low water temperatures cause damage to protein and fat membranes and it is very influential in the seaweed life cycle. Photosynthesis on seaweed generally lasts at a maximum temperature of 23°C–30°C (Msuya, 2011; Ding *et al.*, 2013). Seawater has an enormous support ability to prevent pH fluctuation. The condition of pH that is very acidic or alkaline will endanger the survival of organisms because it led to metabolic and respiration disorders. The pH range at the study site is 7.26–7.63. These values are feasible for seaweed cultivation (Rahman *et al.*, 2019). Based on Noor (2015), a suitable pH range for *K. alvarezii* cultivation is in normal and disposed alkaline level. n. Optimal seaweed growth requires a pH value range of 7.5–8.5 (PI, 2004; BSN, 2015; Radiarta *et al.*, 2018).

The suitability distribution for TSS and substrate type are moderately suitable or more in the study area. A substrate is dominated by reef type almost at the entire study area which is categorized as moderately suitable. The suitable condition is located around the small islands with the sand as the substrate type. Seaweed will grow optimally when the cultivation site is in coral, rubble, sand-based substrate, or a mixture of all three. The most optimum substrate type for seaweed cultivation is a hard substrate like the rocky coral then followed by sandy corals (Parakkasi *et al.*, 2020). Waters with a fractional substrate of coral to sand are usually passed by the current that was suitable for the growth of seaweed. However, cultivating seaweed with a longline method is not overly influenced by the substrate type. For TSS, the eastern part of the study area is highly suitable while the western part is moderately suitable. This indicates that the further distance from the mainland the lower TSS is observed. TSS is insoluble sedimentation which increases the turbidity of water (Nurdin *et al.*, 2019). High turbidity due to the suspended material impedes light penetration into the water column and affects seaweed growth. Thus, TSS indicates water pollution as *K. alvarezii* lives optimally in transparent oligotrophic waters (Hayashi *et al.*, 2011). The TSS levels in the study area range between 3.6–57.6 mg/L which means that the waters are suitable for the growth of *K. alvarezii*. Low TSS concentration is probably due to the absence of re-suspension due to the non-silt/clay of substrate type or no input from coastal sediment. Total dissolved solids (TDS) still meets the seawater quality standard for marine biota according to PI (2004), namely the standard quality value for coral and seagrass life is < 20 mg/L.

For DO and salinity, the suitability distributions vary from less suitable to highly suitable. Based on the DO criterion, the study area is mostly highly suitable for seaweed cultivation, with only 2 spots in the study area which are less suitable. The water circulation among small islands plays an important role in the high DO level. In open water, the low concentration of oxygen in a natural condition is rarely found. The optimal dissolved oxygen for seaweed cultivation is in the range of 3–7 mg/l (Rahman *et al.*, 2019). For salinity, the less suitable classes are spread in some areas and even the class at the eastern part of Sanane Island is not suitable for seaweed cultivation. The stability of the salinity becomes an important factor for seaweed because seaweed is a stenohaline organism that cannot resist the high fluctuation salinity (Reis *et al.*, 2010). The extreme salinity changes can lead to ice-ice disease. Salinity

can affect the osmoregulation process of marine biota including seaweed. However, most of the area is still feasible for seaweed cultivation.

Nitrate and phosphate concentrations as indicators of water fertility are relatively low in this region. This may be related to the absence of big rivers in the small islands of the study area as the nutrient sources. Nitrate is a limiting factor for seaweed growth. Nitrate is one of the nutrients needed for the growth of seaweed. The result shows that nitrate concentrations in the study area range between 0.001-0.12 mg/l which is relatively low. According to Kautsari and Ahdiansyah (2015), the feasible nitrate concentration for seaweed cultivation ranges from 0.01 mg/l to 0.07 mg/l, while the concentration of <0.01 mg/l and > 0.1 mg/l are not suitable for seaweed cultivation. Phosphate can be a limiting factor due to the source limitation. In the study area, phosphate concentration ranges from 0.001 mg/l to 0.062 mg/l which indicates low to insufficient fertility levels. Phosphate concentration for the fertile water should range from 0.051 mg/l to 0.1 mg/l. Besides, phosphate concentration also controls the carrageenan content of seaweed (Patty *et al.*, 2015).

The depth of the study area ranges from 2 m to 39.2 m. The longline method can be applied at the 1 m depth from the lowest tide or below (WWF, 2014). The sufficient depth provides seaweed to be submerged. Nevertheless, noted that the deeper waters will increase the cost of constructing seaweed cultivation. Thus, the areas with a depth of more than 15 m are categorized as not suitable for seaweed cultivation.

The distribution of physical and chemical parameters at the seaweed site is highly determined by the wave and current. These aspects play a pivotal role in the water movement for algae growth (Gaylord *et al.*, 2007; Kregting *et al.*, 2013). The variability of current speed and wave height is distinguished by the 4 different characteristics of the west monsoon; transition 1; east monsoon; and transition 2 based on the results of 2D hydrodynamic modeling (Figure 4 and Figure 5). During west monsoon and transition 1 (east monsoon and transition 2), strong current speed appears at the northwestern (southern) part of the study area. The strongest current speed occurs during the transition 1 season. The pattern of the surface current in the study area is influenced by the current pattern of breaking waves in the small islands waters and the surrounding shallow area; and the influence of wind and tidal conditions in these waters. Based on the criteria shown in Table 2, the study area is categorized as highly suitable (dark blue color) to less suitable (light blue color).

For the wave height, almost entire study areas are categorized as suitable and highly suitable for seaweed cultivation except during east monsoon. During west monsoon transition 1 and transition 2, the blue and dark blue colors mean that the wave height ranges from 0.1 m to 0.3 m. During east monsoon, the wave higher than 0.3 m occurs at almost the entire study area which makes it less suitable for seaweed cultivation.

The protected area is determined by several types of physical conditions such as current velocity, and wave height. The influence of strong current and ocean waves will drag and damage the cultivated seaweed and its system. The maximum value of the distribution of current speeds and wave height for 4 seasons is described in Figure 6. Based on the results in Figure 6, the current speed and wave height around the study of the water are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. The less suitable areas for the current speed (wave height) category only appear mostly in the northeastern (northern) part of the study area. There is no area with a current speed of more than 0.5 m/s and the wave height more than 0.5 m which makes the area is not suitable for seaweed cultivation. The absence of strong current and wave may correspond to the morphology of the study area which is categorized as coral/reef waters with a lot of shallow/reef flat area located within the small islands that protect the area from the high wave and strong current. As mentioned by Sulma *et al.* (2008), the types of protected waters are bays, caps, straits, lagoons, and reef flat. Furthermore, Helmi *et al.* (2017) also stated that the coral reef is a shoreline protector.

3.2. Potential Areas for Seaweed culture Development and Production

Various suitability criteria as limiting factor of site selection for seaweed cultivation include physical and chemical parameters and distance suitability for accessibility and restricted area inside the shipping lanes has been determined into 5 highly suitable classes (HS), suitable (S), moderately suitable (MS), less suitable (LS) and not suitable (N) (Figure 7). Figure 7 shows that most of the study areas are moderately suitable and suitable for seaweed cultivation. The area which is not suitable is only located along the shipping lane. Figure 8 summarizes the areal size of each criteria as follows: the highly suitable (HS) is 83.2 ha (0.42%); suitable (S) is 4,463.08 ha (22.21%); moderately suitable (MS) is 9,983.52 ha (49.68%); less suitable (LS) is an area of 494.84 ha (2.46%); and not suitable (N) is 5,071.15 ha (25.23%). The potential area for seaweed cultivation development reaches 4,546.28 ha as the sum area of highly suitable and suitable criteria. This result is more than obtained by Arifin *et al.* (2014) which only found 1,963.6 ha area which is suitable and highly suitable for *K. alvarezii* cultivation in the same study area. This indicates that our method can optimally classify the potential area for *K. alvarezii* cultivation. Furthermore, this potential area can be utilized for 22,730 cultivation units with seaweed production reaching about 56,825 tons of wet weight per crop, or within a year (2 times of harvesting) can reach about 113,650 tons. This amount is only about one-third of the total seaweed production in Pangkep Regency in 2018. However, it is important to be noted that, there are some limitations in the present study which should be considered. For example, the present study does not include the long-term climate variability which may also influence the physical and chemical parameters in the study area. As reported by Kunarso *et al.* (2019) and Msuya *et al.* (2014). The climate variability influences the oceanographic condition which then affects fisheries activity such as aquaculture and fish catch. The present study also does not consider the risk analysis which can disturb the cultivation process as shown by Castelar *et al.*, (2015). Furthermore, to maintain the carrying capacity of the study area, the area that might be utilized for the expansion of *K. alvarezii* cultivation is about 50% (covering 2,273.14 ha) of the total suitable area accordingly. The result of the present study indicates the effectiveness of geospatial modeling for developing the area of *K. alvarezii* culture which can be applied in the other area.

4. Conclusions

We have conducted a comprehensive geospatial modeling approach for the site selection of *K. alvarezii* cultivation at the Spermonde Islands. Physical and chemical parameters as well as the accessibility and restricted area of shipping lanes were used to define the suitability criteria for *K. alvarezii* cultivation. The result is SST and pH are categorized as highly suitable for *K. alvarezii* cultivation in the whole study area. A substrate is dominated by reef type almost at the whole study area which is categorized as moderately suitable. TSS levels in the study area range between 3.6-57.6 mg/l which means that the waters are suitable for the growth of *K. alvarezii*. For DO and salinity, the suitability distributions vary from less suitable to highly suitable. Nitrate and phosphate concentrations mainly are too low which are not suitable for *K. alvarezii* cultivation. The depth at the study area ranges from 2 m to 39.2 m. The area with a depth more than 15 m is categorized as not suitable for *K. alvarezii* cultivation. The current speed and wave height around the study of the water are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. After summarizing all criteria according to their weighting score, the suitable for seaweed cultivation at the study area is 4,546.28 ha, which is the combination of the suitable (S) area of 4,463.08 ha and highly suitable criteria (HS) with an area of 83.2 ha. The potential production per crop can reach 56,825 tons per cycle and 113,650 tons per year.

Acknowledgment

The author thanked the Ministry of Research Technology and Higher Education of the Republic of Indonesia for the financial support of research. The author thanked the Center for Coastal Rehabilitation and Disaster Mitigation Studies (CoREM), Diponegoro University for the facilities' support on hydro-oceanographic modeling and geospatial data processing. We also thanked the Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University for the surveys and laboratory analysis of water quality data.

References

- Afandi, A. Musadat, F. 2018. Suitability analysis for seaweed culture area at the waters of Kamelanta Village and Panjang Island using Geographical Information System. *Jurnal Akuakultura* 2(1), 69-78. (In Bahasa)
- Arbit, N., Omar, S., Soekendarsi, E., Yasir, I., Tresnati, J., Tuwo, A. 2019. Morphological and genetic analysis of *Gracilaria* sp. cultured in ponds and coastal waters. IOP Conf. Ser.: Earth Env. Sci. 370 012018.
- Arifin T., Bohari, R., Arlyza, I.S. 2014. Analysis of Space Suitability Based on Marine Agriculture in The Small Islands Around Makassar. *Forum geography* 28 (1), 91-102. DOI: <https://doi.org/10.23917/forgeo.v28i1.441>
- Ariyanto, F. Y., Astor, Y., Sidqi, M. 2018. Application of water boundary theory for determining site licenses at the coastal area and small islands in Indonesia. *Proceeding National Seminar of Geomatics 2018*. Bandung, pp. 893-900. (In Bahasa)
- Azhar, R. M., Wurjanto, A., Yuanita, N. 2011. Study of coastal protection of submerged breakwater at Tanjung Kait beach. Master Thesis. Master Program of Water Resource Management, Bandung Institute of Technology, Indonesia.
- Bixler, H. J., Porse, H. 2011. A Decade of Change in the Seaweed Hydrocolloids Industry. *J. Appl. Phycol.* 23, 321-335.
- Castelar, B., de Siqueira, M. F., Sánchez-Tapia, A., Reis, R. P. 2015. Risk Analysis Using Species Distribution Modelling to Support Public Policies for The Alien Alga *Kappaphycus alvarezii* Aquaculture in Brazil. *Aquacult.* 446, 217-226.
- Dee, D. P. *et al.* 2011. The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Q.J.R. Meteorol. Soc.* 137(656), 553–597. doi: 10.1002/qj.828
- DHI. 2010. MIKE-21 and MIKE-3 Flow Model. Hydrodynamic Module Step-by-Step Training Guide. DHI MIKE, Denmark. 96 p.
- Ding L., Ma, Y., Huang, B., Chen, S. 2013. Effects of Seawater Salinity and Temperature on Growth and Pigment Contents in *Hypnea cervicornis*. *J. Agardh (Gigartinales, Rhodophyta)*. *BioMed Res. Int.* 2013, 594308. <https://doi.org/10.1155/2013/594308>
- Gaylord, B., Rosman, J. H., Reed, D. C., Koseff, J. R., Fram, J., Macintyre, S., Arkema, K., McDonald, C., Brzezinski, M. A., Largier, J. L., Monismith, S. G., Raimondi, P. T., Mardian, B. 2007. Spatial Patterns of Flow and Their Modification within and Around A Giant Kelp Forest. *Limnol. Oceanogr.* 52, 1838–1852.
- Hayashi, L., Santos, A. A., Faria, G. S. M., Nunes, B. G., Souza, M. S., Fonseca, A. L. D., Barreto, P. L. M., Oliveira, E. C., Bouzon, Z. L. 2011. *Kappaphycus alvarezii* (Rhodophyta, Areschougaceae) Cultivated in Subtropical Waters in Southern Brazil. *J. Appl. Phycol.* 23, 337–343.
- Helmi, M., Satriadi, A., Suryoputro, A. A. D., Marwoto, J., Setiyono, H. Hariyadi. 2018. Rehabilitation Priority Area Assessment on Death Coral using Cell-Based Modelling Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.* 9(11), 2949-2961.
- Helmi, M., Purwanto, Atmodjo, W., Subardjo, P., Aysira, A. 2017. Benthic Diversity Mapping and Analysis Base on Remote Sensing and Seascape Ecology Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.* 9(10), 227-235.
- Jalil, A. R., Samawi, M. F., Azis, H. Y., Jaya, I., Malik, A., Yunus, I., Sohopi, M. A. A. 2020. Comparison of physical-chemical conditions for seaweed cultivation in the Spermonde Archipelago, Indonesia. *AAFL Bioflux* 13(4), 2071-2082.

- Junaidi, M., Lestary, D. P., Scabra, A. R. 2018. Mariculture Site Selection Based on Environmental Parameters in Tanjung and Gangga Sub-District, North Lombok. In: Proceeding of 1st ICWBB 2018. Mataram University, Mataram, pp. 233-251.
- Kaiser, M. J., Yu, Y., Snyder, B. 2010. Economic Feasibility of Using Offshore Oil and Gas Structures in the Gulf of Mexico for Platform-based Aquaculture. *Mar. Pol.* 34, 699-707.
- Kautsari, N., Ahdiansyah, Y. 2015. Carrying capacity and suitability assessment of Labuhan Terata waters, Sumbawa for the development of seaweed culture. *Jurnal Ilmu Kelautan* 20(4), 233-238. (In Bahasa)
- Kim, J. K., Yarish, C., Hwang, E. K., Park, M., Kim, Y. 2017. Seaweed Aquaculture: Cultivation Technologies, Challenges, and Its Ecosystem Services. *Algae*. 32(1), 1-13.
- Kregting, L.T., Blight, A. B., Elsässer, B., Savidge, G. 2013. The Influence of Water Motion on The Growth Rate of The Kelp *Laminaria Hyperborea*. *J. Exp. Mar. Biol. Ecol.*, 448, 337–345.
- Kunarso, Wirasatriya, A., Irwani, Satriadi, A., Helmi, M., Prayogi, H. Munandar, B. 2019. Impact of Climate Variability on Aquatic Productivity and Fisheries Resources in Jepara. IOP Conf. Ser.: Earth Env. Sci., 246 012021: 1-12.
- Longdill, P. C., Healy, T. R., Black, K. P., 2008. An integrated GIS approach for sustainable aquaculture management area site selection. *Ocean Coast. Manag.* 51, 612-624. doi:10.1016/j.ocecoaman.2008.06.010
- Malczewski, J. 2006. Ordered weighted averaging with fuzzy quantifiers: GIS-based multicriteria evaluation for land-use suitability analysis. *Int. J. Appl. Earth Observ. Geoinf.* 8(4), 270- 277.
- Malczewski, J., Rinner, C. 2015. *Multicriteria Decision Analysis in Geographic Information Science*. Springer, Berlin
- Micael, J., Costa, A.C., Aguiar, P., Medeiros, A., Calado, H. 2015. Geographic Information System in A Multi-criteria tool for Mariculture Site Selection. *Coast. Manag.* 43(1), 52-66.
- Msuya, F. E., Buriyo, A., Omar, I., Pascal, B., Narrain, K., Ravina, J. J. M., Mrabu, E., Wakibia, J. G. 2014. Cultivation and Utilisation of Red Seaweeds in the Western Indian Ocean (WIO) Region. *J. Appl. Phycol.* 26, 699–705.
- Msuya, F. E. 2011. Environmental Changes and Their Impact on Seaweed Farming in Tanzania. *World Aquacult.* 42, 34-37.
- Nirmala, K., Ratnasari, A., Budiman, S. 2014. Site Selection for Seaweed Culture at Gerupuk Bay-West Nusa Tenggara Using Remote Sensing and GIS. *Jurnal Akuakultur Indonesia* 13(1), 73-82.
- Noor, N. M. 2015 The Water Bodies Compatibility Analysis for Culturing Brown Seaweed *Kappapycus alvarezii* in Ketapang Seashore, South Lampung. *Maspari Journal* 7(2), 91-100. (in Bahasa)
- Nuridin, N., Helmi, M., Setiawan, R. Y., Zainudin, M., Rustam, A., Akbar, M. 2019. Landsat Capability to Mapping of Total Solid Suspended (TSS) and Coastal Ecosystem. *ISPRS – Int. Arch. Photogram., Remote Sens. Spatial Inf. Sci.*, 42(4), 497-502.
- Parakkasi, P., Rani, C., Syam, R., Zainuddin, Achmad, M. 2020. Growth Response and Quality of Seaweed *Kappaphycus alvarezii* Cultivated in Various Coastal Ecosystems in the Waters of West Sulawesi, Indonesia. *AACL Bioflux* 13(2), 627-639.
- Patty, S. I., Arfah, H., Abdul, M. S. 2015. The relationship among Nutrient (Phosphate and Nitrate), DO, and pH with the water fertility in Jikumerasa waters, Buru Island. *Jurnal Pesisir dan Laut Tropis* 1(1), 43-50. (In Bahasa)

- Pickering, T. D., Skelton, P. Sulu, J. R. 2007. Intentional Introductions of Commercially Harvested Alien Seaweeds. *Bot. Mar.* 50, 338–350.
- Radiarta, I. N., Saputra, A. Albasri, H. 2012. Feasibility mapping of seaweed culture area (*Kappaphycus alvarezii*) at Bintan Regency, Riau Islands Province using GIS and remote sensing approach. *Akuakultur* 7(1): 145-157. (In Bahasa)
- Radiarta, I. N., Saitoh, S. Yasui, H. 2011. Aquaculture Site Selection for Japanese Kelp (*Laminaria japonica*) in Southern Hokkaido, Japan, Using Satellite Remote Sensing and GIS-based Models. *ICES J. Mar. Sci.* 68(4), 773-780. doi:10.1093/icesjms/fsq163
- Rahman, I. S., Fadjar, M., Tjahjono, A. 2019. The Relationship Water Physical-Chemical Parameters in Seaweed Cultivation *Eucheuma cottonii* with Long-line System. *Int. J. Sci. Tech. Res.* 8(11), 1448-1452.
- Rahmayanti, F., Diana, F., Kusumawati, I. 2018. Suitability analysis for area development of seaweed (*Eucheuma cottonii*) culture in Lhok Bubon waters, Samatiga District, Aceh Barat Regency. *Jurnal Akuakultura* 2(1), 26-32. (In Bahasa).
- Reis, R. P, Loureiro, R. R., Mesquita, F. S. 2010. Do Salinity Affect Growth and Carrageenan Yield of *Kappaphycus alvarezii* (Gigartinales/ Rhodophyta)?. *Aquacult. Res.* 42, 1231-1234.
- Rusliani. 2016. Study of water quality on the growth of seaweed *Gracilaria verrucosa* in a pond with verticulture method. *Octopus J. Fish. Sci.* 5(2), 522-527. (in Bahasa)
- Setiawan, R. Y., Kawamura, H. 2011. Summertime Phytoplankton Bloom in the south Sulawesi Sea. *IEEE J. Sel. Topics Appl. Earth Observ. Remote Sens.* 4(1), 241-244. doi:10.1109/JSTARS.2010.2094604.
- Sousa, F. E. S., Moura, E. A., Marinho-Soriano, E. 2012. Use of Geographic Information Systems (GIS) to Identify Adequate Sites for Cultivation of The Seaweed *Gracilaria birdie* in Rio Grande Do Norte, Northeastern Brazil. *Revista Brasileira de Farmacognosia* 22(4), 868–873.
- Sulma, S., Manoppo, A. K. S., Hartuti, M. 2008. Identification of Suitable Area for Seaweed Culture in Bali Waters Based on Remote Sensing Satellite Data. *Int. J. Remote Sens. Earth Sci.* 5, 57-66.
- Susanto, R. D., Wei, Z., Adi, T. R., Zheng, Q., Fang, G., Bin, F., Supangat, A., Agustyadi, T., Li, S., Trenggono, M., Setiawan, A. 2016. Oceanography Surrounding Krakatau Volcano in the Sunda Strait, Indonesia. *Oceanogr.* 29(2), 228-237. <http://dx.doi.org/10.5670/oceanog.2016.31>.
- Teniwut, W. A., Marimin, Djatna, T. 2019. GIS-Based multi-criteria decision-making model for site selection of seaweed farming information center: A lesson from small islands, Indonesia. *Decis. Sci. Let.* 8(2019), 137-150 doi: 10.5267/j.dsl.2018.8.001
- Utama, R. Z., Handayani, T. 2018. Suitability Between Oceanography and Seaweed (*Eucheuma cottonii*) Cultivation Potential in Tidung Island with Geographic Information System (GIS). *ICENIS: E3S Web of Conference* 73, 12009. 1-5.
- Valderrama, D., Cai, J., Hishamunda, N., Ridler, N., Neish, I. C., Hurtado, A. Q., Msuya, F. E., Krishnan, M., Narayanakumar, R., Kronen, M., Robledo, D., Gasca-Leyva, E., Fraga, J. 2015. The Economics of *Kappaphycus* Seaweed Cultivation in Developing Countries: A Comparative Analysis of Farming Systems. *Aqua. Econ. Manag.* 19, 251–277.
- Wirasatriya, A., Sugianto, D. N., Helmi, M., Setiawan, R. Y. Koch, M. 2019. Distinct Characteristics of SST Variabilities in the Sulawesi Sea and the Northern Part of the Maluku Sea during the Southeast Monsoon. *IEEE J. Sel. Topics Appl. Earth Obs. Remote Sens.* 12(6), 1763-1770.

- WWF-Indonesia. 2014. Better Management Practices: Guideline series of small scale fisheries of seaweed culture - Cottoni (*Kappaphycus alvarezii*), Sacol (*Kappaphycus striatum*) dan Spinosum (*Eucheuma Denticulatum*) Version 1. WWF-Indonesia. South Jakarta. 53 p. (in Bahasa)
- Yulianto, H., Damai, A. A., Delis, P. C., Elisdiana, Y. 2017. Spatial Analysis to Evaluate the Suitability of Seaweed Farming Site in Lampung Bay, Indonesia. *Turkish J. Fish. Aquatic Sci.* 17, 1253-1261.

List of Tables

Table 1. Water quality data collection

No	Variables	Unit	Tools/Methods	Data Retrieval Methods
1	Sea Surface Temperature (SST)	°C	Thermometer	In situ
2	Salinity	Psu	Salinometer	In situ
3	pH	-	<i>Water quality checker</i>	In situ
4	Dissolved oxygen (DO)	ppm	Winkler method	Laboratory analysis
5	Nitrate	mg/l	Ascorbic Acid Method	Laboratory analysis
6	Phosphate	mg/l	Brucine Sulfate Method	Laboratory analysis
7	Total Suspended Solids (TSS)	mg/l	<i>Milipore filter</i>	Laboratory analysis
8	Substrate type	-	Visually observation	In situ
9	Depth	m	Modelled depth from LPI Maps (BIG, 2011) and BATNAS (BIG, 2018) data	Integrating modelled data

Table 2. Table of suitability criteria for seaweed *Kappaphycus alvarezii* cultivation

Variables	Level of conformity (score)					Weight	Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)		
SST (°C)	28-30	24-27 or 31-32	22-23 or >32	21 or >34	<21 or >36	0.5	Modification of Noor (2015); Rahmayanti <i>et al.</i> (2018)
pH	7-8	6.5-7 or 8-8.5	6-6.4 or >8.5	4-6	<4 or >9	0.5	Modification of Kautsari and Ahdiansyah (2015); Noor (2015)
Salinity (psu)	28-31	26-28 or 31-33	<26 or ≥ 34	18-25 or 35-36	<18 or >37	1	Modification of Kautsari and Ahdiansyah (2015); Rahmayanti <i>et al.</i> (2018); Utama and Handayani (2018)
DO (mg/l)	>6	5-6	4-5	2-4	<2	0.5	Modification of Afandi and Musadat (2018)
Nitrate (mg/l)	0.04-0.07	0.02-0.03	0.01 or >0.1	<0.01 or >0.5	>1	1.5	Modification of Kautsari and Ahdiansyah (2015); Afandi and Musadat (2018)
Phosphate (mg/l)	0.051-0.1	0.02-0.04	0.01-<0.02 or >0.2-1	<0.01 or >1	>2	1.5	Modification of Patty <i>et al.</i> (2015)
Depth (m)	3-10	2-3 or 11-13	1-2	>13-15	<1 or >15	0.5	Modification of Tiensongrusme (1990) in Radiarta <i>et al.</i> (2012); Kautsari and Ahdiansyah (2015); Yulianto <i>et al.</i> (2017)
Wave height (m)	0.2-0.3	0.1-0.19	>0.3	<0.1 or >0.4	>0.5	1.5	Modification of Rahmayanti <i>et al.</i> (2018)
Current speed (m/s)	0.25-0.3	0.2-0.24 or 0.31-0.4	0.1-0.19 or >0.4	<0.1 or >0.45	>0.5	1.5	Modification of Utama and Handayani (2018)
TSS (mg/l)	<25	25-50	50-100	100-400	>400	0.5	Modification of Yulianto <i>et al.</i> (2017)
Seabed	Coral, Rubbel	Sand	Reefs	Muddy Sand	Mud	0.5	Modification of Noor (2015); Yulianto <i>et al.</i> (2017)

Note : HS: Highly suitable, S: suitable, MS: moderately suitable, LS: less suitable, and N: not suitable

Table 3. Distance suitability analysis

Parameters	Suitability Level (Score)					Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)	
Shipping lanes (m)	> 500	-	-	-	≤ 500	Ariyanto <i>et al.</i> (2018)
Distance to settlements (km)	4	4-5	5-6	6-7	>7	Modification of Radiarta <i>et al.</i> (2012)

List of Figures




Figure 1. Study area




Figure 2. Research Flowchart




Figure 3. Distribution of physical and chemical parameters




Figure 4. Distribution of current speed (m/s) based on 2D hydrodynamic model in each season




Figure 5. Distribution of significant wave height (m) based on 2D hydrodynamic model in each season




Figure 6. Maximum statistical value of current speed (left) and wave height (right) for all season




Figure 7. Suitability distribution for cultivation area of *K. Alvarezii*




Figure 8. Potential area for *K. alvarezii* cultivation in each



Sarjito Sarjito <sarjito@live.undip.ac.id>

[IK.IJMS] Editor Decision

3 messages

Ilmu Kelautan <ijms.undip@gmail.com>
To: sarjito@live.undip.ac.id

Wed, Jul 27, 2022 at 8:03 AM

Dear
Sarjito

We have reached a decision regarding your submission to ILMU KELAUTAN: Indonesian Journal of Marine Sciences, "Site selection of Kappaphycus alvarezii farming to optimize seaweed production using Geographic Information system with geospatial modeling at Spermonde Islands waters, South Sulawesi, Indonesia".


Our decision is to:
Revision Required

Indonesian Journal of Marine Science
ijms.undip@gmail.com

REVIEW A

1. Hydrodynamics of marine waters has dynamic properties divided into 4 seasonal models, namely: West Monsoon, Transition 1, East Monsoon, and Transition 2. This article will be interesting if each model is made a model for the distribution of Kappaphycus alvarezii cultivation locations.
2. Where are recommendations for selecting and developing Kappaphycus alvarezii cultivation sites? whether in HS, S, MS?

INDONESIAN JOURNAL OF MARINE SCIENCE
<http://ejournal.undip.ac.id/index.php/ijms>

 **47735RV.docx**
2251K

Site selection of *Kappaphycus alvarezii* farming to optimize seaweed production using Geographic Information system with geospatial modeling at Spermonde Islands waters, South Sulawesi, Indonesia

Sarjito Sarjito^{1*}, Hanny Ammaria¹, Muhammad Helmi^{2,3}, Slamet Budi Prayitno¹, Nurjannah Nurdin^{4,5}, Riza Yuliratno Setiawan⁶, Parichat Wetchayont⁷, Anindya Wirasatriya^{2,3}

¹Aquaculture Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro, Indonesia

Jl. Prof. Soedarto, SH., Tembalang, Semarang, Central Java (50275) Indonesia

²Oceanography Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro, Indonesia

³Center for Coastal Rehabilitation and Disaster Mitigation Studies (CoREM), Indonesia

⁴Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University, Indonesia

⁵Marine Science and Fisheries Faculty, Hasanuddin University, Indonesia

⁶Marine Science and Fisheries Faculty, Gadjah Mada University, Indonesia

⁷Department of Geography, Srinakharinwirot University, Thailand

e-mail: sarjito@live.undip.ac.id

Abstract

Indonesia is the largest producer of red algae in the world with one of the major *K. alvarezii* farming areas within the Indonesian seas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province. *K. alvarezii* production in Pangkep Regency increased year by year. However, this production still can be increased by expanding the utilized areas for seaweed farming. Proper site selection is crucial to guarantee the success of the seaweed farming development. We conducted a Geographic Information system (GIS) with geospatial modeling approach for the site selection of *K. alvarezii* cultivation at the Liukang Tuppabiring District, Spermonde Islands by combining various physical and chemical parameters obtained from in-situ data and hydrodynamics modeling. The parameters are Sea Surface Temperature, (SST), salinity pH, substrate type, current speed, wave height, DO, nitrate, phosphate, and Total Suspended Solid (TSS). We also include the accessibility and the restricted area of shipping lanes for determining the area suitability of *K. alvarezii* cultivation. This research demonstrates the cell-based geospatial modeling can be effectively used and found the suitable categorized area for *K. alvarezii* culture of 4,546.28 ha. The Potential productivity of wet-weight seaweed can reach 56,825 tons per cycle or 113,650 tons per year. This result indicates the GIS with geospatial modeling approach for developing the area of *K. alvarezii* culture which can be applied in the other area.

Keywords: Seaweed production, site selection, geospatial, hydrodynamics modeling

Commented [H1]: Suggestions for the title "Identification of Potential Locations for *Kappaphycus Alvarezii* Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia."

Commented [RV2]: Abstract must in 250 words

Commented [H3]: It is better for the background on the existing fundamental problems, "However, the availability of natural seaweed can be threatened along with the increasing demand so that seaweed cultivation is absolutely necessary (Pickering et al., 2007)."

1. Introduction

Indonesia is the largest producer of red algae in the world (83% of global seaweed production) followed by the Philippines and several countries (Valderrama *et al.*, 2015; Kim *et al.*, 2017). Seaweed production in Indonesia comes not only from mariculture, and brackish water ponds but also from natural collections. However, the availability of natural seaweed can be threatened in line with their increasing demand so that seaweed cultivation is completely necessary (Pickering *et al.*, 2007). The increasing demand for seaweed was mostly driven by new markets in China, Eastern Europe, Brazil, and others (Bixler and Porse, 2011). Furthermore, Arbit *et al.* (2019) stated that 99.73% of Indonesia's seaweed production comes from cultivation. Therefore, seaweed farming is the best choice for fulfilling market demand. Moreover, one of the major *K. alvarezii* farming areas within the Indonesian seas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province (Jalil *et al.*, 2020).

Site selection is essential in aquaculture development which depends on the types of commodities and applied cultivation methods (Kaiser *et al.*, 2010; Junaidi *et al.*, 2018). For seaweed, the success of cultivation is intrinsically linked to physical and chemical parameters of seaweed farming areas such as turbidity, temperature, salinity, pH, DO, nitrate and phosphate concentration (Rusliani 2016, Jalil *et al.* 2020). Hydrodynamic aspects should also be considered in the site selection of *K. alvarezii* farming since the cultivation method uses the longline method which is prone to the strong current and wave. The existence of Indonesian Throughflow along the Makassar Strait (Susanto *et al.* 2016) and the occurrence of strong upwelling at the southern part of the South Sulawesi each summer (Setiawan and Kawamura 2011) may influence the hydrodynamics condition in the Spermonde Island. Besides, the existence of small islands among the water area makes the hydrodynamics condition at the Spermonde Islands becomes more complex. Spermonde Islands area also has dense shipping activities. Many shipping lanes have lied among these areas (Figure 1) which are should be avoided in the mariculture site selection. This shows the complex considerations for conducting site selection of seaweed farming at Spermonde Islands.

In dealing with the complex considerations as mentioned above, the use of the geospatial model which is part of Geographical Information Systems (GIS) becomes an effective solution for determining the proper mariculture site selection (Longdill *et al.* 2008; Micael *et al.* 2015). This approach, in general, is an analysis of layers that contains spatial data to evaluate and select a particular area based on its suitability criteria classified according to certain environmental aspects (Malczewski, 2006; Malczewski & Rinner, 2015). GIS utilization in mariculture development has been widely used in various types of commodities including seaweed (e.g., Radiarta *et al.*, 2011; Radiarta *et al.*, 2012; Sousa *et al.*, 2012; Nirmala *et al.*, 2014; Teniwut *et al.*, 2019) which were mainly used in-situ data and remote sensing data for the input of geospatial modeling. The application of GIS for site selection for aquaculture in the Spermonde Islands has conducted by Arifin *et al.* (2014). However, their analysis was only based on in-situ measurement which cannot represent the temporal change of ocean parameters. In the present study, we perform cell-based geospatial modeling for the determination of sustainable site selection of seaweed cultivation, which integrates various physical and chemical parameters obtained not only from in-situ data but also the hydrodynamic model. The combination of in-situ data and the hydrodynamic model becomes the new approach proposed in the present study. Considering the importance of hydrodynamics aspects for the site selection of seaweed culture which is not available in the remote sensing data, we built a 2D hydrodynamic model to represent the hydrodynamic condition in the study area. On the other hand, Indonesia as the maritime continent experiences four-season influenced by monsoon wind (Helmi *et al.*, 2018) which impacts the hydrodynamics conditions. Thus, in-situ and incidental measurements of current and wave cannot represent the hydrodynamics condition in the study area. This issue only can be resolved by performing the hydrodynamics model. Furthermore, we also simulate the potential productivity of *K. alvarezii* at the selected site. This research is expected to be a prototype of a geospatial model to explore the potential aquaculture area in small islands and can be scientific-based information for local governments, researchers, NGOs, and related stakeholders for planning and developing *K. alvarezii* cultivation in the study area.

2. Material and Method

2.1. Study Area

The study area is presented in Figure 1 which is located at the Spermonde islands waters precisely in the Liukang Tupabbiring District, Pangkep Regency, South Sulawesi. The Liukang Tupabbiring District area is 54.44 km² (4.89% of Pangkep Regency) and consists of 20 islands. The population of this district reached 29,680 inhabitants, generally as fishermen and sea farmers. The distribution of field station is located at 4°5'3.98"S - 4°59'51.137"S and 119°18'35.532"E - 119°26'19.849"E. There are 94 sampling points distributed among the island of Bontosua, Sanane, Panjenekang, Pankaiya, Langkadea, Balanglombo, Balangcaddi Island, and reef flat area.

2.2. Water Quality Data Collection

We conducted a field survey for collecting water quality data at 94 points which the depths are more than 2 meters. Survey site determination was using Garmin 76C GPS. The collection of water quality data are presented in Table 1. The in-situ measurements were Sea Surface Temperature, (SST), salinity, pH, and substrate type. For the chemical parameters such as DO, nitrate, phosphate, and Total Suspended Solid (TSS), the water samples were analyzed at Chemical Oceanography Laboratory, Marine Science Department, Hasanuddin University, Indonesia (May 2019). The dynamic and non-dynamic water parameter data were interpolated to obtain a spatial distribution pattern at the time of data retrieval. The data were compiled in a digital form of 5 x 5 meters cell size using spline and classified according to suitability class in Table 2.

2.3. Two dimensions Hydrodynamic Model

Two dimensions (2D) hydrodynamic model was used to simulate current and wave condition at the study area. The data for generating a hydrodynamic model are as follows:

1. High-resolution satellite imagery GeoEye-1 from ESRI online (<https://esriindonesia.co.id/arcgis-online>)
2. Bathymetry data from Indonesia Geospatial Agency (<http://batnas.big.go.id/>)
3. Surface wind and wave height data obtained from 6 hourly ERA-interim data with a grid interval of 0.125° from July 2018 to July 2019 (Dee *et al.* 2011)
4. Tidal elevation forecast from July 2018 to July 2019 (<https://www.dhigroup.com/download/mike-by-dhi-tools/coastandseatools/global-tide-model>)

The current model was constructed by continuity equations and momentum equations with the average of depth. On-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry, wind and tidal dynamics were used as an input for generating current model.

The initial process was making a mesh, and setting the boundary condition of the model using the mathematical continuity equation as follows:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = S$$

The horizontal momentum equation for component x is as follows (DHI, 2010) :

$$\frac{\partial u}{\partial t} + \frac{\partial u^2}{\partial x} + \frac{\partial uv}{\partial y} + \frac{\partial wu}{\partial z} = f_v - g \frac{\partial h}{\partial x} - \frac{1}{r_0} \frac{\partial p_a}{\partial x} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial x} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{xx}}{\partial x} + \frac{\partial S_{xy}}{\partial y} \right) + F_u + \frac{\partial}{\partial z} \left(v_t \frac{\partial u}{\partial z} \right) + u_s S$$

While the horizontal momentum equation for component y is as follows (DHI, 2010):

$$\frac{\partial v}{\partial t} + \frac{\partial v^2}{\partial y} + \frac{\partial uv}{\partial x} + \frac{\partial wv}{\partial z} = -f_u - g \frac{\partial h}{\partial y} - \frac{1}{r_0} \frac{\partial p_a}{\partial y} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial y} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{yx}}{\partial x} + \frac{\partial S_{yy}}{\partial y} \right) + F_v + \frac{\partial}{\partial z} \left(v_t \frac{\partial v}{\partial z} \right) + v_s S$$

where x, y, z = cartesian coordinates of $x, y,$ and z direction; t = time; u = water particle speed of direction x ; v = water particle speed of direction y ; w = water particle speed of direction z ; η = water surface elevation; d = sea depth; h = total depth of the sea; f = coriolis parameter ($f = 2\omega \sin\phi$); Ω = ratio of earth revolution; ϕ = latitude coordinates; g = acceleration of gravity; $S_{xx}, S_{xy}, S_{yx}, S_{yy}$ = component of radiation stress tensor; ν_t = vertical eddy viscosity; p_a = atmospheric pressure; ρ_0 = sea density; S = debits due to sources; u_s = particle velocity of air sources direction x ; and v_s = the particle speed of air sources direction y .

Wave modeling used directional decoupled parametric and fully spectral formulations based on wave eternity movements with the input data of on-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry data, wind speed and direction and wave data. Parameterization was created in the frequency domain by introducing the 0 and 1st moments of the wave spectrum as a variable (Azhar *et al.*, 2011). The arranger equation was the equilibrium wave force in cartesian coordinate formulated as follow (DHI, 2010):

$$\frac{\partial N}{\partial t} + \frac{\partial}{\partial \phi} C\phi N + \frac{\partial}{\partial \lambda} C\lambda N + \frac{\partial}{\partial \sigma} C\sigma N + \frac{\partial}{\partial \theta} C\theta N = \frac{S}{\sigma}$$

Source energy, S , shows the superposition of surface functions in various styles (Azhar *et al.*, 2011).

$$S = S_{in} + S_{nl} + S_{ds} + S_{bot} + S_{surf}$$

where $N_{x,\sigma,\theta,t}$ = power density, ϕ = coordinate spherical, C = speed of propagation of four-dimensional wave group, S = energy source, S_{in} = formation by wind, S_{nl} = energy transfer, non-linear, S_{ds} = wave energy due to white capping, S_{bot} = by bottom friction, S_{surf} = dissipation due to the depth-induced breaking.

We took the maximum value of the current speed and wave height in each grid that comes from 4 seasonal patterns characteristics, that is the west monsoon (December), transition 1 (May), east monsoon (July), and transition 2 seasons (October). The results of the 2D hydrodynamic model form the patterns of current speed and wave height every season were interpolated with a spline method with a cell size of 5 x 5 meters and analyzed using cell statistics to obtain the maximum value of current speed and wave height. The results of the distribution map were classified according to suitability class in table 2.

2.4. Determination of Area Suitability Criteria

The determination of the weighting and scoring was done to provide value in each criterion for supporting the feasibility of seaweed cultivation. The weight determination of each criterion was based on the literature research and later on was used to weight the variables by priority. The variable that is very influential for the growth of seaweed got a high score. The determination was based on the defined suitability matrix as shown in Table 2.

The total score multiplication and weight in table 2 were used to determine the class of water suitability of seaweed cultivation with quantitative analysis using the following equation :

$$Y = \sum a_i \cdot X_n$$

The range of suitability cultivation area of *K. Alvarezii* was adjusted to use the following equation :

$$I = \frac{(\sum a_i \cdot X_n) - (\sum a_i \cdot X_n)_{min}}{k}$$

where, Y = The final value, a_i = weight per parameter, and X_n = score of suitability level, I = interval of suitability class, k = number of suitability class.

Finally, the suitability interval was divided into 5 classes, 1) highly suitable (HS) is in the range of the value 42-50, 2) suitable category (S) is in the range of 33-41, C) moderately suitable (MS) range of 24-32, 4) less suitable (LS) are in the range of 15-23, and 5) not suitable (N) had the value range of < 15. Each parameter in table 2 was transformed in digital format using a spline interpolation method with a cell size of 5 x 5 meters resulting in an index map of water suitability.

2.5. Distance Suitability Analysis

Distance suitability analysis was also carried out as the other limiting factors out of the water parameters. Those were the distance to shipping lanes or sea lanes and the distance to a settlement which are arranged in a digital format with 5 x 5 -meter cell-sized using the buffer and euclidean distance method to consider the accessibility and disruption of marine traffic. These criteria are described in Table 3. The suitability of distance to the settlement was compiled using the data of inhabited island, namely Bontosua, Sanane, Panjenekang, Balanglombo, and Balangcaddi Islands, while the shipping lanes data used was inter-regional shipping routes. The results of the analysis were integrated with a water suitability index map.

Commented [H4]: The distance made by the buffer should be conveyed based on the determination of the distance,

2.6. Spatial Modelling

The spatial modelling in this study used cell-based modelling by integrating all oceanographic and water quality parameters based on the suitability classification according to table 2 with the following modelling algorithms:

$$WS = (T_i \times 0.5) + (pH_i \times 0.5) + (S_i \times 1) + (DO_i \times 0.5) + (N_i \times 1.5) + (F_i \times 1.5) + (B_i \times 0.5) + (HS_i \times 1.5) + (CS_i \times 1.5) + (TSS_i \times 0.5) + (SB_i \times 0.5)$$

with, WS = Water suitability, T_i = temperature index map, pH_i = pH index map S_i = salinity index map, DO_i = Dissolved Oxygen index map, N_i = Nitrate index map, F_i = phosphate index map, B_i = bathymetry index map, HS_i = significant wave height index map, CS_i = current speed index map, TSS_i = TSS index map, SB_i = seabed index map

The result of this modeling is continued by determining the classification of water suitability based on a class of suitability table 2. Then, integrating the results of water suitability above with the result of distance suitability analysis according to table 3 using cell statistics analysis.

Commented [H5]: The source of the algorithm used should be presented,

2.7. Estimating Productivity

After obtaining the distribution of classes areas for all variable, the seaweed production can be estimated only by considering the area which is suitable or highly suitable. Longline method of seaweed cultivation unit was designed as 25 x 50 meter per unit, the distance between the rope was 1 meter and the planting distance of each tied seaweed was 25 cm (WWF, 2014). The initial weight of the seedlings was approximately 100 grams per bunch (Parakkasi *et al.*, 2020) with the wet weight crop was 500 grams per tie. The cultivation of seaweed with a floating rope/longline system produces an excellent product and can gain a weight up to four to five times from the initial weight of the seedlings (100 grams) in a period of cultivation time. The number of the allocated cultivation units was estimated to be 5 units/ha. The distance correction between units is 10 meters in width (twice longer than the average length of the fishing boat) for fishing traffic. This amount represents 62.5% of the maximum number of units i.e., 8 units/ha. To estimated production is the multiplication between the number of cultivation units (from the total suitable area) and once crop yields per unit with the following equations:

$$Ye = N \times Cy$$

with, Ye = estimated yield per cycle (tons), N = number of cultivation unit, Cy = crop yield per cycle (tons). Briefly, the research method is summarized in the flowchart presented in Figure 2.

3. Result and Discussion

3.1. Distribution of Water Quality Parameters

Spatial distributions of the suitability of water quality parameters are shown in Figure 3. It can be seen that the water quality conditions based on the variable of SST, and pH, are highly suitable for seaweed cultivation in the study area. SST is the important limiting factor for marine organisms (Wirasatriya *et al.*, 2019). The range of measured temperature in the study area is very narrow i.e., only around 30°C. According to BSN (2005) the optimal temperature for seaweed culture is 26-32°C, so this temperatures are feasible for this activity. whereas according to Andi *et al.*(2016), *K. alvarezii* grown well at the temperature range of 27°C–30°C. The high temperature of the water causes death on seaweed while low water temperatures cause damage to protein and fat membranes and it is very influential in the seaweed life cycle. Photosynthesis on seaweed generally lasts at a maximum temperature of 23°C-30°C (Msuya, 2011; Ding *et al.*, 2013) Seawater has an enormous support ability to prevent pH fluctuation. The condition of pH that is very acidic or alkaline will endanger the survival of organisms because it led to metabolic and respiration disorders. The pH range at the study site is 7.26-7.63. These values are feasible for seaweed cultivation (Rahman *et al.*, 2019). Based on Noor (2015), a suitable pH range for *K. alvarezii* cultivation is in normal and disposed alkaline level. n. Optimal seaweed growth requires a pH value range of 7.5-8.5 (PI, 2004; BSN, 2015; Radiarta *et al.*, 2018).

Commented [H6]: Is the maximum temperature limit (Msuya, 2011; Ding *et al.*, 2013),

Commented [H7]: Should be directly compared with the appropriate range of pH limits,

The suitability distribution for TSS and substrate type are moderately suitable or more in the study area. A substrate is dominated by reef type almost at the entire study area which is categorized as moderately suitable. The suitable condition is located around the small islands with the sand as the substrate type. Seaweed will grow optimally when the cultivation site is in coral, rubble, sand-based substrate, or a mixture of all three. The most optimum substrate type for seaweed cultivation is a hard substrate like the rocky coral then followed by sandy corals (Parakkasi *et al.*, 2020). Waters with a fractional substrate of coral to sand are usually passed by the current that was suitable for the growth of seaweed. However, cultivating seaweed with a longline method is not overly influenced by the substrate type. For TSS, the eastern part of the study area is highly suitable while the western part is moderately suitable. This indicates that the further distance from the mainland the lower TSS is observed. TSS is insoluble sedimentation which increases the turbidity of water (Nurdin *et al.*, 2019). High turbidity due to the suspended material impedes light penetration into the water column and affects seaweed growth. Thus, TSS indicates water pollution as *K. alvarezii* lives optimally in transparent oligotrophic waters (Hayashi *et al.*, 2011). The TSS levels in the study area range between 3.6-57.6 mg/L which means that the waters are suitable for the growth of *K. alvarezii*. Low TSS concentration is probably due to the absence of re-suspension due to the non-silt/clay of substrate type or no input from coastal sediment. Total dissolved solids (TDS) still meets the seawater quality standard for marine biota according to PI (2004), namely the standard quality value for coral and seagrass life is < 20 mg/L.

Commented [H8]: It is recommended that the results of the TSS measurement are presented in paragraphs, so that the level of conformity is conveyed next,

For DO and salinity, the suitability distributions vary from less suitable to highly suitable. Based on the DO criterion, the study area is mostly highly suitable for seaweed cultivation, with only 2 spots in the study area which are less suitable. The water circulation among small islands plays an important role in the high DO level. In open water, the low concentration of oxygen in a natural condition is rarely found. The optimal dissolved oxygen for seaweed cultivation is in the range of 3-7 mg/l (Rahman *et al.*, 2019). For salinity, the less suitable classes are spread in some areas and even the class at the eastern part of Sanane Island is not suitable for seaweed cultivation. The stability of the salinity becomes an important factor for seaweed because seaweed is a stenohaline organism that cannot resist the high fluctuation salinity (Reis *et al.*, 2010). The extreme salinity changes can lead to ice-ice disease. Salinity

Commented [H9]: It is advisable to provide a source,

can affect the osmoregulation process of marine biota including seaweed. However, most of the area is still feasible for seaweed cultivation.

Commented [H10]: It is advisable to provide a source,

Nitrate and phosphate concentrations as indicators of water fertility are relatively low in this region. This may be related to the absence of big rivers in the small islands of the study area as the nutrient sources. Nitrate is a limiting factor for seaweed growth. Nitrate is one of the nutrients needed for the growth of seaweed. The result shows that nitrate concentrations in the study area range between 0.001-0.12 mg/l which is relatively low. According to Kautsari and Ahdiansyah (2015), the feasible nitrate concentration for seaweed cultivation ranges from 0.01 mg/l to 0.07 mg/l, while the concentration of <0.01 mg/l and > 0.1 mg/l are not suitable for seaweed cultivation. Phosphate can be a limiting factor due to the source limitation. In the study area, phosphate concentration ranges from 0.001 mg/l to 0.062 mg/l which indicates low to insufficient fertility levels. Phosphate concentration for the fertile water should range from 0.051 mg/l to 0.1 mg/l. Besides, phosphate concentration also controls the carrageenan content of seaweed (Patty *et al.*, 2015).

The depth of the study area ranges from 2 m to 39.2 m. The longline method can be applied at the 1 m depth from the lowest tide or below (WWF, 2014). The sufficient depth provides seaweed to be submerged. Nevertheless, noted that the deeper waters will increase the cost of constructing seaweed cultivation. Thus, the areas with a depth of more than 15 m are categorized as not suitable for seaweed cultivation.

The distribution of physical and chemical parameters at the seaweed site is highly determined by the wave and current. These aspects play a pivotal role in the water movement for algae growth (Gaylord *et al.*, 2007; Kregting *et al.*, 2013). The variability of current speed and wave height is distinguished by the 4 different characteristics of the west monsoon; transition 1; east monsoon; and transition 2 based on the results of 2D hydrodynamic modeling (Figure 4 and Figure 5). During west monsoon and transition 1 (east monsoon and transition 2), strong current speed appears at the northwestern (southern) part of the study area. The strongest current speed occurs during the transition 1 season. The pattern of the surface current in the study area is influenced by the current pattern of breaking waves in the small islands waters and the surrounding shallow area; and the influence of wind and tidal conditions in these waters. Based on the criteria shown in Table 2, the study area is categorized as highly suitable (dark blue color) to less suitable (light blue color).

For the wave height, almost entire study areas are categorized as suitable and highly suitable for seaweed cultivation except during east monsoon. During west monsoon transition 1 and transition 2, the blue and dark blue colors mean that the wave height ranges from 0.1 m to 0.3 m. During east monsoon, the wave higher than 0.3 m occurs at almost the entire study area which makes it less suitable for seaweed cultivation.

The protected area is determined by several types of physical conditions such as current velocity, and wave height. The influence of strong current and ocean waves will drag and damage the cultivated seaweed and its system. The maximum value of the distribution of current speeds and wave height for 4 seasons is described in Figure 6. Based on the results in Figure 6, the current speed and wave height around the study of the water are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. The less suitable areas for the current speed (wave height) category only appear mostly in the northeastern (northern) part of the study area. There is no area with a current speed of more than 0.5 m/s and the wave height more than 0.5 m which makes the area is not suitable for seaweed cultivation. The absence of strong current and wave may correspond to the morphology of the study area which is categorized as coral/reef waters with a lot of shallow/reef flat area located within the small islands that protect the area from the high wave and strong current. As mentioned by Sulma *et al.* (2008), the types of protected waters are bays, caps, straits, lagoons, and reef flat. Furthermore, Helmi *et al.* (2017) also stated that the coral reef is a shoreline protector.

3.2. Potential Areas for Seaweed culture Development and Production

Various suitability criteria as limiting factor of site selection for seaweed cultivation include physical and chemical parameters and distance suitability for accessibility and restricted area inside the shipping lanes has been determined into 5 highly suitable classes (HS), suitable (S), moderately suitable (MS), less suitable (LS) and not suitable (N) (Figure 7). Figure 7 shows that most of the study areas are moderately suitable and suitable for seaweed cultivation. The area which is not suitable is only located along the shipping lane. Figure 8 summarizes the areal size of each criteria as follows: the highly suitable (HS) is 83.2 ha (0.42%); suitable (S) is 4,463.08 ha (22.21%); moderately suitable (MS) is 9,983.52 ha (49.68%); less suitable (LS) is an area of 494.84 ha (2.46%); and not suitable (N) is 5,071.15 ha (25.23%). The potential area for seaweed cultivation development reaches 4,546.28 ha as the sum area of highly suitable and suitable criteria. This result is more than obtained by Arifin *et al.* (2014) which only found 1,963.6 ha area which is suitable and highly suitable for *K. alvarezii* cultivation in the same study area. This indicates that our method can optimally classify the potential area for *K. alvarezii* cultivation. Furthermore, this potential area can be utilized for 22,730 cultivation units with seaweed production reaching about 56,825 tons of wet weight per crop, or within a year (2 times of harvesting) can reach about 113,650 tons. This amount is only about one-third of the total seaweed production in Pangkep Regency in 2018. However, it is important to be noted that, there are some limitations in the present study which should be considered. For example, the present study does not include the long-term climate variability which may also influence the physical and chemical parameters in the study area. As reported by Kunarso *et al.* (2019) and Msuya *et al.* (2014). The climate variability influences the oceanographic condition which then affects fisheries activity such as aquaculture and fish catch. The present study also does not consider the risk analysis which can disturb the cultivation process as shown by Castelar *et al.*, (2015). Furthermore, to maintain the carrying capacity of the study area, the area that might be utilized for the expansion of *K. alvarezii* cultivation is about 50% (covering 2,273.14 ha) of the total suitable area accordingly. The result of the present study indicates the effectiveness of geospatial modeling for developing the area of *K. alvarezii* culture which can be applied in the other area.

4. Conclusions

We have conducted a comprehensive geospatial modeling approach for the site selection of *K. alvarezii* cultivation at the Spermonde Islands. Physical and chemical parameters as well as the accessibility and restricted area of shipping lanes were used to define the suitability criteria for *K. alvarezii* cultivation. The result is SST and pH are categorized as highly suitable for *K. alvarezii* cultivation in the whole study area. A substrate is dominated by reef type almost at the whole study area which is categorized as moderately suitable. TSS levels in the study area range between 3.6-57.6 mg/l which means that the waters are suitable for the growth of *K. alvarezii*. For DO and salinity, the suitability distributions vary from less suitable to highly suitable. Nitrate and phosphate concentrations mainly are too low which are not suitable for *K. alvarezii* cultivation. The depth at the study area ranges from 2 m to 39.2 m. The area with a depth more than 15 m is categorized as not suitable for *K. alvarezii* cultivation. The current speed and wave height around the study of the water are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. After summarizing all criteria according to their weighting score, the suitable for seaweed cultivation at the study area is 4,546.28 ha, which is the combination of the suitable (S) area of 4,463.08 ha and highly suitable criteria (HS) with an area of 83.2 ha. The potential production per crop can reach 56,825 tons per cycle and 113,650 tons per year.

Acknowledgment

The author thanked the Ministry of Research Technology and Higher Education of the Republic of Indonesia for the financial support of research. The author thanked the Center for Coastal Rehabilitation and Disaster Mitigation Studies (CoREM), Diponegoro University for the facilities' support on hydro-oceanographic modeling and geospatial data processing. We also thanked the Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University for the surveys and laboratory analysis of water quality data.

References

- Afandi, A. Musadat, F. 2018. Suitability analysis for seaweed culture area at the waters of Kamelanta Village and Panjang Island using Geographical Information System. *Jurnal Akuakultura* 2(1), 69-78. (In Bahasa)
- Arbit, N., Omar, S., Soekendarsi, E., Yasir, I., Tresnati, J., Tuwo, A. 2019. Morphological and genetic analysis of *Gracilaria* sp. cultured in ponds and coastal waters. *IOP Conf. Ser.: Earth Env. Sci.* 370 012018.
- Arifin T., Bohari, R., Arlyza, I.S. 2014. Analysis of Space Suitability Based on Marine Agriculture in The Small Islands Around Makassar. *Forum geography* 28 (1), 91-102. DOI: <https://doi.org/10.23917/forgeo.v28i1.441>
- Ariyanto, F. Y., Astor, Y., Sidqi, M. 2018. Application of water boundary theory for determining site licenses at the coastal area and small islands in Indonesia. *Proceeding National Seminar of Geomatics 2018*. Bandung, pp. 893-900. (In Bahasa)
- Azhar, R. M., Wurjanto, A., Yuanita, N. 2011. Study of coastal protection of submerged breakwater at Tanjung Kait beach. Master Thesis. Master Program of Water Resource Management, Bandung Institute of Technology, Indonesia.
- Bixler, H. J., Porse, H. 2011. A Decade of Change in the Seaweed Hydrocolloids Industry. *J. Appl. Phycol.* 23, 321-335.
- Castelar, B., de Siqueira, M. F., Sánchez-Tapia, A., Reis, R. P. 2015. Risk Analysis Using Species Distribution Modelling to Support Public Policies for The Alien Alga *Kappaphycus alvarezii* Aquaculture in Brazil. *Aquacult.* 446, 217-226.
- Dee, D. P. *et al.* 2011. The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Q.J.R. Meteorol. Soc.* 137(656), 553–597. doi: 10.1002/qj.828
- DHI. 2010. MIKE-21 and MIKE-3 Flow Model. Hydrodynamic Module Step-by-Step Training Guide. DHI MIKE, Denmark. 96 p.
- Ding L., Ma, Y., Huang, B., Chen, S. 2013. Effects of Seawater Salinity and Temperature on Growth and Pigment Contents in *Hypnea cervicornis*. *J. Agardh (Gigartinales, Rhodophyta)*. *BioMed Res. Int.* 2013, 594308. <https://doi.org/10.1155/2013/594308>
- Gaylor, B., Rosman, J. H., Reed, D. C., Koseff, J. R., Fram, J., Macintyre, S., Arkema, K., McDonald, C., Brzezinski, M. A., Largier, J. L., Monismith, S. G., Raimondi, P. T., Mardian, B. 2007. Spatial Patterns of Flow and Their Modification within and Around A Giant Kelp Forest. *Limnol. Oceanogr.* 52, 1838–1852.
- Hayashi, L., Santos, A. A., Faria, G. S. M., Nunes, B. G., Souza, M. S., Fonseca, A. L. D., Barreto, P. L. M., Oliveira, E. C., Bouzon, Z. L. 2011. *Kappaphycus alvarezii* (Rhodophyta, Areschougaceae) Cultivated in Subtropical Waters in Southern Brazil. *J. Appl. Phycol.* 23, 337–343.
- Helmi, M., Satriadi, A., Suryoputro, A. A. D., Marwoto, J., Setiyono, H. Hariyadi. 2018. Rehabilitation Priority Area Assessment on Death Coral using Cell-Based Modelling Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.* 9(11), 2949-2961.
- Helmi, M., Purwanto, Atmodjo, W., Subardjo, P., Aysira, A. 2017. Benthic Diversity Mapping and Analysis Base on Remote Sensing and Seascape Ecology Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.* 9(10), 227-235.
- Jalil, A. R., Samawi, M. F., Azis, H. Y., Jaya, I., Malik, A., Yunus, I., Sohopi, M. A. A. 2020. Comparison of physical-chemical conditions for seaweed cultivation in the Spermonde Archipelago, Indonesia. *AACL Bioflux* 13(4), 2071-2082.

Commented [RV11]: Fill complete name of author

- Junaidi, M., Lestary, D. P., Scabra, A. R. 2018. Mariculture Site Selection Based on Environmental Parameters in Tanjung and Gangga Sub-District, North Lombok. In: Proceeding of 1st ICWBB 2018. Mataram University, Mataram, pp. 233-251.
- Kaiser, M. J., Yu, Y., Snyder, B. 2010. Economic Feasibility of Using Offshore Oil and Gas Structures in the Gulf of Mexico for Platform-based Aquaculture. *Mar. Pol.* 34, 699-707.
- Kautsari, N., Ahdiansyah, Y. 2015. Carrying capacity and suitability assessment of Labuhan Terata waters, Sumbawa for the development of seaweed culture. *Jurnal Ilmu Kelautan* 20(4), 233-238. (In Bahasa)
- Kim, J. K., Yarish, C., Hwang, E. K., Park, M., Kim, Y. 2017. Seaweed Aquaculture: Cultivation Technologies, Challenges, and Its Ecosystem Services. *Algae*. 32(1), 1-13.
- Kregting, L.T., Blight, A. B., Elsässer, B., Savidge, G. 2013. The Influence of Water Motion on The Growth Rate of The Kelp *Laminaria Hyperborea*. *J. Exp. Mar. Biol. Ecol.*, 448, 337–345.
- Kunarso, Wirasatriya, A., Irwani, Satriadi, A., Helmi, M., Prayogi, H. Munandar, B. 2019. Impact of Climate Variability on Aquatic Productivity and Fisheries Resources in Jepara. IOP Conf. Ser.: Earth Env. Sci., 246 012021: 1-12.
- Longdill, P. C., Healy, T. R., Black, K. P., 2008. An integrated GIS approach for sustainable aquaculture management area site selection. *Ocean Coast. Manag.* 51, 612-624. doi:10.1016/j.ocecoaman.2008.06.010
- Malczewski, J. 2006. Ordered weighted averaging with fuzzy quantifiers: GIS-based multicriteria evaluation for land-use suitability analysis. *Int. J. Appl. Earth Observ. Geoinf.* 8(4), 270- 277.
- Malczewski, J., Rinner, C. 2015. *Multicriteria Decision Analysis in Geographic Information Science*. Springer, Berlin
- Micael, J., Costa, A.C., Aguiar, P., Medeiros, A., Calado, H. 2015. Geographic Information System in A Multi-criteria tool for Mariculture Site Selection. *Coast. Manag.* 43(1), 52-66.
- Msuya, F. E., Buriyo, A., Omar, I., Pascal, B., Narrain, K., Ravina, J. J. M., Mrabu, E., Wakibia, J. G. 2014. Cultivation and Utilisation of Red Seaweeds in the Western Indian Ocean (WIO) Region. *J. Appl. Phycol.* 26, 699–705.
- Msuya, F. E. 2011. Environmental Changes and Their Impact on Seaweed Farming in Tanzania. *World Aquacult.* 42, 34-37.
- Nirmala, K., Ratnasari, A., Budiman, S. 2014. Site Selection for Seaweed Culture at Gerupuk Bay-West Nusa Tenggara Using Remote Sensing and GIS. *Jurnal Akuakultur Indonesia* 13(1), 73-82.
- Noor, N. M. 2015 The Water Bodies Compatibility Analysis for Culturing Brown Seaweed *Kappapycus alvarezii* in Ketapang Seashore, South Lampung. *Maspuri Journal* 7(2), 91-100. (in Bahasa)
- Nurdin, N., Helmi, M., Setiawan, R. Y., Zainudin, M., Rustam, A., Akbar, M. 2019. Landsat Capability to Mapping of Total Solid Suspended (TSS) and Coastal Ecosystem. *ISPRS – Int. Arch. Photogram., Remote Sens. Spatial Inf. Sci.*, 42(4), 497-502.
- Parakkasi, P., Rani, C., Syam, R., Zainuddin, Achmad, M. 2020. Growth Response and Quality of Seaweed *Kappaphycus alvarezii* Cultivated in Various Coastal Ecosystems in the Waters of West Sulawesi, Indonesia. *AAFL Bioflux* 13(2), 627-639.
- Patty, S. I., Arfah, H., Abdul, M. S. 2015. The relationship among Nutrient (Phosphate and Nitrate), DO, and pH with the water fertility in Jikumerasa waters, Buru Island. *Jurnal Pesisir dan Laut Tropis* 1(1), 43-50. (In Bahasa)

- Pickering, T. D., Skelton, P. Sulu, J. R. 2007. Intentional Introductions of Commercially Harvested Alien Seaweeds. *Bot. Mar.* 50, 338–350.
- Radiarta, I. N., Saputra, A. Albasri, H. 2012. Feasibility mapping of seaweed culture area (*Kappaphycus alvarezii*) at Bintan Regency, Riau Islands Province using GIS and remote sensing approach. *Akuakultur* 7(1): 145-157. (In Bahasa)
- Radiarta, I. N., Saitoh, S. Yasui, H. 2011. Aquaculture Site Selection for Japanese Kelp (*Laminaria japonica*) in Southern Hokkaido, Japan, Using Satellite Remote Sensing and GIS-based Models. *ICES J. Mar. Sci.* 68(4), 773-780. doi:10.1093/icesjms/fsq163
- Rahman, I. S., Fadjar, M., Tjahjono, A. 2019. The Relationship Water Physical-Chemical Parameters in Seaweed Cultivation *Eucheuma cottonii* with Long-line System. *Int. J. Sci. Tech. Res.* 8(11), 1448-1452.
- Rahmayanti, F., Diana, F., Kusumawati, I. 2018. Suitability analysis for area development of seaweed (*Eucheuma cottonii*) culture in Lhok Bubon waters, Samatiga District, Aceh Barat Regency. *Jurnal Akuakultura* 2(1), 26-32. (In Bahasa).
- Reis, R. P., Loureiro, R. R., Mesquita, F. S. 2010. Do Salinity Affect Growth and Carrageenan Yield of *Kappaphycus alvarezii* (Gigartinales/ Rhodophyta)? *Aquacult. Res.* 42, 1231-1234.
- Rusliani. 2016. Study of water quality on the growth of seaweed *Gracilaria verrucosa* in a pond with verticulture method. *Octopus J. Fish. Sci.* 5(2), 522-527. (in Bahasa)
- Setiawan, R. Y., Kawamura, H. 2011. Summertime Phytoplankton Bloom in the south Sulawesi Sea. *IEEE J. Sel. Topics Appl. Earth Observ. Remote Sens.* 4(1), 241-244. doi:10.1109/JSTARS.2010.2094604.
- Sousa, F. E. S., Moura, E. A., Marinho-Soriano, E. 2012. Use of Geographic Information Systems (GIS) to Identify Adequate Sites for Cultivation of The Seaweed *Gracilaria birdie* in Rio Grande Do Norte, Northeastern Brazil. *Revista Brasileira de Farmacognosia* 22(4), 868–873.
- Sulma, S., Manoppo, A. K. S., Hartuti, M. 2008. Identification of Suitable Area for Seaweed Culture in Bali Waters Based on Remote Sensing Satellite Data. *Int. J. Remote Sens. Earth Sci.* 5, 57-66.
- Susanto, R. D., Wei, Z., Adi, T. R., Zheng, Q., Fang, G., Bin, F., Supangat, A., Agustyadi, T., Li, S., Trenggono, M., Setiawan, A. 2016. Oceanography Surrounding Krakatau Volcano in the Sunda Strait, Indonesia. *Oceanogr.* 29(2), 228-237. <http://dx.doi.org/10.5670/oceanog.2016.31>.
- Teniwut, W. A., Marimin, Djatna, T. 2019. GIS-Based multi-criteria decision-making model for site selection of seaweed farming information center: A lesson from small islands, Indonesia. *Decis. Sci. Let.* 8(2019), 137-150 doi: 10.5267/j.dsl.2018.8.001
- Utama, R. Z., Handayani, T. 2018. Suitability Between Oceanography and Seaweed (*Eucheuma cottonii*) Cultivation Potential in Tidung Island with Geographic Information System (GIS). *ICENIS: E3S Web of Conference* 73, 12009. 1-5.
- Valderrama, D., Cai, J., Hishamunda, N., Ridler, N., Neish, I. C., Hurtado, A. Q., Msuya, F. E., Krishnan, M., Narayanakumar, R., Kronen, M., Robledo, D., Gasca-Leyva, E., Fraga, J. 2015. The Economics of *Kappaphycus* Seaweed Cultivation in Developing Countries: A Comparative Analysis of Farming Systems. *Aqua. Econ. Manag.* 19, 251–277.
- Wirasatriya, A., Sugianto, D. N., Helmi, M., Setiawan, R. Y. Koch, M. 2019. Distinct Characteristics of SST Variabilities in the Sulawesi Sea and the Northern Part of the Maluku Sea during the Southeast Monsoon. *IEEE J. Sel. Topics Appl. Earth Obs. Remote Sens.* 12(6), 1763-1770.

- WWF-Indonesia. 2014. Better Management Practices: Guideline series of small scale fisheries of seaweed culture - Cottoni (*Kappaphycus alvarezii*), Sacol (*Kappaphycus striatum*) dan Spinosum (*Eucheuma Denticulatum*) Version 1. WWF-Indonesia. South Jakarta. 53 p. (in Bahasa)
- Yulianto, H., Damai, A. A., Delis, P. C., Elisdiana, Y. 2017. Spatial Analysis to Evaluate the Suitability of Seaweed Farming Site in Lampung Bay, Indonesia. *Turkish J. Fish. Aquatic Sci.* 17, 1253-1261.

List of Tables

Table 1. Water quality data collection

No	Variables	Unit	Tools/Methods	Data Retrieval Methods
1	Sea Surface Temperature (SST)	°C	Thermometer	In situ
2	Salinity	Psu	Salinometer	In situ
3	pH	-	<i>Water quality checker</i>	In situ
4	Dissolved oxygen (DO)	ppm	Winkler method	Laboratory analysis
5	Nitrate	mg/l	Ascorbic Acid Method	Laboratory analysis
6	Phosphate	mg/l	Brucine Sulfate Method	Laboratory analysis
7	Total Suspended Solids (TSS)	mg/l	<i>Milipore filter</i>	Laboratory analysis
8	Substrate type	-	Visually observation	In situ
9	Depth	m	Modelled depth from LPI Maps (BIG, 2011) and BATNAS (BIG, 2018) data	Integrating modelled data

Table 2. Table of suitability criteria for seaweed *Kappaphycus alvarezii* cultivation


Variables	Level of conformity (score)					Weight	Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)		
SST (°C)	28-30	24-27 or 31-32	22-23 or >32	21 or >34	<21 or >36	0.5	Modification of Noor (2015); Rahmayanti <i>et al.</i> (2018)
pH	7-8	6.5-7 or 8-8.5	6-6.4 or >8.5	4-6	<4 or >9	0.5	Modification of Kautsari and Ahdiansyah (2015); Noor (2015)
Salinity (psu)	28-31	26-28 or 31-33	<26 or ≥ 34	18-25 or 35-36	<18 or >37	1	Modification of Kautsari and Ahdiansyah (2015); Rahmayanti <i>et al.</i> (2018); Utama and Handayani (2018)
DO (mg/l)	>6	5-6	4-5	2-4	<2	0.5	Modification of Afandi and Musadat (2018)
Nitrate (mg/l)	0.04-0.07	0.02-0.03	0.01 or >0.1	<0.01 or >0.5	>1	1.5	Modification of Kautsari and Ahdiansyah (2015); Afandi and Musadat (2018)
Phosphate (mg/l)	0.051-0.1	0.02-0.04	0.01-<0.02 or >0.2-1	<0.01 or >1	>2	1.5	Modification of Patty <i>et al.</i> (2015)
Depth (m)	3-10	2-3 or 11-13	1-2	>13-15	<1 or >15	0.5	Modification of Tiensongrusme (1990) in Radiarta <i>et al.</i> (2012); Kautsari and Ahdiansyah (2015); Yulianto <i>et al.</i> (2017)
Wave height (m)	0.2-0.3	0.1-0.19	>0.3	<0.1 or >0.4	>0.5	1.5	Modification of Rahmayanti <i>et al.</i> (2018)
Current speed (m/s)	0.25-0.3	0.2-0.24 or 0.31-0.4	0.1-0.19 or >0.4	<0.1 or >0.45	>0.5	1.5	Modification of Utama and Handayani (2018)
TSS (mg/l)	<25	25-50	50-100	100-400	>400	0.5	Modification of Yulianto <i>et al.</i> (2017)
Seabed	Coral, Rubbel	Sand	Reefs	Muddy Sand	Mud	0.5	Modification of Noor (2015); Yulianto <i>et al.</i> (2017)

Note : HS: Highly suitable, S: suitable, MS: moderately suitable, LS: less suitable, and N: not suitable

Table 3. Distance suitability analysis

Parameters	Suitability Level (Score)					Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)	
Shipping lanes (m)	> 500	-	-	-	≤ 500	Ariyanto <i>et al.</i> (2018)
Distance to settlements (km)	4	4-5	5-6	6-7	>7	Modification of Radiarta <i>et al.</i> (2012)

List of Figures






Figure 2. Research Flowchart

Commented [RV12]: Text can not read




Figure 3. Distribution of physical and chemical parameters

Commented [H13]: The title of the image is still not informative, it should be given letter numbering to make it easier for users,




Figure 4. Distribution of current speed (m/s) based on 2D hydrodynamic model in each season




Figure 5. Distribution of significant wave height (m) based on 2D hydrodynamic model in each season




Figure 6. Maximum statistical value of current speed (left) and wave height (right) for all season




Figure 7. Suitability distribution for cultivation area of *K. Alvarezii*




Figure 8. Potential area for *K. alvarezii* cultivation in each



Sarjito Sarjito <sarjito@live.undip.ac.id>
To: Ilmu Kelautan <ijms.undip@gmail.com>

Tue, Aug 16, 2022 at 7:17 AM

Dear editor in chief

We have revised our manuscript, 47735RV, with an entitle : **Identification of Potential Locations for Kappaphycus Alvarezii Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia**

We hope our manuscrip could processed further

Thank you for your kindly


I wait a good news from you

Your sincerely

Sarjito

[Quoted text hidden]

2 attachments

 **47735RV 15082022 revised 1.docx**
2271K

 **Rebuttal IJMS for reviwer A and B.docx**
22K

Identification of Potential Locations for Kappaphycus Alvarezii Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia

Sarjito Sarjito^{1*}, Hanny Ammaria¹, Muhammad Helmi^{2,3}, Slamet Budi Prayitno¹, Nurjannah Nurdin^{4,5}, Riza Yuliratno Setiawan⁶, Parichat Wetchayont⁷, Anindya Wirasatriya^{2,3}

¹Aquaculture Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro, Indonesia

Jl. Prof. Soedarto, SH., Tembalang, Semarang, Central Java (50275) Indonesia

²Oceanography Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro, Indonesia

³Center for Coastal Rehabilitation and Disaster Mitigation Studies (CoREM), Indonesia

⁴Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University, Indonesia

⁵Marine Science and Fisheries Faculty, Hasanuddin University, Indonesia

⁶Marine Science and Fisheries Faculty, Gadjah Mada University, Indonesia

⁷Department of Geography, Srinakharinwirot University, Thailand

*e-mail: sarjito@live.undip.ac.id

Commented [H1]: Suggestions for the title "Identification of Potential Locations for Kappaphycus Alvarezii Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia."

Commented [W2R1]: Done

Abstract

Indonesia is the second largest producer of red algae in the world with one of the major *K. alvarezii* farming areas within the Indonesian seas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province. *K. alvarezii* production in Pangkep Regency increased year by year. However, the availability of natural seaweed can be threatened along with the increasing demand so that seaweed cultivation is absolutely necessary. Proper site selection is crucial to guarantee the success of the seaweed farming development. We conducted a Geographic Information system (GIS) with geospatial modeling approach for the site selection of *K. alvarezii* cultivation at the Liukang Tuppabiring District, Spermonde Islands by combining various physical and chemical parameters obtained from in-situ data and hydrodynamics modeling. The parameters are Sea Surface Temperature, (SST), salinity pH, substrate type, current speed, wave height, DO, nitrate, phosphate, and Total Suspended Solid (TSS). We also include the accessibility and the restricted area of shipping lanes for determining the area suitability of *K. alvarezii* cultivation. This research demonstrates the cell-based geospatial modeling can be effectively used and found the suitable categorized area for *K. alvarezii* culture of 4,546.28 ha. The Potential productivity of wet-weight seaweed can reach 56,825 tons per cycle or 113,650 tons per year. This result indicates the GIS with geospatial modeling approach for developing the area of *K. alvarezii* culture which can be applied in the other area.

Commented [RV3]: Abstract must in 250 words

Commented [W4R3]: Abstract maximum in 250 words

Commented [H5]: It is better for the background on the existing fundamental problems, "However, the availability of natural seaweed can be threatened along with the increasing demand so that seaweed cultivation is absolutely necessary (Pickering et al., 2007)."

Commented [W6R5]: done

Keywords: Seaweed production, site selection, geospatial, hydrodynamics modeling

1. Introduction

Indonesia is the largest producer of red algae in the world (83% of global seaweed production) followed by the Philippines and several countries (Valderrama *et al.*, 2015; Kim *et al.*, 2017). Seaweed production in Indonesia comes not only from mariculture, and brackish water ponds but also from natural collections. However, the availability of natural seaweed can be threatened in line with their increasing demand so that seaweed cultivation is completely necessary (Pickering *et al.*, 2007). The increasing demand for seaweed was mostly driven by new markets in China, Eastern Europe, Brazil, and others (Bixler and Porse, 2011). Furthermore, Arbit *et al.* (2019) stated that 99.73% of Indonesia's seaweed production comes from cultivation. Therefore, seaweed farming is the best choice for fulfilling market demand. Moreover, one of the major *K. alvarezii* farming areas within the Indonesian seas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province (Jalil *et al.*, 2020).

Site selection is essential in aquaculture development which depends on the types of commodities and applied cultivation methods (Kaiser *et al.*, 2010; Junaidi *et al.*, 2018). For seaweed, the success of cultivation is intrinsically linked to physical and chemical parameters of seaweed farming areas such as turbidity, temperature, salinity, pH, DO, nitrate and phosphate concentration (Rusliani 2016, Jalil *et al.* 2020). Hydrodynamic aspects should also be considered in the site selection of *K. alvarezii* farming since the cultivation method uses the longline method which is prone to the strong current and wave. The existence of Indonesian Throughflow along the Makassar Strait (Susanto *et al.* 2016) and the occurrence of strong upwelling at the southern part of the South Sulawesi each summer (Setiawan and Kawamura 2011) may influence the hydrodynamics condition in the Spermonde Island. Besides, the existence of small islands among the water area makes the hydrodynamics condition at the Spermonde Islands becomes more complex. Spermonde Islands area also has dense shipping activities. Many shipping lanes have lied among these areas (Figure 1) which are should be avoided in the mariculture site selection. This shows the complex considerations for conducting site selection of seaweed farming at Spermonde Islands.

In dealing with the complex considerations as mentioned above, the use of the geospatial model which is part of Geographical Information Systems (GIS) becomes an effective solution for determining the proper mariculture site selection (Longdill *et al.* 2008; Micael *et al.* 2015). This approach, in general, is an analysis of layers that contains spatial data to evaluate and select a particular area based on its suitability criteria classified according to certain environmental aspects (Malczewski, 2006; Malczewski & Rinner, 2015). GIS utilization in mariculture development has been widely used in various types of commodities including seaweed (e.g., Radiarta *et al.*, 2011; Radiarta *et al.*, 2012; Sousa *et al.*, 2012; Nirmala *et al.*, 2014; Teniwut *et al.*, 2019) which were mainly used in-situ data and remote sensing data for the input of geospatial modeling. The application of GIS for site selection for aquaculture in the Spermonde Islands has conducted by Arifin *et al.* (2014). However, their analysis was only based on in-situ measurement which cannot represent the temporal change of ocean parameters. In the present study, we perform cell-based geospatial modeling for the determination of sustainable site selection of seaweed cultivation, which integrates various physical and chemical parameters obtained not only from in-situ data but also the hydrodynamic model. The combination of in-situ data and the hydrodynamic model becomes the new approach proposed in the present study. Considering the importance of hydrodynamics aspects for the site selection of seaweed culture which is not available in the remote sensing data, we built a 2D hydrodynamic model to represent the hydrodynamic condition in the study area. On the other hand, Indonesia as the maritime continent experiences four-season influenced by monsoon wind (Helmi *et al.*, 2018) which impacts the hydrodynamics conditions. Thus, in-situ and incidental measurements of current and wave cannot represent the hydrodynamics condition in the study area. This issue only can be resolved by performing the hydrodynamics model. Furthermore, we also simulate the potential productivity of *K. alvarezii* at the selected site. This research is expected to be a prototype of a geospatial model to explore the potential aquaculture area in small islands and can be scientific-based information for local governments, researchers, NGOs, and related stakeholders for planning and developing *K. alvarezii* cultivation in the study area.

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: Fig.

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

2. Material and Method

2.1. Study Area

The study area is presented in [Figure 1](#), which is located at the Spermonde islands waters precisely in the Liukang Tupabbiring District, Pangkep Regency, South Sulawesi. The Liukang Tupabbiring District area is 54.44 km² (4.89% of Pangkep Regency) and consists of 20 islands. The population of this district reached 29,680 inhabitants, generally as fishermen and sea farmers. The distribution of field station is located at 4°5'3.98"S - 4°59'51.137"S and 119°18'35.532"E - 119°26'19.849"E. There are 94 sampling points distributed among the island of Bontosua, Sanane, Panjenekang, Pankaiya, Langkadea, Balanglombo, Balangcaddi Island, and reef flat area.

Deleted: Fig.

2.2. Water Quality Data Collection

We conducted a field survey for collecting water quality data at 94 points which the depths are more than 2 meters. Survey site determination was using Garmin 76C GPS. The collection of water quality data are presented in Table 1. The in-situ measurements were Sea Surface Temperature, (SST), salinity, pH, and substrate type. For the chemical parameters such as DO, nitrate, phosphate, and Total Suspended Solid (TSS), the water samples were analyzed at Chemical Oceanography Laboratory, Marine Science Department, Hasanuddin University, Indonesia (May 2019). The dynamic and non-dynamic water parameter data were interpolated to obtain a spatial distribution pattern at the time of data retrieval. The data were compiled in a digital form of 5 x 5 meters cell size using spline and classified according to suitability class in Table 2.

2.3. Two dimensions Hydrodynamic Model

Two dimensions (2D) hydrodynamic model was used to simulate current and wave condition at the study area. The data for generating a hydrodynamic model are as follows:

1. High-resolution satellite imagery GeoEye-1 from ESRI online (<https://esriindonesia.co.id/arcgis-online>)
2. Bathymetry data from Indonesia Geospatial Agency (<http://batnas.big.go.id/>)
3. Surface wind and wave height data obtained from 6 hourly ERA-interim data with a grid interval of 0.125° from July 2018 to July 2019 (Dee *et al.* 2011)
4. Tidal elevation forecast from July 2018 to July 2019 (<https://www.dhigroup.com/download/mike-by-dhi-tools/coastandseatools/global-tide-model>)

Deleted: et al

The current model was constructed by continuity equations and momentum equations with the average of depth. On-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry, wind and tidal dynamics were used as an input for generating current model.

The initial process was making a mesh, and setting the boundary condition of the model using the mathematical continuity equation as follows:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = S$$

The horizontal momentum equation for component x is as follows (DHI, 2010) :

$$\frac{\partial u}{\partial t} + \frac{\partial u^2}{\partial x} + \frac{\partial uv}{\partial y} + \frac{\partial wu}{\partial z} = f_v - g \frac{\partial h}{\partial x} - \frac{1}{r_0} \frac{\partial p_a}{\partial x} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial x} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{xx}}{\partial x} + \frac{\partial S_{xy}}{\partial y} \right) + F_u + \frac{\partial}{\partial z} \left(v_t \frac{\partial u}{\partial z} \right) + u_s S$$

While the horizontal momentum equation for component y is as follows (DHI, 2010):

$$\frac{\partial v}{\partial t} + \frac{\partial v^2}{\partial y} + \frac{\partial uv}{\partial x} + \frac{\partial wv}{\partial z} = -f_u - g \frac{\partial h}{\partial y} - \frac{1}{r_0} \frac{\partial p_a}{\partial y} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial y} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{yx}}{\partial x} + \frac{\partial S_{yy}}{\partial y} \right) + F_v + \frac{\partial}{\partial z} \left(v_t \frac{\partial v}{\partial z} \right) + v_s S$$

where x, y, z = cartesian coordinates of x, y, and z direction; t = time; u = water particle speed of direction x; v = water particle speed of direction y; w = water particle speed of direction z; η = water surface elevation; d = sea depth; h = total depth of the sea; f = coriolis parameter (f = 2ω sinØ); Ω = ratio of earth revolution; Ø = latitude coordinates; g = acceleration of gravity; S_{xx}, S_{xy}, S_{yx}, S_{yy} = component of radiation stress tensor; ν_t = vertical eddy viscosity; p_a = atmospheric pressure; ρ₀ = sea density; S = debits due to sources; u_s = particle velocity of air sources direction x; and v_s = the particle speed of air sources direction y.

Wave modeling used directional decoupled parametric and fully spectral formulations based on wave eternity movements with the input data of on-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry data, wind speed and direction and wave data. Parameterization was created in the frequency domain by introducing the 0 and 1st moments of the wave spectrum as a variable (Azhar *et al.*, 2011). The arranger equation was the equilibrium wave force in cartesian coordinate formulated as follow (DHI, 2010):

$$\frac{\partial N}{\partial t} + \frac{\partial}{\partial \phi} C \phi N + \frac{\partial}{\partial \lambda} C \lambda N + \frac{\partial}{\partial \sigma} C \sigma N + \frac{\partial}{\partial \theta} C \theta N = \frac{S}{\sigma}$$

Source energy, S, shows the superposition of surface functions in various styles (Azhar *et al.*, 2011).

$$S = S_{in} + S_{nl} + S_{ds} + S_{bot} + S_{surf}$$

where $N_{x,\sigma,\theta,t}$ = power density, φλ = coordinate spherical, C = speed of propagation of four-dimensional wave group, S = energy source, S_{in} = formation by wind, S_{nl} = energy transfer, non-linear, S_{ds} = wave energy due to white capping, S_{bot} = by bottom friction, S_{surf} = dissipation due to the depth-induced breaking.

We took the maximum value of the current speed and wave height in each grid that comes from 4 seasonal patterns characteristics, that is the west monsoon (December), transition 1 (May), east monsoon (July), and transition 2 seasons (October). The results of the 2D hydrodynamic model form the patterns of current speed and wave height every season were interpolated with a spline method with a cell size of 5 x 5 meters and analyzed using cell statistics to obtain the maximum value of current speed and wave height. The results of the distribution map were classified according to suitability class in table 2.

2.4. Determination of Area Suitability Criteria

The determination of the weighting and scoring was done to provide value in each criterion for supporting the feasibility of seaweed cultivation. The weight determination of each criterion was based on the literature research and later on was used to weight the variables by priority. The variable that is very influential for the growth of seaweed got a high score. The determination was based on the defined suitability matrix as shown in Table 2.

The total score multiplication and weight in table 2 were used to determine the class of water suitability of seaweed cultivation with quantitative analysis using the following equation :

$$Y = \sum a_i \cdot X_n$$

The range of suitability cultivation area of *K. Alvarezii* was adjusted to use the following equation :

$$I = \frac{(\sum a_i \cdot X_n) - (\sum a_i \cdot X_n)_{min}}{k}$$

where, Y = The final value, a_i = weight per parameter, and X_n = score of suitability level, I = interval of suitability class, k = number of suitability class.

Deleted: et al

Deleted: et al

Finally, the suitability interval was divided into 5 classes, 1) highly suitable (HS) is in the range of the value 42-50, 2) suitable category (S) is in the range of 33-41, C) moderately suitable (MS) range of 24-32, 4) less suitable (LS) are in the range of 15-23, and 5) not suitable (N) had the value range of < 15. Each parameter in table 2 was transformed in digital format using a spline interpolation method with a cell size of 5 x 5 meters resulting in an index map of water suitability.

2.5. Distance Suitability Analysis

Distance suitability analysis was also carried out as the other limiting factors out of the water parameters. Those were the distance to shipping lanes or sea lanes and the distance to a settlement which are arranged in a digital format with 5 x 5 -meter cell-sized using the buffer and euclidean distance method to consider the accessibility and disruption of marine traffic. These criteria are described in Table 3. The suitability of distance to the settlement was compiled using the data of inhabited island, namely Bontosua, Sanane, Panjenekang, Balanglombo, and Balangcaddi Islands, while the shipping lanes data used was inter-regional shipping routes. The results of the analysis were integrated with a water suitability index map.

Commented [H7]: The distance made by the buffer should be conveyed based on the determination of the distance,

Commented [W8R7]: done

2.6. Spatial Modelling

The spatial modelling in this study used cell-based modelling by integrating all oceanographic and water quality parameters based on the suitability classification according to table 2 with the following modelling algorithms develop from Danish Hydraulic Institute (2010):

$$WS = (T_i \times 0.5) + (pH_i \times 0.5) + (S_i \times 1) + (DO_i \times 0.5) + (N_i \times 1.5) + (F_i \times 1.5) + (B_i \times 0.5) + (HS_i \times 1.5) + (CS_i \times 1.5) + (TSS_i \times 0.5) + (SB_i \times 0.5)$$

with, WS = Water suitability, T_i = temperature index map, pH_i = pH index map S_i = salinity index map, DO_i = Dissolved Oxygen index map, N_i = Nitrate index map, F_i = phosphate index map, B_i = bathymetry index map, HS_i = significant wave height index map, CS_i = current speed index map, TSS_i = TSS index map, SB_i = seabed index map

The result of this modeling is continued by determining the classification of water suitability based on a class of suitability table 2. Then, integrating the results of water suitability above with the result of distance suitability analysis according to table 3 using cell statistics analysis.

Commented [H9]: The source of the algorithm used should be presented,

Commented [W10R9]: done

2.7. Estimating Productivity

After obtaining the distribution of classes areas for all variable, the seaweed production can be estimated only by considering the area which is suitable or highly suitable. Longline method of seaweed cultivation unit was designed as 25 x 50 meter per unit, the distance between the rope was 1 meter and the planting distance of each tied seaweed was 25 cm (WWF, 2014). The initial weight of the seedlings was approximately 100 grams per bunch (Parakkasi *et al.*, 2020) with the wet weight crop was 500 grams per tie. The cultivation of seaweed with a floating rope/longline system produces an excellent product and can gain a weight up to four to five times from the initial weight of the seedlings (100 grams) in a period of cultivation time. The number of the allocated cultivation units was estimated to be 5 units/ha. The distance correction between units is 10 meters in width (twice longer than the average length of the fishing boat) for fishing traffic. This amount represents 62.5% of the maximum number of units i.e., 8 units/ha. To estimated production is the multiplication between the number of cultivation units (from the total suitable area) and once crop yields per unit with the following equations:

Deleted: *et al*

$$Ye = N \times Cy$$

with, Ye = estimated yield per cycle (tons), N = number of cultivation unit, Cy = crop yield per cycle (tons). Briefly, the research method is summarized in the flowchart presented in [Figure 2](#).

3. Result and Discussion

3.1. Distribution of Water Quality Parameters

Spatial distributions of the suitability of water quality parameters are shown in [Figure 3](#). It can be seen that the water quality conditions based on the variable of SST, and pH, are highly suitable for seaweed cultivation in the study area. SST is the important limiting factor for marine organisms (Wirasatriya *et al.*, 2019). The range of measured SST in the study area is very narrow i.e., only around 30°C. **According to BSN (2015) the optimal temperature for this activity is 26-32°C, so the temperature in this site still suitable for seaweed cultivation.** The high temperature of the water causes death on seaweed while low water temperatures cause damage to protein and fat membranes and it is very influential in the seaweed life cycle. Photosynthesis on seaweed generally lasts at a maximum temperature of 23°C-30°C (Msuya, 2011; Ding *et al.*, 2013). Seawater has an enormous support ability to prevent pH fluctuation. The condition of pH that is very acidic or alkaline will endanger the survival of organisms because it led to metabolic and respiration disorders. The pH range at the study site is 7.26-7.63. **Whereas, optimal seaweed growth requires a pH value range of 7.5-8.5 (BSN, 2015; Radiarta *et al.*, 2018).** Moreover, Noor (2015) also stated that a suitable pH range for *K. alvarezii* cultivation is in normal and disposed alkaline level. **Therefore, these values of pH of this area are feasible for seaweed cultivation (Rahman *et al.*, 2019).**

The suitability distribution for TSS and substrate type are moderately suitable or more in the study area. A substrate is dominated by reef type almost at the entire study area which is categorized as moderately suitable. The suitable condition is located around the small islands with the sand as the substrate type. Seaweed will grow optimally when the cultivation site is in coral, rubble, sand-based substrate, or a mixture of all three. The most optimum substrate type for seaweed cultivation is a hard substrate like the rocky coral then followed by sandy corals (Parakkasi *et al.*, 2020). Waters with a fractional substrate of coral to sand are usually passed by the current that was suitable for the growth of seaweed. However, cultivating seaweed with a longline method is not overly influenced by the substrate type. For TSS, the eastern part of the study area is highly suitable while the western part is moderately suitable. This indicates that the further distance from the mainland the lower TSS is observed. TSS is insoluble sedimentation which increases the turbidity of water (Nurdin *et al.*, 2019). High turbidity due to the suspended material impedes light penetration into the water column and affects seaweed growth. Thus, TSS indicates water pollution as *K. alvarezii* lives optimally in transparent oligotrophic waters (Hayashi *et al.*, 2011). **According to Ministry of Environment R.I (1998), the recommended TSS value is <25mg/L, although up to 80 mg/L is allowed for marine cultivation.** The TSS levels in the study area range between 3.6-57.6 mg/L which means that the waters are suitable for the growth of *K. alvarezii*. TSS concentration is probably due to the absence of re-suspension due to the non-silt/clay of substrate type or no input from coastal sediment.

For DO and salinity, the suitability distributions vary from less suitable to highly suitable. Based on the DO criterion, the study area is mostly highly suitable for seaweed cultivation, with only 2 spots in the study area which are less suitable. The water circulation among small islands plays an important role in the high DO level. In open water, the low concentration of oxygen in a natural condition is rarely found (Eriksson, and Rosenberg, 2013). The optimal dissolved oxygen for seaweed cultivation is in the range of 3-7 mg/l (Rahman *et al.*, 2019). For salinity, the less suitable classes are spread in some areas and even the class at the eastern part of Sanane Island is not suitable for seaweed cultivation. The stability of the salinity becomes an important factor for seaweed because seaweed is a stenohaline organism that cannot resist the high fluctuation salinity (Reis *et al.*, 2010; Aris *et al.*, 2021). The extreme salinity changes can lead to ice-ice disease. Salinity can affect the osmoregulation process of marine biota including seaweed (Aris *et al.*, 2021). **Due to *K. alvarezii* can grow optimally in the range of**

Deleted: Fig.

Deleted: Fig.

Deleted: *et al*

Commented [H11]: Is the maximum temperature limit (Msuya, 2011; Ding *et al.*, 2013),

Deleted: *et al*

Commented [H12]: Should be directly compared with the appropriate range of pH limits,

Commented [W13R12]: done

Deleted: *et al*

Deleted: *et al*

Deleted: *et al*

Deleted: *et al*

Commented [H14]: It is recommended that the results of the TSS measurement are presented in paragraphs, so that the level of conformity is conveyed next,

Deleted: *et al*

Deleted: *et al*

Commented [H15]: It is advisable to provide a source,

salinity values 29 - 34 ppt (Doty,1987), therefore, most of the area is still feasible for seaweed cultivation.

Nitrate and phosphate concentrations as indicators of water fertility are relatively low in this region. This may be related to the absence of big rivers in the small islands of the study area as the nutrient sources. Nitrate is a limiting factor for seaweed growth. Nitrate is one of the nutrients needed for the growth of seaweed. The result shows that nitrate concentrations in the study area range between 0.001-0.12 mg/l which is relatively low. According to Kautsari and Ahdiansyah (2015), the feasible nitrate concentration for seaweed cultivation ranges from 0.01 mg/l to 0.07 mg/l, while the concentration of <0.01 mg/l and > 0.1 mg/l are not suitable for seaweed cultivation. Phosphate can be a limiting factor due to the source limitation. In the study area, phosphate concentration ranges from 0.001 mg/l to 0.062 mg/l which indicates low to insufficient fertility levels. Phosphate concentration for the fertile water should range from 0.051 mg/l to 0.1 mg/l. Besides, phosphate concentration also controls the growth (Nursidi *et al.*, 2017) and the carrageenan content of seaweed (Patty *et al.*, 2015).

Deleted: et al

The depth of the study area ranges from 2 m to 39.2 m. The longline method can be applied at the 1 m depth from the lowest tide or below (WWF, 2014). The sufficient depth provides seaweed to be submerged. Nevertheless, noted that the deeper waters will increase the cost of constructing seaweed cultivation. Thus, the areas with a depth of more than 15 m are categorized as not suitable for seaweed cultivation.

The distribution of physical and chemical parameters at the seaweed site is highly determined by the wave and current. These aspects play a pivotal role in the water movement for algae growth (Gaylord *et al.*, 2007; Kregting *et al.*, 2013; Nursidi *et al.*, 2017). The variability of current speed and wave height is distinguished by the 4 different characteristics of the west monsoon; transition 1; east monsoon; and transition 2 based on the results of 2D hydrodynamic modeling (Figure 4 and Figure 5). During west monsoon and transition 1 (east monsoon and transition 2), strong current speed appears at the northwestern (southern) part of the study area. The strongest current speed occurs during the transition 1 season. The pattern of the surface current in the study area is influenced by the current pattern of breaking waves in the small islands waters and the surrounding shallow area; and the influence of wind and tidal conditions in these waters. Based on the criteria shown in Table 2, the study area is categorized as highly suitable (dark blue color) to less suitable (light blue color).

Deleted: et al

Deleted: Fig.

Deleted: Fig.

For the wave height, almost entire study areas are categorized as suitable and highly suitable for seaweed cultivation except during east monsoon. During west monsoon transition 1 and transition 2, the blue and dark blue colors mean that the wave height ranges from 0.1 m to 0.3 m. During east monsoon, the wave higher than 0.3 m occurs at almost the entire study area which makes it less suitable for seaweed cultivation.

The protected area is determined by several types of physical conditions such as current velocity, and wave height. The influence of strong current and ocean waves will drag and damage the cultivated seaweed and its system. The maximum value of the distribution of current speeds and wave height for 4 seasons is described in Figure 6. Based on the results in Figure 6, the current speed and wave height around the study of the water are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. The less suitable areas for the current speed (wave height) category only appear mostly in the northeastern (northern) part of the study area. There is no area with a current speed of more than 0.5 m/s and the wave height more than 0.5 m which makes the area is not suitable for seaweed cultivation. The absence of strong current and wave may correspond to the morphology of the study area which is categorized as coral/reef waters with a lot of shallow/reef flat area located within the small islands that protect the area from the high wave and strong current. As mentioned by Sulma *et al.* (2008), the types of protected waters are bays, caps, straits, lagoons, and reef flat. Furthermore, Helmi *et al.* (2017) also stated that the coral reef is a shoreline protector.

Deleted: Fig.

Deleted: Fig.

Deleted: et al

Deleted: et al

3.2. Potential Areas for Seaweed culture Development and Production

Various suitability criteria as limiting factor of site selection for seaweed cultivation include physical and chemical parameters and distance suitability for accessibility and

restricted area inside the shipping lanes has been determined into 5 highly suitable classes (HS), suitable (S), moderately suitable (MS), less suitable (LS) and not suitable (N) (Figure 7). Figure 7 shows that most of the study areas are moderately suitable and suitable for seaweed cultivation. The area which is not suitable is only located along the shipping lane. Figure 8 summarizes the areal size of each criteria as follows: the highly suitable (HS) is 83.2 ha (0.42%); suitable (S) is 4,463.08 ha (22.21%); moderately suitable (MS) is 9,983.52 ha (49.68%); less suitable (LS) is an area of 494.84 ha (2.46%); and not suitable (N) is 5,071.15 ha (25.23%). The potential area for seaweed cultivation development reaches 4,546.28 ha as the sum area of highly suitable and suitable criteria. This result is more than obtained by Arifin *et al.* (2014) which only found 1,963.6 ha area which is suitable and highly suitable for *K. alvarezii* cultivation in the same study area. This indicates that our method can optimally classify the potential area for *K. alvarezii* cultivation. Furthermore, this potential area can be utilized for 22,730 cultivation units with seaweed production reaching about 56,825 tons of wet weight per crop, or within a year (2 times of harvesting) can reach about 113,650 tons. This amount is only about one-third of the total seaweed production in Pangkep Regency in 2018. However, it is important to be noted that, there are some limitations in the present study which should be considered. For example, the present study does not include the long-term climate variability which may also influence the physical and chemical parameters in the study area. As reported by Kunarso *et al.* (2019) and Msuya *et al.* (2014), the climate variability influences the oceanographic condition which then affects fisheries activity such as aquaculture and fish catch. The present study also does not consider the risk analysis which can disturb the cultivation process as shown by Castelar *et al.*, (2015). Furthermore, to maintain the carrying capacity of the study area, the potential area that might be utilized for the expansion of *K. alvarezii* cultivation is about 50% (covering 2,273.14 ha) of the total suitable area accordingly. The result of the present study indicates the effectiveness of **Geographic Information Systems with** geospatial modeling for developing the area of *K. alvarezii* culture which can be applied in the other area.

4. Conclusions

We have conducted a comprehensive geospatial modeling approach for the site selection of *K. alvarezii* cultivation at the Spermonde Islands. Physical and chemical parameters as well as the accessibility and restricted area of shipping lanes were used to define the suitability criteria for *K. alvarezii* cultivation. The result is SST and pH are categorized as highly suitable for *K. alvarezii* cultivation in the whole study area. A substrate is dominated by reef type almost at the whole study area which is categorized as moderately suitable. TSS levels in the study area range between 3.6-57.6 mg/l which means that the waters are suitable for the growth of *K. alvarezii*. For DO and salinity, the suitability distributions vary from less suitable to highly suitable. Nitrate and phosphate concentrations mainly are too low which are not suitable for *K. alvarezii* cultivation. The depth at the study area ranges from 2 m to 39.2 m. The area with a depth more than 15 m is categorized as not suitable for *K. alvarezii* cultivation. The current speed and wave height around the study of the water are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. After summarizing all criteria according to their weighting score, the suitable for seaweed cultivation at the study area is 4,546.28 ha, which is the combination of the suitable (S) area of 4,463.08 ha and highly suitable criteria (HS) with an area of 83.2 ha. The potential production per crop can reach 56,825 tons per cycle and 113,650 tons per year.

Acknowledgment

The author thanked the Ministry of Research Technology and Higher Education of the Republic of Indonesia for the financial support of research. The author thanked the Center for Coastal Rehabilitation and Disaster Mitigation Studies (CoREM), Diponegoro University for the facilities' support on hydro-oceanographic modeling and geospatial data processing. We also thanked the Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University for the surveys and laboratory analysis of water quality data.

Deleted: Fig.

Deleted: Fig.

Deleted: Fig.

Deleted: et al

Deleted: et al

Deleted: et al

Deleted: et al

References

- Afandi, A. Musadat, F. 2018. Suitability analysis for seaweed culture area at the waters of Kamelanta Village and Panjang Island using Geographical Information System. *Jurnal Akuakultura* 2(1), 69-78. (In Bahasa)
- Arbit, N., Omar, S., Soekendarsi, E., Yasir, I., Tresnati, J., Tuwo, A. 2019. Morphological and genetic analysis of *Gracilaria* sp. cultured in ponds and coastal waters. IOP Conf. Ser.: Earth Env. Sci. 370 012018.
- Arifin T., Bohari, R., Arlyza, I.S. 2014. Analysis of Space Suitability Based on Marine Agriculture in The Small Islands Around Makassar. *Forum geography* 28 (1), 91-102. DOI: <https://doi.org/10.23917/forgeo.v28i1.441>
- Aris, M., Muchdar, F., Labenua, R. 2021. Study of Seaweed *Kappaphycus alvarezii* Explants Growth in the Different Salinity Concentrations. *Jurnal Ilmiah Perikanan Dan Kelautan*, 13(1), 97–105. <https://doi.org/10.20473/jipk.v13i1.19842>
- Ariyanto, F. Y., Astor, Y., Sidqi, M. 2018. Application of water boundary theory for determining site licenses at the coastal area and small islands in Indonesia. *Proceeding National Seminar of Geomatics 2018*. Bandung, pp. 893-900. (In Bahasa)
- Azhar, R. M., Wurjanto, A., Yuanita, N. 2011. Study of coastal protection of submerged breakwater at Tanjung Kait beach. Master Thesis. Master Program of Water Resource Management, Bandung Institute of Technology, Indonesia.
- Bixler, H. J., Porse, H. 2011. A Decade of Change in the Seaweed Hydrocolloids Industry. *J. Appl. Phycol.* 23, 321-335.
- Castelar, B., de Siqueira, M. F., Sánchez-Tapia, A., Reis, R. P. 2015. Risk Analysis Using Species Distribution Modelling to Support Public Policies for The Alien Alga *Kappaphycus alvarezii* Aquaculture in Brazil. *Aquacult.* 446, 217-226.
- Danish Hydraulic Institute (DHI). 2010. MIKE-21 and MIKE-3 Flow Model. Hydrodynamic Module Step-by-Step Training Guide. DHI MIKE, Denmark. 96 p.
- Dee, D.P., Uppala, S.M., Simmons A.J., Berrisford, P., Poli, P., Kobayashi S., Andrae U., , M. A. Balmaseda, M.A., Balsamo G., Bauer, P., Bechtold, P. Beljaars, A.C.M., van de Berg L., Bidlot, J., Bormann, N., Delsol,C., Dragani, R., Fuentes,M., Geer, A.J.,, L. Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen L., Kállberg, P., Köhler M., Matricardi M, McNally, A.P., Monge-Sanz, B.M., Morcrette, J.J., Park, B.K., Peubey C, P. de Rosnay, P., Tavolato,C.,, Thépaut, J.N., Vitart , F., 2011.The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Q.J.R. Meteorol. Soc.*, 137(656), 553–597. doi: 10.1002/qj.828
- Doty, M.S., 1987. The production and use of *Eucaema*. in: Doty, M.A., Caddy, J.F, Santilices, B. (Eds), *Case Studies of Seven Commercial Seaweed Resources*, FAO Fish, Tech, Pap, 281,1987 p. 123- 161
- Ding L., Ma, Y., Huang, B., Chen, S. 2013. Effects of Seawater Salinity and Temperature on Growth and Pigment Contents in *Hypnea cervicornis*. *J. Agardh (Gigartinales, Rhodophyta)*. *BioMed Res. Int.* 2013, 594308. <https://doi.org/10.1155/2013/594308>
- Eriksson, R.J.H, and Rosenberg, H.R., 2013. *Managing Ocean Environments in a Changing Climate Sustainability and Economic Perspectives: Chapter 4.*

Oksigen - Hypoxia, Pages 67-96. <https://doi.org/10.1016/B978-0-12-407668-6.00004-5>

- Gaylord, B., Rosman, J. H., Reed, D. C., Koseff, J. R., Fram, J., Macintyre, S., Arkema, K., McDonald, C., Brzezinski, M. A., Largier, J. L., Monismith, S. G., Raimondi, P. T., Mardian, B. 2007. Spatial Patterns of Flow and Their Modification within and Around A Giant Kelp Forest. *Limnol. Oceanogr.* 52, 1838–1852.
- Hayashi, L., Santos, A. A., Faria, G. S. M., Nunes, B. G., Souza, M. S., Fonseca, A. L. D., Barreto, P. L. M., Oliveira, E. C., Bouzon, Z. L. 2011. *Kappaphycus alvarezii* (Rhodophyta, Areschougaceae) Cultivated in Subtropical Waters in Southern Brazil. *J. Appl. Phycol.* 23, 337–343.
- Helmi, M., Satriadi, A., Suryoputro, A. A. D., Marwoto, J., Setiyono, H. Hariyadi. 2018. Rehabilitation Priority Area Assessment on Death Coral using Cell-Based Modelling Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.* 9(11), 2949-2961.
- Helmi, M., Purwanto, Atmodjo, W., Subardjo, P., Aysira, A. 2017. Benthic Diversity Mapping and Analysis Base on Remote Sensing and Seascape Ecology Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.* 9(10), 227-235.
- Jalil, A. R., Samawi, M. F., Azis, H. Y., Jaya, I., Malik, A., Yunus, I., Sohopy, M. A. A. 2020. Comparison of physical-chemical conditions for seaweed cultivation in the Spermonde Archipelago, Indonesia. *AAFL Bioflux* 13(4), 2071-2082.
- Junaidi, M., Lestary, D. P., Scabra, A. R. 2018. Mariculture Site Selection Based on Environmental Parameters in Tanjung and Gangga Sub-District, North Lombok. In: *Proceeding of 1st ICWBB 2018*. Mataram University, Mataram, pp. 233-251.
- Kaiser, M. J., Yu, Y., Snyder, B. 2010. Economic Feasibility of Using Offshore Oil and Gas Structures in the Gulf of Mexico for Platform-based Aquaculture. *Mar. Pol.* 34, 699-707.
- Kautsari, N., Ahdiansyah, Y. 2015. Carrying capacity and suitability assessment of Labuhan Terata waters, Sumbawa for the development of seaweed culture. *Jurnal Ilmu Kelautan* 20(4), 233-238. (In Bahasa)
- Kim, J. K., Yarish, C., Hwang, E. K., Park, M., Kim, Y. 2017. Seaweed Aquaculture: Cultivation Technologies, Challenges, and Its Ecosystem Services. *Algae.* 32(1), 1-13.
- Kregting, L.T., Blight, A. B., Elsässer, B., Savidge, G. 2013. The Influence of Water Motion on The Growth Rate of The Kelp *Laminaria Hyperborea*. *J. Exp. Mar. Biol. Ecol.*, 448, 337–345.
- Kunarso, Wirasatriya, A., Irwani, Satriadi, A., Helmi, M., Prayogi, H. Munandar, B. 2019. Impact of Climate Variability on Aquatic Productivity and Fisheries Resources in Jepara. *IOP Conf. Ser.: Earth Env. Sci.*, 246 012021: 1-12.
- Longdill, P. C., Healy, T. R., Black, K. P., 2008. An integrated GIS approach for sustainable aquaculture management area site selection. *Ocean Coast. Manag.* 51, 612-624. doi:10.1016/j.ocecoaman.2008.06.010
- Malczewski, J. 2006. Ordered weighted averaging with fuzzy quantifiers: GIS-based multicriteria evaluation for land-use suitability analysis. *Int. J. Appl. Earth Observ. Geoinf.* 8(4), 270- 277.
- Malczewski, J., Rinner, C. 2015. *Multicriteria Decision Analysis in Geographic Information Science*. Springer, Berlin
- Micael, J., Costa, A.C., Aguiar, P., Medeiros, A., Calado, H. 2015. Geographic Information System in A Multi-criteria tool for Mariculture Site Selection. *Coast. Manag.* 43(1), 52-66.

- Ministry of Environment R.I, Decree of the State Minister for Population and Environment Number 02 of 1988 concerning Guidelines for Determining Environmental Quality Standards for Marine Biota (Aquaculture). Jakarta, 1988, p. 31.
- Msuya, F. E., Buriyo, A., Omar, I., Pascal, B., Narrain, K., Ravina, J. J. M., Mrabu, E., Wakibia, J. G. 2014. Cultivation and Utilisation of Red Seaweeds in the Western Indian Ocean (WIO) Region. *J. Appl. Phycol.* 26, 699–705.
- Msuya, F. E. 2011. Environmental Changes and Their Impact on Seaweed Farming in Tanzania. *World Aquacult.* 42, 34-37.
- National Standardization Agency (BSN), 2015. **Good Aquaculture Practices (CBIB). 2 : Seaweed.** SNI, 8228.2.:2015. National Standardization Agency (in Bahasa)
- Nirmala, K., Ratnasari, A., Budiman, S. 2014. Site Selection for Seaweed Culture at Gerupuk Bay-West Nusa Tenggara Using Remote Sensing and GIS. *Jurnal Akuakultur Indonesia* 13(1), 73-82.
- Noor, N. M. 2015 The Water Bodies Compatibility Analysis for Culturing Brown Seaweed *Kappaphycus alvarezii* in Ketapang Seashore, South Lampung. *Maspari Journal* 7(2), 91-100. (in Bahasa)
- Nurdin, N., Helmi, M., Setiawan, R. Y., Zainudin, M., Rustam, A., Akbar, M. 2019. Landsat Capability to Mapping of Total Solid Suspended (TSS) and Coastal Ecosystem. *ISPRS – Int. Arch. Photogram., Remote Sens. Spatial Inf. Sci.*, 42(4), 497-502.
- Nursidi, Ali S. A., Anshary H., Tahya A. M., 2017 **Environmental parameters and specific growth of *Kappaphycus alvarezii* in Saugi Island, South Sulawesi Province, Indonesia.** *AAFL Bioflux* 10(4):698-702.
- Parakkasi, P., Rani, C., Syam, R., Zainuddin, Achmad, M. 2020. Growth Response and Quality of Seaweed *Kappaphycus alvarezii* Cultivated in Various Coastal Ecosystems in the Waters of West Sulawesi, Indonesia. *AAFL Bioflux* 13(2), 627-639.
- Patty, S. I., Arfah, H., Abdul, M. S. 2015. The relationship among Nutrient (Phosphate and Nitrate), DO, and pH with the water fertility in Jikumerasa waters, Buru Island. *Jurnal Pesisir dan Laut Tropis* 1(1), 43-50. (In Bahasa)
- Pickering, T. D., Skelton, P. Sulu, J. R. 2007. Intentional Introductions of Commercially Harvested Alien Seaweeds. *Bot. Mar.* 50, 338–350.
- Radiarta, I.N., Erlania, Haryadi, J., 2018. **Analysis of land suitability and carrying capacity of waters for the development of seaweed cultivation in Simeulue Regency; Aceh.** *Jurnal Segara*, 14 (1) : 11 – 22. *Jurnal Segara*, 14 (1) : 11 – 22 (In Bahasa)
- Radiarta, I. N., Saitoh, S. Yasui, H. 2011. Aquaculture Site Selection for Japanese Kelp (*Laminaria japonica*) in Southern Hokkaido, Japan, Using Satellite Remote Sensing and GIS-based Models. *ICES J. Mar. Sci.* 68(4), 773-780. doi:10.1093/icesjms/fsq163
- Radiarta, I. N., Saputra, A. Albasri, H. 2012. Feasibility mapping of seaweed culture area (*Kappaphycus alvarezii*) at Bintan Regency, Riau Islands Province using GIS and remote sensing approach. *Akuakultur* 7(1): 145-157. (In Bahasa)
- Rahman, I. S., Fadjar, M., Tjahjono, A. 2019. The Relationship Water Physical-Chemical Parameters in Seaweed Cultivation *Eucheuma cottonii* with Long-line System. *Int. J. Sci. Tech. Res.* 8(11), 1448-1452.
- Rahmayanti, F., Diana, F., Kusumawati, I. 2018. Suitability analysis for area development of seaweed (*Eucheuma cottonii*) culture in Lhok Bubon waters, Samatiga District, Aceh Barat Regency. *Jurnal Akuakultura* 2(1), 26-32. (In Bahasa).

- Reis, R. P., Loureiro, R. R., Mesquita, F. S. 2010. Do Salinity Affect Growth and Carrageenan Yield of *Kappaphycus alvarezii* (Gigartinales/ Rhodophyta)? *Aquacult. Res.* 42, 1231-1234.
- Rusliani. 2016. Study of water quality on the growth of seaweed *Gracilaria verrucosa* in a pond with verticulture method. *Octopus J. Fish. Sci.* 5(2), 522-527. (in Bahasa)
- Setiawan, R. Y., Kawamura, H. 2011. Summertime Phytoplankton Bloom in the south Sulawesi Sea. *IEEE J. Sel. Topics Appl. Earth Observ. Remote Sens.* 4(1), 241-244. doi:10.1109/JSTARS.2010.2094604.
- Sousa, F. E. S., Moura, E. A., Marinho-Soriano, E. 2012. Use of Geographic Information Systems (GIS) to Identify Adequate Sites for Cultivation of The Seaweed *Gracilaria birdie* in Rio Grande Do Norte, Northeastern Brazil. *Revista Brasileira de Farmacognosia* 22(4), 868–873.
- Sulma, S., Manoppo, A. K. S., Hartuti, M. 2008. Identification of Suitable Area for Seaweed Culture in Bali Waters Based on Remote Sensing Satellite Data. *Int. J. Remote Sens. Earth Sci.* 5, 57-66.
- Susanto, R. D., Wei, Z., Adi, T. R., Zheng, Q., Fang, G., Bin, F., Supangat, A., Agustyadi, T., Li, S., Trenggono, M., Setiawan, A. 2016. Oceanography Surrounding Krakatau Volcano in the Sunda Strait, Indonesia. *Oceanogr.* 29(2), 228-237. <http://dx.doi.org/10.5670/oceanog.2016.31>.
- Teniwut, W. A., Marimin, Djatna, T. 2019. GIS-Based multi-criteria decision-making model for site selection of seaweed farming information center: A lesson from small islands, Indonesia. *Decis. Sci. Let.* 8(2019), 137-150 doi: 10.5267/j.dsl.2018.8.001
- Utama, R. Z., Handayani, T. 2018. Suitability Between Oceanography and Seaweed (*Eucheuma cottonii*) Cultivation Potential in Tidung Island with Geographic Information System (GIS). *ICENIS: E3S Web of Conference* 73, 12009. 1-5.
- Valderrama, D., Cai, J., Hishamunda, N., Ridler, N., Neish, I. C., Hurtado, A. Q., Msuya, F. E., Krishnan, M., Narayanakumar, R., Kronen, M., Robledo, D., Gasca-Leyva, E., Fraga, J. 2015. The Economics of *Kappaphycus* Seaweed Cultivation in Developing Countries: A Comparative Analysis of Farming Systems. *Aqua. Econ. Manag.* 19, 251–277.
- Wirasatriya, A., Sugianto, D. N., Helmi, M., Setiawan, R. Y. Koch, M. 2019. Distinct Characteristics of SST Variabilities in the Sulawesi Sea and the Northern Part of the Maluku Sea during the Southeast Monsoon. *IEEE J. Sel. Topics Appl. Earth Obs. Remote Sens.* 12(6), 1763-1770.
- WWF-Indonesia. 2014. Better Management Practices: Guideline series of small scale fisheries of seaweed culture - Cottoni (*Kappaphycus alvarezii*), Sacol (*Kappaphycus striatum*) dan Spinosum (*Eucheuma Denticulatum*) Version 1. WWF-Indonesia. South Jakarta. 53 p. (in Bahasa)
- Yulianto, H., Damai, A. A., Delis, P. C., Elisdiana, Y. 2017. Spatial Analysis to Evaluate the Suitability of Seaweed Farming Site in Lampung Bay, Indonesia. *Turkish J. Fish. Aquatic Sci.* 17, 1253-1261.

List of Tables

Table 1. Water quality data collection

No	Variables	Unit	Tools/Methods	Data Retrieval Methods
1	Sea Surface Temperature (SST)	°C	Thermometer	In situ
2	Salinity	Psu	Salinometer	In situ
3	pH	-	<i>Water quality checker</i>	In situ
4	Dissolved oxygen (DO)	ppm	Winkler method	Laboratory analysis
5	Nitrate	mg/l	Ascorbic Acid Method	Laboratory analysis
6	Phosphate	mg/l	Brucine Sulfate Method	Laboratory analysis
7	Total Suspended Solids (TSS)	mg/l	<i>Milipore filter</i>	Laboratory analysis
8	Substrate type	-	Visually observation	In situ
9	Depth	m	Modelled depth from LPI Maps (BIG, 2011) and BATNAS (BIG, 2018) data	Integrating modelled data

Table 2. Table of suitability criteria for seaweed *Kappaphycus alvarezii* cultivation

Variables	Level of conformity (score)					Weight	Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)		
SST (°C)	28-30	24-27 or 31-32	22-23 or >32	21 or >34	<21 or >36	0.5	Modification of Noor (2015); Rahmayanti <i>et al.</i> (2018)
pH	7-8	6.5-7 or 8-8.5	6-6.4 or >8.5	4-6	<4 or >9	0.5	Modification of Kautsari and Ahdiansyah (2015); Noor (2015)
Salinity (psu)	28-31	26-28 or 31-33	<26 or ≥ 34	18-25 or 35-36	<18 or >37	1	Modification of Kautsari and Ahdiansyah (2015); Rahmayanti <i>et al.</i> (2018); Utama and Handayani (2018)
DO (mg/l)	>6	5-6	4-5	2-4	<2	0.5	Modification of Afandi and Musadat (2018)
Nitrate (mg/l)	0.04-0.07	0.02-0.03	0.01 or >0.1	<0.01 or >0.5	>1	1.5	Modification of Kautsari and Ahdiansyah (2015); Afandi and Musadat (2018)
Phosphate (mg/l)	0.051-0.1	0.02-0.04	0.01-<0.02 or >0.2-1	<0.01 or >1	>2	1.5	Modification of Patty <i>et al.</i> (2015)
Depth (m)	3-10	2-3 or 11-13	1-2	>13-15	<1 or >15	0.5	Modification of Tiensongrusme (1990) in Radiarta <i>et al.</i> (2012); Kautsari and Ahdiansyah (2015); Yulianto <i>et al.</i> (2017)
Wave height (m)	0.2-0.3	0.1-0.19	>0.3	<0.1 or >0.4	>0.5	1.5	Modification of Rahmayanti <i>et al.</i> (2018)
Current speed (m/s)	0.25-0.3	0.2-0.24 or 0.31-0.4	0.1-0.19 or >0.4	<0.1 or >0.45	>0.5	1.5	Modification of Utama and Handayani (2018)
TSS (mg/l)	<25	25-50	50-100	100-400	>400	0.5	Modification of Yulianto <i>et al.</i> (2017)
Seabed	Coral, Rubbel	Sand	Reefs	Muddy Sand	Mud	0.5	Modification of Noor (2015); Yulianto <i>et al.</i> (2017)

Note : HS: Highly suitable, S: suitable, MS: moderately suitable, LS: less suitable, and N: not suitable


Table 3. Distance suitability analysis

Parameters	Suitability Level (Score)					Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)	
Shipping lanes (m)	> 500	-	-	-	≤ 500	Ariyanto <i>et al.</i> (2018)
Distance to settlements (km)	4	4-5	5-6	6-7	>7	Modification of Radiarta <i>et al.</i> (2012)

Deleted: et al

Deleted: et al

List of Figures






Figure 2. Research Flowchart

Commented [RV16]: Text can not read

Commented [W17R16]: done




Figure 3. Distribution of physical and chemical parameters: a) SST (Sea surface temperature); b). Depth; c). Salinity; d).Nitrate; e). Phosphate; f).; DO (dissolved Oxygen); g).TSS; h: seabed type; i): pH




Figure 4. Distribution of current speed (m/s) based on 2D hydrodynamic model in each season




Figure 5. Distribution of significant wave height (m) based on 2D hydrodynamic model in each season




Figure 6. Maximum statistical value of current speed (left) and wave height (right) for all season




Figure 7. Suitability distribution for cultivation area of *K. Alvarezii*




Figure 8. Potential area for *K. alvarezii* cultivation in each

REBUTAL IJMS

Title : Identification of Potential Locations for Kappaphycus Alvarezii Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia

Authors : Sarjito Sarjito^{1*}, Hanny Ammaria¹, Muhammad Helmi^{2,3}, Slamet Budi Prayitno¹, Nurjannah Nurdin^{4,5}, Riza Yuliratno Setiawan⁶, Parichat Wetchayont⁷, Anindya Wirasatriya^{2,3}

	Comment /Suggestion of the reviews	Corrections
	Review A	
1.	Hydrodynamics of marine waters has dynamic properties divided into 4 seasonal models, namely: West Monsoon, Transition 1, East Monsoon, and Transition 2. This article will be interesting if each model is made a model for the distribution of Kappaphycus alvarezii cultivation locations.	Thank you very much for your suggestion. However, in our opinion it is not applicable since in the Kappaphycus alvarezii cultivation, it is not efficient if should move the location following the changing season. In our analysis we have already considered the influence of season by combining the model of dynamic properties in each season (Figure 4 and Figure.5).
2.	Where are recommendations for selecting and developing Kappaphycus alvarezii cultivation sites? whether in HS, S, MS?	Thank you very much for your recommendation. However, in our article have already mentioned about these. The area of HS, S and MS has already clearly shown in Figure 7.
	Review B	
1.	Suggestions for the title "Identification of Potential Locations for Kappaphycus Alvarezii Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia.	Thank you for your excellent suggestion. We already changes the title : Identification of Potential Locations for Kappaphycus Alvarezii Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia.
2.	Abstract must in 250 words	Thank you very much for your suggestions. However the maximum abstract is 250 words
3.	It is better for the background on the existing fundamental problems, "However, the availability of natural seaweed can be threatened along with the increasing demand so that seaweed cultivation is absolutely necessary (Pickering et al., 2007)."	Thank you very much for your suggestions, we have already canged; However, the availability of natural seaweed can be threatened along with the increasing demand so that seaweed cultivation is absolutely necessary.

4.	Deleted et al. in the hole article.	Thank you very much for your corrections. We have done in this article
5.	The distance made by the buffer should be conveyed based on the determination of the distance,	Thank you very much for your suggestion. Buffer distance method is one of distance determination method in GIS. We used the buffer distance method as one of methods for determining distance from.shipping lane and settlement.
6.	The source of the algorithm used should be presented !	Done algorithms develop from Danish Hidraulic Institute (2010):
7.	Is the maximum temperature limit (Msuya, 2011; Ding et al., 2013),	It is not necessary . We have added the proper references : According to BSN (2015) the optimal temperature for this activity is 26-32°C, so the temperature in this site still suitable for seaweed cultivation.
8.	Should be directly compared with the appropriate range of pH limits,	Thank you For your good suggestion. So we changed the sentence : The pH range at the study site is 7.26-7.63. Whereas, optimal seaweed growth requires a pH value range of 7.5-8.5 (BSN, 2015; Radiarta et al., 2018). Moreover, Noor (2015) also stated that a suitable pH range for <i>K. alvarezii</i> cultivation is in normal and disposed alkaline level. Therefore ,these values of pH of this area are feasible for seaweed cultivation (Rahmancet al., 2019).
10.	It is recommended that the results of the TSS measurement are presented in paragraphs, so that the level of conformity is conveyed next,	Thank you verymuch for your good suggestion So,we canged in red transparent oligotrophic waters (Hayashi et al., 2011). According to Ministry of Environment R.I (1998), the recommended TSS value is <25mg/L, although up to 80 mg/L is allowed for marine cultivation. TSS levels in the study area range between 3.6-57.6 mg/L which means that the waters are suitable for the growth of <i>K. alvarezii</i> .
11.	The title of the image is still not informative, it should be given letter numbering to make it easier for users,	Thank you for your correction We changed in Red

		<p>Figure 3. Distribution of physical and chemical parameters : a) SST (Sea surface temperature); b). Depth; c). Salinity; d).Nitrate; e). Phosphate; f).; DO (dissolved Oxygen); g).TSS; h: seabed type; i): pH</p>
--	--	--



Sarjito Sarjito <sarjito@live.undip.ac.id>

[IK.IJMS] [ID-47735] Revised Version Acknowledgement

3 messages

Ambariyanto <ijms@live.undip.ac.id>
Reply-To: sarjito - sarjito <sarjito@live.undip.ac.id>
To: sarjito - sarjito <sarjito@live.undip.ac.id>

Tue, Aug 16, 2022 at 7:33 AM

sarjito - sarjito:

Thank you for submitting the revision of manuscript, "Site selection of Kappaphycus alvarezii farming to optimize seaweed production using Geographic Information system with geospatial modeling at Spermonde Islands waters, South Sulawesi, Indonesia" to ILMU KELAUTAN: Indonesian Journal of Marine Sciences. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL: <https://ejournal.undip.ac.id/index.php/ijms/author/submission/47735>
Username: sarjito
Editor: Indonesian Journal of Marine Science

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Ambariyanto
ILMU KELAUTAN: Indonesian Journal of Marine Sciences

INDONESIAN JOURNAL OF MARINE SCIENCE
<http://ejournal.undip.ac.id/index.php/ijms>

Ambariyanto <ijms@live.undip.ac.id>
Reply-To: sarjito - sarjito <sarjito@live.undip.ac.id>
To: sarjito - sarjito <sarjito@live.undip.ac.id>

Tue, Aug 16, 2022 at 7:36 AM

[Quoted text hidden]

Ambariyanto <ijms@live.undip.ac.id>
Reply-To: sarjito - sarjito <sarjito@live.undip.ac.id>
To: sarjito - sarjito <sarjito@live.undip.ac.id>

Tue, Aug 16, 2022 at 7:36 AM

[Quoted text hidden]



Sarjito Sarjito <sarjito@live.undip.ac.id>

[IK.IJMS] Editor Decision

3 messages

Ilmu Kelautan <ijms.undip@gmail.com>
To: sarjito@live.undip.ac.id

Tue, Aug 30, 2022 at 8:03 AM

Dear
Sarjito

We have reached a decision regarding your submission to ILMU KELAUTAN: Indonesian Journal of Marine Sciences, "Identification of Potential Location for Kapphyscus alverazi Optimization of Seaweed Production Based on Geographic Information Systems In Spermonde Archipelago Waters, South Sulawesi, Indonesia".

Our decision is to:
Accepted

Indonesian Journal of Marine Science
ijms.undip@gmail.com



Sarjito Sarjito <sarjito@live.undip.ac.id>

[IK.IJMS] Proof Layout - 47735

2 messages

Ilmu Kelautan <ijms.undip@gmail.com>
To: sarjito@live.undip.ac.id

Thu, Sep 8, 2022 at 11:10 AM

Dear
Sarjito
Aquaculture Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro

Please check the manuscript proof. Utilize the file contained in this email attachment to make necessary corrections to the manuscript.

Send the cost of publication (Rp. 2.500.000,-) to Widianingsih
Bank : Bank Nasional Indonesia (BNI)
Branch : UNDIP Semarang, Indonesia
Account No : 0162816376

Manuscript revise and proof of payment are sent before September 11, 2022 to publish the manuscript in September 2022.

Thank you for your contribution to this journal.

Sincerely
IJMS Editor



Virus-free. www.avast.com



47735-152785-2-ED proof.docx
1804K

Identification of Potential Locations for *Kappaphycus alvarezii* Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia

Sarjito Sarjito^{1*}, Hanny Ammaria¹, Muhammad Helmi^{2,3}, Slamet Budi Prayitno¹, Nurjannah Nurdin^{4,5}, Riza Yuliratno Setiawan⁶, Parichat Wetchayont⁷, Anindya Wirasatriya^{2,3}

¹Aquaculture Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro

²Oceanography Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro

³Center for Coastal Rehabilitation and Disaster Mitigation Studies

Jl. Prof.H. Soedarto, S.H. Tembalang, Semarang, Central Java, 50275 Indonesia

⁴Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University

⁵Marine Science and Fisheries Faculty, Hasanuddin University

Jl. Perintis Kemerdekaan Km.10 Tamalanrea, Makassar, Sulawesi Selatan Indonesia

⁶Marine Science and Fisheries Faculty, Gadjah Mada University

Bulaksumur, Caturtunggal, Depok, Sleman, Daerah Istimewa Yogyakarta 55281 Indonesia

⁷Department of Geography, Srinakharinwirot University

Sukhumvit23, Wattana,, Bangkok, 10110, Thailand

Email: sarjito@live.undip.ac.id

Abstract

Indonesia is the second largest producer of red algae in the world with one of the major *Kappaphycus alvarezii* farming areas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province. *K. alvarezii* production in Pangkep Regency increased every year. However, the availability of natural seaweed can be threatened along with the increasing demand so that seaweed cultivation is necessary. Proper site selection is crucial to guarantee the success of the seaweed farming development. We conducted a Geographic Information system (GIS) with geospatial modeling approach for the site selection of *K. alvarezii* cultivation at the Liukang Tuppabiring District, Spermonde Islands by combining various physical and chemical parameters obtained from in-situ data and hydrodynamics modeling. The parameters are Sea Surface Temperature, (SST), salinity pH, substrate type, current speed, wave height, DO, nitrate, phosphate, and Total Suspended Solid (TSS). We also include the accessibility and the restricted area of shipping lanes for determining the area suitability of *K. alvarezii* cultivation. This research demonstrates the cell-based geospatial modeling can be effectively used and found the suitable categorized area for *K. alvarezii* culture of 4,546.28 ha. The potential productivity of wet-weight seaweed can reach 56,825 tons per cycle or 113,650 tons per year. This result indicates the GIS with geospatial modeling approach for developing the area of *K. alvarezii* culture which can be applied in the other area.

Commented [A1]: Avoid using "we"

Commented [A2]: Make this abstract to 250 words

Keywords: Seaweed production, site selection, geospatial, hydrodynamics modeling

Introduction

Indonesia is the largest producer of red algae in the world (83% of global seaweed production) followed by the Philippines and several countries (Valderrama *et al.*, 2015; Kim *et al.*, 2017). Seaweed production in Indonesia comes not only from mariculture, and brackish water ponds but also from natural collections. However, the availability of natural seaweed can be threatened in line with their increasing demand so that seaweed cultivation is completely necessary (Pickering *et al.*, 2007). The increasing demand for seaweed was mostly driven by new markets in China, Eastern Europe, Brazil, and others (Bixler and Porse, 2011). Furthermore, Arbit

et al. (2019) stated that 99.73% of Indonesia's seaweed production comes from cultivation. Therefore, seaweed farming is the best choice for fulfilling market demand.

Site selection is essential in aquaculture development which depends on the species of commodities and applied cultivation methods (Kaiser *et al.*, 2010; Junaidi *et al.*, 2018). For seaweed, the success of cultivation is intrinsically linked to physical and chemical parameters of seaweed farming areas such as turbidity, temperature, salinity, pH, DO, nitrate and phosphate concentration (Rusliani, 2016; Jalil *et al.*, 2020). Hydrodynamic aspects should also be considered in

the site selection of *K. alvarezii* farming since the cultivation method uses the longline method which is prone to strong current and wave. One of major *K. alvarezii*'s farming areas within the Indonesian seas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province (Jalil *et al.*, 2020). The existence of Indonesian Troughflow along the Makassar Strait (Susanto *et al.*, 2016) and the occurrence of strong upwelling at the southern part of the South Sulawesi during each summer (Setiawan and Kawamura 2011) may influence the hydrodynamics condition in the Spermonde Island. Besides, the existence of small islands among the water area makes the hydrodynamics condition at the Spermonde Islands becomes more complex. Spermonde Islands area also has dense shipping activities. Many shipping lanes have lied among these areas which are should be avoided in the mariculture site selection. This shows the complex considerations for conducting site selection of seaweed farming at Spermonde Islands.

In dealing with the complex considerations as mentioned above, the use of the geospatial model which is part of Geographical Information Systems (GIS) becomes an effective solution for determining the proper mariculture site selection (Longdill *et al.*, 2008; Micael *et al.*, 2015). This approach, in general, is an analysis of layers that contains spatial data to evaluate and select a particular area based on its suitability criteria classified according to certain environmental aspects (Malczewski, 2006; Malczewski & Rinner, 2015). GIS utilization in mariculture development has been widely used in various types of commodities including seaweed (e.g., Radiarta *et al.*, 2011; Radiarta *et al.*, 2012; Sousa *et al.*, 2012; Nirmala *et al.*, 2014; Teniwut *et al.*, 2019) which were mainly used in-situ data and remote sensing data for the input of geospatial modeling. The application of GIS for site selection for aquaculture in the Spermonde Islands has conducted by Arifin *et al.* (2014). However, their analysis was only based on in-situ measurement which cannot represent the temporal change of ocean parameters. In the present study, cell-based geospatial modeling was performed for the determination of sustainable site selection of seaweed cultivation, which integrates various physical and chemical parameters obtained from in-situ data and the hydrodynamic model. The combination of both data becomes the new approach proposed in the present study. Considering that the hydrodynamics aspects for the site selection of seaweed culture is not available in the remote sensing data, a 2D hydrodynamic model was built to represent the hydrodynamic condition in the study area. On the other hand, Indonesia as the maritime continent experiences four-season influenced by monsoon wind (Helmi *et al.*, 2018) which impacts

the hydrodynamics conditions. Thus, in-situ and incidental measurements of current and wave cannot represent the hydrodynamics condition in the study area. This issue only can be resolved by performing the hydrodynamics model. Furthermore, the potential productivity of *K. alvarezii* at the selected site is also simulated. This research is expected to become a prototype of a geospatial model to explore the potential aquaculture area in small islands and can be a scientific-based information for local governments, researchers, NGOs, and related stakeholders for planning and developing *K. alvarezii* cultivation in the study area.

Material and Method

The study area is presented in Figure 1. which is located at the Spermonde islands waters precisely in the Liukang Tupabbiring District, Pangkep Regency, South Sulawesi. The Liukang Tupabbiring District area is 54.44 km² (4.89% of Pangkep Regency) and consists of 20 islands. The population of this district reached 29,680 inhabitants, generally fishermen and sea farmers. The field stations are located at 4°5'3.98"S - 4°59'51.137"S and 119°18'35.532"E - 119°26'19.849"E. There are 94 sampling points distributed among the island of Bontosua, Sanane, Panjenekang, Pankaiya, Langkadea, Balanglombo, Balangcaddi Island, and reef flat area.

A field survey was conducted for collecting water quality data at 94 points in which the depths are more than 2 meters. Survey site determination was using Garmin 76C GPS. The water quality data collected are presented in Table 1. The in-situ measurements were Sea Surface Temperature, (SST), salinity, pH, and substrate type. For the chemical parameters, such as DO, nitrate, phosphate, and Total Suspended Solid (TSS), the water samples were taken and analyzed at Chemical Oceanography Laboratory, Marine Science Department, Hasanuddin University, Indonesia (May 2019). The dynamic and non-dynamic water parameter data were interpolated to obtain a spatial distribution pattern at the time of data retrieval. The data were compiled in a digital form of 5 x 5 meters cell size using spline and classified according to suitability class in Table 2.

Two dimensions (2D) hydrodynamic model was used to simulate current and wave condition at the study area. The data for generating a hydrodynamic model are as follows: (1) High-resolution satellite imagery GeoEye-1 from ESRI online (<https://esriindonesia.co.id/arcgis-online>); (2) Bathymetry data from Indonesia Geospatial Agency (<http://batnas.big.go.id/>); (3) Surface wind and wave height data obtained from 6 hourly ERA-interim data

with a grid interval of 0.125° from July 2018 to July 2019 (Dee et al., 2011); (4) Tidal elevation forecast




Figure 1. Study area

Table 1. Water quality data collection

No	Variables	Unit	Tools/Methods	Data Retrieval Methods
1	Sea Surface Temperature (SST)	°C	Thermometer	In situ
2	Salinity	Psu	Salinometer	In situ
3	pH	-	Water quality checker	In situ
4	Dissolved oxygen (DO)	ppm	Winkler method	Laboratory analysis
5	Nitrate	mg.L ⁻¹	Ascorbic Acid Method	Laboratory analysis
6	Phosphate	mg.L ⁻¹	Brucine Sulfate Method	Laboratory analysis
7	Total Suspended Solids (TSS)	mg.L ⁻¹	Millipore filter	Laboratory analysis
8	Substrate type	-	Visually observation	In situ
9	Depth	m	Modelled depth from LPI Maps (BIG, 2011) and BATNAS (BIG, 2018) data	Integrating modelled data

from July 2018 to July 2019 (<https://www.dhigroup.com/download/mike-by-dhi-tools/coastandseatools/global-tide-model>). The current model was constructed by continuity equations and momentum equations with the average of depth. On-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry, wind and tidal dynamics were used as an input for generating current model. The initial process was making a mesh, and setting the boundary condition of the model using the mathematical continuity equation as follows:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = S$$

The horizontal momentum equation for component x is as follows (DHI, 2010):

$$\frac{\partial u}{\partial t} + \frac{\partial u^2}{\partial x} + \frac{\partial vu}{\partial y} + \frac{\partial wu}{\partial z} = fv - g \frac{\partial h}{\partial x} - \frac{1}{r_0} \frac{\partial p_a}{\partial x} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial x} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{xx}}{\partial x} + \frac{\partial S_{xy}}{\partial y} \right) + F_u + \frac{\partial}{\partial z} \left(v_t \frac{\partial u}{\partial z} \right) + u_s S$$

While the horizontal momentum equation for component y is as follows (DHI, 2010):

$$\frac{\partial v}{\partial t} + \frac{\partial v^2}{\partial y} + \frac{\partial uv}{\partial x} + \frac{\partial wv}{\partial z} = fu - g \frac{\partial h}{\partial y} - \frac{1}{r_0} \frac{\partial p_a}{\partial y} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial y} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{yx}}{\partial x} + \frac{\partial S_{yy}}{\partial y} \right) + F_v + \frac{\partial}{\partial z} \left(v_t \frac{\partial v}{\partial z} \right) + v_s S$$

Note: x, y, z = cartesian coordinates of x, y, and z direction; t = time; u = water particle speed of direction x; v = water particle speed of direction y; w = water particle speed of direction z; η = water surface elevation; d = sea depth; h = total depth of the sea; f = coriolis parameter (f = 2ω sinθ); Ω = ratio of earth revolution; θ = latitude coordinates; g = acceleration of gravity; S_{xx}, S_{xy}, S_{yx}, S_{yy} = component of radiation stress tensor; v_t = vertical eddy viscosity; p_a = atmospheric pressure; ρ₀ = sea density; S = debits due to sources; u_s = particle velocity of air sources direction x; and v_s = the particle speed of air sources direction y.

Wave modeling used directional decoupled parametric and fully spectral formulations based on wave eternity movements with the input data of on-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry data, wind speed and direction and wave data. Parameterization was created in the frequency domain by introducing the 0 and 1st moments of the wave spectrum as a variable (Azhar *et al.*, 2011). The arrange equation was the equilibrium wave force in cartesian coordinate formulated as follow (DHI, 2010):

$$\frac{\partial N}{\partial t} + \frac{\partial}{\partial \phi} C\phi N + \frac{\partial}{\partial \lambda} C\lambda N + \frac{\partial}{\partial \sigma} C\sigma N + \frac{\partial}{\partial \theta} C\theta N = \frac{S}{\sigma}$$

Source energy, S, shows the superposition of surface functions in various styles (Azhar *et al.*, 2011).

$$S = S_{in} + S_{nl} + S_{ds} + S_{bot} + S_{surf}$$

Note : $N_{x,\sigma,\theta,t}$ = power density, φλ = coordinate spherical, C = speed of propagation of four-dimensional wave group, S = energy source, S_{in} = formation by wind, S_{nl} = energy transfer, non-linear, S_{ds} = wave energy due to white capping, S_{bot} = by bottom friction, S_{surf} = dissipation due to the depth-induced breaking.

The maximum value of the current speed and wave height were taken in each grid that comes from 4 seasonal patterns characteristics, that is the west monsoon (December), transition 1 (May), east monsoon (July), and transition 2 seasons (October). The results of the 2D hydrodynamic model form the patterns of current speed and wave height every season were interpolated with a spline method with a cell size of 5 x 5 meters and analyzed using cell statistics to obtain the maximum value of current

speed and wave height. The results of the distribution map were classified according to suitability class in Table 2.

Determination of Area Suitability Criteria

The determination of the weighting and scoring was done to provide value in each criterion for supporting the feasibility of seaweed cultivation. The weight determination of each criterion was based on the literature research and later on was used to weight the variables by priority. The variable that is very influential for the growth of seaweed got a high score. The determination was based on the defined suitability matrix as shown in Table 2. The total score multiplication and weight in table 2 were used to determine the class of water suitability of seaweed cultivation with quantitative analysis using the following equation :

$$Y = \sum ai . Xn$$

The range of suitability cultivation area of *K. Alvarezii* was adjusted to use the following equation :

$$I = \frac{(\sum ai . Xn) - (\sum ai . Xn)_{min}}{k}$$

Note: Y = The final value, ai = weight per parameter, and X_n = score of suitability level, I = interval of suitability class, k = number of suitability class.

Finally, the suitability interval was divided into 5 classes, i.e. 1) highly suitable (HS) is in the range of the value 42-50, 2) suitable (S) is in the range of 33-41; C) moderately suitable (MS) is in the range of 24-32; 4) less suitable (LS) are in the range of 15-23; and 5) not suitable (N) had the value of <15. Each parameter in Table 2 was transformed in digital format using a spline interpolation method with a cell size of 5 x 5 meters resulting in an index map of water suitability.

Distance Suitability Analysis

Distance suitability analysis was also carried out as the other limiting factors out of the water parameters. Those were the distance to shipping lanes or sea lanes and the distance to a settlement which are arranged in a digital format of 5 x 5-meter cell-sized using the buffer and the euclidean distance method to consider the accessibility and disruption of marine traffic. These criteria are described in Table 3. The suitability of distance to the settlement was compiled using the data of inhabited island, namely Bontosua, Sanane, Panjenekang, Balanglombo, and Balangcaddi Islands, while the shipping lanes data used was inter-regional shipping routes. The results of the

analysis were integrated with a water suitability index map.

Spatial Modelling

The spatial modelling in this study used cell-based modelling by integrating all oceanographic and water quality parameters based on the suitability classification according to Table 2 with the following modelling algorithms develop from Danish Hydraulic Institute (2010):

$$WS = (T_i \times 0.5) + (pH_i \times 0.5) + (S_i \times 1) + (DO_i \times 0.5) + (N_i \times 1.5) + (F_i \times 1.5) + (B_i \times 0.5) + (HS_i \times 1.5) + (CS_i \times 1.5) + (TSS_i \times 0.5) + (SB_i \times 0.5)$$

Note: WS= Water suitability, T_i = temperature index map, pH_i = pH index map S_i = salinity index map, DO_i = Dissolved Oxygen index map, N_i = Nitrate index map, F_i = phosphate index map, B_i = bathymetry index map, HS_i = significant wave height index map, CS_i = current speed index map, TSS_i = TSS index map, SB_i = seabed index map

Table 2. Table of suitability criteria for seaweed *Kappaphycus alvarezii* cultivation

Variables	Level of conformity (score)					Weight	Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)		
SST (°C)	28-30	24-27 or 31-32	22-23 or >32	21 or >34	<21 or >36	0.5	Modification of Noor (2015); Rahmayanti et al. (2018)
pH	7-8	6.5-7 or 8-8.5	6-6.4 or >8.5	4-6	<4 or >9	0.5	Modification of Kautsari and Ahdiansyah (2015); Noor (2015)
Salinity (psu)	28-31	26-28 or 31-33	<26 or ≥ 34	18-25 or 35-36	<18 or >37	1	Modification of Kautsari and Ahdiansyah (2015); Rahmayanti et al. (2018); Utama and Handayani (2018)
DO (mg.L ⁻¹)	>6	5-6	4-5	2-4	<2	0.5	Modification of Afandi and Musadat (2018)
Nitrate (mg.L ⁻¹)	0.04-0.07	0.02-0.03	0.01 or >0.1	<0.01 or >0.5	>1	1.5	Modification of Kautsari and Ahdiansyah (2015); Afandi and Musadat (2018)
Phosphate (mg.L ⁻¹)	0.051-0.1	0.02-0.04	0.01-<0.02 or >0.2-1	<0.01 or >1	>2	1.5	Modification of Patty et al. (2015)
Depth (m)	3-10	2-3 or 11-13	1-2	>13-15	<1 or >15	0.5	Modification of Tiensongrusmee (1990) in Radiarta et al. (2012); Kautsari and Ahdiansyah (2015); Yulianto et al. (2017)
Wave height (m)	0.2-0.3	0.1-0.19	>0.3	<0.1 or >0.4	>0.5	1.5	Modification of Rahmayanti et al. (2018)
Current speed (m.s ⁻¹)	0.25-0.3	0.2-0.24 or 0.31-0.4	0.1-0.19 or >0.4	<0.1 or >0.45	>0.5	1.5	Modification of Utama and Handayani (2018)
TSS (mg.L ⁻¹)	<25	25-50	50-100	100-400	>400	0.5	Modification of Yulianto et al. (2017)
Seabed	Coral, Rubble	Sand	Reefs	Muddy Sand	Mud	0.5	Modification of Noor (2015); Yulianto et al. (2017)

The result of this modeling is continued by determining the classification of water suitability based on a class of suitability presented in Table 2. Then, integrating the results of water suitability above with the result of distance suitability analysis was conducted according to Table 3 using cell statistics analysis.

Estimating Productivity

After obtaining the distribution of classes areas for all variable, the seaweed production is estimated only by considering the area which is suitable or highly suitable. Longline method of seaweed cultivation unit was designed as 25 × 50 meter per unit, the distance between the rope was 1 meter and the planting distance of each tied seaweed was 25 cm (WWF, 2014). The initial weight of the seedlings was approximately 100 grams per bunch (Parakkasi et al., 2020) with the wet weight crop was 500 grams per tie. The cultivation of seaweed with a floating

Commented [A3]: Do not use secondary reference

Note : HS: Highly suitable, S: suitable, MS: moderately suitable, LS: less suitable, and N: not suitable

rope/longline system produces an excellent product and can gain a weight up to four to five times from the initial weight of the seedlings (100 g) in a period of cultivation time. The number of the allocated cultivation units was estimated to be 5 units.ha⁻¹. The distance correction between units is 10 m in width (twice longer than the average length of the fishing boat) for fishing traffic. This amount represents 62.5% of the maximum number of units i.e., 8 units.ha⁻¹. To estimated production is the multiplication between the number of cultivation units (from the total suitable area) and once crop yields per unit with the following equations:

$$Ye = N \times Cy$$

Note: Ye= estimated yield per cycle (tons), N= number of cultivation unit, Cy= crop yield per cycle (tons). Briefly, the research method is summarized in the flowchart presented in Figure 2.

Result and Discussion

Distribution of water quality parameters

Spatial distributions of the suitability of water quality parameters are shown in Figure 3. It can be seen that the water quality conditions based on the

variable of SST, and pH, are highly suitable for seaweed cultivation in the study area. SST is the important limiting factor for marine organisms (Wirasatriya *et al.*, 2019). The range of measured SST in the study area is very narrow i.e., only around 30°C. According to BSN (2015) the optimal temperature for this activity is 26-32°C, so the temperature in this site is still suitable for seaweed cultivation. The high temperature of the water causes death on seaweed while low water temperatures cause damage to protein and fat membranes and is very influential in the seaweed life cycle. Photosynthesis on seaweed generally lasts at a maximum temperature of 23°C-30°C (Msuya, 2011; Ding *et al.*, 2013). Seawater has an enormous support ability to prevent pH fluctuation. The condition of pH that is very acidic or alkaline will endanger the survival of organisms because it led to metabolic and respiration disorders. The pH range at the study site is 7.26-7.63. Whereas, optimal seaweed growth requires a pH value range of 7.5-8.5 (BSN, 2015; Radiarta *et al.*, 2018). Moreover, Noor (2015) stated that a suitable pH range for *K. alvarezii* cultivation is in normal and disposed alkaline level. Therefore, these values of pH of this area are feasible for seaweed cultivation (Rahman *et al.*, 2019).




Figure 2. Research Flowchart

Table 3. Distance suitability analysis

Parameters	Suitability Level (Score)					Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)	
Shipping lanes (m)	> 500	-	-	-	≤ 500	Ariyanto <i>et al.</i> (2018)

Commented [RV4]: Replace use high resolution

Distance to settlements (km)	4	4-5	5-6	6-7	>7	Modification of Radiarta et al. (2012)
------------------------------	---	-----	-----	-----	----	--




Figure 3. Distribution of physical and chemical parameters: a) SST (Sea surface temperature); b). Depth; c). Salinity; d).Nitrate; e). Phosphate; f). DO (Dissolved Oxygen); g).TSS; h: seabed type; i): pH

The suitability distribution for TSS and substrate type are moderately suitable or more in the study area. A substrate is dominated by reef almost at entire study area which is categorized as moderately suitable. The suitable condition is located around the small islands with the sand as the substrate type. Seaweed will grow optimally when the cultivation site is in coral, rubble, sand-based substrate, or a mixture of all those three. The most optimum substrate type for seaweed cultivation is a hard substrate like the rocky coral then followed by sandy corals (Parakkasi et al., 2020). Waters with a fractional substrate of coral to sand are usually passed by the current that was suitable for the growth of seaweed. However, cultivating seaweed with a longline method is not overly influenced by the substrate type. For TSS, the eastern part of the study area is highly suitable while the western part is moderately suitable. This indicates that the further distance from the mainland the lower TSS is observed. TSS is insoluble sedimentation which increases the turbidity of water (Nurdin et al., 2019). High turbidity due to the suspended material impedes light penetration into

water column and affects seaweed growth. Thus, TSS indicates water pollution as *K. alvarezii* lives optimally in transparent oligotrophic waters (Hayashi et al., 2011). According to Ministry of Environment R.I (1998), the recommended TSS value is <25 mg.L⁻¹, although up to 80 mg.L⁻¹ is allowed for marine cultivation. The TSS levels in the study area range between 3.6-57.6 mg.L⁻¹ which means that the waters are suitable for the growth of *K. alvarezii*. TSS concentration is probably due to the absence of re-suspension due to the non-silt/clay of substrate type or no input from coastal sediment.

For DO and salinity values, the suitability distributions vary from less suitable to highly suitable. Based on the DO criterion, the study area is mostly highly suitable for seaweed cultivation, with only 2 spots in the study area which are less suitable. The water circulation among small islands plays an important role in high DO level. In open water, the low concentration of oxygen in a natural condition is rarely found (Diaz et al., 2013). The optimal dissolved oxygen for seaweed cultivation is in the range of 3-7 mg.L⁻¹ (Rahman et al., 2019). For

Commented [A5]: Rewrite please, so clearly understood

salinity, the less suitable classes are spread in some areas and even the class at the eastern part of Sanane Island is not suitable for seaweed cultivation. The stability of salinity becomes an important factor for seaweed because seaweed is a stenohaline organism that cannot resist the high fluctuation salinity (Reis *et al.*, 2010; Aris *et al.*, 2021). The extreme salinity changes can lead to ice-ice disease. Salinity can affect the osmoregulation process of marine biota including seaweed (Aris *et al.*, 2021). Because *K. alvarezii* can grow optimally in the range of salinity values of 29-34 ppt (Doty, 1987), therefore, most of the area is still feasible for seaweed cultivation.

Nitrate and phosphate concentrations as indicators of water fertility are relatively low in this region. This may be related to the absence of big rivers in the small islands of the study area as the nutrient sources. Nitrate is a limiting factor for seaweed growth and one of the nutrients needed for the growth of seaweed. The result shows that nitrate concentrations in the study area range between 0.001-0.12 mg.L⁻¹ which is relatively low. According to Kautsari and Ahdiansyah (2015), the feasible nitrate concentration for seaweed cultivation ranges from 0.01 mg.L⁻¹ to 0.07 mg.L⁻¹, while the concentration of <0.01 mg.L⁻¹ and > 0.1 mg.L⁻¹ are not suitable for seaweed cultivation. Phosphate can be a limiting factor due to the source limitation. In the study area, phosphate concentration ranges from 0.001 mg.L⁻¹ to 0.062 mg.L⁻¹ which indicates low to insufficient fertility levels. Phosphate concentration for the fertile water should range from 0.051 mg.L⁻¹ to 0.1 mg.L⁻¹. Besides, phosphate concentration also controls the growth (Nursidi *et al.*, 2017) and the carrageenan content of seaweed (Patty *et al.*, 2015).

The depth of the study area ranges from 2 m to 39.2 m. The longline method can be applied at the 1 m depth from the lowest tide or below (WWF, 2014). The sufficient depth provides seaweed to be submerged. Nevertheless, noted that the deeper waters will increase the cost of constructing seaweed cultivation. Thus, the areas with a depth of more than 15 m are categorized as not suitable for seaweed cultivation.

The distribution of physical and chemical parameters at the seaweed site is highly determined by the wave and current. These aspects play a pivotal role in the water movement for algae growth (Gaylord *et al.*, 2007; Kregting *et al.*, 2013; Nursidi *et al.*, 2017). The variability of current speed and wave height is distinguished by four different characteristics of the west monsoon; transition 1; east monsoon; and transition 2 based on the results of 2D hydrodynamic modeling (Figure 4 and Figure

5). During west monsoon and transition 1 (east monsoon and transition 2), strong current speed appears at the northwestern (southern) part of the study area. The strongest current speed occurs during the transition 1 season. The pattern of the surface current in the study area is influenced by the current pattern of breaking waves in the small islands waters and the surrounding shallow area; and the influence of wind and tidal conditions in these waters. Based on the criteria shown in Table 2, the study area is categorized as highly suitable (dark blue color) to less suitable (light blue color). For the wave height, almost entire study areas are categorized as suitable and highly suitable for seaweed cultivation except during east monsoon. During west monsoon transition 1 and transition 2, the blue and dark blue colors mean that the wave height ranges from 0.1 m to 0.3 m. During east monsoon, the wave higher than 0.3 m occurs at almost the entire study area which makes it less suitable for seaweed cultivation.

The protected area is determined by several types of physical conditions such as current velocity, and wave height. The influence of strong current and ocean waves will drag and damage the cultivated seaweed and its system. The maximum value of the distribution of current speeds and wave height for 4 seasons is described in Figure 6. Based on the results in Figure 6, the current speed and wave height around the study of the water are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. The less suitable areas for the current speed (wave height) category appear mostly in the northeastern (northern) part of the study area. There is no area with a current speed of more than 0.5 m.s⁻¹ and the wave height more than 0.5 m which makes the area is not suitable for seaweed cultivation. The absence of strong current and wave may correspond to the morphology of the study area which is categorized as coral waters with a lot of shallow/reef flat area located within the small islands that protect the area from the high wave and strong current. As mentioned by Sulma *et al.* (2008), the types of protected waters are bays, caps, straits, lagoons, and reef flat. Furthermore, Helmi *et al.* (2017) also stated that the coral reef is a shoreline protector.

Potential areas for seaweed culture development and production

Various suitability criteria as limiting factor of site selection for seaweed cultivation include physical and chemical parameters and distance suitability for accessibility and restricted area inside the shipping lanes has been determined into 5 highly suitable classes (HS), suitable (S), moderately suitable (MS), less suitable (LS) and not suitable (N).

Commented [A6]: What rivers?




Figure 4. Distribution of current speed ($m \cdot s^{-1}$) based on 2D hydrodynamic model in each season




Figure 5. Distribution of significant wave height (m) based on 2D hydrodynamic model in each season




Figure 6. Maximum statistical value of current speed (left) and wave height (right) for all season




Figure 7. Suitability distribution for cultivation area of *K. alvarezii*




Figure 8. Potential area for *K. alvarezii* cultivation in each

Figure 7 shows that most of the study areas are moderately suitable and suitable for seaweed cultivation. The area which is not suitable is only

located along the shipping lane. Figure 8 summarizes the area size of each criteria as follows: the highly suitable (HS) is 83.2 ha (0,42%); suitable

(S) is 4,463.08 ha (22.21%); moderately suitable (MS) is 9,983.52 ha (49.68%); less suitable (LS) is an area of 494.84 ha (2.46%); and not suitable (N) is 5,071.15 ha (25.23%). The potential area for seaweed cultivation development reaches 4,546.28 ha as the sum area of highly suitable and suitable criteria. This result is more than obtained by Arifin *et al.* (2014) which only found 1,963.6 ha area suitable and highly suitable for *K. alvarezii* cultivation in the same study area. This indicates that our method can optimally classify the potential area for *K. alvarezii* cultivation. Furthermore, this potential area can be utilized for 22,730 cultivation units with seaweed production reaching about 56,825 tons of wet weight per crop, or within a year (2 times of harvesting) can reach about 113,650 tons. This amount is only about one-third of the total seaweed production in Pangkep Regency in 2018. However, it is important to be noted that, there are some limitations in the present study which should be considered. For example, the present study did not include the longterm climate variability which may also influence the physical and chemical parameters in the study area. As reported by Msuya *et al.* (2014) and Kunarso *et al.* (2019), the climate variability influences the oceanographic condition which then affects fisheries activity such as aquaculture and fish catch. The present study also did not consider the risk analysis which was able to disturb the cultivation process as shown by Castelar *et al.*, (2015). Furthermore, to maintain the carrying capacity of the study area, the potential area that might be utilized for the expansion of *K. alvarezii* cultivation is about 50% (covering 2,273.14 ha) of the total suitable area accordingly. The result of the present study indicates the effectiveness of Geographic Information Systems with geospatial modeling for developing the area of *K. alvarezii* culture which can be applied in the other area.

Conclusions

Physical and chemical parameters as well as the accessibility and restricted area of shipping lanes were used to define the suitability criteria for *K. alvarezii* cultivation. The result is SST and pH are categorized as highly suitable for *K. alvarezii* cultivation in the whole study area. A substrate is dominated by reef type almost at the whole study area which is categorized as moderately suitable. TSS levels in the study area range between 3.6-57.6 mg.L⁻¹ which means that the waters are suitable for the growth of *K. alvarezii*. For DO and salinity, the suitability distributions vary from less suitable to highly suitable. Nitrate and phosphate concentrations mainly are too low which are not suitable for *K. alvarezii* cultivation. The depth at the study area ranges from 2 m to 39.2 m. The area with a depth more than 15 m is categorized as not

suitable for *K. alvarezii* cultivation. The current speed and wave height around the study area are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. After summarizing all criteria according to their weighting score, the suitable for seaweed cultivation at the study area is 4,546.28 ha, which is the combination of the suitable (S) area of 4,463.08 ha and highly suitable criteria (HS) with an area of 83.2 ha. The potential production per crop can reach 56,825 tons per cycle and 113,650 tons per year.

Acknowledgment

The author thanked the Ministry of Research Technology and Higher Education of the Republic of Indonesia for the financial support of research. The author thanked the Center for Coastal Rehabilitation and Disaster Mitigation Studies (CoREM), Diponegoro University for the facilities' support on hydro-oceanographic modeling and geospatial data processing. We also thanked the Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University for the surveys and laboratory analysis of water quality data.

References

- Afandi, A. & Musadat, F. 2018. Suitability analysis for seaweed culture area at the waters of Kamelanta Village and Panjang Island using Geographical Information System. *J. Akuakultura* 2(1): 69-78. <https://doi.org/10.35308/ja.v2i1.789>
- Arbit, N., Omar, S., Soekendarsi, E., Yasir, I., Tresnati, J. & Tuwo, A. 2019. Morphological and genetic analysis of *Gracilaria* sp. cultured in ponds and coastal waters. *IOP Conf. Ser.: Earth Env. Sci.* 370: 012018. <https://doi.org/10.1088/1755-1315/370/1/012018>
- Arifin, T., Bohari, R. & Arlyza, I.S. 2014. Analysis of space suitability based on marine agriculture in the small islands around Makassar. *Forum geography*, 28(1): 91-102. <https://doi.org/10.23917/forgeo.v28i1.441>
- Aris, M., Muchdar, F. & Labenua, R. 2021. Study of seaweed *Kappaphycus alvarezii* explants growth in the different salinity concentrations. *J. Ilmiah Perikanan Kelautan*, 13(1): 97-105. <https://doi.org/10.20473/jipk.v13i1.19842>
- Ariyanto, F.Y., Astor, Y. & Sidqi, M. 2018. Application of water boundary theory for determining site licenses at the coastal area and small islands

- in Indonesia. *Proc. Nat. Seminar of Geomatics 2018*. Bandung, pp. 893-900. <https://doi.org/10.24895/SNG.2018.3-0.1055>
- Azhar, R.M., Wurjanto, A., & Yuanita, N. 2011. Study of coastal protection of submerged breakwater at Tanjung Kait beach. Master Thesis. Master Program of Water Resource Management, Bandung Institute of Technology, Indonesia.
- Bixler, H.J. & Porse, H. 2011. A Decade of Change in the Seaweed Hydrocolloids Industry. *J. Appl. Phycol.* 23: 321-335. <https://doi.org/10.1007/s10811-010-9529-3>
- Castelar, B., de Siqueira, M.F., Sánchez-Tapia, A. & Reis, R.P. 2015. Risk Analysis Using Species Distribution Modelling to Support Public Policies for The Alien Alga *Kappaphycus alvarezii* Aquaculture in Brazil. *Aquacult.* 446: 217-226. <https://doi.org/10.1016/j.aquaculture.2015.05.012>
- Danish Hydraulic Institute (DHI). 2010. MIKE-21 and MIKE-3 Flow Model. Hydrodynamic Module Step-by-Step Training Guide. DHI MIKE, Denmark. 96 p
- Dee, D.P., Uppala, S.M., Simmons A.J., Berrisford, P., Poli, P., Kobayashi S., Andrae U., Balmaseda, M.A., Balsamo G., Bauer, P., Bechtold, P. Beljaars, A.C.M., van de Berg L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen L., Kållberg, P., Köhler M., Matricardi M, McNally, A.P., Monge-Sanz, B.M., Morcrette, J.J., Park, B.K., Peubey C.P. de Rosnay, P., Tavolato, C., Thépaut, J.N., & Vitart, F., 2011. The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Q.J.R. Meteorol. Soc.*, 137(656): 553-597. <https://doi.org/10.1002/qj.828>
- Doty, M.S. 1987. The production and use of *Eucheuma*. in: Doty, M.A., Caddy, J.F., & Santilices, B. (Eds), Case Studies of Seven Commercial Seaweed Resources, FAO Fish, Tech, Pap, 281: 123- 161
- Ding L., Ma, Y., Huang, B., & Chen, S. 2013. Effects of Seawater Salinity and Temperature on Growth and Pigment Contents in *Hypnea cervicornis*. *J. Agardh (Gigartinales, Rhodophyta)*. *BioMed. Res. Int.* 594308: 10p. <https://doi.org/10.1155/2013/594308>
- Diaz, R.J., Eriksson-Hagg, H. & Rosenberg, R. 2013. Managing Ocean Environments in a Changing Climate Sustainability and Economic Perspectives: Chapter 4. Oxygen - Hypoxia, Pages 67-96. <https://doi.org/10.1016/B978-0-12-407668-6.00004-5>
- Gaylord, B., Rosman, J.H., Reed, D. C., Koseff, J.R., Fram, J., Macintyre, S., Arkema, K., McDonald, C., Brzezinski, M. A., Largier, J.L., Monismith, S.G., Raimondi, P.T. & Mardian, B. 2007. Spatial Patterns of Flow and Their Modification within and Around A Giant Kelp Forest. *Limnol. Oceanogr.*, 52: 1838-1852. <https://doi.org/10.4319/lo.2007.52.5.1838>
- Hayashi, L., Santos, A.A., Faria, G.S.M., Nunes, B.G., Souza, M.S., Fonseca, A.L.D., Barreto, P.L.M., Oliveira, E.C. & Bouzon, Z.L. 2011. *Kappaphycus alvarezii* (Rhodophyta, Areschougaceae) Cultivated in Subtropical Waters in Southern Brazil. *J. Appl. Phycol.*, 23: 337-343. <https://doi.org/10.1007/s10811-010-9543-5>
- Helmi, M., Satriadi, A., Suryoputro, A.A.D., Marwoto, J. & Hariyadi, H.S. 2018. Rehabilitation Priority Area Assessment on Death Coral using Cell-Based Modelling Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.*, 9(11): 2949-2961.
- Helmi, M., Purwanto, Atmodjo, W., Subardjo, P. & Aysira, A. 2017. Benthic Diversity Mapping and Analysis Base on Remote Sensing and Seascape Ecology Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.*, 9(10): 227-235.
- Jalil, A.R., Samawi, M.F., Azis, H.Y., Jaya, I., Malik, A., Yunus, I. & Sohopi, M.A.A. 2020. Comparison of physical-chemical conditions for seaweed cultivation in the Spermonde Archipelago, Indonesia. *AAAL Bioflux*, 13(4), 2071-2082.
- Junaidi, M., Lestary, D.P. & Scabra, A.R. 2018. Mariculture Site Selection Based on Environmental Parameters in Tanjung and Gangga Sub-District, North Lombok. In: Proceeding of 1st ICWBB 2018. Mataram University, Mataram, pp. 233-251.
- Kaiser, M.J., Yu, Y. & Snyder, B. 2010. Economic Feasibility of Using Offshore Oil and Gas Structures in the Gulf of Mexico for Platform-based Aquaculture. *Mar. Pol.*, 34: 699-707. <https://doi.org/10.1016/j.marpol.2010.01.002>

- Kautsari, N. & Ahdiansyah, Y. 2015. Carrying capacity and suitability assessment of Labuhan Terata waters, Sumbawa for the development of seaweed culture. *Ilmu Kelautan: Indonesian Journal of Marine Science*, 20(4): 233-238. <https://doi.org/10.14710/ik.ijms.20.4.23328>
- Kim, J.K., Yarish, C., Hwang, E.K., Park, M. & Kim, Y. 2017. Seaweed Aquaculture: Cultivation Technologies, Challenges, and Its Ecosystem Services. *Algae*, 32(1): 1-13. <https://doi.org/10.4490/algae.2017.32.3.3>
- Kregting, L.T., Blight, A.B., Elsässer, B. & Savidge, G. 2013. The Influence of Water Motion on The Growth Rate of The Kelp *Laminaria hyperborea*. *J. Exp. Mar. Biol. Ecol.*, 448: 337-345. <https://doi.org/10.1016/j.jembe.2013.07.017>
- Kunarno, Wirasatriya, A., Irwani, Satriadi, A., Helmi, M., Prayogi, H. & Munandar, B. 2019. Impact of Climate Variability on Aquatic Productivity and Fisheries Resources in Jepara. *IOP Conf. Ser.: Earth Env. Sci.*, 246: 012021. <https://doi.org/10.1088/1755-1315/246/1/012021>
- Longdill, P.C., Healy, T.R. & Black, K.P. 2008. An integrated GIS approach for sustainable aquaculture management area site selection. *Ocean Coast. Manag.*, 51: 612-624. <https://doi.org/10.1016/j.ocecoaman.2008.06.010>
- Malczewski, J. 2006. Ordered weighted averaging with fuzzy quantifiers: GIS-based multicriteria evaluation for land-use suitability analysis. *Int. J. Appl. Earth Observ. Geoinf.*, 8(4): 270-277. <https://doi.org/10.1016/j.jag.2006.01.003>
- Malczewski, J. & Rinner, C. 2015. Multicriteria Decision Analysis in Geographic Information Science. Springer, Berlin. <https://doi.org/10.1007/978-3-540-74757-4>
- Micael, J., Costa, A.C., Aguiar, P., Medeiros, A. & Calado, H. 2015. Geographic Information System in A Multi-criteria tool for Mariculture Site Selection. *Coast. Manag.*, 43(1): 52-66. <https://doi.org/10.1080/08920753.2014.985178>
- Ministry of Environment R.I, 1988. Decree of the State Minister for Population and Environment Number 02 of 1988 concerning Guidelines for Determining Environmental Quality Standards for Marine Biota (Aquaculture). Jakarta, 31 pp.
- Msuya, F.E., Buriyo, A., Omar, I., Pascal, B., Narrain, K., Ravina, J.J.M., Mrabu, E. & Wakibia, J.G. 2014. Cultivation and Utilisation of Red Seaweeds in the Western Indian Ocean (WIO) Region. *J. Appl. Phycol.*, 26: 699-705.
- Msuya, F.E. 2011. Environmental Changes and Their Impact on Seaweed Farming in Tanzania. *World Aquacult.*, 42: 34-37. <https://doi.org/10.1007/s10811-013-0086-4>
- National Standardization Agency (BSN), 2015. Good Aquaculture Practices (CBIB). 2: Seaweed. SNI 8228.2.:2015. National Standardization Agency
- Nirmala, K., Ratnasari, A. & Budiman, S. 2014. Site Selection for Seaweed Culture at Gerupuk Bay-West Nusa Tenggara Using Remote Sensing and GIS. *J. Akuakultur Indonesia*, 13(1): 73-82. <https://doi.org/10.19027/jai.13.73-82>
- Noor, N.M. 2015 The Water Bodies Compatibility Analysis for Culturing Brown Seaweed *Kappapycus alvarezii* in Ketapang Seashore, South Lampung. *Maspari J.*, 7(2): 91-100.
- Nurdin, N., Helmi, M., Setiawan, R.Y., Zainudin, M., Rustam, A. & Akbar, M. 2019. Landsat Capability to Mapping of Total Solid Suspended (TSS) and Coastal Ecosystem. *Int. Arch. Photogram., Remote Sens. Spatial Inf. Sci.*, 42(4): 497-502. <https://doi.org/10.5194/isprs-archives-XLII-4-W16-497-2019>
- Nursidi, Ali S.A., Anshary H. & Tahya A.M., 2017 Environmental parameters and specific growth of *Kappaphycus alvarezii* in Saugi Island, South Sulawesi Province, Indonesia. *AAAL Bioflux*, 10(4): 698-702.
- Parakkasi, P., Rani, C., Syam, R., Zainuddin & Achmad, M. 2020. Growth Response and Quality of Seaweed *Kappaphycus alvarezii* Cultivated in Various Coastal Ecosystems in the Waters of West Sulawesi, Indonesia. *AAAL Bioflux*, 13(2): 627-639.
- Patty, S.J., Arfah, H. & Abdul, M.S. 2015. The relationship among Nutrient (Phosphate and Nitrate), DO, and pH with the water fertility in Jikumerasa waters, Buru Island. *J. Pesisir dan Laut Tropis*, 1(1): 43-50. <https://doi.org/10.35800/jplt.3.1.2015.9578>
- Pickering, T.D., Skelton, P. & Sulu, J.R. 2007. Intentional Introductions of Commercially Harvested Alien Seaweeds. *Bot. Mar.* 50: 338-350. <https://doi.org/10.1515/BOT.2007.039>

- Radiarta, I.N., Erlania & Haryadi, J., 2018. Analysis of land suitability and carrying capacity of waters for the development of seaweed cultivation in Simeulue Regency; Aceh. *J. Segara*, 14(1): 11-22. <https://doi.org/10.15578/segara.v14i1.6626>
- Radiarta, I.N., Saitoh, S. & Yasui, H. 2011. Aquaculture Site Selection for Japanese Kelp (*Laminaria japonica*) in Southern Hokkaido, Japan, Using Satellite Remote Sensing and GIS-based Models. *ICES J. Mar. Sci.*, 68(4): 773-780. <https://doi.org/10.1093/icesjms/fsq163>
- Radiarta, I.N., Saputra, A. & Albasri, H. 2012. Feasibility mapping of seaweed culture area (*Kappaphycus alvarezii*) at Bintan Regency, Riau Islands Province using GIS and remote sensing approach. *Akuakultur*, 7(1): 145-157. <https://doi.org/10.15578/jra.7.1.2012.145-157>
- Rahman, I.S., Fadjar, M. & Tjahjono, A. 2019. The Relationship Water Physical-Chemical Parameters in Seaweed Cultivation *Euचेuma cottonii* with Long-line System. *Int. J. Sci. Tech. Res.*, 8(11): 1448-1452.
- Rahmayanti, F., Diana, F. & Kusumawati, I. 2018. Suitability analysis for area development of seaweed (*Euचेuma cottonii*) culture in Lhok Bubon waters, Samatiga District, Aceh Barat Regency. *J. Akuakultura*, 2(1): 26-32. <https://doi.org/10.35308/ja.v2i1.773>
- Reis, R.P., Loureiro, R.R. & Mesquita, F.S. 2010. Do Salinity Affect Growth and Carrageenan Yield of *Kappaphycus alvarezii* (Gigartinales/Rhodophyta)? *Aquacult. Res.*, 42: 1231-1234. <https://doi.org/10.1111/j.1365-2109.2010.02699.x>
- Rusliani. 2016. Study of water quality on the growth of seaweed *Gracilaria verrucosa* in a pond with verticulture method. *Octopus J. Fish. Sci.*, 5(2): 522-527.
- Setiawan, R.Y. & Kawamura, H. 2011. Summertime Phytoplankton Bloom in the south Sulawesi Sea. *IEEE J. Sel. Topics Appl. Earth Observ. Remote Sens.* 4(1): 241-244. <https://doi.org/10.1109/JSTARS.2010.2094604>.
- Sousa, F.E.S., Moura, E.A., Marinho-Soriano, E. 2012. Use of Geographic Information Systems (GIS) to Identify Adequate Sites for Cultivation of The Seaweed *Gracilaria birdie* in Rio Grande Do Norte, Northeastern Brazil. *Revista Brasileira de Farmacognosia* 22(4): 868-873.
- <https://doi.org/10.1590/S0102-695X2012005000087>
- Sulma, S., Manoppo, A.K.S. & Hartuti, M. 2008. Identification of Suitable Area for Seaweed Culture in Bali Waters Based on Remote Sensing Satellite Data. *Int. J. Remote Sens. Earth Sci.*, 5: 57-66. <https://doi.org/10.30536/ijreses.2008.v5.a1229>
- Susanto, R.D., Wei, Z., Adi, T.R., Zheng, Q., Fang, G., Bin, F., Supangat, A., Agustyadi, T., Li, S., Trenggono, M. & Setiawan, A. 2016. Oceanography Surrounding Krakatau Volcano in the Sunda Strait, Indonesia. *Oceanogr.* 29(2): 228-237. <https://doi.org/10.5670/oceanog.2016.31>.
- Teniwut, W.A., Marimin & Djatna, T. 2019. GIS-Based multi-criteria decision-making model for site selection of seaweed farming information center: A lesson from small islands, Indonesia. *Decis. Sci. Let.*, 8: 137-150 <https://doi.org/10.5267/j.dsl.2018.8.001>
- Utama, R.Z. & Handayani, T. 2018. Suitability Between Oceanography and Seaweed (*Euचेuma cottonii*) Cultivation Potential in Tidung Island with Geographic Information System (GIS). *ICENIS: E3S Web of Conference*, 73: p.12009. <https://doi.org/10.1051/e3sc onf/20187312009>
- Valderrama, D., Cai, J., Hishamunda, N., Ridler, N., Neish, I. C., Hurtado, A. Q., Msuya, F.E., Krishnan, M., Narayanakumar, R., Kronen, M., Robledo, D., Gasca-Leyva, E. & Fraga, J. 2015. The Economics of *Kappaphycus* Seaweed Cultivation in Developing Countries: A Comparative Analysis of Farming Systems. *Aqua. Econ. Manag.*, 19: 251-277. <https://doi.org/10.1080/13657305.2015.1024348>
- Wirasatriya, A., Sugianto, D.N., Helmi, M., Setiawan, R.Y. & Koch, M. 2019. Distinct Characteristics of SST Variabilities in the Sulawesi Sea and the Northern Part of the Maluku Sea during the Southeast Monsoon. *IEEE J. Sel. Topics Appl. Earth Obs. Remote Sens.*, 12(6): 1763-1770.
- WWF-Indonesia. 2014. Better Management Practices: Guideline series of small scale fisheries of seaweed culture - Cottoni (*Kappaphycus alvarezii*), Sacol (*Kappaphycus striatum*) dan Spinosum (*Euचेuma denticulatum*) Version 1. WWF-Indonesia. South Jakarta. 53 p.

Sarjito Sarjito <sarjito@live.undip.ac.id>
To: Ilmu Kelautan <ijms.undip@gmail.com>

Sat, Sep 10, 2022 at 2:28 PM

Dear Editor in chief IJMS,

Thank you very much for giving us the opportunity to publish our article no 47735 RV with entitle : Identification of Potential Location for *Kapphyscus alverazi* Optimization of Seaweed Production Based on Geographic Information Systems In Spermonde Archipelago Waters, South Sulawesi,Indonesia,

We send the manuscript revised and proof of publication fee transfer .

Thank you for your kindness

Sincerely

Sarjito Sarjito

[Quoted text hidden]

2 attachments



47735-152785-2-ED proof Read revised 2.docx
1881K



Proof of Transfer publication fee.pdf
38K

Identification of Potential Locations for *Kappaphycus alvarezii* Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia

Sarjito Sarjito^{1*}, Hanny Ammaria¹, Muhammad Helmi^{2,3}, Slamet Budi Prayitno¹, Nurjannah Nurdin^{4,5}, Riza Yuliratno Setiawan⁶, Parichat Wetchayont⁷, Anindya Wirasatriya^{2,3}

¹Aquaculture Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro

²Oceanography Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro

³Center for Coastal Rehabilitation and Disaster Mitigation Studies

Jl. Prof.H. Soedarto, S.H. Tembalang, Semarang, Central Java, 50275 Indonesia

⁴Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University

⁵Marine Science and Fisheries Faculty, Hasanuddin University

Jl. Perintis Kemerdekaan Km.10 Tamalanrea, Makassar, Sulawesi Selatan Indonesia

⁶Marine Science and Fisheries Faculty, Gadjah Mada University

Bulaksumur, Caturtunggal, Depok, Sleman, Daerah Istimewa Yogyakarta 55281 Indonesia

⁷Department of Geography, Srinakharinwirot University

Sukhumvit23, Wattana,, Bangkok, 10110, Thailand

Email: sarjito@live.undip.ac.id

Abstract

Indonesia is the second largest producer of red algae in the world with one of the major *Kappaphycus alvarezii* farming areas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province. *K. alvarezii* production in Pangkep Regency increased every year. However, the availability of natural seaweed can be threatened along with the increasing demand so that seaweed cultivation is necessary. Proper site selection is crucial to guarantee the success of the seaweed farming development. A Geographic Information system (GIS) with geospatial modeling approach was applied for identification of the site selection for *K. alvarezii* cultivation at the Liukang Tuppabiring District, Spermonde Islands by combining various physical and chemical parameters obtained from in-situ data and hydrodynamics modeling. The parameters are Sea Surface Temperature, salinity, pH, substrate type, current speed, wave height, DO, nitrate, phosphate, and Total Suspended Solid. This research also include the accessibility and the restricted area of shipping lanes for determining the area suitability of *K. alvarezii* cultivation. This research demonstrates a Geographic Information system with the cell-based geospatial modeling can be effectively used and found the the suitable categorized area for *K. alvarezii* is 4,546.28 ha, which is the combination of the suitable area of 4,463.08 ha and highly suitable criteria with an area of 83.2 ha. The potential productivity of wet-weight seaweed can reach 56,825 tons per cycle or 113,650 tons per year. This result indicates the GIS with geospatial modeling approach for developing the area of *K. alvarezii* culture which can be applied in the other area.

Keywords: Seaweed production, site selection, geospatial, hydrodynamics modeling

Introduction

Indonesia is the largest producer of red algae in the world (83% of global seaweed production) followed by the Philippines and several countries (Valderrama et al., 2015; Kim et al., 2017). Seaweed production in Indonesia comes not only from mariculture, and brackish water ponds but also from natural collections. However, the availability of natural seaweed can be threatened in line with their increasing demand so that seaweed cultivation is completely necessary (Pickering et al., 2007). The increasing demand for seaweed was mostly driven by new markets in China, Eastern Europe, Brazil, and

others (Bixler and Porse, 2011). Furthermore, Arbit et al. (2019) stated that 99.73% of Indonesia's seaweed production comes from cultivation. Therefore, seaweed farming is the best choice for fulfilling market demand.

Site selection is essential in aquaculture development which depends on the species of commodities and applied cultivation methods (Kaiser et al., 2010; Junaidi et al., 2018). For seaweed, the success of cultivation is intrinsically linked to physical and chemical parameters of seaweed farming areas such as turbidity, temperature, salinity, pH, DO, nitrate and phosphate concentration (Rusliani, 2016;

Commented [A1]: Make this abstract to 250 words

Commented [W2R1]: Done

Jalil *et al.*, 2020). Hydrodynamic aspects should also be considered in the site selection of *K. alvarezii* farming since the cultivation method uses the longline method which is prone to strong current and wave. One of major *K. alvarezii*'s farming areas within the Indonesian seas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province (Jalil *et al.*, 2020). The existence of Indonesian Troughflow along the Makassar Strait (Susanto *et al.*, 2016) and the occurrence of strong upwelling at the southern part of the South Sulawesi during each summer (Setiawan and Kawamura 2011) may influence the hydrodynamics condition in the Spermonde Island. Besides, the existence of small islands among the water area makes the hydrodynamics condition at the Spermonde Islands becomes more complex. Spermonde Islands area also has dense shipping activities. Many shipping lanes have lied among these areas which are should be avoided in the mariculture site selection. This shows the complex considerations for conducting site selection of seaweed farming at Spermonde Islands.

In dealing with the complex considerations as mentioned above, the use of the geospatial model which is part of Geographical Information Systems (GIS) becomes an effective solution for determining the proper mariculture site selection (Longdill *et al.*, 2008; Micael *et al.*, 2015). This approach, in general, is an analysis of layers that contains spatial data to evaluate and select a particular area based on its suitability criteria classified according to certain environmental aspects (Malczewski, 2006; Malczewski & Rinner, 2015). GIS utilization in mariculture development has been widely used in various types of commodities including seaweed (e.g., Radiarta *et al.*, 2011; Radiarta *et al.*, 2012; Sousa *et al.*, 2012; Nirmala *et al.*, 2014; Teniwut *et al.*, 2019) which were mainly used in-situ data and remote sensing data for the input of geospatial modeling. The application of GIS for site selection for aquaculture in the Spermonde Islands has conducted by Arifin *et al.* (2014). However, their analysis was only based on in-situ measurement which cannot represent the temporal change of ocean parameters. In the present study, cell-based geospatial modeling was performed for the determination of sustainable site selection of seaweed cultivation, which integrates various physical and chemical parameters obtained from in-situ data and the hydrodynamic model. The combination of both data becomes the new approach proposed in the present study. Considering that the hydrodynamics aspects for the site selection of seaweed culture is not available in the remote sensing data, a 2D hydrodynamic model was built to represent the hydrodynamic condition in the study area. On the other hand, Indonesia as the maritime continent experiences four-season influenced by monsoon wind (Helmi *et al.*, 2018) which impacts the

hydrodynamics conditions. Thus, in-situ and incidental measurements of current and wave cannot represent the hydrodynamics condition in the study area. This issue only can be resolved by performing the hydrodynamics model. Furthermore, the potential productivity of *K. alvarezii* at the selected site is also simulated. This research is expected to become a prototype of a geospatial model to explore the potential aquaculture area in small islands and can be a scientific-based information for local governments, researchers, NGOs, and related stakeholders for planning and developing *K. alvarezii* cultivation in the study area.

Material and Method

The study area is presented in Figure 1. which is located at the Spermonde islands waters precisely in the Liukang Tupabbiring District, Pangkep Regency, South Sulawesi. The Liukang Tuppabiring District area is 54.44 km² (4.89% of Pangkep Regency) and consists of 20 islands. The population of this district reached 29,680 inhabitants, generally fishermen and sea farmers. The field stations are located at 4°5'3.98"S - 4°59'51.137"S and 119°18'35.532"E - 119°26'19.849"E. There are 94 sampling points distributed among the island of Bontosua, Sanane, Panjenekang, Pankaiya, Langkadea, Balanglombo, Balangcaddi Island, and reef flat area.

A field survey was conducted for collecting water quality data at 94 points in which the depths are more than 2 meters. Survey site determination was using Garmin 76C GPS. The water quality data collected are presented in Table 1. The in-situ measurements were Sea Surface Temperature, (SST), salinity, pH, and substrate type. For the chemical parameters, such as DO, nitrate, phosphate, and Total Suspended Solid (TSS), the water samples were taken and analyzed at Chemical Oceanography Laboratory, Marine Science Department, Hasanuddin University, Indonesia (May 2019). The dynamic and non-dynamic water parameter data were interpolated to obtain a spatial distribution pattern at the time of data retrieval. The data were compiled in a digital form of 5 x 5 meters cell size using spline and classified according to suitability class in Table 2.

Two dimensions (2D) hydrodynamic model was used to simulate current and wave condition at the study area. The data for generating a hydrodynamic model are as follows: (1) High-resolution satellite imagery GeoEye-1 from ESRI online (<https://esriindonesia.co.id/arcgis-online>); (2) Bathymetry data from Indonesia Geospatial Agency (<http://batnas.big.go.id/>); (3) Surface wind and wave height data obtained from 6 hourly ERA-interim data

with a grid interval of 0.125° from July 2018 to July 2019 (Dee et al., 2011); (4) Tidal elevation forecast




Figure 1. Study area

Table 1. Water quality data collection

No	Variables	Unit	Tools/Methods	Data Retrieval Methods
1	Sea Surface Temperature (SST)	°C	Thermometer	In situ
2	Salinity	Psu	Salinometer	In situ
3	pH	-	Water quality checker	In situ
4	Dissolved oxygen (DO)	ppm	Winkler method	Laboratory analysis
5	Nitrate	mg.L ⁻¹	Ascorbic Acid Method	Laboratory analysis
6	Phosphate	mg.L ⁻¹	Brucine Sulfate Method	Laboratory analysis
7	Total Suspended Solids (TSS)	mg.L ⁻¹	Milipore filter	Laboratory analysis
8	Substrate type	-	Visually observation	In situ
9	Depth	m	Modelled depth from LPI Maps (BIG, 2011) and BATNAS (BIG, 2018) data	Integrating modelled data

from July 2018 to July 2019 (<https://www.dhigroup.com/download/mike-by-dhi-tools/coastandseetools/global-tide-model>). The current model was constructed by continuity equations and momentum equations with the average of depth. On-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry, wind and tidal dynamics were used as an input for generating current model. The initial process was making a mesh, and setting the boundary

condition of the model using the mathematical continuity equation as follows:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = S$$

The horizontal momentum equation for component x is as follows (DHI, 2010) :

$$\frac{\partial u}{\partial t} + \frac{\partial u^2}{\partial x} + \frac{\partial uv}{\partial y} + \frac{\partial wu}{\partial z} = f_v \cdot g \frac{\partial h}{\partial x} - \frac{1}{r_0} \frac{\partial p_a}{\partial x} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial x} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{xx}}{\partial x} + \frac{\partial S_{xy}}{\partial y} \right) + F_u + \frac{\partial}{\partial z} \left(v_t \frac{\partial u}{\partial z} \right) + u_s S$$

While the horizontal momentum equation for component y is as follows (DHI, 2010):

$$\frac{\partial v}{\partial t} + \frac{\partial v^2}{\partial y} + \frac{\partial uv}{\partial x} + \frac{\partial wv}{\partial z} = f_u \cdot g \frac{\partial h}{\partial y} - \frac{1}{r_0} \frac{\partial p_a}{\partial y} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial y} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{yx}}{\partial x} + \frac{\partial S_{yy}}{\partial y} \right) + F_v + \frac{\partial}{\partial z} \left(v_t \frac{\partial v}{\partial z} \right) + v_s S$$

Note: x, y, z = cartesian coordinates of x, y, and z direction; t = time; u = water particle speed of direction x; v = water particle speed of direction y; w = water particle speed of direction z; η = water surface elevation; d = sea depth; h = total depth of the sea; f = coriolis parameter (f = 2ω sinθ); Ω = ratio of earth revolution; θ = latitude coordinates; g = acceleration of gravity; S_{xx}, S_{xy}, S_{yx}, S_{yy} = component of radiation stress tensor; v_t = vertical eddy viscosity; p_a = atmospheric pressure; ρ₀ = sea density; S = debits due to sources; u_s = particle velocity of air sources direction x; and v_s = the particle speed of air sources direction y.

Wave modeling used directional decoupled parametric and fully spectral formulations based on wave eternity movements with the input data of on-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry data, wind speed and direction and wave data. Parameterization was created in the frequency domain by introducing the 0 and 1st moments of the wave spectrum as a variable (Azhar *et al.*, 2011). The arranger equation was the equilibrium wave force in cartesian coordinate formulated as follow (DHI, 2010):

$$\frac{\partial N}{\partial t} + \frac{\partial}{\partial \phi} C \phi N + \frac{\partial}{\partial \lambda} C \lambda N + \frac{\partial}{\partial \sigma} C \sigma N + \frac{\partial}{\partial \theta} C \theta N = \frac{S}{\sigma}$$

Source energy, S, shows the superposition of surface functions in various styles (Azhar *et al.*, 2011).

$$S = S_{in} + S_{nl} + S_{ds} + S_{bot} + S_{surf}$$

Note : N_{x,σ,θ,t} = power density, φλ = coordinate spherical, C = speed of propagation of four-dimensional wave group, S = energy source, S_{in} = formation by wind, S_{nl} = energy transfer, non-linear, S_{ds} = wave energy due to white capping, S_{bot} = by bottom friction, S_{surf} = dissipation due to the depth-induced breaking.

The maximum value of the current speed and wave height were taken in each grid that comes from 4 seasonal patterns characteristics, that is the west

monsoon (December), transition 1 (May), east monsoon (July), and transition 2 seasons (October). The results of the 2D hydrodynamic model form the patterns of current speed and wave height every season were interpolated with a spline method with a cell size of 5 x 5 meters and analyzed using cell statistics to obtain the maximum value of current speed and wave height. The results of the distribution map were classified according to suitability class in Table 2.

Determination of Area Suitability Criteria

The determination of the weighting and scoring was done to provide value in each criterion for supporting the feasibility of seaweed cultivation. The weight determination of each criterion was based on the literature research and later on was used to weight the variables by priority. The variable that is very influential for the growth of seaweed got a high score. The determination was based on the defined suitability matrix as shown in Table 2. The total score multiplication and weight in table 2 were used to determine the class of water suitability of seaweed cultivation with quantitative analysis using the following equation:

$$Y = \sum ai \cdot Xn$$

The range of suitability cultivation area of *K. Alvarezii* was adjusted to use the following equation :

$$I = \frac{(\sum ai \cdot Xn) - (\sum ai \cdot Xn)_{min}}{k}$$

Note: Y = The final value, ai = weight per parameter, and X_n = score of suitability level, I = interval of suitability class, k = number of suitability class.

Finally, the suitability interval was divided into 5 classes, i.e. 1) highly suitable (HS) is in the range of the value 42-50, 2) suitable (S) is in the range of 33-41; C) moderately suitable (MS) is in the range of 24-32; 4) less suitable (LS) are in the range of 15-23; and 5) not suitable (N) had the value of <15. Each parameter in Table 2 was transformed in digital format using a spline interpolation method with a cell size of 5 x 5 meters resulting in an index map of water suitability.

Distance Suitability Analysis

Distance suitability analysis was also carried out as the other limiting factors out of the water parameters. Those were the distance to shipping lanes or sea lanes and the distance to a settlement which are arranged in a digital format of 5 x 5 -meter cell-sized using the buffer and the euclidean distance method to consider the accessibility and disruption of

marine traffic. These criteria are described in Table 3. The suitability of distance to the settlement was compiled using the data of inhabited island, namely Bontosua, Sanane, Panjenekang, Balanglombo, and Balangcaddi Islands, while the shipping lanes data used was inter-regional shipping routes. The results of the analysis were integrated with a water suitability index map.

Spatial Modelling

The spatial modelling in this study used cell-based modelling by integrating all oceanographic and water quality parameters based on the suitability classification according to Table 2 with the following modelling algorithms develop from Danish Hydraulic Institute (2010):

$$WS = (T_i \times 0.5) + (pH_i \times 0.5) + (S_i \times 1) + (DO_i \times 0.5) + (N_i \times 1.5) + (F_i \times 1.5) + (B_i \times 0.5) + (HS_i \times 1.5) + (CS_i \times 1.5) + (TSS_i \times 0.5) + (SB_i \times 0.5)$$

Note: WS= Water suitability, T_i = temperature index map, pH_i = pH index map, S_i = salinity index map, DO_i = Dissolved Oxygen index map, N_i = Nitrate index map,

F_i = phosphate index map, B_i = bathymetry index map, HS_i = significant wave height index map, CS_i = current speed index map, TSS_i = TSS index map, SB_i = seabed index map

The result of this modeling is continued by determining the classification of water suitability based on a class of suitability presented in Table 2. Then, integrating the results of water suitability above with the result of distance suitability analysis was conducted according to Table 3 using cell statistics analysis.

Estimating Productivity

After obtaining the distribution of classes areas for all variable, the seaweed production is estimated only by considering the area which is suitable or highly suitable. Longline method of seaweed cultivation unit was designed as 25 × 50 meter per unit, the distance between the rope was 1 meter and the planting distance of each tied seaweed was 25 cm (WWF, 2014). The initial weight of the seedlings was approximately 100 grams per bunch (Parakkasi et al., 2020) with the wet weight crop was 500 grams per tie. The cultivation of seaweed with a floating

Table 2. Table of suitability criteria for seaweed *Kappaphycus alvarezii* cultivation

Variables	Level of conformity (score)					Weight	Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)		
SST (°C)	28-30	24-27 or 31-32	22-23 or >32	21 or >34	<21 or >36	0.5	Modification of Noor (2015); Rahmayanti et al. (2018)
pH	7-8	6.5-7 or 8-8.5	6-6.4 or >8.5	4-6	<4 or >9	0.5	Modification of Kautsari and Ahdiansyah (2015); Noor (2015)
Salinity (psu)	28-31	26-28 or 31-33	<26 or ≥ 34	18-25 or 35-36	<18 or >37	1	Modification of Kautsari and Ahdiansyah (2015); Rahmayanti et al. (2018); Utama and Handayani (2018)
DO (mg.L ⁻¹)	>6	5-6	4-5	2-4	<2	0.5	Modification of Afandi and Musadat (2018)
Nitrate (mg.L ⁻¹)	0.04-0.07	0.02-0.03	0.01 or >0.1	<0.01 or >0.5	>1	1.5	Modification of Kautsari and Ahdiansyah (2015); Afandi and Musadat (2018)
Phosphate (mg.L ⁻¹)	0.051-0.1	0.02-0.04	0.01-0.02 or >0.2-1	<0.01 or >1	>2	1.5	Modification of Patty et al. (2015)
Depth (m)	3-10	2-3 or 11-13	1-2	>13-15	<1 or >15	0.5	Modification of Radiarta et al. (2012); Kautsari and Ahdiansyah (2015); Yulianto et al. (2017)
Wave height (m)	0.2-0.3	0.1-0.19	>0.3	<0.1 or >0.4	>0.5	1.5	Modification of Rahmayanti et al. (2018)
Current speed (m.s ⁻¹)	0.25-0.3	0.2-0.24 or 0.31-0.4	0.1-0.19 or >0.4	<0.1 or >0.45	>0.5	1.5	Modification of Utama and Handayani (2018)
TSS (mg.L ⁻¹)	<25	25-50	50-100	100-400	>400	0.5	Modification of Yulianto et al. (2017)

Commented [A3]: Do not use secondary reference

Commented [W4R3]: Tiensongrusmee (1990) was deleted

Seabed	Coral, Rubble	Sand	Reefs	Muddy Sand	Mud	0.5	Modification of Noor (2015); Yulianto <i>et al.</i> (2017)
--------	---------------	------	-------	------------	-----	-----	--

Note : HS: Highly suitable, S: suitable, MS: moderately suitable, LS: less suitable, and N: not suitable

rope/longline system produces an excellent product and can gain a weight up to four to five times from the initial weight of the seedlings (100 g) in a period of cultivation time. The number of the allocated cultivation units was estimated to be 5 units.ha⁻¹. The distance correction between units is 10 m in width (twice longer than the average length of the fishing boat) for fishing traffic. This amount represents 62.5% of the maximum number of units i.e., 8 units.ha⁻¹. To estimated production is the multiplication between the number of cultivation units (from the total suitable area) and once crop yields per unit with the following equations:

$$Ye = N \times Cy$$

Note: Ye= estimated yield per cycle (tons), N= number of cultivation unit, Cy= crop yield per cycle (tons). Briefly, the research method is summarized in the flowchart presented in Figure 2.

Result and Discussion

Distribution of water quality parameters

Spatial distributions of the suitability of water quality parameters are shown in Figure 3. It can be

seen that the water quality conditions based on the variable of SST, and pH, are highly suitable for seaweed cultivation in the study area. SST is the important limiting factor for marine organisms (Wirasatriya *et al.*, 2019). The range of measured SST in the study area is very narrow *i.e.*, only around 30°C. According to BSN (2015) the optimal temperature for this activity is 26-32°C, so the temperature in this site is still suitable for seaweed cultivation. The high temperature of the water causes death on seaweed while low water temperatures cause damage to protein and fat membranes and is very influential in the seaweed life cycle. Photosynthesis on seaweed generally lasts at a maximum temperature of 23°C-30°C (Msuya, 2011; Ding *et al.*, 2013). Seawater has an enormous support ability to prevent pH fluctuation. The condition of pH that is very acidic or alkaline will endanger the survival of organisms because it led to metabolic and respiration disorders. The pH range at the study site is 7.26-7.63. Whereas, optimal seaweed growth requires a pH value range of 7.5-8.5 (BSN, 2015; Radiarta *et al.*, 2018). Moreover, Noor (2015) stated that a suitable pH range for *K. alvarezii* cultivation is in normal and disposed alkaline level. Therefore, these values of pH of this area are feasible for seaweed cultivation (Rahman *et al.*, 2019).




Figure 2. Research Flowchart

Table 3. Distance suitability analysis

Parameters	Suitability Level (Score)					Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)	
Shipping lanes (m)	> 500	-	-	-	≤ 500	Ariyanto <i>et al.</i> (2018)
Distance to settlements (km)	4	4-5	5-6	6-7	>7	Modification of Radiarta <i>et al.</i> (2012)

Commented [RV5]: Replace use high resolution

Commented [W6R5]: Done




Figure 3. Distribution of physical and chemical parameters: a) SST (Sea surface temperature); b). Depth; c). Salinity; d).Nitrate; e). Phosphate; f). DO (Dissolved Oxygen); g).TSS; h: seabed type; i): pH

The suitability distribution for TSS and substrate type are moderately suitable or more in the study area. A substrate is dominated by reef almost at entire study area which is categorized as moderately suitable. The suitable condition is located around the small islands with the sand as the substrate type. Seaweed will grow optimally when the cultivation site is in coral, rubble, sand-based substrate, or a mixture of all those three. The most optimum substrate type for seaweed cultivation is a hard substrate like the rocky coral then followed by sandy corals (Parakkasi *et al.*, 2020). Waters with a fractional substrate of coral to sand are usually passed by the current that was suitable for the growth of seaweed. However, cultivating seaweed with a longline method is not overly influenced by the substrate type. For TSS, the eastern part of the study area is highly suitable while the western part is moderately suitable. This indicates that the further distance from the mainland the lower TSS is observed. TSS is insoluble sedimentation which increases the turbidity of water (Nurdin *et al.*, 2019). High turbidity due to the suspended material impedes light penetration into water column and affects

seaweed growth. Thus, TSS indicates water pollution as *K. alvarezii* lives optimally in transparent oligotrophic waters (Hayashi *et al.*, 2011). According to Ministry of Environment R.I (1998), the recommended TSS value is $<25 \text{ mg.L}^{-1}$, although up to 80 mg.L^{-1} is allowed for marine cultivation. The TSS levels in the study area range between $3.6\text{-}57.6 \text{ mg.L}^{-1}$ which means that the waters are suitable for the growth of *K. alvarezii*. TSS concentration is probably due to the absence of re-suspension from bottom substrate and no input of the coastal sediment.

For DO and salinity values, the suitability distributions vary from less suitable to highly suitable. Based on the DO criterion, the study area is mostly highly suitable for seaweed cultivation, with only 2 spots in the study area which are less suitable. The water circulation among small islands plays an important role in high DO level. In open water, the low concentration of oxygen in a natural condition is rarely found (Diaz *et al.*, 2013). The optimal dissolved oxygen for seaweed cultivation is in the range of $3\text{-}7 \text{ mg.L}^{-1}$ (Rahman *et al.*, 2019). For salinity, the less suitable classes are spread in some areas and even

Commented [A7]: Rewrite please, so clearly understood

Commented [W8R7]: done

the class at the eastern part of Sanane Island is not suitable for seaweed cultivation. The stability of salinity becomes an important factor for seaweed because seaweed is a stenohaline organism that cannot resist the high fluctuation salinity (Reis *et al.*, 2010; Aris *et al.*, 2021). The extreme salinity changes can lead to ice-ice disease. Salinity can affect the osmoregulation process of marine biota including seaweed (Aris *et al.*, 2021). Because *K. alvarezii* can grow optimally in the range of salinity values of 29-34 ppt (Doty, 1987), therefore, most of the area is still feasible for seaweed cultivation.

Nitrate and phosphate concentrations as indicators of water fertility are relatively low in this region. This may be related to the absence of big rivers in the small islands of the study area as the nutrient sources. Nitrate is a limiting factor for seaweed growth and one of the nutrients needed for the growth of seaweed. The result shows that nitrate concentrations in the study area range between 0.001-0.12 mg.L⁻¹ which is relatively low. According to Kautsari and Ahadiansyah (2015), the feasible nitrate concentration for seaweed cultivation ranges from 0.01 mg.L⁻¹ to 0.07 mg.L⁻¹, while the concentration of <0.01 mg.L⁻¹ and > 0.1 mg.L⁻¹ are not suitable for seaweed cultivation. Phosphate can be a limiting factor due to the source limitation. In the study area, phosphate concentration ranges from 0.001 mg.L⁻¹ to 0.062 mg.L⁻¹ which indicates low to insufficient fertility levels. Phosphate concentration for the fertile water should range from 0.051 mg.L⁻¹ to 0.1 mg.L⁻¹. Besides, phosphate concentration also controls the growth (Nursidi *et al.*, 2017) and the carrageenan content of seaweed (Patty *et al.*, 2015).

The depth of the study area ranges from 2 m to 39.2 m. The longline method can be applied at the 1 m depth from the lowest tide or below (WWF, 2014). The sufficient depth provides seaweed to be submerged. Nevertheless, noted that the deeper waters will increase the cost of constructing seaweed cultivation. Thus, the areas with a depth of more than 15 m are categorized as not suitable for seaweed cultivation.

The distribution of physical and chemical parameters at the seaweed site is highly determined by the wave and current. These aspects play a pivotal role in the water movement for algae growth (Gaylord *et al.*, 2007; Kregting *et al.*, 2013; Nursidi *et al.*, 2017). The variability of current speed and wave height is distinguished by four different characteristics of the west monsoon; transition 1; east monsoon; and transition 2 based on the results of 2D hydrodynamic modeling (Figure 4 and Figure 5). During west monsoon and transition 1 (east monsoon

and transition 2), strong current speed appears at the northwestern (southern) part of the study area. The strongest current speed occurs during the transition 1 season. The pattern of the surface current in the study area is influenced by the current pattern of breaking waves in the small islands waters and the surrounding shallow area; and the influence of wind and tidal conditions in these waters. Based on the criteria shown in Table 2, the study area is categorized as highly suitable (dark blue color) to less suitable (light blue color). For the wave height, almost entire study areas are categorized as suitable and highly suitable for seaweed cultivation except during east monsoon. During west monsoon transition 1 and transition 2, the blue and dark blue colors mean that the wave height ranges from 0.1 m to 0.3 m. During east monsoon, the wave higher than 0.3 m occurs at almost the entire study area which makes it less suitable for seaweed cultivation.

The protected area is determined by several types of physical conditions such as current velocity, and wave height. The influence of strong current and ocean waves will drag and damage the cultivated seaweed and its system. The maximum value of the distribution of current speeds and wave height for 4 seasons is described in Figure 6. Based on the results in Figure 6, the current speed and wave height around the study of the water are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. The less suitable areas for the current speed (wave height) category appear mostly in the northeastern (northern) part of the study area. There is no area with a current speed of more than 0.5 m.s⁻¹ and the wave height more than 0.5 m which makes the area is not suitable for seaweed cultivation. The absence of strong current and wave may correspond to the morphology of the study area which is categorized as coral waters with a lot of shallow/reef flat area located within the small islands that protect the area from the high wave and strong current. As mentioned by Sulma *et al.* (2008), the types of protected waters are bays, caps, straits, lagoons, and reef flat. Furthermore, Helmi *et al.* (2017) also stated that the coral reef is a shoreline protector.

Potential areas for seaweed culture development and production

Various suitability criteria as limiting factor of site selection for seaweed cultivation include physical and chemical parameters and distance suitability for accessibility and restricted area inside the shipping lanes has been determined into 5 highly suitable classes (HS), suitable (S), moderately suitable (MS), less suitable (LS) and not suitable (N).

Commented [A9]: What rivers?

Commented [W10R9]: There is no big river in this area !!!




Figure 4. Distribution of current speed (m.s⁻¹) based on 2D hydrodynamic model in each season




Figure 5. Distribution of significant wave height (m) based on 2D hydrodynamic model in each season




Figure 6. Maximum statistical value of current speed (left) and wave height (right) for all season




Figure 7. Suitability distribution for cultivation area of *K. alvarezii*




Figure 8. Potential area for *K. alvarezii* cultivation in each

Figure 7 shows that most of the study areas are moderately suitable and suitable for seaweed cultivation. The area which is not suitable is only

located along the shipping lane. Figure 8 summarizes the area size of each criteria as follows: the highly suitable (HS) is 83.2 ha (0,42%); suitable (S) is

4,463.08 ha (22.21%); moderately suitable (MS) is 9,983.52 ha (49.68%); less suitable (LS) is an area of 494.84 ha (2.46%); and not suitable (N) is 5,071.15 ha (25.23%). The potential area for seaweed cultivation development reaches 4,546.28 ha as the sum area of highly suitable and suitable criteria. This result is more than obtained by Arifin *et al.* (2014) which only found 1,963.6 ha area suitable and highly suitable for *K. alvarezii* cultivation in the same study area. This indicates that our method can optimally classify the potential area for *K. alvarezii* cultivation. Furthermore, this potential area can be utilized for 22,730 cultivation units with seaweed production reaching about 56,825 tons of wet weight per crop, or within a year (2 times of harvesting) can reach about 113,650 tons. This amount is only about one-third of the total seaweed production in Pangkep Regency in 2018. However, it is important to be noted that, there are some limitations in the present study which should be considered. For example, the present study did not include the longterm climate variability which may also influence the physical and chemical parameters in the study area. As reported by Msuya *et al.* (2014) and Kunarso *et al.* (2019), the climate variability influences the oceanographic condition which then affects fisheries activity such as aquaculture and fish catch. The present study also did not consider the risk analysis which was able to disturb the cultivation process as shown by Castelar *et al.*, (2015). Furthermore, to maintain the carrying capacity of the study area, the potential area that might be utilized for the expansion of *K. alvarezii* cultivation is about 50% (covering 2,273.14 ha) of the total suitable area accordingly. The result of the present study indicates the effectiveness of Geographic Information Systems with geospatial modeling for developing the area of *K. alvarezii* culture which can be applied in the other area.

Conclusions

Physical and chemical parameters as well as the accessibility and restricted area of shipping lanes were used to define the suitability criteria for *K. alvarezii* cultivation. The result is SST and pH are categorized as highly suitable for *K. alvarezii* cultivation in the whole study area. A substrate is dominated by reef type almost at the whole study area which is categorized as moderately suitable. TSS levels in the study area range between 3.6-57.6 mg.L⁻¹ which means that the waters are suitable for the growth of *K. alvarezii*. For DO and salinity, the suitability distributions vary from less suitable to highly suitable. Nitrate and phosphate concentrations mainly are too low which are not suitable for *K. alvarezii* cultivation. The depth at the study area ranges from 2 m to 39.2 m. The area with a depth more than 15 m is categorized as not suitable for *K. alvarezii* cultivation. The current speed and wave

height around the study area are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. After summarizing all criteria according to their weighting score, the suitable for seaweed cultivation at the study area is 4,546.28 ha, which is the combination of the suitable (S) area of 4,463.08 ha and highly suitable criteria (HS) with an area of 83.2 ha. The potential production per crop can reach 56,825 tons per cycle and 113,650 tons per year.

Acknowledgment

The author thanked the Ministry of Research Technology and Higher Education of the Republic of Indonesia for the financial support of research. The author thanked the Center for Coastal Rehabilitation and Disaster Mitigation Studies (CoREM), Diponegoro University for the facilities support on hydro-oceanographic modeling and geospatial data processing. We also thanked the Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University for the surveys and laboratory analysis of water quality data.

References

- Afandi, A. & Musadat, F. 2018. Suitability analysis for seaweed culture area at the waters of Kamelanta Village and Panjang Island using Geographical Information System. *J. Akuakultura* 2(1): 69-78. <https://doi.org/10.35308/ja.v2i1.789>
- Arbit, N., Omar, S., Soekendarsi, E., Yasir, I., Tresnati, J. & Tuwo, A. 2019. Morphological and genetic analysis of *Gracilaria* sp. cultured in ponds and coastal waters. *IOP Conf. Ser.: Earth Env. Sci.* 370: 012018. <https://doi.org/10.1088/1755-1315/370/1/012018>
- Arifin, T., Bohari, R. & Arlyza, I.S. 2014. Analysis of space suitability based on marine agriculture in the small islands around Makassar. *Forum geography*, 28(1): 91-102. <https://doi.org/10.23917/forgeo.v28i1.441>
- Aris, M., Muchdar, F. & Labenua, R. 2021. Study of seaweed *Kappaphycus alvarezii* explants growth in the different salinity concentrations. *J. Ilmiah Perikanan Kelautan*, 13(1): 97-105. <https://doi.org/10.20473/jipk.v13i1.19842>
- Ariyanto, F.Y., Astor, Y. & Sidqi, M. 2018. Application of water boundary theory for determining site licenses at the coastal area and small islands in Indonesia. *Proc. Nat. Seminar of Geomatics*

2018. Bandung, pp. 893-900. <https://doi.org/10.24895/SNG.2018.3.0.1055>
- Azhar, R.M., Wurjanto, A., & Yuanita, N. 2011. Study of coastal protection of submerged breakwater at Tanjung Kait beach. Master Thesis. Master Program of Water Resource Management, Bandung Institute of Technology, Indonesia.
- Bixler, H.J. & Porse, H. 2011. A Decade of Change in the Seaweed Hydrocolloids Industry. *J. Appl. Phycol.* 23: 321-335. <https://doi.org/10.1007/s10811-010-9529-3>
- Castelar, B., de Siqueira, M.F., Sánchez-Tapia, A. & Reis, R.P. 2015. Risk Analysis Using Species Distribution Modelling to Support Public Policies for The Alien Alga *Kappaphycus alvarezii* Aquaculture in Brazil. *Aquacult.* 446: 217-226. <https://doi.org/10.1016/j.aquaculture.2015.05.012>
- Danish Hydraulic Institute (DHI). 2010. MIKE-21 and MIKE-3 Flow Model. Hydrodynamic Module Step-by-Step Training Guide. DHI MIKE, Denmark. 96 p
- Dee, D.P., Uppala, S.M., Simmons A.J., Berrisford, P., Poli, P., Kobayashi S., Andrae U., Balmaseda, M.A., Balsamo G., Bauer, P., Bechtold, P., Beljaars, A.C.M., van de Berg L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen L., Kållberg, P., Köhler M., Matricardi M, McNally, A.P., Monge-Sanz, B.M., Morcrette, J.J., Park, B.K., Peubey C.P. de Rosnay, P., Tavolato, C., Thépaut, J.N., & Vitart, F., 2011. The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Q.J.R. Meteorol. Soc.*, 137(656): 553-597. <https://doi.org/10.1002/qj.828>
- Doty, M.S. 1987. The production and use of *Eucheuma*. in: Doty, M.A., Caddy, J.F., & Santilices, B. (Eds), Case Studies of Seven Commercial Seaweed Resources, FAO Fish, Tech, Pap, 281: 123- 161
- Ding L., Ma, Y., Huang, B., & Chen, S. 2013. Effects of Seawater Salinity and Temperature on Growth and Pigment Contents in *Hypnea cervicornis*. *J. Agardh (Gigartinales, Rhodophyta)*. *BioMed. Res. Int.* 594308: 10p. <https://doi.org/10.1155/2013/594308>
- Diaz, R.J., Eriksson-Hagg, H. & Rosenberg, R. 2013. Managing Ocean Environments in a Changing Climate Sustainability and Economic Perspectives: Chapter 4. Oxygen-Hypoxia, Pages 67-96. <https://doi.org/10.1016/B978-0-12-407668-6.00004-5>
- Gaylord, B., Rosman, J.H., Reed, D. C., Koseff, J.R., Fram, J., Macintyre, S., Arkema, K., McDonald, C., Brzezinski, M. A., Largier, J.L., Monismith, S.G., Raimondi, P.T. & Mardian, B. 2007. Spatial Patterns of Flow and Their Modification within and Around A Giant Kelp Forest. *Limnol. Oceanogr.*, 52: 1838-1852. <https://doi.org/10.4319/lo.2007.52.5.1838>
- Hayashi, L., Santos, A.A., Faria, G.S.M., Nunes, B.G., Souza, M.S., Fonseca, A.L.D., Barreto, P.L.M., Oliveira, E.C. & Bouzon, Z.L. 2011. *Kappaphycus alvarezii* (Rhodophyta, Areschougaceae) Cultivated in Subtropical Waters in Southern Brazil. *J. Appl. Phycol.*, 23: 337-343. <https://doi.org/10.1007/s10811-010-9543-5>
- Helmi, M., Satriadi, A., Suryoputro, A.A.D., Marwoto, J. & Hariyadi, H.S. 2018. Rehabilitation Priority Area Assessment on Death Coral using Cell-Based Modelling Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.*, 9(11): 2949-2961.
- Helmi, M., Purwanto, Atmodjo, W., Subardjo, P. & Aysira, A. 2017. Benthic Diversity Mapping and Analysis Base on Remote Sensing and Seascape Ecology Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.*, 9(10): 227-235.
- Jalil, A.R., Samawi, M.F., Azis, H.Y., Jaya, I., Malik, A., Yunus, I. & Sohopi, M.A.A. 2020. Comparison of physical-chemical conditions for seaweed cultivation in the Spermonde Archipelago, Indonesia. *AAFL Bioflux*, 13(4), 2071-2082.
- Junaidi, M., Lestary, D.P. & Scabra, A.R. 2018. Mariculture Site Selection Based on Environmental Parameters in Tanjung and Gangga Sub-District, North Lombok. In: Proceeding of 1st ICWBB 2018. Mataram University, Mataram, pp. 233-251.
- Kaiser, M.J., Yu, Y. & Snyder, B. 2010. Economic Feasibility of Using Offshore Oil and Gas Structures in the Gulf of Mexico for Platform-based Aquaculture. *Mar. Pol.*, 34: 699-707. <https://doi.org/10.1016/j.marpol.2010.01.002>
- Kautsari, N. & Ahdiansyah, Y. 2015. Carrying capacity and suitability assessment of Labuhan Terata waters, Sumbawa for the development of seaweed culture. *Ilmu Kelautan: Indonesian Journal of Marine Science*, 20(4): 233-238. <https://doi.org/10.14710/ik.ijms.20.4.23328>

- Kim, J.K., Yarish, C., Hwang, E.K., Park, M. & Kim, Y. 2017. Seaweed Aquaculture: Cultivation Technologies, Challenges, and Its Ecosystem Services. *Algae*, 32(1): 1-13. <https://doi.org/10.4490/algae.2017.32.3.3>
- Kregting, L.T., Blight, A.B., Elsässer, B. & Savidge, G. 2013. The Influence of Water Motion on The Growth Rate of The Kelp *Laminaria hyperborea*. *J. Exp. Mar. Biol. Ecol.*, 448: 337-345. <https://doi.org/10.1016/j.jembe.2013.07.017>
- Kunarso, Wirasatriya, A., Irwani, Satriadi, A., Helmi, M., Prayogi, H. & Munandar, B. 2019. Impact of Climate Variability on Aquatic Productivity and Fisheries Resources in Jepara. *IOP Conf. Ser.: Earth Env. Sci.*, 246: 012021. <https://doi.org/10.1088/1755-1315/246/1/012021>
- Longdill, P.C., Healy, T.R. & Black, K.P. 2008. An integrated GIS approach for sustainable aquaculture management area site selection. *Ocean Coast. Manag.*, 51: 612-624. <https://doi.org/10.1016/j.ocecoaman.2008.06.010>
- Malczewski, J. 2006. Ordered weighted averaging with fuzzy quantifiers: GIS-based multicriteria evaluation for land-use suitability analysis. *Int. J. Appl. Earth Observ. Geoinf.*, 8(4): 270-277. <https://doi.org/10.1016/j.jag.2006.01.003>
- Malczewski, J. & Rinner, C. 2015. Multicriteria Decision Analysis in Geographic Information Science. Springer, Berlin. <https://doi.org/10.1007/978-3-540-74757-4>
- Micael, J., Costa, A.C., Aguiar, P., Medeiros, A. & Calado, H. 2015. Geographic Information System in A Multi-criteria tool for Mariculture Site Selection. *Coast. Manag.*, 43(1): 52-66. <https://doi.org/10.1080/08920753.2014.985178>
- Ministry of Environment R.I, 1988. Decree of the State Minister for Population and Environment Number 02 of 1988 concerning Guidelines for Determining Environmental Quality Standards for Marine Biota (Aquaculture). Jakarta, 31 pp.
- Msuya, F.E., Buriyo, A., Omar, I., Pascal, B., Narrain, K., Ravina, J.J.M., Mrabu, E. & Wakibia, J.G. 2014. Cultivation and Utilisation of Red Seaweeds in the Western Indian Ocean (WIO) Region. *J. Appl. Phycol.*, 26: 699-705.
- Msuya, F.E. 2011. Environmental Changes and Their Impact on Seaweed Farming in Tanzania. *World Aquacult.*, 42: 34-37. <https://doi.org/10.1007/s10811-013-0086-4>
- National Standardization Agency (BSN), 2015. Good Aquaculture Practices (CBIB). 2: Sea weed. SNI 8228.2.:2015. National Standardization Agency
- Nirmala, K., Ratnasari, A. & Budiman, S. 2014. Site Selection for Seaweed Culture at Gerupuk Bay-West Nusa Tenggara Using Remote Sensing and GIS. *J. Akuakultur Indonesia*, 13(1): 73-82. <https://doi.org/10.19027/jai.13.73-82>
- Noor, N.M. 2015 The Water Bodies Compatibility Analysis for Culturing Brown Seaweed *Kappaphycus alvarezii* in Ketapang Seashore, South Lampung. *Maspari J.*, 7(2): 91-100.
- Nurdin, N., Helmi, M., Setiawan, R.Y., Zainudin, M., Rustam, A. & Akbar, M. 2019. Landsat Capability to Mapping of Total Solid Suspended (TSS) and Coastal Ecosystem. *Int. Arch. Photogram., Remote Sens. Spatial Inf. Sci.*, 42(4): 497-502. <https://doi.org/10.5194/isprs-archives-XLII-4-W16-497-2019>
- Nursidi, Ali S.A., Anshary H. & Tahya A.M., 2017 Environmental parameters and specific growth of *Kappaphycus alvarezii* in Saugi Island, South Sulawesi Province, Indonesia. *AAAL Bioflux*, 10(4): 698-702.
- Parakkasi, P., Rani, C., Syam, R., Zainuddin & Achmad, M. 2020. Growth Response and Quality of Seaweed *Kappaphycus alvarezii* Cultivated in Various Coastal Ecosystems in the Waters of West Sulawesi, Indonesia. *AAAL Bioflux*, 13(2): 627-639.
- Patty, S.I., Arfah, H. & Abdul, M.S. 2015. The relationship among Nutrient (Phosphate and Nitrate), DO, and pH with the water fertility in Jikumerasa waters, Buru Island. *J. Pesisir dan Laut Tropis*, 1(1): 43-50. <https://doi.org/10.35800/jplt.3.1.2015.9578>
- Pickering, T.D., Skelton, P. & Sulu, J.R. 2007. Intentional Introductions of Commercially Harvested Alien Seaweeds. *Bot. Mar.* 50: 338-350. <https://doi.org/10.1515/BOT.2007.039>
- Radiarta, I.N., Erlania & Haryadi, J., 2018. Analysis of land suitability and carrying capacity of waters for the development of seaweed cultivation in Simeulue Regency; Aceh. *J. Segara*, 14(1): 11-22. <https://doi.org/10.15578/segara.v14i1.6626>

- Radiarta, I.N., Saitoh, S. & Yasui, H. 2011. Aquaculture Site Selection for Japanese Kelp (*Laminaria japonica*) in Southern Hokkaido, Japan, Using Satellite Remote Sensing and GIS-based Models. *ICES J. Mar. Sci.*, 68(4): 773-780. <https://doi.org/10.1093/icesjms/fsq163>
- Radiarta, I.N., Saputra, A. & Albasri, H. 2012. Feasibility mapping of seaweed culture area (*Kappaphycus alvarezii*) at Bintan Regency, Riau Islands Province using GIS and remote sensing approach. *Akuakultur*, 7(1): 145-157. <https://doi.org/10.15578/jra.7.1.2012.145-157>
- Rahman, I.S., Fadjar, M. & Tjahjono, A. 2019. The Relationship Water Physical-Chemical Parameters in Seaweed Cultivation *Eucheuma cottonii* with Long-line System. *Int. J. Sci. Tech. Res.*, 8(11): 1448-1452.
- Rahmayanti, F., Diana, F. & Kusumawati, I. 2018. Suitability analysis for area development of seaweed (*Eucheuma cottonii*) culture in Lhok Bubon waters, Samatiga District, Aceh Barat Regency. *J. Akuakultura*, 2(1): 26-32. <https://doi.org/10.35308/ja.v2i1.773>
- Reis, R.P, Loureiro, R.R. & Mesquita, F.S. 2010. Do Salinity Affect Growth and Carrageenan Yield of *Kappaphycus alvarezii* (Gigartinales/Rhodophyta)? *Aquacult. Res.*, 42: 1231-1234. <https://doi.org/10.1111/j.1365-2109.2010.02699.x>
- Rusliani. 2016. Study of water quality on the growth of seaweed *Gracilaria verrucosa* in a pond with verticulture method. *Octopus J. Fish. Sci.*, 5(2): 522-527.
- Setiawan, R.Y. & Kawamura, H. 2011. Summertime Phytoplankton Bloom in the south Sulawesi Sea. *IEEE J. Sel. Topics Appl. Earth Observ. Remote Sens.* 4(1): 241-244. <https://doi.org/10.1109/JSTARS.2010.2094604>.
- Sousa, F.E.S., Moura, E.A., Marinho-Soriano, E. 2012. Use of Geographic Information Systems (GIS) to Identify Adequate Sites for Cultivation of The Seaweed *Gracilaria birdie* in Rio Grande Do Norte, Northeastern Brazil. *Revista Brasileira de Farmacognosia* 22(4): 868-873. <https://doi.org/10.1590/S0102-695X2012005000087>
- Sulma, S., Manoppo, A.K.S. & Hartuti, M. 2008. Identification of Suitable Area for Seaweed Culture in Bali Waters Based on Remote Sensing Satellite Data. *Int. J. Remote Sens. Earth Sci.*, 5: 57-66. <https://doi.org/10.30536/ijreses.2008.v5.a1229>
- Susanto, R.D., Wei, Z., Adi, T.R., Zheng, Q., Fang, G., Bin, F., Supangat, A., Agustiadi, T., Li, S., Trenggono, M. & Setiawan, A. 2016. Oceanography Surrounding Krakatau Volcano in the Sunda Strait, Indonesia. *Oceanogr.* 29(2): 228-237. <https://doi.org/10.5670/oceanog.2016.31>.
- Teniwut, W.A., Marimin & Djatna, T. 2019. GIS-Based multi-criteria decision-making model for site selection of seaweed farming information center: A lesson from small islands, Indonesia. *Decis. Sci. Let.*, 8: 137-150 <https://doi.org/10.5267/j.dsl.2018.8.001>
- Utama, R.Z. & Handayani, T. 2018. Suitability Between Oceanography and Seaweed (*Eucheuma cottonii*) Cultivation Potential in Tidung Island with Geographic Information System (GIS). *ICENIS: E3S Web of Conference*, 73: p.12009. <https://doi.org/10.1051/e3sconf/20187312009>
- Valderrama, D., Cai, J., Hishamunda, N., Ridler, N., Neish, I. C., Hurtado, A. Q., Msuya, F.E., Krishnan, M., Narayanakumar, R., Kronen, M., Robledo, D., Gasca-Leyva, E. & Fraga, J. 2015. The Economics of Kappaphycus Seaweed Cultivation in Developing Countries: A Comparative Analysis of Farming Systems. *Aqua. Econ. Manag.*, 19: 251-277. <https://doi.org/10.1080/13657305.2015.1024348>
- Wirasatriya, A., Sugianto, D.N., Helmi, M., Setiawan, R.Y. & Koch, M. 2019. Distinct Characteristics of SST Variabilities in the Sulawesi Sea and the Northern Part of the Maluku Sea during the Southeast Monsoon. *IEEE J. Sel. Topics Appl. Earth Obs. Remote Sens.*, 12(6): 1763-1770.
- WWF-Indonesia. 2014. Better Management Practices: Guideline series of small scale fisheries of seaweed culture - Cottoni (*Kappaphycus alvarezii*), Sacol (*Kappaphycus striatum*) dan Spinosum (*Eucheuma denticulatum*) Version 1. WWF-Indonesia. South Jakarta. 53 p.
- Yulianto, H., Damai, A.A., Delis, P.C. & Elisdiana, Y. 2017. Spatial Analysis to Evaluate the Suitability of Seaweed Farming Site in Lampung Bay, Indonesia. *Turkish J. Fish. Aquatic Sci.* 17: 1253-1261. https://doi.org/10.4194/1303-2712-v17_6_18

Yulianto, H., Damai, A.A., Delis, P.C. & Elisdiana, Y.
2017. Spatial Analysis to Evaluate the
Suitability of Seaweed Farming Site in Lampung
Bay, Indonesia. *Turkish J. Fish. Aquatic Sci.* 17:
1253-1261. [https://doi.org/10.4194/1303-
2712-v17_6_18](https://doi.org/10.4194/1303-2712-v17_6_18)

Proof of Transfer Publication Fee



TANGGAL JAM ATM
10/09/22 06:54:22 40017501
JL PROF SUDIARTO NO 13
TVR :
NO KARTU :XXXXXXXXXXXX1147
NO REFF :8945

TRANSFER LINK

DARI :BTN
NAMA :SARJITO

TUJUAN :BNI
NAMA :- WIDIANINGSIH
REK :0162816376
REFF :

JUMLAH: RP. 2,500,000.00

DEMI KEAMANAN ANDA
SIMPAN RESI INI SEBAGAI
BUKTI SAH TRANSAKSI ANDA

Identification of Potential Locations for *Kappaphycus alvarezii* Cultivation for Optimization of Seaweed Production Based on Geographic Information Systems in Spermonde Archipelago Waters, South Sulawesi, Indonesia

Sarjito Sarjito^{1*}, Hanny Ammaria¹, Muhammad Helmi^{2,3}, Slamet Budi Prayitno¹, Nurjannah Nurdin^{4,5}, Riza Yuliratno Setiawan⁶, Parichat Wetchayont⁷, Anindya Wirasatriya^{2,3}

¹Aquaculture Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro

²Oceanography Department, Faculty of Fisheries and Marine Sciences, Universitas Diponegoro

³Center for Coastal Rehabilitation and Disaster Mitigation Studies

Jl. Prof.H. Soedarto, S.H. Tembalang, Semarang, Central Java, 50275 Indonesia

⁴Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University

⁵Marine Science and Fisheries Faculty, Hasanuddin University

Jl. Perintis Kemerdekaan Km.10 Tamalanrea, Makassar, Sulawesi Selatan Indonesia

⁶Marine Science and Fisheries Faculty, Gadjah Mada University

Bulaksumur, Caturtunggal, Depok, Sleman, Daerah Istimewa Yogyakarta 55281 Indonesia

⁷Department of Geography, Srinakharinwirot University

Sukhumvit23, Wattana, Bangkok, 10110, Thailand

Email: sarjito@live.undip.ac.id

Abstract

Indonesia is the second largest producer of red algae in the world with one of the major *Kappaphycus alvarezii* farming areas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province. *K. alvarezii* production in Pangkep Regency increased every year. However, the availability of natural seaweed can be threatened along with the increasing demand so that seaweed cultivation is necessary. Proper site selection is crucial to guarantee the success of the seaweed farming development. A Geographic Information system (GIS) with geospatial modeling approach was applied for identification of the site selection for *K. alvarezii* cultivation at the Liukang Tuppabiring District, Spermonde Islands by combining various physical and chemical parameters obtained from in-situ data and hydrodynamics modeling. The parameters are Sea Surface Temperature, salinity, pH, substrate type, current speed, wave height, DO, nitrate, phosphate, and Total Suspended Solid. This research also include the accessibility and the restricted area of shipping lanes for determining the area suitability of *K. alvarezii* cultivation. This research demonstrates a Geographic Information system with the cell-based geospatial modeling can be effectively used and found the the suitable categorized area for *K. alvarezii* is 4,546.28 ha, which is the combination of the suitable area of 4,463.08 ha and highly suitable criteria with an area of 83.2 ha. The potential productivity of wet-weight seaweed can reach 56,825 tons per cycle or 113,650 tons per year. This result suggests the GIS with geospatial modeling approach for developing the area of *K. alvarezii* culture can be applied in other area.

Keywords: Seaweed production, site selection, geospatial, hydrodynamics modeling

Introduction

Indonesia is the largest producer of red algae in the world (83% of global seaweed production) followed by the Philippines and several countries (Valderrama *et al.*, 2015; Kim *et al.*, 2017). Seaweed production in Indonesia comes not only from mariculture, and brackish water ponds but also from natural collections. However, the availability of natural seaweed can be threatened in line with their increasing demand so that seaweed cultivation is completely necessary (Pickering *et al.*, 2007). The increasing demand for seaweed was mostly driven by new markets in China, Eastern Europe, Brazil, and

others (Bixler and Porse, 2011). Furthermore, Arbit *et al.* (2019) stated that 99.73% of Indonesia's seaweed production comes from cultivation. Therefore, seaweed farming is the best choice for fulfilling market demand.

Site selection is essential in aquaculture development which depends on the species of commodities and applied cultivation methods (Kaiser *et al.*, 2010; Junaidi *et al.*, 2018). For seaweed, the success of cultivation is intrinsically linked to physical and chemical parameters of seaweed farming areas such as turbidity, temperature, salinity, pH, DO, nitrate and phosphate concentration (Rusliani, 2016;

Jalil *et al.*, 2020). Hydrodynamic aspects should also be considered in the site selection of *K. alvarezii* farming since the cultivation method uses the longline method which is prone to strong current and wave. One of major *K. alvarezii*'s farming areas within the Indonesian seas is in Spermonde Islands, Pangkep Regency, South Sulawesi Province (Jalil *et al.*, 2020). The existence of Indonesian Troughflow along the Makassar Strait (Susanto *et al.*, 2016) and the occurrence of strong upwelling at the southern part of the South Sulawesi during each summer (Setiawan and Kawamura 2011) may influence the hydrodynamics condition in the Spermonde Island. Besides, the existence of small islands among the water area makes the hydrodynamics condition at the Spermonde Islands becomes more complex. This area also has dense shipping activities. Many shipping lanes have lied among these areas which are should be avoided in the mariculture site selection. This shows the complex considerations for conducting site selection of seaweed farming at Spermonde Islands.

In dealing with the complex considerations as mentioned above, the use of the geospatial model which is part of Geographical Information Systems (GIS) becomes an effective solution for determining the proper mariculture site selection (Longdill *et al.*, 2008; Micael *et al.*, 2015). This approach, in general, is an analysis of layers that contains spatial data to evaluate and select a particular area based on its suitability criteria classified according to certain environmental aspects (Malczewski, 2006; Malczewski and Rinner, 2015). GIS utilization in mariculture development has been widely used in various types of commodities including seaweed (e.g., Radiarta *et al.*, 2011; Radiarta *et al.*, 2012; Sousa *et al.*, 2012; Nirmala *et al.*, 2014; Teniwut *et al.*, 2019) which were mainly used in-situ data and remote sensing data for the input of geospatial modeling. The application of GIS for site selection for aquaculture in the Spermonde Islands has conducted by Arifin *et al.* (2014). However, their analysis was only based on in-situ measurement which cannot represent the temporal change of ocean parameters. In the present study, cell-based geospatial modeling was performed for the determination of sustainable site selection of seaweed cultivation, which integrates various physical and chemical parameters obtained from in-situ data and the hydrodynamic model. The combination of both data becomes the new approach proposed in the present study. Considering that the hydrodynamics aspects for the site selection of seaweed culture is not available in the remote sensing data, a 2D hydrodynamic model was built to represent the hydrodynamic condition in the study area. On the other hand, Indonesia as the maritime continent experiences four-season influenced by monsoon wind (Helmi *et al.*, 2018) which impacts the

hydrodynamics conditions. Thus, in-situ and incidental measurements of current and wave cannot represent the hydrodynamics condition in the study area. This issue only can be resolved by performing the hydrodynamics model. Furthermore, the potential productivity of *K. alvarezii* at the selected site is also simulated. This research is expected to become a prototype of a geospatial model to explore the potential aquaculture area in small islands and can be a scientific-based information for local governments, researchers, NGOs, and related stakeholders for planning and developing *K. alvarezii* cultivation in the study area.

Material and Methods

The study area is presented in Figure 1. which is located at the Spermonde islands waters precisely in the Liukang Tupabbiring District, Pangkep Regency, South Sulawesi. The Liukang Tupabbiring District area is 54.44 km² (4.89% of Pangkep Regency) and consists of 20 islands. The population of this district reached 29,680 inhabitants, generally fishermen and sea farmers. The field stations are located at 4°5'3.98"S - 4°59'51.137"S and 119°18'35.532"E - 119°26'19.849"E. There are 94 sampling points distributed among the island of Bontosua, Sanane, Panjenekang, Pankaiya, Langkadea, Baranglampo, Barangcaddi Island, and reef flat area.

A field survey was conducted for collecting water quality data at 94 points in which the depths are more than 2 meters. Survey site determination was using Garmin 76C GPS. The water quality data collected are presented in Table 1. The in-situ measurements were Sea Surface Temperature (SST), salinity, pH, and substrate type. For the chemical parameters, such as DO, nitrate, phosphate, and Total Suspended Solid (TSS), the water samples were taken and analyzed. The dynamic and non-dynamic water parameter data were interpolated to obtain a spatial distribution pattern at the time of data retrieval. The data were compiled in a digital form of 5 x 5 meters cell size using spline and classified according to suitability class in Table 2.

Two dimensions (2D) hydrodynamic model was used to simulate current and wave condition at the study area. The data for generating a hydrodynamic model are as follows: (1) High-resolution satellite imagery GeoEye-1 from ESRI online (<https://esriindonesia.co.id/arcgis-online>); (2) Bathymetry data from Indonesia Geospatial Agency (<http://batnas.big.go.id/>); (3) Surface wind and wave height data obtained from 6 hourly ERA-interim data with a grid interval of 0.125° from July 2018 to July (Dee *et al.*, 2011); (4) Tidal elevation forecast from July 2018 to July 2019 (<https://www.dhigroup.com/download/mike-by-dhi-tools/coastandseatools/global-tide-model>).




Figure 1. Study area

Table 1. Water quality data collection

No	Variables	Unit	Tools/Methods	Data Retrieval Methods
1	Sea Surface Temperature (SST)	°C	Thermometer	In situ
2	Salinity	Psu	Salinometer	In situ
3	pH	-	Water quality checker	In situ
4	Dissolved oxygen (DO)	ppm	Winkler method	Laboratory analysis
5	Nitrate	mg.L ⁻¹	Ascorbic Acid Method	Laboratory analysis
6	Phosphate	mg.L ⁻¹	Brucine Sulfate Method	Laboratory analysis
7	Total Suspended Solids (TSS)	mg.L ⁻¹	Milipore filter	Laboratory analysis
8	Substrate type	-	Visually observation	In situ
9	Depth	m	Modelled depth from LPI Maps (BIG, 2011) and BATNAS (BIG, 2018) data	Integrating modelled data

The current model was constructed by continuity equations and momentum equations with the average of depth. On-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry, wind and tidal dynamics were used as an input for generating current model. The initial process was making a mesh, and setting the boundary condition of the model using the mathematical continuity equation as follows:

$$\frac{\partial u}{\partial x} + \frac{\partial v}{\partial y} + \frac{\partial w}{\partial z} = S$$

The horizontal momentum equation for component x is as follows (DHI, 2010) :

$$\frac{\partial u}{\partial t} + \frac{\partial u^2}{\partial x} + \frac{\partial uv}{\partial y} + \frac{\partial wu}{\partial z} = f_v \cdot g \frac{\partial h}{\partial x} - \frac{1}{r_0} \frac{\partial p_a}{\partial x} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial x} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{xx}}{\partial x} + \frac{\partial S_{xy}}{\partial y} \right) + F_u + \frac{\partial}{\partial z} \left(v_t \frac{\partial u}{\partial z} \right) + u_s S$$

While the horizontal momentum equation for component y is as follows (DHI, 2010):

$$\frac{\partial v}{\partial t} + \frac{\partial v^2}{\partial y} + \frac{\partial uv}{\partial x} + \frac{\partial wv}{\partial z} = f_u \cdot g \frac{\partial h}{\partial y} - \frac{1}{r_0} \frac{\partial p_a}{\partial y} - \frac{g}{r_0} \int_z^h \frac{\partial r}{\partial y} dz - \frac{1}{r_0 h} \left(\frac{\partial S_{yx}}{\partial x} + \frac{\partial S_{yy}}{\partial y} \right) + F_v + \frac{\partial}{\partial z} \left(v_t \frac{\partial v}{\partial z} \right) + v_s S$$

Note: x, y, z = cartesian coordinates of $x, y,$ and z direction; t = time; u = water particle speed of direction x ; v = water particle speed of direction y ; w = water particle speed of direction z ; η = water surface elevation; d = sea depth; h = total depth of the sea; f = coriolis parameter ($f= 2\omega \sin\theta$); Ω = ratio of earth revolution; θ = latitude coordinates; g = acceleration of gravity; $S_{xx}, S_{xy}, S_{yy}, S_{yy}$ = component of radiation stress tensor; ν_t = vertical eddy viscosity; p_a = atmospheric pressure; ρ_0 = sea density; S = debits due to sources; u_s = particle velocity of air sources direction x ; and v_s = the particle speed of air sources direction y .

Wave modeling used directional decoupled parametric and fully spectral formulations based on wave eternity movements with the input data of on-screen digitized coastline from high-resolution satellite imagery GeoEye-1, bathymetry data, wind speed and direction and wave data. Parameterization was created in the frequency domain by introducing the 0 and 1st moments of the wave spectrum as a variable (Azhar et al., 2011). The arranger equation was the equilibrium wave force in cartesian coordinate formulated as follow (DHI, 2010):

$$\frac{\partial N}{\partial t} + \frac{\partial}{\partial \phi} C\phi N + \frac{\partial}{\partial \lambda} C\lambda N + \frac{\partial}{\partial \sigma} C\sigma N + \frac{\partial}{\partial \theta} C\theta N = \frac{S}{\sigma}$$

Source energy, S , shows the superposition of surface functions in various styles (Azhar et al., 2011).

$$S=S_{in}+S_{nl}+S_{ds}+S_{bot}+S_{surf}$$

Note: $N_{x,\sigma,\theta,t}$ = power density, $\phi\lambda$ = coordinate spherical, C = speed of propagation of four-dimensional wave group, S = energy source, S_{in} = formation by wind, S_{nl} = energy transfer, non-linear, S_{ds} = wave energy due to white capping, S_{bot} = by bottom friction, S_{surf} = dissipation due to the depth-induced breaking.

The maximum value of the current speed and wave height were taken in each grid that comes from 4 seasonal patterns characteristics, that is the west monsoon (December), transition 1 (May), east monsoon (July), and transition 2 seasons (October). The results of the 2D hydrodynamic model form the patterns of current speed and wave height every season were interpolated with a spline method with a cell size of 5 x 5 meters and analyzed using cell statistics to obtain the maximum value of current speed and wave height. The results of the distribution map were classified according to suitability class in Table 2.

Determination of area suitability criteria

The determination of the weighting and scoring was done to provide value in each criterion for supporting the feasibility of seaweed cultivation. The weight determination of each criterion was based on the literature research and later on was used to weight the variables by priority. The variable that is very influential for the growth of seaweed got a high score. The determination was based on the defined suitability matrix as shown in Table 2. The total score multiplication and weight in table 2 were used to determine the class of water suitability of seaweed cultivation with quantitative analysis using the following equation:

$$Y = \sum a_i . X_n$$

The range of suitability cultivation area of *K. Alvarezii* was adjusted to use the following equation:

$$I = \frac{(\sum a_i . X_n) - (\sum a_i . X_n)_{min}}{k}$$

Note: Y = The final value, a_i = weight per parameter, and X_n = score of suitability level, I = interval of suitability class, k = number of suitability class.

Finally, the suitability interval was divided into 5 classes, i.e. 1) highly suitable (HS) is in the range of the value 42-50, 2) suitable (S) is in the range of 33-41; C) moderately suitable (MS) is in the range of 24-32; 4) less suitable (LS) are in the range of 15-23; and 5) not suitable (N) had the value of <15. Each parameter in Table 2 was transformed in digital format using a spline interpolation method with a cell size of 5 x 5 meters resulting in an index map of water suitability.

Distance suitability analysis

Distance suitability analysis was also carried out as the other limiting factors out of the water parameters. Those were the distance to shipping lanes or sea lanes and the distance to a settlement which are arranged in a digital format of 5 x 5 -meter cell-sized using the buffer and the euclidean distance method to consider the accessibility and disruption of marine traffic. These criteria are described in Table 3. The suitability of distance to the settlement was compiled using the data of inhabited island, namely Bontosua, Sanane, Panjenekang, Baranglombo, and Baranggaddi Islands, while the shipping lanes data used was inter-regional shipping routes. The results of the analysis were integrated with a water suitability index map.

Spatial modelling

The spatial modelling in this study used cell-based modelling by integrating all oceanographic and

water quality parameters based on the suitability classification according to Table 2 with the following modelling algorithms develop from Danish Hydraulic Institute (2010):

$$WS = (T_i \times 0.5) + (pH_i \times 0.5) + (S_i \times 1) + (DO_i \times 0.5) + (N_i \times 1.5) + (F_i \times 1.5) + (B_i \times 0.5) + (HS_i \times 1.5) + (CS_i \times 1.5) + (TSS_i \times 0.5) + (SB_i \times 0.5)$$

Note: WS= Water suitability, T_i = temperature index map, pH_i = pH index map S_i = salinity index map, DO_i = Dissolved Oxygen index map, N_i = Nitrate index map, F_i = phosphate index map, B_i = bathymetry index map, HS_i = significant wave height index map, CS_i = current speed index map, TSS_i = TSS index map, SB_i = seabed index map

The result of this modeling is continued by determining the classification of water suitability based on a class of suitability presented in Table 2. Then, integrating the results of water suitability above with the result of distance suitability analysis was

conducted according to Table 3 using cell statistics analysis.

Estimating productivity

After obtaining the distribution of classes areas for all variable, the seaweed production is estimated only by considering the area which is suitable or highly suitable. Longline method of seaweed cultivation unit was designed as 25 × 50 meter per unit, the distance between the rope was 1 meter and the planting distance of each tied seaweed was 25 cm (WWF, 2014). The initial weight of the seedlings was approximately 100 grams per bunch (Parakkasi et al., 2020) with the wet weight crop was 500 grams per tie. The cultivation of seaweed with a floating rope/longline system produces an excellent product and can gain a weight up to four to five times from the initial weight of the seedlings (100 g) in a period of cultivation time. The number of the allocated cultivation units was estimated to be 5 units.ha⁻¹. The distance correction between units is 10 m in width

Table 2. Table of suitability criteria for seaweed *Kappaphycus alvarezii* cultivation

Variables	Level of conformity (score)					Weight	Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)		
SST (°C)	28-30	24-27 or 31-32	22-23 or >32	21 or >34	<21 or >36	0.5	Modification of Noor (2015); Rahmayanti et al. (2018)
pH	7-8	6.5-7 or 8-8.5	6-6.4 or >8.5	4-6	<4 or >9	0.5	Modification of Kautsari and Ahdiansyah (2015); Noor (2015)
Salinity (psu)	28-31	26-28 or 31-33	<26 or ≥ 34	18-25 or 35-36	<18 or >37	1	Modification of Kautsari and Ahdiansyah (2015); Rahmayanti et al. (2018); Utama and Handayani (2018)
DO (mg.L ⁻¹)	>6	5-6	4-5	2-4	<2	0.5	Modification of Afandi and Musadat (2018)
Nitrate (mg.L ⁻¹)	0.04-0.07	0.02-0.03	0.01 or >0.1	<0.01 or >0.5	>1	1.5	Modification of Kautsari and Ahdiansyah (2015); Afandi and Musadat (2018)
Phosphate (mg.L ⁻¹)	0.051-0.1	0.02-0.04	0.01-<0.02 or >0.2-1	<0.01 or >1	>2	1.5	Modification of Patty et al. (2015)
Depth (m)	3-10	2-3 or 11-13	1-2	>13-15	<1 or >15	0.5	Modification of Radiarta et al. (2012); Kautsari and Ahdiansyah (2015); Yulianto et al. (2017)
Wave height (m)	0.2-0.3	0.1-0.19	>0.3	<0.1 or >0.4	>0.5	1.5	Modification of Rahmayanti et al. (2018)
Current speed (m.s ⁻¹)	0.25-0.3	0.2-0.24 or 0.31-0.4	0.1-0.19 or >0.4	<0.1 or >0.45	>0.5	1.5	Modification of Utama and Handayani (2018)
TSS (mg.L ⁻¹)	<25	25-50	50-100	100-400	>400	0.5	Modification of Yulianto et al. (2017)
Seabed	Coral, Rubble	Sand	Reefs	Muddy Sand	Mud	0.5	Modification of Noor (2015); Yulianto et al. (2017)

Note : HS: Highly suitable, S: suitable, MS: moderately suitable, LS: less suitable, and N: not suitable

(twice longer than the average length of the fishing boat) for fishing traffic. This amount represents 62.5% of the maximum number of units i.e., 8 units.ha⁻¹. To estimated production is the multiplication between the number of cultivation units (from the total suitable area) and once crop yields per unit with the following equations:

$$Ye = N \times Cy$$

Note: Ye= estimated yield per cycle (tons), N= number of cultivation unit, Cy= crop yield per cycle (tons). Briefly, the research method is summarized in the flowchart presented in Figure 2.

Result and Discussion

Distribution of water quality parameters

Spatial distributions of the suitability of water quality parameters are shown in Figure 3. It can be seen that the water quality conditions based on the

variable of SST, and pH, are highly suitable for seaweed cultivation in the study area. SST is the important limiting factor for marine organisms (Wirasatriya *et al.*, 2019). The range of measured SST in the study area is very narrow i.e., only around 30°C. According to BSN (2015) the optimal temperature for this activity is 26-32°C, so the temperature in this site is still suitable for seaweed cultivation. The high temperature of the water causes death on seaweed while low water temperatures cause damage to protein and fat membranes and is very influential in the seaweed life cycle. Photosynthesis on seaweed generally lasts at a maximum temperature of 23°C-30°C (Msuya, 2011; Ding *et al.*, 2013). Seawater has an enormous support ability to prevent pH fluctuation. The condition of pH that is very acidic or alkaline will endanger the survival of organisms because it led to metabolic and respiration disorders. The pH range at the study site is 7.26-7.63. Whereas, optimal seaweed growth requires a pH value range of 7.5-8.5 (BSN, 2015; Radiarta *et al.*, 2018). Moreover, Noor (2015) stated that a suitable pH range for *K. alvarezii*




Figure 2. Research Flowchart

Table 3. Distance suitability analysis

Parameters	Suitability Level (Score)					Sources
	HS (5)	S (4)	MS (3)	LS (2)	N (1)	
Shipping lanes (m)	> 500	-	-	-	≤ 500	Ariyanto <i>et al.</i> (2018)
Distance to settlements (km)	4	4-5	5-6	6-7	>7	Modification of Radiarta <i>et al.</i> (2012)




Figure 3. Distribution of physical and chemical parameters: a) SST (Sea surface temperature); b). Depth; c). Salinity; d).Nitrate; e). Phosphate; f). DO (Dissolved Oxygen); g).TSS; h: seabed type; i): pH

cultivation is in normal and disposed alkaline level. Therefore, these values of pH of this area are feasible for seaweed cultivation (Rahman et al., 2019).

The suitability distribution for TSS and substrate type are moderately suitable or more in the study area. A substrate is dominated by reef almost at entire study area which is categorized as moderately suitable. The suitable condition is located around the small islands with the sand as the substrate type. Seaweed will grow optimally when the cultivation site is in coral, rubble, sand-based substrate, or a mixture of all those three. The most optimum substrate type for seaweed cultivation is a hard substrate like the rocky coral then followed by sandy corals (Parakkasi et al., 2020). Waters with a fractional substrate of coral to sand are usually passed by the current that was suitable for the growth of seaweed. However, cultivating seaweed with a longline method is not overly influenced by the substrate type. For TSS, the eastern part of the study area is highly suitable while the western part is

moderately suitable. This indicates that the further distance from the mainland the lower TSS is observed. TSS is insoluble sedimentation which increases the turbidity of water (Nurdin et al., 2019). High turbidity due to the suspended material impedes light penetration into water column and affects seaweed growth. Thus, TSS indicates water pollution as *K. alvarezii* lives optimally in transparent oligotrophic waters (Hayashi et al., 2011). According to Ministry of Environment R.I (1998), the recommended TSS value is <math><25 \text{ mg.L}^{-1}</math>, although up to 80 mg.L^{-1} is allowed for marine cultivation. The TSS levels in the study area range between 3.6-57.6 mg.L^{-1} which means that the waters are suitable for the growth of *K. alvarezii*. TSS concentration is probably due to the absence of re-suspension from bottom substrate and no input of the coastal sediment.

For DO and salinity values, the suitability distributions vary from less suitable to highly suitable. Based on the DO criterion, the study area is mostly highly suitable for seaweed cultivation, with only 2

spots in the study area which are less suitable. The water circulation among small islands plays an important role in high DO level. In open water, the low concentration of oxygen in a natural condition is rarely found (Diaz *et al.*, 2013). The optimal dissolved oxygen for seaweed cultivation is in the range of 3-7 mg.L⁻¹ (Rahman *et al.*, 2019). For salinity, the less suitable classes are spread in some areas and even the class at the eastern part of Sanane Island is not suitable for seaweed cultivation. The stability of salinity becomes an important factor for seaweed because seaweed is a stenohaline organism that cannot resist the high fluctuation salinity (Reis *et al.*, 2010; Aris *et al.*, 2021). The extreme salinity changes can lead to ice-ice disease. Salinity can affect the osmoregulation process of marine biota including seaweed (Aris *et al.*, 2021). Because *K. alvarezii* can grow optimally in the range of salinity values of 29-34 ppt (Doty, 1987), therefore, most of the area is still feasible for seaweed cultivation.

Nitrate and phosphate concentrations as indicators of water fertility are relatively low in this region. This may be related to the absence of big rivers in the small islands of the study area as the nutrient sources. Nitrate is a limiting factor for seaweed growth and one of the nutrients needed for the growth of seaweed. The result shows that nitrate concentrations in the study area range between 0.001-0.12 mg.L⁻¹ which is relatively low. According to Kautsari and Ahdiansyah (2015), the feasible nitrate concentration for seaweed cultivation ranges from 0.01 mg.L⁻¹ to 0.07 mg.L⁻¹, while the concentration of <0.01 mg.L⁻¹ and > 0.1 mg.L⁻¹ are not suitable for seaweed cultivation. Phosphate can be a limiting factor due to the source limitation. In the study area, phosphate concentration ranges from 0.001 mg.L⁻¹ to 0.062 mg.L⁻¹ which indicates low to insufficient fertility levels. Phosphate concentration for the fertile water should range from 0.051 mg.L⁻¹ to 0.1 mg.L⁻¹. Besides, phosphate concentration also controls the growth (Nursidi *et al.*, 2017) and the carrageenan content of seaweed (Patty *et al.*, 2015).

The depth of the study area ranges from 2 m to 39.2 m. The longline method can be applied at the 1 m depth from the lowest tide or below (WWF, 2014). The sufficient depth provides seaweed to be submerged. Nevertheless, noted that the deeper waters will increase the cost of constructing seaweed cultivation. Thus, the areas with a depth of more than 15 m are categorized as not suitable for seaweed cultivation.

The distribution of physical and chemical parameters at the seaweed site is highly determined by the wave and current. These aspects play a pivotal role in the water movement for algae growth (Gaylord *et al.*, 2007; Kregting *et al.*, 2013; Nursidi *et al.*,

2017). The variability of current speed and wave height is distinguished by four different characteristics of the west monsoon; transition 1; east monsoon; and transition 2 based on the results of 2D hydrodynamic modeling (Figure 4 and Figure 5). During west monsoon and transition 1 (east monsoon and transition 2), strong current speed appears at the northwestern (southern) part of the study area. The strongest current speed occurs during the transition 1 season. The pattern of the surface current in the study area is influenced by the current pattern of breaking waves in the small islands waters and the surrounding shallow area; and the influence of wind and tidal conditions in these waters. Based on the criteria shown in Table 2, the study area is categorized as highly suitable (dark blue color) to less suitable (light blue color). For the wave height, almost entire study areas are categorized as suitable and highly suitable for seaweed cultivation except during east monsoon. During west monsoon transition 1 and transition 2, the blue and dark blue colors mean that the wave height ranges from 0.1 m to 0.3 m. During east monsoon, the wave higher than 0.3 m occurs at almost the entire study area which makes it less suitable for seaweed cultivation.

The protected area is determined by several types of physical conditions such as current velocity, and wave height. The influence of strong current and ocean waves will drag and damage the cultivated seaweed and its system. The maximum value of the distribution of current speeds and wave height for 4 seasons is described in Figure 6. Based on the results in Figure 6, the current speed and wave height around the study of the water are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. The less suitable areas for the current speed (wave height) category appear mostly in the northeastern (northern) part of the study area. There is no area with a current speed of more than 0.5 m.s⁻¹ and the wave height more than 0.5 m which makes the area is not suitable for seaweed cultivation. The absence of strong current and wave may correspond to the morphology of the study area which is categorized as coral waters with a lot of shallow/reef flat area located within the small islands that protect the area from the high wave and strong current. As mentioned by Sulma *et al.* (2008), the types of protected waters are bays, caps, straits, lagoons, and reef flat. Furthermore, Helmi *et al.* (2017) also stated that the coral reef is a shoreline protector.

Potential areas for seaweed culture development and production

Various suitability criteria as limiting factor of site selection for seaweed cultivation include physical and chemical parameters and distance suitability for accessibility and restricted area inside the shipping




Figure 4. Distribution of current speed ($m \cdot s^{-1}$) based on 2D hydrodynamic model in each season




Figure 5. Distribution of significant wave height (m) based on 2D hydrodynamic model in each season

lanes has been determined into 5 highly suitable classes (HS), suitable (S), moderately suitable (MS), less suitable (LS) and not suitable (N).

Figure 7 shows that most of the study areas are moderately suitable and suitable for seaweed

cultivation. The area which is not suitable is only located along the shipping lane. Figure 8 summarizes the area size of each criteria as follows: the highly suitable (HS) is 83.2 ha (0.42%); suitable (S) is 4,463.08 ha (22.21%); moderately suitable (MS) is 9,983.52 ha (49.68%); less suitable (LS) is an area of




Figure 6. Maximum statistical value of current speed (left) and wave height (right) for all season




Figure 7. Suitability distribution for cultivation area of *K. alvarezii*




Figure 8. Potential area for *K. alvarezii* cultivation in each

494.84 ha (2.46%); and not suitable (N) is 5,071.15 ha (25.23%). The potential area for seaweed cultivation development reaches 4,546.28 ha as the

sum area of highly suitable and suitable criteria. This result is more than obtained by Arifin *et al.* (2014) which only found 1,963.6 ha area suitable and highly

suitable for *K. alvarezii* cultivation in the same study area. This indicates that our method can optimally classify the potential area for *K. alvarezii* cultivation. Furthermore, this potential area can be utilized for 22,730 cultivation units with seaweed production reaching about 56,825 tons of wet weight per crop, or within a year (2 times of harvesting) can reach about 113,650 tons. This amount is only about one-third of the total seaweed production in Pangkep Regency in 2018. However, it is important to be noted that, there are some limitations in the present study which should be considered. For example, the present study did not include the longterm climate variability which may also influence the physical and chemical parameters in the study area. As reported by Msuya *et al.* (2014) and Kunarso *et al.* (2019), the climate variability influences the oceanographic condition which then affects fisheries activity such as aquaculture and fish catch. The present study also did not consider the risk analysis which was able to disturb the cultivation process as shown by Castelar *et al.* (2015). Furthermore, to maintain the carrying capacity of the study area, the potential area that might be utilized for the expansion of *K. alvarezii* cultivation is about 50% (covering 2,273.14 ha) of the total suitable area accordingly. The result of the present study indicates the effectiveness of Geographic Information Systems with geospatial modeling for developing the area of *K. alvarezii* culture which can be applied in the other area.

Conclusions

Physical and chemical parameters as well as the accessibility and restricted area of shipping lanes were used to define the suitability criteria for *K. alvarezii* cultivation. The result is SST and pH are categorized as highly suitable for *K. alvarezii* cultivation in the whole study area. A substrate is dominated by reef type almost at the whole study area which is categorized as moderately suitable. TSS levels in the study area range between 3.6-57.6 mg.L⁻¹ which means that the waters are suitable for the growth of *K. alvarezii*. For DO and salinity, the suitability distributions vary from less suitable to highly suitable. Nitrate and phosphate concentrations mainly are too low which are not suitable for *K. alvarezii* cultivation. The depth at the study area ranges from 2 m to 39.2 m. The area with a depth more than 15 m is categorized as not suitable for *K. alvarezii* cultivation. The current speed and wave height around the study area are mostly at the range of moderately suitable to highly suitable for seaweed cultivation. After summarizing all criteria according to their weighting score, the suitable for seaweed cultivation at the study area is 4,546.28 ha, which is the combination of the suitable (S) area of 4,463.08 ha and highly suitable criteria (HS) with an area of

83.2 ha. The potential production per crop can reach 56,825 tons per cycle and 113,650 tons per year.

Acknowledgment

The author thanked the Ministry of Research Technology and Higher Education of the Republic of Indonesia for the financial support of research. The author thanked the Center for Coastal Rehabilitation and Disaster Mitigation Studies (CoREM), Diponegoro University for the facilities support on hydro-oceanographic modeling and geospatial data processing. We also thanked the Research and Development Center for Marine, Coast, and Small Island, Hasanuddin University for the surveys and laboratory analysis of water quality data.

References

- Afandi, A. & Musadat, F. 2018. Suitability analysis for seaweed culture area at the waters of Kamelanta Village and Panjang Island using Geographical Information System. *J. Akuakultura* 2(1): 69-78. <https://doi.org/10.35308/ja.v2i1.789>
- Arbit, N., Omar, S., Soekendarsi, E., Yasir, I., Tresnati, J. & Tuwo, A. 2019. Morphological and genetic analysis of *Gracilaria* sp. cultured in ponds and coastal waters. *IOP Conf. Ser.: Earth Env. Sci.* 370: 012018. <https://doi.org/10.1088/1755-1315/370/1/012018>
- Arifin, T., Bohari, R. & Arlyza, I.S. 2014. Analysis of space suitability based on marine agriculture in the small islands around Makassar. *Forum geography*, 28(1): 91-102. <https://doi.org/10.23917/forgeo.v28i1.441>
- Aris, M., Muchdar, F. & Labenua, R. 2021. Study of seaweed *Kappaphycus alvarezii* explants growth in the different salinity concentrations. *J. Ilmiah Perikanan Kelautan*, 13(1): 97-105. <https://doi.org/10.20473/jipk.v13i1.19842>
- Ariyanto, F.Y., Astor, Y. & Sidqi, M. 2018. Application of water boundary theory for determining site licenses at the coastal area and small islands in Indonesia. *Proc. Nat. Seminar of Geomatics 2018*. Bandung, pp. 893-900. <https://doi.org/10.24895/SNG.2018.3-0.1055>
- Azhar, R.M., Wurjanto, A., & Yuanita, N. 2011. Study of coastal protection of submerged breakwater at Tanjung Kait beach. Master Thesis. Master Program of Water Resource Management, Bandung Institute of Technology, Indonesia.

- Bixler, H.J. & Porse, H. 2011. A Decade of Change in the Seaweed Hydrocolloids Industry. *J. Appl. Phycol.* 23: 321-335. <https://doi.org/10.1007/s10811-010-9529-3>
- Castelar, B., de Siqueira, M.F., Sánchez-Tapia, A. & Reis, R.P. 2015. Risk Analysis Using Species Distribution Modelling to Support Public Policies for The Alien Alga *Kappaphycus alvarezii* Aquaculture in Brazil. *Aquacult.* 446: 217-226. <https://doi.org/10.1016/j.aquaculture.2015.05.012>
- Danish Hydraulic Institute (DHI). 2010. MIKE-21 and MIKE-3 Flow Model. Hydrodynamic Module Step-by-Step Training Guide. DHI MIKE, Denmark. 96 p
- Dee, D.P., Uppala, S.M., Simmons A.J., Berrisford, P., Poli, P., Kobayashi S., Andrae U., Balmaseda, M.A., Balsamo G., Bauer, P., Bechtold, P. Beljaars, A.C.M., van de Berg L., Bidlot, J., Bormann, N., Delsol, C., Dragani, R., Fuentes, M., Geer, A.J., Haimberger, L., Healy, S.B., Hersbach, H., Hólm, E.V., Isaksen L., Kållberg, P., Köhler M., Matricardi M, McNally, A.P., Monge-Sanz, B.M., Morcrette, J.J., Park, B.K., Peubey C.P. de Rosnay, P., Tavolato, C., Thépaut, J.N., & Vitart, F., 2011. The ERA-Interim reanalysis: Configuration and performance of the data assimilation system. *Q.J.R. Meteorol. Soc.*, 137(656): 553–597. <https://doi.org/10.1002/qj.828>
- Doty, M.S. 1987. The production and use of *Eucheuma*. in: Doty, M.A., Caddy, J.F, & Santilices, B. (Eds), Case Studies of Seven Commercial Seaweed Resources, FAO Fish, Tech, Pap, 281: 123- 161
- Ding L., Ma, Y., Huang, B., & Chen, S. 2013. Effects of Seawater Salinity and Temperature on Growth and Pigment Contents in *Hypnea cervicornis*. *J. Agardh (Gigartinales, Rhodophyta). BioMed. Res. Int.* 594308: 10p. <https://doi.org/10.1155/2013/594308>
- Diaz, R.J., Eriksson-Hagg, H. & Rosenberg, R. 2013. Managing Ocean Environments in a Changing Climate Sustainability and Economic Perspectives: Chapter 4. Oxygen - Hypoxia, Pages 67-96. <https://doi.org/10.1016/B978-0-12-407668-6.00004-5>
- Gaylord, B., Rosman, J.H., Reed, D. C., Koseff, J.R., Fram, J., Macintyre, S., Arkema, K., McDonald, C., Brzezinski, M. A., Largier, J.L., Monismith, S.G., Raimondi, P.T. & Mardian, B. 2007. Spatial Patterns of Flow and Their Modification within and Around A Giant Kelp Forest. *Limnol. Oceanogr.*, 52: 1838–1852. <https://doi.org/10.4319/lo.2007.52.5.1838>
- Hayashi, L., Santos, A.A., Faria, G.S.M., Nunes, B.G., Souza, M.S., Fonseca, A.L.D., Barreto, P.L.M., Oliveira, E.C. & Bouzon, Z.L. 2011. *Kappaphycus alvarezii* (Rhodophyta, Areschougaceae) Cultivated in Subtropical Waters in Southern Brazil. *J. Appl. Phycol.*, 23: 337–343. <https://doi.org/10.1007/s10811-010-9543-5>
- Helmi, M., Satriadi, A., Suryoputro, A.A.D., Marwoto, J. & Hariyadi, H.S. 2018. Rehabilitation Priority Area Assessment on Death Coral using Cell-Based Modelling Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.*, 9(11): 2949-2961.
- Helmi, M., Purwanto, Atmodjo, W., Subardjo, P. & Aysira, A. 2017. Benthic Diversity Mapping and Analysis Base on Remote Sensing and Seascape Ecology Approach at Parang Islands, Karimunjawa National Park, Indonesia. *Int. J. Civil Eng. Tech.*, 9(10): 227-235.
- Jalil, A.R., Samawi, M.F., Azis, H.Y., Jaya, I., Malik, A., Yunus, I. & Sohopi, M.A.A. 2020. Comparison of physical-chemical conditions for seaweed cultivation in the Spermonde Archipelago, Indonesia. *AAFL Bioflux*, 13(4), 2071-2082.
- Junaidi, M., Lestary, D.P. & Scabra, A.R. 2018. Mariculture Site Selection Based on Environmental Parameters in Tanjung and Gangga Sub-District, North Lombok. In: Proceeding of 1st ICWBB 2018. Mataram University, Mataram, pp. 233-251.
- Kaiser, M.J., Yu, Y. & Snyder, B. 2010. Economic Feasibility of Using Offshore Oil and Gas Structures in the Gulf of Mexico for Platform-based Aquaculture. *Mar. Pol.*, 34: 699-707. <https://doi.org/10.1016/j.marpol.2010.01.002>
- Kautsari, N. & Ahdiansyah, Y. 2015. Carrying capacity and suitability assessment of Labuhan Terata waters, Sumbawa for the development of seaweed culture. *Ilmu Kelautan: Indonesian Journal of Marine Science*, 20(4): 233-238. <https://doi.org/10.14710/ik.ijms.20.4.23328>
- Kim, J.K., Yarish, C., Hwang, E.K., Park, M. & Kim, Y. 2017. Seaweed Aquaculture: Cultivation Technologies, Challenges, and Its Ecosystem Services. *Algae*, 32(1): 1-13. <https://doi.org/10.4490/algae.2017.32.3.3>

- Kregting, L.T., Blight, A.B., Elsässer, B. & Savidge, G. 2013. The Influence of Water Motion on The Growth Rate of The Kelp *Laminaria hyperborea*. *J. Exp. Mar. Biol. Ecol.*, 448: 337–345. <https://doi.org/10.1016/j.jembe.2013.07.017>
- Kunarso, Wirasatriya, A., Irwani, Satriadi, A., Helmi, M., Prayogi, H. & Munandar, B. 2019. Impact of Climate Variability on Aquatic Productivity and Fisheries Resources in Jepara. *IOP Conf. Ser.: Earth Env. Sci.*, 246: 012021. <https://doi.org/10.1088/1755-1315/246/1/012021>
- Longdill, P.C., Healy, T.R. & Black, K.P. 2008. An integrated GIS approach for sustainable aquaculture management area site selection. *Ocean Coast. Manag.*, 51: 612-624. <https://doi.org/10.1016/j.ocecoaman.2008.06.010>
- Malczewski, J. 2006. Ordered weighted averaging with fuzzy quantifiers: GIS-based multicriteria evaluation for land-use suitability analysis. *Int. J. Appl. Earth Observ. Geoinf.*, 8(4): 270-277. <https://doi.org/10.1016/j.jag.2006.01.003>
- Malczewski, J. & Rinner, C. 2015. Multicriteria Decision Analysis in Geographic Information Science. Springer, Berlin. <https://doi.org/10.1007/978-3-540-74757-4>
- Micael, J., Costa, A.C., Aguiar, P., Medeiros, A. & Calado, H. 2015. Geographic Information System in A Multi-criteria tool for Mariculture Site Selection. *Coast. Manag.*, 43(1): 52-66. <https://doi.org/10.1080/08920753.2014.985178>
- Ministry of Environment R.I, 1988. Decree of the State Minister for Population and Environment Number 02 of 1988 concerning Guidelines for Determining Environmental Quality Standards for Marine Biota (Aquaculture). Jakarta, 31 pp.
- Msuya, F.E., Buriyo, A., Omar, I., Pascal, B., Narrain, K., Ravina, J.J.M., Mrabu, E. & Wakibia, J.G. 2014. Cultivation and Utilisation of Red Seaweeds in the Western Indian Ocean (WIO) Region. *J. Appl. Phycol.*, 26: 699–705.
- Msuya, F.E. 2011. Environmental Changes and Their Impact on Seaweed Farming in Tanzania. *World Aquacult.*, 42: 34-37. <https://doi.org/10.1007/s10811-013-0086-4>
- National Standardization Agency (BSN), 2015. Good Aquaculture Practices (CBIB). 2: Sea weed. SNI 8228.2.:2015. National Standardization Agency
- Nirmala, K., Ratnasari, A. & Budiman, S. 2014. Site Selection for Seaweed Culture at Gerupuk Bay-West Nusa Tenggara Using Remote Sensing and GIS. *J. Akuakultur Indonesia*, 13(1): 73-82. <https://doi.org/10.19027/jai.13.73-82>
- Noor, N.M. 2015 The Water Bodies Compatibility Analysis for Culturing Brown Seaweed *Kappapycus alvarezii* in Ketapang Seashore, South Lampung. *Maspari J.*, 7(2): 91-100.
- Nurdin, N., Helmi, M., Setiawan, R.Y., Zainudin, M., Rustam, A. & Akbar, M. 2019. Landsat Capability to Mapping of Total Solid Suspended (TSS) and Coastal Ecosystem. *Int. Arch. Photogram., Remote Sens. Spatial Inf. Sci.*, 42(4): 497-502. <https://doi.org/10.5194/isprs-archives-XLII-4-W16-497-2019>
- Nursidi, Ali S.A., Anshary H. & Tahya A.M., 2017 Environmental parameters and specific growth of *Kappaphycus alvarezii* in Saugi Island, South Sulawesi Province, Indonesia. *AACL Bioflux*, 10(4): 698-702.
- Parakkasi, P., Rani, C., Syam, R., Zainuddin & Achmad, M. 2020. Growth Response and Quality of Seaweed *Kappaphycus alvarezii* Cultivated in Various Coastal Ecosystems in the Waters of West Sulawesi, Indonesia. *AACL Bioflux*, 13(2): 627-639.
- Patty, S.I., Arfah, H. & Abdul, M.S. 2015. The relationship among Nutrient (Phosphate and Nitrate), DO, and pH with the water fertility in Jikumerasa waters, Buru Island. *J. Pesisir dan Laut Tropis*, 1(1): 43-50. <https://doi.org/10.35800/jplt.3.1.2015.9578>
- Pickering, T.D., Skelton, P. & Sulu, J.R. 2007. Intentional Introductions of Commercially Harvested Alien Seaweeds. *Bot. Mar.* 50: 338–350. <https://doi.org/10.1515/BOT.2007.039>
- Radiarta, I.N., Erlania & Haryadi, J., 2018. Analysis of land suitability and carrying capacity of waters for the development of seaweed cultivation in Simeulue Regency; Aceh. *J. Segara*, 14(1): 11–22. <https://doi.org/10.15578/segara.v14i1.6626>
- Radiarta, I.N., Saitoh, S. & Yasui, H. 2011. Aquaculture Site Selection for Japanese Kelp (*Laminaria japonica*) in Southern Hokkaido, Japan, Using Satellite Remote Sensing and GIS-based Models. *ICES J. Mar. Sci.*, 68(4): 773-780. <https://doi.org/10.1093/icesjms/fsq163>

- Radiarta, I.N., Saputra, A. & Albasri, H. 2012. Feasibility mapping of seaweed culture area (*Kappaphycus alvarezii*) at Bintan Regency, Riau Islands Province using GIS and remote sensing approach. *Akuakultur*, 7(1): 145-157. <https://doi.org/10.15578/jra.7.1.2012.145-157>
- Rahman, I.S., Fadjar, M. & Tjahjono, A. 2019. The Relationship Water Physical-Chemical Parameters in Seaweed Cultivation *Eucheuma cottonii* with Long-line System. *Int. J. Sci. Tech. Res.*, 8(11): 1448-1452.
- Rahmayanti, F., Diana, F. & Kusumawati, I. 2018. Suitability analysis for area development of seaweed (*Eucheuma cottonii*) culture in Lhok Bubon waters, Samatiga District, Aceh Barat Regency. *J. Akuakultura*, 2(1): 26-32. <https://doi.org/10.35308/ja.v2i1.773>
- Reis, R.P, Loureiro, R.R. & Mesquita, F.S. 2010. Do Salinity Affect Growth and Carrageenan Yield of *Kappaphycus alvarezii* (Gigartinales/Rhodophyta)? *Aquacult. Res.*, 42: 1231-1234. <https://doi.org/10.1111/j.1365-2109.2010.02699.x>
- Rusliani. 2016. Study of water quality on the growth of seaweed *Gracilaria verrucosa* in a pond with verticulture method. *Octopus J. Fish. Sci.*, 5(2): 522-527.
- Setiawan, R.Y. & Kawamura, H. 2011. Summertime Phytoplankton Bloom in the south Sulawesi Sea. *IEEE J. Sel. Topics Appl. Earth Observ. Remote Sens.* 4(1): 241-244. <https://doi.org/10.1109/JSTARS.2010.2094604>.
- Sousa, F.E.S., Moura, E.A., Marinho-Soriano, E. 2012. Use of Geographic Information Systems (GIS) to Identify Adequate Sites for Cultivation of The Seaweed *Gracilaria birdie* in Rio Grande Do Norte, Northeastern Brazil. *Revista Brasileira de Farmacognosia* 22(4): 868-873. <https://doi.org/10.1590/S0102-695X2012005000087>
- Sulma, S., Manoppo, A.K.S. & Hartuti, M. 2008. Identification of Suitable Area for Seaweed Culture in Bali Waters Based on Remote Sensing Satellite Data. *Int. J. Remote Sens. Earth Sci.*, 5: 57-66. <https://doi.org/10.30536/j.ijreses.2008.v5.a1229>
- Susanto, R.D., Wei, Z., Adi, T.R., Zheng, Q., Fang, G., Bin, F., Supangat, A., Agustiadi, T., Li, S., Trenggono, M. & Setiawan, A. 2016. Oceanography Surrounding Krakatau Volcano in the Sunda Strait, Indonesia. *Oceanogr.* 29(2): 228-237. <https://doi.org/10.5670/oceanog.2016.31>.
- Teniwut, W.A., Marimin & Djatna, T. 2019. GIS-Based multi-criteria decision-making model for site selection of seaweed farming information center: A lesson from small islands, Indonesia. *Decis. Sci. Let.*, 8: 137-150 <https://doi.org/10.5267/j.dsl.2018.8.001>
- Utama, R.Z. & Handayani, T. 2018. Suitability Between Oceanography and Seaweed (*Eucheuma cottonii*) Cultivation Potential in Tidung Island with Geographic Information System (GIS). *ICENIS: E3S Web of Conference*, 73: p.12009. <https://doi.org/10.1051/e3sconf/20187312009>
- Valderrama, D., Cai, J., Hishamunda, N., Ridler, N., Neish, I. C., Hurtado, A. Q., Msuya, F.E., Krishnan, M., Narayanakumar, R., Kronen, M., Robledo, D., Gasca-Leyva, E. & Fraga, J. 2015. The Economics of *Kappaphycus* Seaweed Cultivation in Developing Countries: A Comparative Analysis of Farming Systems. *Aqua. Econ. Manag.*, 19: 251-277. <https://doi.org/10.1080/13657305.2015.1024348>
- Wirasatriya, A., Sugianto, D.N., Helmi, M., Setiawan, R.Y. & Koch, M. 2019. Distinct Characteristics of SST Variabilities in the Sulawesi Sea and the Northern Part of the Maluku Sea during the Southeast Monsoon. *IEEE J. Sel. Topics Appl. Earth Obs. Remote Sens.*, 12(6): 1763-1770.
- WWF-Indonesia. 2014. Better Management Practices: Guideline series of small scale fisheries of seaweed culture - Cottoni (*Kappaphycus alvarezii*), Sacol (*Kappaphycus striatum*) dan Spinosum (*Eucheuma denticulatum*) Version 1. WWF-Indonesia. South Jakarta. 53 p.
- Yulianto, H., Damai, A.A., Delis, P.C. & Elisdiana, Y. 2017. Spatial Analysis to Evaluate the Suitability of Seaweed Farming Site in Lampung Bay, Indonesia. *Turkish J. Fish. Aquatic Sci.* 17: 1253-1261. <https://doi.org/10.4194/1303-2712-v17618>