

CERTIFICATE


Awarded to

Wiludjeng Roessali


as

Presenter

in International Conference on Sustainable Agriculture for Rural Development 2018
(ICSARD 2018) on 23 - 24 October 2018 at Java Heritage Hotel, Purwokerto, Indonesia



Dr. Ir. Anisur Rosyad, M.S.
Dean of Faculty of Agriculture
Jenderal Soedirman University



Handwritten signature of Susanto Budi Sulisty.

Susanto Budi Sulisty, S.TP., M.Si., Ph.D
Chairman of ICSARD 2018



< Back to results | < Previous 8 of 16 Next >

Download Print E-mail Save to PDF Save to list More... >

IOP Conference Series: Earth and Environmental Science • Open Access • Volume 250, Issue 1 • 5 April 2019 • Article number 012072 • International Conference on Sustainable Agriculture for Rural Development 2018, ICSARD 2018 • Purwokerto • 23 October 2018 through 24 October 2018 • Code 147383

Document type

Conference Paper • Bronze Open Access

Source type

Conference Proceedings

ISSN

17551307

DOI

10.1088/1755-1315/250/1/012072

View more >

The adoption behaviour and its influenced factors of true shallot seed technology in Central Java

[Roessali W.](#) ; [Purbajanti E.D.](#); [Dalmyiatun T.](#)

Save all to author list

^a Faculty of Animal and Agricultural Sciences, Diponegoro University Tembalang, Campus Semarang, Central Java, 50275, Indonesia

2⁷⁸th percentile Citations in Scopus | 1.32 FWCI | 15 Views count | [View all metrics >](#)

[View PDF](#) Full text options > Export >

Abstract

Indexed keywords

SciVal Topics

Metrics

Abstract

True shallot seed (TSS) technology is one of the way to increase shallots productivity. Its ability to produce high amount of onion requires a willingness to carry out good cultivation practices. This study was aimed to analyze the behavior of adoption of TSS technology and to analysis factors that influence the behavior of farmers adoption. The research was carried out in two regions, namely Wanasari District, Brebes Regency and Klambu District, Grobogan Regency which were the centers of shallot production. One hundred respondents were chosen with 50 farmers in each region. The data were analyzed using descriptive method and multiple regression. The result showed that the adoption rate of TSS technology in Wanasari District was higher than in Klambu District. The farmers age, education level, land acreage, farmers income, number of families, land status, and regions were significantly influence the farmers behavior in adopting TSS technology. © 2019 Published under licence by IOP Publishing Ltd.

Cited by 2 documents

Farmer Perceptions and Preferences to Shallot Farming Technology Introduction in Grobogan District, Central Java, Indonesia

Sahara, D. , Citra Kusumasari, A. , Hermawan, A. (2021) *E3S Web of Conferences*

Source Identification and Spatial Distribution of Heavy Metal Concentrations in Shallot Fields in Brebes Regency, Central Java, Indonesia

Dewi, T. , Martono, E. , Hanudin, E. (2021) *Applied and Environmental Soil Science*

[View all 2 citing documents](#)

Inform me when this document is cited in Scopus:

[Set citation alert >](#)

Related documents

Growth and yields performance of true shallot seed (TSS) in dry land of Sigi district

Saidah , Muchtar , Wahyuni, A.N. (2020) *IOP Conference Series: Earth and Environmental Science*

Perception and innovativeness level of farmers on the integration system of rice and cattle farming

Mulyono, J. , Suryana, A.T. , Suryana, E.A. (2021) *IOP Conference Series: Earth and Environmental Science*

The Effect of True Shallot Seed (TSS) Varieties and Population on Growth and Bulb Yield

Devy, N.F. , Hardiyanto, H. , Syah, J.A. (2021) *Indian Journal of Agricultural Research*

[View all related documents based on references](#)

Find more related documents in Scopus based on:

[Authors >](#) [Keywords >](#)

Source details

IOP Conference Series: Earth and Environmental Science

Scopus coverage years: from 2010 to Present

ISSN: 1755-1307 E-ISSN: 1755-1315

Subject area: Earth and Planetary Sciences: General Earth and Planetary Sciences
Environmental Science: General Environmental Science

Source type: Conference Proceeding

[View all documents >](#) [Set document alert](#) [Save to source list](#)

CiteScore 2021
0.6

SJR 2021
0.202

SNIP 2021
0.409

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology x

CiteScore 2021 counts the citations received in 2018-2021 to articles, reviews, conference papers, book chapters and data papers published in 2018-2021, and divides this by the number of publications published in 2018-2021. [Learn more >](#)

CiteScore 2021 v

$$0.6 = \frac{45,063 \text{ Citations 2018 - 2021}}{74,324 \text{ Documents 2018 - 2021}}$$

Calculated on 05 May, 2022

CiteScoreTracker 2022 i

$$0.8 = \frac{55,106 \text{ Citations to date}}{73,273 \text{ Documents to date}}$$

Last updated on 06 December, 2022 • Updated monthly

CiteScore rank 2021 i

Category	Rank	Percentile
Earth and Planetary Sciences	#153/191	20th
General Earth and Planetary Sciences		
Environmental Science	#191/228	16th
General Environmental Science		

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site &](#)

PAPER • OPEN ACCESS

International Conference on Sustainable Agriculture for Rural Development 2018 (ICSARD 2018)

To cite this article: 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **250** 011001

View the [article online](#) for updates and enhancements.

You may also like

- [Conference Committee](#)
- [11th Joint Conference on Chemistry in Conjunction with the 4th Regional Biomaterials Scientific Meeting](#)
- [International Conference on Food Science and Engineering 2016](#)

ECS Toyota Young Investigator Fellowship



For young professionals and scholars pursuing research in batteries, fuel cells and hydrogen, and future sustainable technologies.

At least one \$50,000 fellowship is available annually.
More than \$1.4 million awarded since 2015!



Application deadline: January 31, 2023

Learn more. Apply today!

PREFACE

On behalf of the Organizing Committee, I would like to express my sincere gratitude for your contributions and participations in the International Conference on Sustainable Agriculture for Rural Development 2018 (ICSARD 2018) which was held in Purwokerto, Indonesia, 23-24 October 2018.

Agriculture is a vital sector since it provides foods and raw materials related to food for human life. Agriculture gives employment opportunities to a very large proportion of communities. It also has a great significance in global trade as well as foreign exchange resources, which in turn might improve a country's GNP value. Agriculture is at a crossroad. It has to find ways to feed the world while being environmentally, socially and economically sustainable. The International Conference on Sustainable Agriculture for Rural Development 2018 (ICSARD 2018), therefore, was aimed to promote scientific and educational activities towards the advancement of knowledge by improving the theory and practice of various disciplines and areas of sustainable agriculture. This conference addressed the food system activities of processing, distributing and consuming food, as well as food production from crop and livestock; the availability, access, utilization and stability dimensions of agriculture; and the synergies and trade-offs between economic, environmental, health and social objectives and outcomes. The conference became an excellent opportunity for academic researchers, industry professionals, government delegates and students to interact and share their experiences and knowledge on cutting-edge developments in the fields of Agro-technology, Soil Science, Agronomy, Horticulture, Plant Protection, Plant Breeding and Biotechnology, Agroecology, Food Science and Technology, Agricultural and Biosystems Engineering, as well as Socio-economics of Agriculture and Agribusiness.

ICSARD 2018 was organized by the Faculty of Agriculture, Jenderal Soedirman University, Indonesia. The conference was joined by eight keynote speakers who are leading experts from reputable organizations, i.e. Prof. Dr. Robert Edwin Paull (University of Hawai'i at Manoa), Prof. Dr. Shao Hui Zheng (Saga University, Japan), Dr. Tuyen Chan Kha (Nong Lam University, Vietnam), Prof. Dr. Rindit Pambayun (Sriwijaya University, Indonesia), Prof. Lilik Soetiarso, Ph.D (Gadjah Mada University, Indonesia), Prof. Dr. Ahmad Yunus (Sebelas Maret University, Indonesia), Prof. Loekas Soesanto, Ph.D (Jenderal Soedirman University, Indonesia), and Prof. Totok Agung DH, Ph.D (Jenderal Soedirman University, Indonesia). Participants of this conference came from Sudan, Ecuador, Japan, Malaysia, Thailand, and Indonesia.

As the general chair of this conference, I realize that the success of the conference depends ultimately on the many people who have worked with us in planning and organizing both the technical program and supporting social arrangements. In particular, we thank the Program Chairs for their wise advice and brilliant suggestion on organizing the technical program and also the Scientific Committee for their thorough and timely reviewing of the papers. Recognition should go to the Organizing Committee members who have all worked extremely hard for the details of important aspects of the conference programs and social activities. We would also like to thank our special honorary guests and plenary speakers for their dedication to this event.

I hope that this publication can bring beneficial contributions to the development of knowledge particularly in the field of sustainable agriculture.

Susanto Budi Sulisty, Ph.D.
Jenderal Soedirman University, Purwokerto, Indonesia
General Chair of ICSARD 2018 Conference Committee
February 2019, Purwokerto, Indonesia



Committee of ICSARD 2018

ORGANIZING COMMITTEE

General Chair	: Susanto Budi Sulistyono
Vice General Chair	: Karseno
General Secretary	: Budi Dharmawan
Treasurer	: Woro Sri Suharti
Program Chair	: Suprayogi
Scientific Chair	: Rifda Naufalin
Proceedings/Publication Chair	: Arief Sudarmaji
Sponsorship Chair	: Budi Sustriawan
Arrangement Chairs	: Santi Dwi Astuti Purwanto Ahmad Rizqul Karim Nur Prihatiningsih Retno Setyawati

STEERING COMMITTEE

1. Suwanto
2. Anisur Rosyad
3. Heru Adi Djatmiko
4. Totok Agung
5. Loekas Soesanto
6. Robert Edwin Paull
7. Shao Hui Zheng
8. Tuyen Chan Kha
9. Rindit Pambayun
10. Lilik Soetiarso
11. Ahmad Yunus

PROGRAM COMMITTEE MEMBERS (in alphabetical order):

Abdul Mukhlis R	Erminawati	Prita Sari Dewi
Afik Hardanto	Ervina Mela Dewi	Purwoko Hari K
Agus Margiwiyatno	Fatichin	Ratna Satriani
Agus Priyono	Furqon	Rian Fedrianto
Agus Riyanto	Gunawan Wijonarko	Riana Listanti
Agus Sarjito	Hidayah Dwiyanti	Ropiudin
Ahadiyah Yugi	Ida Widyawati	Rostaman
Ahmad Arijal Lutfi	Ike Sitoresmi	Rumpoko Wicaksono
Ahmad Sujoko	Imastini Dinuriah	Sakhidin
Ali Maksum	Indah Widyarini	Saparso
Alpha Nadeira	Ismangil	Sapto Nugoho
Andi Nugroho	Kardino	Septi Dewi Yanti



Anita Widhiani
Anung Slamet Dwi P
Ardiansyah
Asna Mustofa
Budi Prakoso
Budiman
Condro Wibowo
Djeimy Kusnaman
Dyah Etika
Dyah Susanti
Eni Sumarni

Kharisun
Krissandi Wijaya
Kusmanto Edy Sularso
Masrukhi
Mohamad Nazarudin B
Mujiono
Noor Farid
Nur Aini
NurWijayanti
Parno
Poppy Arsil

Sidharta Sahirman
Siswantoro
Suswati
Suyono
Tamad
Tarso
Tatang Widjojoko
Tri Hidayatun
Wilujeng Trisasiwi
Yoga

Table of contents

Volume 250

2019

[◀ Previous issue](#) [Next issue ▶](#)

International Conference on Sustainable Agriculture for Rural Development 2018 (ICSARD 2018) 23–24 October 2018, Purwokerto, Indonesia

Accepted papers received: 01 March 2019

Published online: 05 April 2019

[Open all abstracts](#)

Preface

OPEN ACCESS 011001

International Conference on Sustainable Agriculture for Rural Development 2018 (ICSARD 2018)

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 011002

Committee of ICSARD 2018

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 011003

Peer review statement

[+ Open abstract](#) [View article](#) [PDF](#)

Papers

OPEN ACCESS 012001

Foliar iron application on growth and yield of tomato

A T Sakya and Sulandjari

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012002

Morphological characterization of local durian as parent tree in Bitingan District, Rembang

A D Sawitri, E Yuniastuti and Nandariyah

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS 012003

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our [Privacy and Cookies policy.](#)



Effectivity and compatibility of *Azotobacter* and *Bacillus* for biological control agents of fusarium wilt on banana seedlings

A Proboningrum, Hadiwiyono, S Widono and Sholahuddin

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012004

Morphophysiology and the yield of two types of moringa (*Moringa oleifera* Lamk) cultivated in two different regions in Madura

C Wasonowati, E Sulistyarningsih, D Indradewa and B Kurniasih

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012005

Readiness of farmers' institutions in facing environmental disruption: The case of participation of farmers as water users in maintaining trash-polluted irrigation in Indonesia

D Sulaeman, S S Arif and Sudarmadji

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012006

NPK fertilizer on peanut and maize cultivar under agroforestry system

D Purnomo, Mth S Budiati and AT Sakya

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012007

Colchicine concentration of *Gloriosa superba* seed at different locations and seasons

F Rahmawati, Sugiyarto and A Yunus

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012008

Vermicompost and arbuscular mycorrhiza application on growth and yield of *Artemisia annua* in the low land

Z L Gaibi, F Roviqowati, A Yunus and E Purwanto

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012009

Content of dietary fiber and vitamin A in slice jam of Tamarillo (*Solanum betaceum* Cav.) and watermelon albedo as complementary foods for school-age children

I Nuraeni, B Sustriawan and A Proverawati

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012010

Growth responses of corn cultivars on weed and nitrogen application

M Rahayu, P Yudono, D Indradewa and E Hanudin

[+ Open abstract](#) [View article](#) [PDF](#)



-
- OPEN ACCESS** 012066
Physical quality improvement of coffee Robusta (*Coffea robusta Lindl*) in Argopuro Mountain, Jember
N Kuswardhani, M Misbahudin and B H Purnomo
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012067
Measuring the coefficient of unit surface conductance of steel balls for frying without cooking oil by using dimensional analysis
Siswantoro, A Margiwiyatno and H Dwiyanti
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012068
Study of interaction between genetic source, harvest time and storage time in some mutant varieties of sweet sorghum to support future bioindustry development
A Nur, S B Santoso, K Syahrudin, R Efendy and E G Lestari
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012069
Strategic ways to develop new variety of upland rice: case study of "Inpago Unsoed 1" in Central Java, Indonesia
B Dharmawan, A R Karim and U Nurdiani
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012070
Comparison of methodologies for determination total humidity in hard corn (*Zea mays L.*)
H Palacios, D Villamarín, J Velásquez and W Vásquez
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012071
Comparison of methodologies for determination total humidity in two types of Andean corn (*Zea mays L.*)
H Palacios, M Tamayo, H Terán, J Velásquez and W Vásquez
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012072
The adoption behaviour and its influenced factors of true shallot seed technology in Central Java
W Roessali, E D Purbajanti and T Dalmiyatun
[+](#) [Open abstract](#) [View article](#) [PDF](#)
-
- OPEN ACCESS** 012073
Formulation of protection policy for sustainable cropland agriculture
This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy.
Wahyuningrat, S Rosyadi, T Haryanto and M Setiansah



[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012113

Pollinator diversity and soybean productivity with flowering plant (Crotalaria and Rosella)

Sholahuddin, R Wijayanti, R B Arniputri, Supriyadi and D Widyaningrum

[+ Open abstract](#) [View article](#) [PDF](#)

OPEN ACCESS

012114

The utilization of phenol in betel leaf (*piper betle*) as feed to reduce *Aeromonas hydrophilla* infection in dumbo catfish (*Clarias gariepinus*) and gourami (*Ospronemus gouramy* Lac.) fish cultivation

H Soeprapto and P Syarif

[+ Open abstract](#) [View article](#) [PDF](#)

JOURNAL LINKS

[Journal home](#)

[Journal scope](#)

[Information for organizers](#)

[Information for authors](#)

[Contact us](#)

[Reprint services from Curran Associates](#)



Comparison of methodologies for determination total humidity in hard corn (*Zea mays* L.)

H Palacios¹, D Villamarín¹, J Velásquez² and W Vásquez¹

¹ Facultad de Ingeniería Agroindustrial de la Universidad de Las Américas, Quito, Pichincha EC170125, Ecuador.

² Instituto Nacional de Investigaciones Agropecuarias, Quito, Pichincha EC170401, Ecuador

E-mail : hector.palacios@udla.edu.ec

Abstract. The importance of humidity moisture determination for storage and harvesting lies in the quality and safety of raw materials, process control and food preservation. There is research that comparing methodologies with significant differences, but none is applied in to hard corn, so in this paper the following objectives are established: 1) Develop standards of hard corn with seven degrees of humidity to ensure homogeneity. 2) Compare six different methodologies for determining of determination of humidity in the ranges of 10-16%. A homogeneity test was performed using the equation of Gough, 1975 and a statistical (DBCA). After verifying that the batches of corn were homogeneous within and between the bags, we proceeded to compare five methodologies in which where an analysis of variance was applied, followed by a Tukey test at 5%. For the homogeneity results inside and between the covers, the lots were homogeneous. For the results of the comparison of methodologies, significant statistical differences were found in the application of one method over the other, finally then, it was made with regression equations the corrections of the moisture results obtained from the different methodologies in relation to the ISO reference method.

1. Introduction

Corn, also known by the scientific name of *Zea mays* L. comes from the Greek Zeo, which means to live and Mahiz which means grain. It is part of the family of grasses and its origin is attributed to Mexico. It is commercialized throughout the world, the world production is approximately 800 million tons per year and is the most cultivated cereal in the world, due to its relevancy in human and animal food in addition to the production of alcohol and starch [1–4]. In Ecuador was reported a production of approximately 1.4 million tons in a cultivated area of 358 thousand hectares, with an average yield calculated of 400 kg per hectare [5].

The great demand of the agroindustry, together with the advances of techniques of sowing, mainly in the use of quality seeds, has stimulated that Ecuador has increased in 188% its maize production, which has meant a reduction of the import of approximately 80%. It is also important to consider that the production cost of 1 ton of corn in Ecuador is approximately 900 dollars, which includes the activities of land preparation, seeding, fertilization, agricultural work, harvesting and land rent [6].

Based on the morphological characteristics in Ecuador, 29 maize races are recognized, of which 13 are not cultivated in the Andean regions. The main challenges in the coastal and Amazonian regions to increase yields in the corn crop are nutrient impoverishment, compaction and salinization [7].



Phenolic acids as plant growth inhibitors from *Tridax procumbens* L.

Y Andriana^{1,2}, T N Quy¹ and T D Xuan¹

¹ Graduate School for International Development and Cooperation, Hiroshima University
Kagamiyama, Higashi-Hiroshima, Hiroshima, 739-8529 Japan

² Development Center for Appropriate Technology, Indonesian Institute of Sciences
Jl. KS. Tubun No. 5 Subang, Jawa Barat, 41213 Indonesia

E-mail: d162930@hiroshima-u.ac.jp, yusuf.andriana@lipi.go.id

Abstract. This study examined inhibitory effects of phenolic acids from *Tridax procumbens* L. on germination and growth of *Raphanus sativus* (radish). Four phenolic acids from this plant, namely benzoic acid, ellagic acid, vanillin, and ferulic acid were identified and quantified from a fraction of ethyl acetate extract by high performance liquid chromatography. These phenolic acids were subjected to germination and growth assays. Vanillin was the major constituents (364.689 µg/mg fraction) of ethyl acetate extract of this plant followed by benzoic, ellagic, and ferulic acids (69.888, 17.589, and 3.590 µg/mg fraction, respectively). In germination and growth assays, benzoic acid presented the most inhibitory effects (IC₅₀ of germination = 5.148 mM) on radish germination and growth. This compound decreased root elongation and shoot growth with IC₅₀ values of 0.947 mM and 3.452 mM respectively. This study revealed that *T. procumbens* possessed phenolic acids that can be utilized as plant growth inhibitors. Benzoic acid might play a role in the phytotoxicity of this plant. However, further trials on evaluating how this plant releases benzoic acid in soil, and examining the influence on more plants in its vicinity are needed to clarify potential use of this plant in weed control of agricultural practice.

1. Introduction

Tridax procumbens L. is an invasive plant species and belongs to Asteraceae family. It is widely spread in more than 80 countries in the world. This plant commonly known as “coat button” or “tridax daisy” (USA) or “gletang” (Indonesia) [1]. *T. procumbens* has been reported to be a weed for more than 30 crops [2]. In some countries, it has been used as a traditional medicine. In India, *T. procumbens* was used for wound healing [3], hepatoprotective [4], anti-inflammatory [5], and antimicrobial agents [6]. In Guatemala and Togo, this plant has been used to treat gastrointestinal disorders, protozoal infections, and liver diseases [7-8].

Primary studies conducted on *T. procumbens* have focused on its pharmaceutical properties. Extracts of this plant showed antioxidant [9], antidiabetic [10], anti-inflammatory [11], and anti-hyperglycemic activities [12]. For pest control, extract of this plant was reported to have vigorous larvicidal activities against *Anopheles subpictus* & *Culex tritaeniorhynchus* [13].

Allelopathic activity of *T. procumbens* has been studied sporadically. The inhibitory effects of this plant on some indicator plants such as *Raniceps ranninus*, *Sorghum bicolor*, *Lactuca sativa*, and *Alium cepa* have been reported [14,15], but the identification of allelopathic substances has not been yet



Relationship between chlorophyll content and soil plant analytical development values in two cultivars of fig (*Ficus carica* L.) as brassinolide effect at an open field

Z M Zulkarnaini, S Z Sakimin, M T M Mohamed and H Z E Jaafar

Department of Crop Science, Faculty of Agriculture, University Putra Malaysia, 43400 Serdang, Selangor, Malaysia.

E-mail: zuliasm1@gmail.com

Abstract. Relationship between chlorophyll and relative chlorophyll values obtained using soil plant analytical development (SPAD)-502 meter after brassinolide treatment were determined in two varieties of fig (*Ficus carica* L.) at an open field conditions using fresh weight basis. The experiment was arranged as split plot randomized complete block design (SRCBD) with four replications, and each experimental unit containing three plants. A different fig sample was considered as the main plot and brassinolide concentrations as sub-plots. There were four plants as destructive samples observed monthly for each replication. The correlation of SPAD value and the content of chlorophyll a, chlorophyll b and total chlorophyll content were significantly correlated in fig leaves. At IBT variety, the optimal mathematic function for SPAD value and chlorophyll a, chlorophyll b and total chlorophyll were $Y=30.209e^{-0.081x}$ ($r=0.6623$), $Y=388.42x^{1.676}$ ($r=0.3145$), and $Y=29.738e^{-0.072x}$ ($r=0.5380$) respectively. At MD variety, the optimal function models were $Y=7.4524e^{-0.032x}$ ($r=0.5318$) for chlorophyll a, $y=-0.0025x^2+0.1268x+0.2896$ ($r=0.3407$) for chlorophyll b and $Y=-0.0052x^2+0.1961x+3.1882$ ($r=0.5129$) for total chlorophyll. All these data proved SPAD-502 can be an effective tool used for rapid and nondestructive estimation of leaf chlorophyll content in fig.

1. Introduction

Chlorophyll is a main material for photosynthesis, found in plants, algae, and bacteria that gives them their green color and it also enables them to absorb the light for photosynthesis. The contents of chlorophyll reflect leaf photosynthesis ability and plant health condition [1, 2, 3, 4]. Traditional way to measure chlorophyll content usually need to extract leaf tissue with organic solvents such as acetone, ethanol, N, N-dimethyl formamide [5, 6, 7, 8]. Although this laboratory analysis method is relatively accurate, however, extraction is laborious, destructive, time-consuming, and expensive [1, 4]. In the meantime, significant pigment losses may occur during the extraction and dilution and resulted a high variability [9].

SPAD-502 chlorophyll meter (Minolta Inc, USA) is easy to use, non-destructive, and hand-held spectral device, portable diagnostic tool that measures the greenness or the relative chlorophyll content of leaves [10, 11, 12, 13, 14]. By measuring the leaf transmittance in two wave bands (400-500 nm and 600-700 nm), this device quantifies the relative amount of chlorophyll with a reading in arbitrary unit (SPAD-502 Chlorophyll Index) that is proportional to the leaf chlorophyll concentration [15, 16], which provides a substantial saving in time, space, resources and suitable for small plot areas.



The effect of lactic acid bacteria-containing calf starter on Holstein calf performance

S Mukodiningsih¹, J Achmadi¹, F Wahyono¹, T R Taswanda², Sugiono³ and S J Ohh⁴

¹ Department of Animal Science Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang, Central Java 50275, Indonesia

² Dinas Ketahanan Pangan Kabupaten Sukabumi, West Java-Indonesia

³ BBPTU-HPT Baturraden, Central Java-Indonesia

⁴ Collage of Animal Life Sciences, Kangwon National University, Chuncheon- **South Korea**

E-mail: mukodiningsih@gmail.com

Abstract. Newborn calve has an undeveloped rumen and tend to easily diarrhea caused by *Escherichia coli* from environment and cause death. Feeding calf starter (CS) contains probiotic after birth can promote rumen development and suppress *E. coli*. This experiment aimed to evaluate the effect of different feeding calf starter on Friesian Holstein (FH) calf performance. The treatments were the use of different calf starter, namely calf starter from farmer (CS-F) and calf starter contains fermented cabbage waste (CS-L). The materials were CS contains fermented waste cabbage, CS from farmer and 30 FH calves aged 7-14 days and 41.14 ± 3.28 kg initial body weight. Calf starters were fed twice per day after giving milk, forage and water given *ad libitum*. The parameters measured were feed intake, feces consistency and daily gain. The data were analyzed with t- test. The result indicated that there were no different ($P > 0.05$) on feed intake and average daily gain, but CS contains fermented waste cabbage produced better feces consistency than CS from farmer ($P < 0.05$). Feeding calf starter with addition of fermented cabbage waste produced good calve performance.

1. Introduction

Rumen is the main stomach for ruminants that develop optimally at the age of 2-6 weeks depend on feeding as soon as after birth. Solid feed consisting of calf starter (CS) and fiber sources can be given to calves starting at 1 week to accelerate rumen development [1]. The nutritional requirements for calves after birth to weaning are sourced from 40% of solid feed and 60% of milk [2]. Calf starter added with 5% molasses given to FH calves aged 2-6 weeks can accelerate rumen development [3]. One indicator of rumen development is the presence of microbes which ferment feed in the rumen, because at birth the rumen is sterile [1]. The rumen microbes come from the environment both from feed and milk given during maintenance after birth. The most dominant microbes in the rumen are bacteria. One method to accelerate the development of rumen microbes is by providing feed contains source of lactic acid bacteria.

Lactic acid bacteria can affect the pH of the rumen, by accelerating the growth of lactate-utilizing bacteria in the rumen [4]. Lactate-utilizing bacteria can reduce the production of lactic acid accumulated

