



Source details

Jurnal Teknologi (Sciences and Engineering)

Open Access ⓘ

Scopus coverage years: from 2006 to 2007, from 2010 to 2020

Publisher: Penerbit Universiti Teknologi Malaysia

ISSN: 0127-9696 E-ISSN: 2180-3722

Subject area: Engineering: General Engineering

CiteScore 2019

1.1



Add CiteScore to your site

SJR 2019

0.183



SNIP 2019

0.417



[View all documents >](#)

[Set document alert](#)

[Save to source list](#) [Journal Homepage](#)

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

i Improved CiteScore methodology



CiteScore 2019 counts the citations received in 2016-2019 to articles, reviews, conference papers, book chapters and data papers published in 2016-2019, and divides this by the number of publications published in 2016-2019. [Learn more >](#)

CiteScore 2019 ▼

$$1.1 = \frac{1,885 \text{ Citations 2016 - 2019}}{1,663 \text{ Documents 2016 - 2019}}$$

Calculated on 06 May, 2020

CiteScoreTracker 2020 ⓘ

$$1.3 = \frac{709 \text{ Citations to date}}{560 \text{ Documents to date}}$$

Last updated on 07 December, 2020 • Updated monthly

CiteScore rank 2019 ⓘ

Category	Rank	Percentile
Engineering		
General Engineering	#160/299	46th

[View CiteScore methodology >](#) [CiteScore FAQ >](#)



About Scopus

- [What is Scopus](#)
- [Content coverage](#)
- [Scopus blog](#)
- [Scopus API](#)
- [Privacy matters](#)

Language

- [日本語に切り替える](#)
- [切换到简体中文](#)
- [切换到繁體中文](#)
- [Русский язык](#)

Customer Service

- [Help](#)
- [Contact us](#)

The background of the journal cover is a deep blue color. It features several faint, light blue technical motifs: a globe at the top, a circuit board pattern with various components and lines, and a large gear on the right side. The text is centered and rendered in a clean, white, sans-serif font.

JURNAL TEKNOLOGI

SCIENCES & ENGINEERING

ISSN : 2180-3722





THE 2nd INTERNATIONAL SYMPOSIUM ON AQUATIC PRODUCTS PROCESSING AND HEALTH

“Sustaining Fish Processing Industry to Support Global Maritime Axis”

Semarang, 13-15th September 2015

Directorate General of Fisheries Product Processing and Marketing, Ministry of Marine Affairs and Fisheries and Department of Fish Products Technology, Faculty of Fisheries and Marine Science, Diponegoro University in collaboration with Indonesian Fisheries Products Processing Society will organize The 2nd International Symposium on Aquatic Products Processing and Health and Exhibition (ISAPPROSH 2) on 13-15 September 2015 at Diponegoro University, Semarang, Central Java, Indonesia. The symposium bring together national and international participant from : 1). Scientists, 2). Policy Makers, 3). Practitioners, 4). Private Sectors, 5). Students.

Topics

1. Development of fishery product handling, processing, and preservation
2. Technological innovation and post-harvest equipment
3. Quality management and product safety
4. Biotechnology aquatic in nutraceutical and functional foods
5. Marketing and management on aquatic processing industry
6. Aquatic enzyme and bacteria for fishery products and health
7. Sustainable environmental management
8. Sanitation and public health

KEYNOTE SPEAKER

1. H.E. Joko Widodo (President of the Republic of Indonesia)
2. H.E. Madam Susi Pudjiastuti (Minister of Marine Affairs and Fisheries)
3. H.E. Mohammad Nasir (Minister of Research, Technology and Higher Education)

INVITED SPEAKER

1. Director of Directorate General of Processing and Marketing of Fisheries Product (Ministry of Marine Affairs and Fisheries)
2. Prof. Kazuo Miyashita (Hokkaido University - Japan / President of Japan Oil Chemist Society (JOCS))
3. Prof. Mohammad Shafiur Rohman (Sultan Qaboos University - Sultanate of Oman)
4. Prof. Sootawat Benjakul (Prince of Songkla University - Thailand)
5. Prof. Toru Suzuki (Tokyo University of Marine Science and Technology - Japan)
6. Prof. Irwandi Jaswir (International Islamic University of Malaysia (IIUM)- Malaysia / Director of International Institute for Halal Research and Training (INHART)
7. Prof. Dr. Ocky Karna Radjasa, MSc (Head of Diponegoro University's Research and Public Service Institute - Diponegoro University - Indonesia)
8. Dr. Ir. Widodo Farid Ma'ruf, MSc (President of Seaweeds Commission of Indonesia / Diponegoro University - Indonesia)
9. Prof. Daniel Khan (Grimsby University - United Kingdom)
10. Dr. Klervi Le Lann (Lab. LEMAR - IUEM, Universite de Bretagne Occidentale, Brest France)
11. Mohammad Novi Saputro (Kelola Mina Laut Foods)

ONLINE REGISTRATION AND ABSTRACT SUBMISSION

Online registration, extended abstract and scientific article submission, and confirmation proofs of payment could be done at the ISAPPROSH 2 website : www.isapprosh2undip.com
For detail information and registration please visit our website.

PUBLICATION

All accepted paper will be published in Knowledge E Digital Libraries-Proceeding and selected paper at Aquatic Procedia-Elsevier.

IMPORTANT DATE AND REGISTRATION

Stage	Deadline
First circular (online)	1 March to 1 June 2015
Second circular	1 June to 1 August 2015
Extended abstract submission	1 March to 1 August 2015
Notification of Acceptance	15 August 2015
Full paper (camera ready) submission ¹	31 August 2015
Registration Fee Payment	20 August to 7 September 2015
Symposium	13-15 September 2015

REGISTRATION FEE

Categories	Early Bird (before 1 June 2015)	Normal (after 1 June 2015)
Indonesia	IDR	IDR
Presenter	500.000	750.000
Students (Student ID)	300.000	500.000
Non-presenting delegate	300.000	500.000
Accompanying person	300.000	500.000
International	US\$	US\$
Presenter	150	200
Students (Student ID)	75	100
Accompanying person	50	75

Please transfer the registration fee to:
Bank account No: 1360006644659 (bank Mandiri)
Holder : ESCROW
Rekening Post Kredit FPIK UNDIP

Hosted by :



Supported by :



Contact Us
Secretariat of ISAPPROSH 2 2015
Department of Fish Products Technology, Faculty of Fisheries and Marine Science,
Diponegoro University, Jl. Prof. Soedarto, SH Tembalang Semarang INDONESIA 50275.
Phone: +6281215290286; Email: isapprosh2undip@gmail.com
Website: www.isapprosh2undip.com

[Home](#) / [Archives](#) /

Vol. 78 No. 4-2: Sustaining Fish Processing Industry to Support Global Maritime Axis

Vol. 78 No. 4-2: Sustaining Fish Processing Industry to Support Global Maritime Axis

Chief Editor: Roy Hendroko Setyobudi (Jakarta, IDN) Â Editorial board: Daniel Khan (London, GBR), Fronthea Swastawati (Semarang, IDN), Maizirwan Mel (Kuala Lumpur, MYS), Mohamad Shafiur Rahman (Muscat, OMN), Nathalie Bourgongnoun (Brest, FRA), Praptiningsih Gamawati Adinurani (Madiun, IDN), Yuzo Shioi (Shizuoka, JPN)

Published: 2016-04-18

[Home](#) / [Archives](#) /

Vol. 78 No. 4-2: Sustaining Fish Processing Industry to Support Global Maritime Axis

Vol. 78 No. 4-2: Sustaining Fish Processing Industry to Support Global Maritime Axis

Chief Editor: Roy Hendroko Setyobudi (Jakarta, IDN) Â Editorial board: Daniel Khan (London, GBR), Fronthea Swastawati (Semarang, IDN), Maizirwan Mel (Kuala Lumpur, MYS), Mohamad Shafiu Rahman (Muscat, OMN), Nathalie Bourgongnoun (Brest, FRA), Praptiningsih Gamawati Adinurani (Madiun, IDN), Yuzo Shioi (Shizuoka, JPN)

Published: 2016-04-18

Science and Engineering

EXTRACTION OF SNAKEHEAD FISH [*Ophiocephalus striatus* (Bloch, 1793)] INTO FISH PROTEIN CONCENTRATE AS ALBUMIN SOURCE USING VARIOUS SOLVENT

Abdul Rasyid Romadhoni, Eddy Afrianto, Rusky Intan Pratama

[PDF](#)

BLOOD GLUCOSE LEVEL AND LIPID PROFILE OF ALLOXAN-INDUCED DIABETIC RATS TREATED WITH NA-ALGINATE FROM SEAWEED *Turbinaria ornata* (Turner) J.Agardh

Amir Husni, Setyaning Pawestri, Alim Isnansetyo

[PDF](#)

ANALYSIS OF BIOACTIVE COMPOUNDS OF *Caulerpa racemosa*, *Sargassum* sp. AND *Gracillaria verrucosa* USING DIFFERENT SOLVENTS

Arham Rusli, Metusalach Metusalach, Mulyati Muhammad Tahir, Salengke Salengke, Syamsuar Syamsuar

[PDF](#)

CARCASS QUALITY OF RAW AND SMOKED FISH FILLETS PREPARED FROM CAGE RAISED RIVER CATFISH (*Hemibagrus nemurus* Valenciennes, 1840) FED HIGH PROTEIN-LOW ENERGY AND LOW PROTEIN-HIGH ENERGY DIETS

Bustari Hasan, Indra Suharman, Desmelati Desmelati, Dian Iriani

[PDF](#)

STUDY OF GROWTH AND N, P CONTENT OF MICROALGAE *Chlorella vulgaris* CULTIVATED IN DIFFERENT CULTURE MEDIA AND LIGHT INTENSITY

Dedi Edwin Satriaaji, Muhammad Zainuri, Ita Widowati

 PDF

HIGHLY COMMERCIAL FISHERIES TAWAR FISH: MOLECULAR ANALYSIS DNA MITOCHONDRIAL COI GENE SEQUENCE AND PROXIMATE ANALYSIS FROM MALACCA STRAIT, RIAU

Delianis Pringgenis, Ragil Susilowati

 PDF

THE DIETARY PROTEIN REQUIREMENT IN FORMULATED FEED FOR SPECIFIC GROWTH, DIGESTIBILITY AND SURVIVAL OF SPIRAL BABYLON (*Babylonia spirata*)

Diana Rachmawati, Istiyanto Samidjan

 PDF

ANTIOXIDANT ACTIVITIES OF PHYCOCYANIN MICROCAPSULES USING MALTODEXTRIN AND CARRAGEENAN AS COATING MATERIALS

Eko Nurcahya Dewi, Lukita Purnamayati, Retno Ayu Kurniasih

 PDF

COMPARATION OF SPINY LOBSTER (*Panulirus Sp.*) POPULATIONS FROM BANTUL AND CILACAP, CENTRAL JAVA, INDONESIA

Florencius Eko Dwi Haryono, Sahala Hutabarat, Johannes Hutabarat, Ambariyanto Ambariyanto

 PDF

CRUDE CATHEPSIN ACTIVITY AND QUALITY CHARACTERISTIC OF SMOKED CATFISH [*Pangasius pangasius* (Hamilton, 1882)] PROCESSED BY DIFFERENT SMOKING TEMPERATURE

Fronthea Swastawati, Ahmad Niâ€™matullah Al Baarri, Tri Winarni Agustini, Eko Nurcahya Dewi, Ima Wijayanti, Dwi Yanuar Budi Prasetyo, Daniel Khan

 PDF

MOLECULAR CHARACTERIZATION OF *Dunaliella salina* AND *Chlorella vulgaris* FUSANT USING 18SrDNA GENE

Hermin Pancasakti Kusumaningrum, Muhammad Zainuri

 PDF

THE INFLUENCE ANALYSIS OF DIFFERENTLY CONSTRUCTED FOLDED TRAPS AND TYPES OF BAITS TO CATCH CRABS [*Portunus pelagicus*, (Linnaeus,1758)] IN REMBANG SEA WATERS

Herry Boesono, Sansan Sansan, Agus Suherman

[Home](#) / [Archives](#) /

[Vol. 78 No. 4-2: Sustaining Fish Processing Industry to Support Global Maritime Axis](#) /

[Science and Engineering](#)

APPLICATION OF *Spirulina platensis* ON ICE CREAM AND SOFT CHEESE WITH RESPECT TO THEIR NUTRITIONAL AND SENSORY PERSPECTIVES

Tri Winarni Agustini

Fisheries Department, Diponegoro University, Jl. Prof. H. Sudharto, S.H, Tembalang, Semarang, 50275, Indonesia

Widodo Farid Maâ€™ruf

Fisheries Department, Diponegoro University, Jl. Prof. H. Sudharto, S.H, Tembalang, Semarang, 50275, Indonesia

Widayat Widayat

Department of Chemical Engineering, Faculty of Engineering, Diponegoro University

Meiny Suzery

Department of Chemistry, Faculty of Science and Mathematics, Diponegoro University

Hadiyanto Hadiyanto

Department of Chemical Engineering, Faculty of Engineering, Diponegoro University

Soottawat Benjakul

Department of Food Technology, Faculty of Agroindustry, Prince of Songkhla University, Hat Yai, Songkhla 90112, [Thailand](#)

DOI: <https://doi.org/10.11113/jt.v78.8216>

Keywords: Ice cream, physical properties, sensory, soft cheese, *Spirulina platensis* (Gomont) Geitler

Abstract

Application of *Spirulina platensis* (Gomont) Geitler into food product can be used for producing functional food and improve its nutritional value. However, some bioactive compounds containing in *S. platensis* are heat sensitive, therefore processing techniques need to be strictly considered. It is necessary to observe the application of *S. platensis* powder into different products of ice cream and soft cheese in which the application of *S. platensis* was in relatively low temperature to protect its bioactive compounds from damage. *S. platensis* contains approximately 55 % to 70% of protein and its utilization on food product can be expected to improve the nutritional value. Innovation technique to produce such kind of product should respect to its acceptance by panelist using sensory test. Therefore, the objective of this research was to find out maximum concentration of *S. platensis* that can be

[Home](#) / [Archives](#) /

[Vol. 78 No. 4-2: Sustaining Fish Processing Industry to Support Global Maritime Axis](#) /

[Science and Engineering](#)

ENVIRONMENTAL ASSESSMENT OF POLY CULTURE FARMING PRACTICE BASED ON MACROBENTHIC ASSEMBLAGES: A STUDY CASE AT COASTAL AREA OF KALIWUNGU, KENDAL (CENTRAL JAVA, INDONESIA)

Sapto Purnomo Putro

Center of Marine Ecology and Biomonitoring for Sustainable Aquaculture (Ce-MEBSA), Diponegoro University, Tembalang, Semarang 50275, Indonesia

Widowati Widowati

Department of Mathematics, Faculty of Sciences and Mathematics, Diponegoro University, Tembalang, Semarang 50275, Indonesia

Ibni Jeudi Febria

Center of Marine Ecology and Biomonitoring for Sustainable Aquaculture (Ce-MEBSA), Diponegoro University, Tembalang, Semarang 50275, Indonesia

Fuad Muhammad

Center of Marine Ecology and Biomonitoring for Sustainable Aquaculture (Ce-MEBSA), Diponegoro University, Tembalang, Semarang 50275, Indonesia

Suhartana Suhartana

Department of Chemistry, Faculty of Sciences and Mathematics, Diponegoro University, Tembalang, Semarang 50275, Indonesia

Suminto Suminto

Department of Fishery, Faculty of Fishery and Marine Science, Diponegoro University, Tembalang, Semarang 50275, Indonesia

Agung Sudaryono

Department of Fishery, Faculty of Fishery and Marine Science, Diponegoro University, Tembalang, Semarang 50275, Indonesia

Sunshuke Koshio

Department of Fisheries, Faculty of Fisheries, Kagoshima University, 1-21-24 Korimoto Kagoshima, Kagoshima Prefecture 890-8580, [Japan](#)

DOI: <https://doi.org/10.11113/jt.v78.8206>

Keywords: Environmental disturbance, macrobenthos, moderately disturbed area, polychaete, polyculture

[Home](#) / [Archives](#) /

[Vol. 78 No. 4-2: Sustaining Fish Processing Industry to Support Global Maritime Axis](#) /

[Science and Engineering](#)

CARBON DIOXIDE CAPTURE EFFICIENCY USING ALGAE BIOLOGICAL ABSORBENT AND SOLID ADSORBENT FOR BIOGAS PURIFICATION

Praptiningsih Gamawati Adinurani

Faculty of Agrotechnology University of Merdeka, Jl. Serayu, PO. Box 12, Madiun 63131, Indonesia

Roy Hendroko Setyobudi

Indonesian Association of Bioenergy Scientist and Technologist, BPPT Building II, 22nd floor Jl. MH. Thamrin No. 8 Jakarta 10340

Satriyo Krido Wahono

Technical Implementation Unit for Development of Chemical Engineering Processes, Indonesian Institute of Sciences, 55861, Yogyakarta, Indonesia

Maizirwan Mel

Faculty of Engineering, International Islamic University Malaysia (IIUM), Gombak, 50728 Kuala Lumpur, Malaysia

Anggi Nindita

Department of Agronomy and Horticulture, Faculty of Agriculture, Bogor Agricultural University, Jl. Meranti, Kampus IPB Darmaga, Bogor 16680, Indonesia

Endang Purbajanti

Agricultural and Animal Science Faculty, Diponegoro University, Jln. Prof Sudharto SH, Tembalang Semarang, 50275, Indonesia

Soni Sisbudi Harsono

Faculty of Agricultural Technology, University of Jember (UNEJ), Jl. Kalimantan 23 Jember 68121 Indonesia

Andoniana Rakoto Malala

Centre de Formation et d` Application du Machinisme Agricole (CFAMA)Route Betafo Ambaniandrefana - B.p. 109, Antsirabe, **Madagascar**

Salafudin Salafudin

Departement of Chemical Engineering, ITENAS Jl. PHH Mustafa No.23, Bandung 40123, Indonesia

Andi Sasmito

Alumnus Department of Biology, Faculty of Mathematics and Natural Sciences Bogor Agricultural University, Jl. Agatis, Kampus IPB Darmaga, Bogor 16880, Indonesia

DOI: <https://doi.org/10.11113/jt.v78.8201>

Keywords: Biogas, biological purification, biomethane, solid adsorbent purification

[Home](#) / [Archives](#) /

[Vol. 78 No. 4-2: Sustaining Fish Processing Industry to Support Global Maritime Axis](#) /

[Science and Engineering](#)

THE INNOVATION OF VULNERABLE FISHERIES USING ECOSYSTEM-BASED FISHERY MANAGEMENT APPROACH: A TEST CASE IN KARIMUNJAWA ECOSYSTEM, CENTRAL JAVA, INDONESIA

Mayanggita Kirana

aFaculty of Economics and Business, Universitas Diponegoro, Jl. Prof. Soedarto, SH Tembalang, Semarang 50275, Indonesia

Indah Susilowati

Faculty of Economics and Business, Universitas Diponegoro, Jl. Prof. Soedarto, SH Tembalang, Semarang 50275, Indonesia

Kuperan Viswanathan

Othman Yeop Abdullah Graduate School of Business, Universiti Utara Malaysia, 06010 UUM Sintok, Kedah Darul Aman, [Malaysia](#)

DOI: <https://doi.org/10.11113/jt.v78.8194>

Keywords: Central Java, ecosystem, fisheries, management

Abstract

The sustainability of marine ecosystem has become a major concern the government; however, the implementation of sustainability-based fisheries management has not been fully carried out and well controlled. Therefore, having a concept of ecosystem-based fisheries management (EBFM) is essential in protecting it preserved. The aim of this study was to analyze the implementation of EBFM in Karimunjawa ecosystem, Central Java, Indonesia. The analysis of this study was based on the primary data collected from fishermen and stakeholders using in-depth interviews, and the secondary data gathered from stakeholders of Karimunjawa documentation. Meta-analysis with triangulation was invoked in this study. The result showed that the vulnerability of marine ecosystem, particularly fisheriesâ€™ resource in the pilot project is in progress. The conventional approach has not yet succeeded in managing fisheriesâ€™ resource in terms of sustainability attributes. Moreover, the EBFM has not yet proven to be a suitable approach for some reasons; although, this concept is very promising in encouraging a new paradigm for sustainable management in Indonesia with a protocol concept. This initial finding needs to be furthered in order to explore other aspects of development.

SAND DOLLARS DISTRIBUTION PATTERN AND ABUNDANCE AT THE COAST OF CEMARA KECIL ISLAND, KARIMUNJAWA, JEPARA, INDONESIA

Article history

Received

5 November 2015

Received in revised form

21 December 2015

Accepted

20 February 2016

Suryanti*, Max Rudolf Muskananfola, Kevin E. Simanjuntak

Aquatic Resource Management Study Program Faculty of Fisheries and Marine Science, Diponegoro University, Jl. Prof. Soedarto S.H., Tembalang, Semarang 50275, Indonesia

*Corresponding author
suryantidr@gmail.com

Graphical abstract



Abstract

This research is aimed to know the distribution patterns and abundance of *sand dollars* [*Clypeaster subdepressus* (Gray, 1840)] at different spatial distances from the coast of Cemara Kecil Island Karimunjawa, Indonesia. The data were collected using purposive sampling and analyzed using cluster analysis PAST-3. The results indicate that the abundance of *sand dollar* amounts to 2 200 individuals, and the existing species are *Laganum centrale*, *L. laganum*, *L. fudsiyama*, *L. depressum*, *L. retinens*, *L. boschi*, *L. depressum tonganense*, *L. dickersoni*, *L. decagonale rectum*, *L. joubini* and *L. decagonale*. The cluster analysis using PAST-3 reveals that there are clumped distributions and uniform distributions of *L. centrale* and *Laganum laganum*. At a distance of 23 m to 38 m from the coastline the animals form clumped distributions.

Keywords: Abundance, sand dollars [*Clypeaster subdepressus*(Gray, 1840)], spatial distribution pattern

© 2016 Penerbit UTM Press. All rights reserved

1.0 INTRODUCTION

Sand dollars [*Clypeaster subdepressus* (Gray, 1840)] belong to the class of echinoidea of the phylum echinoderms and have irregular anatomy, flattened and have no arms. In general, sand dollars live in the area of intertidal to sub-tidal zone, buried in mud or fine sand [1]. This animal is a deposit feeder, eating the residuals of living organisms found in a substrate and ingesting planktons and algae, or passively collects microorganisms and organic matters trapped by the spines in particular with aboral sucker [2]. The body of sand dollars is wrapped in structured shells (test), consisting of dials that form a box like a hard shell. Usually there are 10 double columns of plates and five pierced for the vessel-like feet which are slender out through shells. The mouth of sand dollars is located in the middle and the interior of the mouth or teeth moor and form a structure called Aristotle's *Lantern* [3].

Cemara Kecil Island is one of the islands of Karimunjawa islands, Jepara, Central Java, Indonesia. Cemara Kecil Island is uninhabited and is the area included in tourism zone as it has abundant potential natural resources such as seagrass beds, coral reefs, mangrove ecosystem and various kinds of echinoderms, one of which is sand dollars [4, 5].

Geographically, Cemara Kecil Island is located at 110° 22' 38" to 110° 22' 44" EL and 5° 49' 51" to 5° 50' 2" SL covering an area of 1.5 ha. Cemara Kecil Island is a beach with white sand with some kinds of seagrass beds, dead coral fractions, mangrove and some coral reefs [6, 7].

Sand dollars have important roles in ecology as the basic ecosystems in terms of cleaning environment as they usually live in fine sand habitats. The abundance and the spatial distribution of this unique biota in Pulau Cemara Kecil coastal waters therefore need to be analyzed. The present work studies the abundance and the spatial distribution of sand dollars at different

distances from the coastline of Cemara Kecil Island in Karimunjawa, Jepara. The study was conducted from March 2015 to April 2015 and an analysis was made utilizing some statistical method.

2.0 EXPERIMENT

Materials used in this research are sand dollars found in Cemara Kecil Island Karimunjawa Jepara, while the data were collected using purposive sampling and the descriptive-qualitative method. Purposive sampling method is the determination of recipient with some certain consideration by researchers based on features or properties of the population which are already known [8, 9].

2.1 The Sampling Location

Sampling was conducted at the sand zone of the lowest low tide spanning 50 m along the coastline. The location was then divided into sections by drawing seven virtual lines each separated 5 m from the next line, and repeated twice as shown in Figure 1.

2.2 Data Analysis

The data obtained from the survey were then statistically analyzed using multivariate statistical analysis with cluster analysis with the help of PAST-3 / Paleontological Statistics_3. Program Multivariate analysis is a statistical analysis best used to analyze data that arise from more than one variable in the experimental design [10, 11].

2.3 Relative Abundance and the Distribution Pattern

The Relative Abundance of sand dollars is the number of percent composition of sand dollars relative to the total number of organism in the area.

$$RA = P_i \times 100 \% \tag{1}$$

Where :

RA = Relative Abundance;

P_i = Probabilities of species i in the total organisms

The distribution pattern of sand dollars at certain distances from the coastline can be obtained from the mean values and the variance [12].

The mean value and the variance were obtained using the formula (2):

$$V = \sqrt{\frac{2}{n-1}} \quad \text{and} \quad m = \frac{n}{N} \tag{2}$$

Where:

V= variance; n = number of individuals, m = mean; N = sample size

The distribution of organisms in nature is grouped into three:

V = m indicates random distribution

V > m indicates clumped distribution

V < m indicates uniform distribution




Figure 1 Sampling with Quadrant Transect

3.0 RESULTS AND DISCUSSION

3.1 Result

The cluster analysis using PAST-3 program reveals that sand dollars in Cemara Kecil Island (presented in Table 1 and Figure 3) are mostly clumped; some,

such as *Laganum centrale* and *Laganum laganum* are uniformly distributed. At a distance of 23 m to 38 m along the coast sand dollars are found clumped in sandy mud substrate. The clumped and clustered distributional patterns of sand dollars are the most common in nature (Figure 2).

Table 1 Abundance and distribution patterns of sand dollars in Cemara Kecil Island

No	Species	Individu	RA (%)	(v)	(m)	DP
1	<i>L. centrale</i> (H.L. Clark, 1925)	384	17.5	0.07	0.17	Uniform
2	<i>L. laganum</i> (Leske, 1778)	661	30.0	0.06	0.30	Uniform
3	<i>L. fudsiyama</i> (Mortensen, 1948)	130	5.9	0.12	0.06	Clumped
4	<i>L. depressum</i> (L. Agassiz, 1841)	125	5.7	0.13	0.06	Clumped
5	<i>L. retinens</i> (Koehler, 1922)	145	6.6	0.12	0.07	Clumped
6	<i>L. boschi</i> (Jeannet & Martin, 1937)	124	5.6	0.13	0.06	Clumped
7	<i>L. depressum tonganense</i> (L. Agassiz, 1841)	141	6.4	0.12	0.06	Clumped
8	<i>L. dickersoni</i> (Israelsky, 1933)	149	6.8	0.12	0.07	Clumped
9	<i>L. decagonale rectum</i> (Gregory, 1892)	110	5.0	0.14	0.05	Clumped
10	<i>L. joubini</i> (Koehler, 1922)	121	5.5	0.13	0.06	Clumped
11	<i>L. decagonale</i> (Blainville, 1827)	110	5.0	0.14	0.05	Clumped
Σ		2 200	100			
H'		2.149				
E		0.896				

Note : RA : Relative Abundance; DP: Distribution Pattern; V: Variance; m: Mean Value




Figure 2 Geometry map showing sand dollars distribution and abundance in Cemara Kecil Island using PAST-3 (scale 1:5)




Figure 3 Sand dollars found on the Cemara Kecil Island, relative abundance and distribution pattern

3.2 Discussion

Sand dollars are abundantly found in the coastal area of Cemara Kecil Island which is known for its vast sloping white sand zone. This is naturally dominated by fine sand substrate in addition to some coral and seagrass beds. This type of substrate may become the underlying condition for the sand dollars abundance.

The abundance of sand dollars on Cemara Kecil Island Karimunjawa was obtained from the analysis

using PAST-3, and is classified into three groups based on the distance to the coastline:

- a. The distance of 0 m to 22 m
 The number of sand dollars within a distance of 0 m to 22 m from the coastline was found to be 632, which can be classified as medium abundance. This may be due to the coarse substrate from coral and shell fractions of some crustacean and bivalve. This finding is in line with a study discovering that the distribution of local sea fauna

is influenced by the habitats, a base substrate, the depth and zoning. Sand dollars of *Laganum laganum* kind was found in a depth of 1 m to 2 m [13].

- b. The distance of 23 m to 38 m
The abundance of sand dollars at a distance of 23 m to 38 m from the coast belongs to the highest category as many as 1 103 individuals were found. This is because sand dollars bury themselves in a substrate of fine sand or sandy mud and obtain the food by swallowing sand which is in the surrounding area. This finding is in line with a study indicating that sand dollars obtain the food by swallowing sand around them [14]. Sand dollars are deposit feeder organisms. Other study shows that sand dollars are organisms that prefer basic sandy mud substrate [15].
- c. The distance of 39 m to 50 m
The abundance of sand dollars at a distance of 39 m to 50 m from coastline belongs to the lowest category with 465 individuals. This is because the more distant from the coastline, the deeper the area whilst sand dollars have tendency to be buried in shallow water. Therefore sand dollars are found more in abundance in sandy areas and sometimes stranded on the coast [16].

The outer margin of these populations is well-defined in 4 m to 12 m of water, and here the largest individuals and greatest densities occur. Most of our observations are of these populations. The other type of distribution is similar, except that it extends into deeper water, and below about 10 m to 15 m; individuals become progressively smaller with depth [17].

3.2.1 Distributional Pattern

The distributional patterns of sand dollars on Cemara Kecil Island are dominated by clumped distributions, and some are random and uniform as in the case of *Laganum centrale* and *Laganum laganum*. The density of the sand dollars and organic matter from sediment is a major factor that regulates the density of sand dollars and muddy sand substrates as there is enough food and place to reproduce. When the density is high, sand dollars are less packed. On the other hand, when the density is low, the amount of organic matter, plankton and algae are eaten more. The proportions can be set by changing the organic content of the sediment [1, 13].

The types of clumped or clustered distributional pattern are the most commonly found in nature. This is because the sand dollars (*Laganum* sp.) tend to find a place suitable for them which then determines the distributional patterns [12].

The diversity index (H') of sand dollars is 2.149 which belongs to high category. According to Odum [12] the diversity index (H') 1.6 to 3.0 indicates high diversity. The high value of diversity can be acquired

if all individuals collected are samples of different species, and on the other hand, diversity is low if there is only one kind of species in the samples [18, 19].

The calculated index of uniformity (e) is 0.896 indicating that the uniformity of sand dollars of the island is high. Index of uniformity of the population (e) having the value > 0.6 is classified as high [12].

3.2.2 Water Quality Parameter

Water temperature on the island of Cemara Kecil is around 29 °C to 30 °C. In terms of the temperature range of the water, it can be seen that this water temperature meets the requirements for the survival of sand dollars. This is further strengthened by Page [19] who claimed that sand dollars can still live in water temperature range between 26 °C to 30 °C [19].

The salinity on the sampling site ranges from 32 ‰ to 33 ‰, which is suitable for sand dollars and also other marine biota. This is in line with the findings of a study that states that the ideal sea water salinity range is from 0 ‰ to 33 ‰. Salinity level change can make a significant difference to the properties of marine water and marine biota [3].

Sand dollars have a radially symmetrical body which is divided into five sections. The movement is slow with the vessel-like feet to regulate water pressure. Usually the animals bury themselves in the sandy substrate. According to Hawkes [20], the speed of either direct or indirect current affects the substrate base where the sand dollars community can be found.

The research site is mainly sandy and sand dollars can be found in the sandy substrates when the water recedes. Hawkes states that the substrate directly influences the composition and distribution of benthos, and serves as the habitat and the source of nutrients for the animals [20].

4.0 CONCLUSION

The composition of sand dollars on the island of Cemara Kecil consists of 11 patterns. The highest relative abundance is *L. laganum* (30 %) and the lowest is *L. decagonale rectum* (5 %) and *L. decagonale* (5 %).

The distributional patterns of sand dollars in Cemara Kecil Island are uniform distribution for *Laganum central* and *Laganum laganum*, while the rest of the animal belongs to clumped distribution.

Acknowledgement

The authors would like to thank the Staff of Balai Taman Nasional Karimunjawa and Ign. Boedi Hendarto for the valuable input and discussion in data processing.

References

- [1] Telford, M. 1981. A Hydrodynamic Interpretation of Sand Dollar Morphology. *Bulletin Marine Science*. 58(1): 605-622.
- [2] Dian R. A, R. Hartati and, A. Ambariyanto. 2005. *Identifikasi Sand Dollar dan Karakteristik Habitatnya di Pulau Cemara Besar, Kepulauan Karimunjawa Jepara*. [Identification Sand Dollars and the Characteristics of Their Habitat in Pulau Cemara Besar Karimunjawa Islands]. *Jurnal Ilmu Kelautan*. 10(1): 1-10. [Bahasa Indonesia].
- [3] Romimohtarto, K and S. Juwana. 2007. *Biologi Laut*. [Marine Biology]. Jakarta: Djambatan. [Bahasa Indonesia].
- [4] Sya'rani, L and A. Suryanto. 2006. *Gambaran Umum Kepulauan Karimunjawa*. [Overview Karimunjawa Island]. Semarang: Unssula Pres. [Bahasa Indonesia].
- [5] Suryanti, S and R. Ruswahyuni. 2014. *Perbedaan Kelimpahan Bulu Babi (Echinoidea) pada Ekosistem Karang dan Lamun di Pancuran Belakang, Karimunjawa, Jepara*. [The Difference in Abundance of Echinoideas on Coral Ecosystem and Seagrass Beds in Pancuran Belakang, Karimunjawa, Jepara]. *Jurnal Saintek Perikanan*. 10(1): 63-67. [Bahasa Indonesia].
- [6] Supono, S. 2009. *Aspek Biologi dan Ekologi Sand Dollar (Dendraster excentricus)*. [Biological and Ecological Aspects of Sand Dollars (*Dendraster excentricus*)]. *Jurnal OSEANA*. 34(4): 43-51. [Bahasa Indonesia].
- [7] Balai Taman Nasional Karimunjawa. 2008. *Data Base Karimunjawa Islands*. Semarang. Central Java. [Bahasa Indonesia].
- [8] Palys, T. 2008. Purposive Sampling. In Given, L. M. (ed.). *The Sage Encyclopedia of Qualitative Research Methods. Vol 2*. Los Angeles: Sage Publisher.
- [9] Fachrul, M. 2007. *Metode Sampling Bioekologi*. [Sampling Method Bioecology]. Jakarta: PT Bumi Aksara. [Bahasa Indonesia].
- [10] Hammer, Ø. and H. Bucher. 2006. Generalized Ammonoid Hydrostatics Modelling, with Application to *Intornites* and Intraspecific Variation in *Amaltheus*. *Paleontological Research*. 10: 91-96.
- [11] Hammer, Ø., D. A. T. Harper, and P. D. Ryan. 2001. PAST: Paleontological Statistics Software Package for Education and Data Analysis. *Palaeontologia Electronica*. 4(1): 1-9.
- [12] Odum. 1993. *Fundamental of Ecology*. Yogyakarta: Gadjah Mada University Press.
- [13] Aziz, A. 1981. *Fauna Echinodermata dari Terumbu Karang Pulau Pari, Pulau-pulau Seribu*. [Fauna of Echinoderms of Coral Reefs, Pari Island, Pulau-pulau Seribu]. *Oseanologi di Indonesia*. 14(1): 41-50. [Bahasa Indonesia].
- [14] Brotowidjoyo, M. D. 1994. *Zoologi Dasar* [Fundamental of Zoology]. Yogyakarta: Erlangga.
- [15] Merrill, R. J. and E. S. Hobson. 1970. Field Observations of *Dendraster excentricus*, a Sand Dollars of Western North America. *American Midland Naturalist*. 83(2): 595-624.
- [16] Frodrie. 2015. Feeding and Growth. Science. [Online]. From: <http://www.asnailsodyssey.com>. [Accessed on 15 July 2015].
- [17] Smith, A. L. 1981. Comparison of Macrofaunal Invertebrates in Sand Dollar (*Dendraster excentricus*) Beds and in Adjacent Areas Free of Sand Dollars. *Marine Biology*. 65(2): 191-198.
- [18] Taghon, G. L. 1982. Optimal Foraging by Deposit-Feeding Invertebrates: Roles of Particle Size and Organic Coating. *Oecologia*. 52(3): 295-304.
- [19] Page, C. 2000. The Common Sand Dollar. [Online]. From: <http://octopus.gma.org/Tidings/sanddollar.html>. [Accessed on 15 April 2015].
- [20] Hawkes, T. 1978. *Structuralism and Semiotics*. Great Britain: Richard Clay Ltd, Bungay, Suffolk.



CERTIFICATE

Is hereby presented to

Suryanti

AS **Presenter** AT

THE 2nd INTERNATIONAL SYMPOSIUM
ON AQUATIC PRODUCTS PROCESSING AND HEALTH (ISAPPROSH)

Diponegoro University, Semarang, Indonesia. September 13-15, 2015

Director Generale
of Marine and Fisheries Products
Competitiveness Enhancement

Ir. R. Nilanto Perbowo, M.Sc



Head of Indonesian
Fisheries Product Processing
Society

Prof. Dr. Hari Eko Irianto, M.Sc