

KORESPONDENSI PAPER

JUDUL : NITROGEN CONCENTRATION IN CORAL REEF: THE CONTRIBUTION OF CORAL FORMS AND NITROGEN

JURNAL : SAINTEK PERIKANAN - UNIVERSITAS DIPONEGORO

Status : Jurnal Nasional Terakreditasi SINTA 2

No	AKTIVITAS	TANGGAL	HALAMAN
1	Submission Artikel	10 Mei 2020	1-7
2	Pengiriman Hasil Review	2 Juni 2020	8 -14
3	Pengiriman hasil review OJS	2 Juni 2020	15 - 18
3	Hasil Perbaikan Artikel	3 Juni 2020	19 - 24
4	Letter of Acceptend	15 Juni 2020	25
5	Copy Editing Review Request	22 Juni 2020	26
6	Artikel Publish	04 Agustus 2020	27
7	Final Artikel	Vol. 16 No. 2 Agustus 2020	28 - 32

Submission Artikel

Search mail

4 of 1,437

[IJFST] Submission Acknowledgement Inbox x

Saintek Perikanan : Indonesian Journal of Fisheries Science via ejournal.undip.ac.id
to S.PI

Sun, May 10, 2020, 11:30 AM

Be careful with this message

This may be a spoofed message. The message claims to have been sent from your account, but Gmail couldn't verify the actual source. Avoid clicking links or replying with sensitive information, unless you are sure you actually sent this message. (No need to reset your password, the real sender does not actually have access to your account!)

Report spam Looks safe

Suryanti Suryanti :

Thank you for submitting the manuscript, "NITROGEN CONCENTRATION IN CORAL REEF: THE CONTRIBUTION OF CORAL FORMS AND NITROGEN FIXING BACTERIA IN KARIMUNJAWA ISLANDS" to Saintek Perikanan : Indonesian Journal of Fisheries Science and Technology. With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL: <https://ejournal.undip.ac.id/index.php/saintek/author/submission/40087>
Username: drsuryanti

Nitrogen Concentration In Coral Reef: The Contribution of Coral Forms and Nitrogen Fixing Bacteria In Karimunjawa Islands

Suryanti Suryanti^{1*}, Supriharyono Supriharyono¹, Sigit Febrianto¹, Muslidin Aini²

¹ Department of Aquatic Resources, Faculty of Fisheries and Marine Science, Universitas Diponegoro, Semarang, Indonesia

Jl. Prof. Soedarto, SH Tembalang, Semarang, Central Java, Indonesia 50275

²Research Center, Coastal Environmental and Fisheries, Jl. RA Kartini no 23, Rakam, Selong, Lombok Timur, NTB. *Indonesia* . 83619

* Corresponding author's e-mail: suryantidr@gmail.com

Abstract.

Coral is ecosystem in the coastal area which plays an important role in controlling nutrient cycle in the sea. This research aimed to study the the nitrogen concentration in coral and to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in coral. The research was carried out by field observation, by taking coral reef samples and analyzed in the laboratory for its nitrogen concentration and number of nitrogen fixing bacteria. Data analysis was carried out by ANOVA and regression. The analysis resulted that there was a significant difference of nitrogen concentration in massive and branching corals. However, there is no significant correlatio between the sampling location and types of coral on the density of nitrogen fixing bacteria, both *Nitrosomonas* sp. and *Nitrobacter* sp. Regression analysis showed that the concentration of nitrogen in coral tissue was not significantly related to the density of nitrogen fixing bacteria. The result of this research implies that the concentration of nitrogen in the coral tissue may be affected by many other factors.

Keywords: bacteria, coral, fixation, nitrogen, tissue

Introduction

As the development of coastal area increase, the problems related to the coastal ecosystem arises. This includes the coral reef ecosystem which has the most important role in supporting marine biodiversity. Coral also provides micro-biome which shows the relationship among organisms in the coral surface. Various processes occur in the micro-biome, such as sulphur cycling, nitrogen fixing, as well as the it defensive system (Krediet, Ritchie, Paul, & Teplitski, 2013).

Coral is a rocky organism that usually lives in the shallow marine ecosystem ranging from the tropical to sub-tropical area. Ecologically, shallow coral reefs are considered as the huge source of benthic nitrogen fixation (Diaz, 2017). The biodiversity in coral reef ecosystem is supported by the coral associated organisms, such as sulphur recycling bacteria or nitrogen fixing bacteria. All of them existed in the coral ecosystem and provide defense toward various threats.

As a living organism, coral absorbs various minerals to grow. One of the important nutrient for coral is nitrogen. Nitrogen for coral is mainly provided by zooxanthelae which which lives mutualistically with coral polyp (Krediet et al., 2013). Coral polyp is where various ecological processes occur. Various physiological response due to the arising stress in the environment is detected and responded in the polyp (Hohn & Merico, 2012).

Symbiodinium is the algae species which has mutualistic association with coral. However, the micro-biome in coral is just recognized recently. The micro-biome is the system which provide support to the organization of coral ecoystems which involved various microorganisms (Rädecker, Pogoreutz, Voolstra, Wiedenmann, & Wild, 2015). There are many function that the microbiome provide. Among the functions the microbiome provides, nitrogen cycling is considered as the most important process for coral sustainability.

There are several studies which found that coral polyp also contain nitrogen fixing bacteria. Nitrogen is an important element required by any plant to grow. However, in some cases the availability of nitrogen in the coral reef ecosystem is limited. In this condition, nitrogen fixing is the main source for coral to support its growth (Putnam, Barott, Ainsworth, & Gates, 2017). The nitrogen fixing bacteria are frequently found in coral mucus (Lema, Willis, & Bourne, 2012).

The nitrogen fixing bacteria is commonly called diazotrophs. Nitrogen fixing bacteria are mostly found in a reef building symbiotic coral as the source of nitrogen for coral's growth (Wang et al., 2015). The bacteria favors the nitrogen supply to the coral organism directly through bacterivory, through polyp and zooxanthellae (Diaz, 2017). The species of nitrogen fixing bacteria is varied among coral species and locations. At least there are 23 species of nitrogen fixing bacteria. Nitrosomonas and Nitrobacter are frequently used as bioremediator for aquaculture. It is mostly functioned as the remediator for ammonia and nitrite (Barik, Ram, Halder, & Vardya, 2018).

Critical concern on the microbiological role on the coral reef ecosystem arises the question whether the associated microorganism also affect the nutrient uptake by coral reef. Least information is known regarding the processes and impact of the nitrogen fixing bacteria to the livelihood of coral community. This research aimed to study the nitrogen concentration in coral and to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in coral.

Methods

The research was carried out in June – July 2019 in the area of Karimunjawa National Park. The research was carried out by field observation. The research stations were located in Karimunjawa island, Menjangan Kecil island, and Cemara Kecil island, while laboratory analysis was in the *Tropical Marine Biotechnology (TMB), Faculty of Fisheries and Marine Science Diponegoro University Semarang*.

There were three sampling stations occupied in the research. Each station consisted of three sampling spots. The objects of the research were including the concentration of Nitrogen in coral organisms and the density of nitrogen fixing bacteria. Coral samples were obtained from two groups of coral organisms, including the massive and branching corals. Thus, the research design applied 3 x 2 factorial treatment.

The coral was sampled with the length of 10 cm. The coral samples were stored in a cool box and identified in the laboratory for its density of nitrogen fixing bacteria and the concentration of nitrogen. The bacteria were isolated from the coral and placed in the enrichment media (nutrient agar). The bacteria collected as the object of the research was Nitrosomonas sp. and Nitrobacter sp.

Data analysis was carried out with ANOVA and regression. Multivariate ANOVA was carried out to analyze the difference of nitrogen concentration and the abundance of nitrogen fixing bacteria. Regression was carried out to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in the coral. The regression analysis was carried out through multiple regression to analyze the simultaneous effect of nitrogen fixing bacteria, and curve estimation to analyze the effect of nitrogen fixing bacteria separately. Analysis was carried out by SPSS.

Results

The result of the laboratory analysis showed that there was variation on the concentration of nitrogen in coral samples. Generally, the concentration of nitrogen obtained from massive coral was lower than that in the branching coral. The concentration of nitrogen in Cemara Kecil island was also higher than in Karimunjawa and Menjangan Kecil for respective types of coral. Detailed result of laboratory analysis and the statistical analysis for nitrogen concentration is presented in Table 1.

Table 1. Average concentration of nitrogen in coral (mg/kg) from each sample

No.	Location	Types of Coral		Total by Location
		Massive	Branching	
1.	Karimunjawa	61.380 ^{*a}	231.527 ^{‡a}	146.453 ^{‡p}
2.	Menjangan Kecil	73.662 ^{*a}	197.390 ^{‡a}	135.526 ^{‡p}

3.	Cemara Kecil	160.223 ^{‡a}	221.486 ^{‡a}	190.855 ^{‡p}
	<i>Total by Types of Coral</i>	98.422 ^{*x}	216.801 ^{*y}	

Notation: * = very low

‡ = low

‡ = moderate

Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Table 1 shows that the average concentration of nitrogen in the coral was ranging from very low to moderate. However, moderate nitrogen concentration was obtained from branching coral, while very low concentration was obtained from massive coral. The concentration of nitrogen is considered as very low within the range $< 0.1\%$, low within the range $0.1 - 0.2$ and moderate within the range $0.2 - 0.5$ (Wahyuni, Kusnadi, & Honorita, 2017). Thus, it can be noted that massive coral has a lower accumulation of nitrogen.

Statistical data analysis showed that the nitrogen concentration was affected by the types of coral. Massive coral tended to have lower nitrogen concentration than the branching coral. However, other treatments did not show any significant effect on the nitrogen concentration. However, the difference in nitrogen concentration affected by location, and types of coral altogether was noticeable. Data analysis resulted F value for location was 2.311 ($p = 0.142$), types of coral showed the value of 28.285 ($p = 0.000$), and combination of both 2.009 ($p = 0.251$).

Table 2. Average density of nitrogen fixing bacteria *Nitrosomonas* sp. (cfu/g) from each sample

No.	Location	Types of Coral		
		Massive	Branching	Total
1.	Karimunjawa	$58.27 \times 10^{5(a)}$	$39.93 \times 10^{5(a)}$	$49.10 \times 10^{5(p)}$
2.	Menjangan Kecil	$11.30 \times 10^{6(a)}$	$15.00 \times 10^{6(a)}$	$13.15 \times 10^{6(p)}$
3.	Cemara Kecil	$11.07 \times 10^{6(a)}$	$64.33 \times 10^{5(a)}$	$87.50 \times 10^{5(p)}$
	<i>Total</i>	$93.98 \times 10^{5(x)}$	$84.76 \times 10^{5(x)}$	

Notation: Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Statistical analysis showed that there was no significant effect of the source of coral samples and the types of coral to the density of *Nitrosomonas* sp. Data analysis with univariate ANOVA showed that the F value obtained from the location, types of coral and combination of both were 2.847 ($p = 0.097$), 0.107 ($p = 0.749$) and 0.753 ($p = 0.492$) respectively. Thus, it can be suggested that massive coral provided better support to the nitrogen fixing bacteria. The environmental condition of Menjangan Kecil island was also more supportive than the other two islands.

The average density of nitrogen fixing bacteria *Nitrobacter* sp. is the last parameter observed in this research. Assessment on the average density of *Nitrobacter* sp. showed that there was slight difference between the massive and branching corals. However, branching coral had higher *Nitrobacter* sp. density even though the difference was not significant. Noticeable difference was obtained from the bacterial density in Karimunjawa island. The difference was obtained not only from the total average density, but also in the density of respective massive and branching coral types. Table 3 presents the detailed *Nitrobacter* sp. density of the samples obtained from the research location.

Table 3. Average density of nitrogen fixing bacteria *Nitrobacter* sp. (cfu/gr) from each sample

No.	Location	Types of Coral		
		Massive	Branching	Total
1.	Karimunjawa	$96.57 \times 10^{5(a)}$	$59.43 \times 10^{5(a)}$	$78.00 \times 10^{5(p)}$
2.	Menjangan Kecil	$15.00 \times 10^{6(a)}$	$19.67 \times 10^{6(a)}$	$17.33 \times 10^{6(p)}$

3.	Cemara Kecil	15.33x10 ^{6(a)}	16.90x10 ^{6(a)}	16.12x10 ^{6(p)}
	Total	13.33x10 ^{6(x)}	14.17x10 ^{6(x)}	

Notation: Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Table 3 shows that the density of *Nitrobacter* sp. in Karimunjawa National Park was higher than the *Nitrosomonas* sp. However, there was no significant difference in the density of nitrogen fixing bacteria *Nitrobacter* sp. both partially and simultaneously. Statistical analysis with ANOVA showed the F value of coral source, types of coral and combination of both were as much as 1.877 ($p = 0.195$), 0.037 ($p = 0.851$) and 0.313 ($p = 0.737$) respectively.

Among the results from the assessment of nitrogen concentration as well as the density of *Nitrosomonas* sp. and *Nitrobacter* sp., it can be suggested that Karimunjawa always had the lowest value. This could be due to the environmental condition of the island. However, further indepth research is needed to study the possible causes.

The result of statistical tests showed that the effect of nitrogen fixing bacteria showed that the concentration of nitrogen in the coral was not related to the density of *Nitrosomonas* sp. nor *Nitrobacter* sp. Both multiple regression and curve estimation analysis did not show the significance of the relationship. Thus, it can be concluded that the concentration of nitrogen in coral is affected by some other factor, or involves some other factors as moderator or as cooperating factor.

Discussion

The accumulation of nitrogen in massive and branching coral is suggested as the effect of growth rate between the two. Nitrogen is required for coral metabolism. However, the environmental condition significantly affect the concentration of nitrogen. Shaded environment tends to cause the increase of the C/N ratio in coral. Moreover, the concentration of nitrogen is decreased along with the increasing depth (Alamaru, Loya, Brokovich, Yam, & Shemesh, 2009).

The size of coral is related to its capability to tolerate environmental stress (Edmunds & Burgess, 2016). However, each coral species has its defensive method to overcome the environmental stress. The response of coral toward environmental disturbance also involved the metabolic system provided by coral's microstructure (Delgadillo-Nuño, Liñán-Cabello, Reyes-Gómez, & Soriano-Santiago, 2014).

Assessment of the abundance of nitrifying bacteria *Nitrosomonas* sp. showed that massive coral had higher average density. The collected data as presented in Table 2 showed that higher density of *Nitrosomonas* sp. in massive coral was obtained from all locations. Furthermore, the density of bacteria from the coral sample obtained from Menjangan Kecil island was the highest, while Karimunjawa provided the lowest density.

The sustainability of coral organisms relies on the nutrient supply. The symbiotic between coral and the symbiodinium is highly reliable to the availability of nitrogen (Rädecker et al., 2015). The significant difference between nitrogen concentration in massive and branching coral resulted in this research can be caused by the absorption rate rather than the nitrification process. Higher nitrogen concentration in the branching corals may be caused by polyp abundance. Branching coral tended to have larger surface due to its three dimensional form, while the massive coral only has the top surface. Thus, since the sample was collected in the same size, the abundance of coral polyp in the branching coral was more than the massive coral. Thus, it affected the nitrogen absorption.

The abundance of nitrogen fixing bacteria indicates its nitrification rate provided by the community (Zulfarina, Rusmana, Mubarik, & Santosa, 2017). It also indicates the potential nitrogen produced from the nitrification process. However, the research found that there was no significant effect of the density of *Nitrosomonas* sp. and *Nitrobacter* sp. to the concentration of nitrogen in coral tissue. The nitrification process occurred in the polyp surface, but it mainly affects the nitrogen availability in the environment. Unfortunately, the high nitrification rate is not followed by higher nitrogen uptake by coral organism.

The difference in the density of nitrogen fixing bacteria is dominantly caused by the environmental condition. One of the affecting factors of nitrifying bacteria is the depth of which the bacteria sampled. The abundance of nitrifying bacteria is higher in the shallow water (Smith, Damashek, Chavez, & Francis, 2016). Unfortunately, measurement on the depth of the sampling location was not carried out.

The bacteria is sensitive to the change of environmental condition, such a temperature (Diaz, 2017). At the research location, Karimunjawa island is the most exploited area. Thus, anthropologic aspects may play a role in changing the coral waters. However, this should be assured by investigate the condition of coral reef ecosystem. Any environmental stress may break the symbiosis of coral and algae (zooxanthellae) (Wooldridge, 2013). This may affect the other associated microorganisms in the polyp.

Nitrosomonas and Nitrobacter are frequently used as bioremediator in aquaculture. However, the preferences of the two is not much understood. Related to the nitrogen fixing process in Karimunjawa National Park, Nitrosomonas and Nitrobacter might have an important role in remediating coastal environment. The density of both bacterias may be related to the high input of ammonia since Karimunjawa island has been much exploited for tourism activities. Nitrosomonas and Nitrobacter mainly act as bioremediator of both pollutant (Barik et al., 2018).

Even though nitrogen fixing bacteria exists in the coral mucus or tissue, but it's suggested that the bacteria are associated with Symbiodinium rather than directly to the coral (Lema et al., 2012). This explains why the Nitrosomonas sp. and Nitrobacter sp. did not influence the concentration of nitrogen significantly.

The role of nitrogen fixing bacteria is obvious in the nitrogen limited environment (Putnam et al., 2017). However, since the effect of Nitrosomonas sp. and Nitrobacter sp. in Karimunjawa National Park remains insignificant, it can be concluded that the source of nitrogen is sufficient. Or otherwise, since the coral growth requires nitrogen, the coral can not accumulate the nitrogen it consumed.

Conclusion


Based on the analysis result, the concentration of nitrogen is higher in branching coral as the impact of its higher growth rate. However, the concentration of nitrogen in coral tissue is not affected by the abundance of nitrogen fixing bacteria.

References

- Alamaru, A., Loya, Y., Brokovich, E., Yam, R., & Shemesh, A. (2009). Carbon and nitrogen utilization in two species of Red Sea corals along a depth gradient: Insights from stable isotope analysis of total organic material and lipids. *Geochimica et Cosmochimica Acta*, 73, 5333–5342.
- Barik, P., Ram, R., Haldar, C., & Vardia, H. K. (2018). Study on nitrifying bacteria as bioremediator of ammonia in simulated aquaculture system. *Journal of Entomology and Zoology Studies*, 6, 1200–1206.
- Delgadillo-Nuño, M. A., Liñán-Cabello, M. A., Reyes-Gómez, J., & Soriano-Santiago, O. (2014). Response to pH stress in the reef-building coral *Pocillopora capitata* (Anthozoa: Scleractinia). *Revista de Biología Marina y Oceanografía*, 49, 449–459.
- Diaz, L. (2017). Response of the coral associated nitrogen fixing bacteria toward elevated water temperature. *Journal of Water Resources and Ocean Science*, 6, 98–109.
- Edmunds, P. J., & Burgess, S. C. (2016). Size-dependent physiological responses of the branching coral *Pocillopora verrucosa* to elevated temperature and PCO₂. *The Journal of Experimental Biology*, 219, 3896–3906.
- Hohn, S., & Merico, A. (2012). Modelling coral polyp calcification in relation to ocean acidification. *Biogeosciences*, 9, 4441–4454.
- Krediet, C. J., Ritchie, K. B., Paul, V. J., & Teplitski, M. (2013). Coral-associated micro-organisms and their roles in promoting coral health and thwarting diseases. *Proceedings of the Royal Society B: Biological Sciences*, 280, 1755.
- Lema, K. A., Willis, B. L., & Bourne, D. G. (2012). Corals form characteristic associations with symbiotic nitrogen-fixing bacteria. *Applied and Environmental Microbiology*, 78, 3136–3144.

- Putnam, H. M., Barott, K. L., Ainsworth, T. D., & Gates, R. D. (2017). The vulnerability and resilience of reef-building corals. *Current Biology*, *27*, R528–R540.
- Rädecker, N., Pogoreutz, C., Voolstra, C. R., Wiedenmann, J., & Wild, C. (2015). Nitrogen cycling in corals: the key to understanding holobiont functioning? *Trends in Microbiology*, *23*, 490–497.
- Smith, J. M., Damashek, J., Chavez, F. P., & Francis, C. A. (2016). Factors influencing nitrification rates and the abundance and transcriptional activity of ammonia-oxidizing microorganisms in the dark northeast Pacific Ocean. *Limnology and Oceanography*, *61*, 596–609.
- Wahyuni, T., Kusnadi, H., & Honorita, B. (2017). Status unsur hara karbon organik dan nitrogen tanah sawah tiga Kabupaten di Provinsi Bengkulu. In S. Herlinda (Ed.), *Prosiding Seminar Nasional Lahan Suboptimal* (pp. 978–979). Palembang.
- Wang, X. T., Sigman, D. M., Cohen, A. L., Sinclair, D. J., Sherrell, R. M., Weigand, M. A., ... Ren, H. (2015). Isotopic composition of skeleton-bound organic nitrogen in reef-building symbiotic corals: A new method and proxy evaluation at Bermuda. *Geochimica et Cosmochimica Acta*, *148*, 179–190.
- Wooldridge, S. A. (2013). Breakdown of the coral-algae symbiosis: towards formalising a linkage between warm-water bleaching thresholds and the growth rate of the intracellular zooxanthellae. *Biogeosciences*, *10*, 1647–1658.
- Zulfarina, Rusmana, I., Mubarik, N. R., & Santosa, D. A. (2017). The abundance of nitrogen fixing, nitrifying, denitrifying and ammonifying bacteria in the soil of tropical rainforests and oil palm plantations in Jambi. *Makara Journal of Science*, *21*, 187–194.

HASIL REVIEW



The screenshot shows an email client interface. At the top, there is a search bar with the text "Search mail" and a hamburger menu icon. Below the search bar is a toolbar with various icons for email actions like back, forward, delete, and search. The email title is "[IJFST] Hasil Review Artikel" with a yellow envelope icon and "Inbox x". The sender is "Saintek jurnal" with the email address "saintekjurnal@gmail.com" and a profile picture. The recipient is "to suryanti" and the location is "di Tempat". The date and time are "Tue, Jun 2, 2020, 10:15 AM". The main body of the email contains a review comment in Indonesian, followed by the journal's contact information. At the bottom, there are two attachments.

Search mail

[IJFST] Hasil Review Artikel Inbox x

Saintek jurnal <saintekjurnal@gmail.com>
to suryanti
di Tempat

Tue, Jun 2, 2020, 10:15 AM

Dengan Hormat, berikut kami kirimkan hasil review artikel saudara dengan judul "NITROGEN CONCENTRATION IN CORAL REEF: THE CONTRIBUTION OF CORAL FORMS AND NITROGEN FIXING BACTERIA IN KARIMUNJAWA ISLANDS". Mohon artikel dapat diperbaiki sesuai dengan sarat dari reviewer dan dikirim kembali ke Redaksi Saintek perikanan. Terima kasih

Saintek Perikanan: Indonesian Journal of Fisheries Science and Technology
Fakultas Perikanan dan Ilmu Kelautan
Universitas Diponegoro
Jl. Prof. H. Soedarto SH Tembalang Semarang
Telp/hp: (024) 76480685/ 085536649493

2 Attachments

Nitrogen Concentration In Coral Reef: The Contribution of Coral Forms and Nitrogen Fixing Bacteria In Karimunjawa Islands

Suryanti Suryanti^{1*}, Supriharyono Supriharyono¹, Sigit Febrianto¹, Muslidin Aini²

¹ Department of Aquatic Resources, Faculty of Fisheries and Marine Science, Universitas Diponegoro, Semarang, Indonesia

Jl. Prof. Soedarto, SH Tembalang, Semarang, Central Java, Indonesia 50275

²Research Center, Coastal Environmental and Fisheries, Jl. RA Kartini no 23, Rakam, Selong, Lombok Timur, NTB. Indonesia . 83619

* Corresponding author's e-mail: suryantidr@gmail.com

Abstract.

Coral is ecosystem in the coastal area which plays an important role in controlling nutrient cycle in the sea. This research aimed to study the the nitrogen concentration in coral and to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in coral. The research was carried out by field observation, by taking coral reef samples and analyzed in the laboratory for its nitrogen concentration and number of nitrogen fixing bacteria. Data analysis was carried out by ANOVA and regression. The analysis resulted that there was a significant difference of nitrogen concentration in massive and branching corals. However, there is no significant correlatio between the sampling location and types of coral on the density of nitrogen fixing bacteria, both Nitrosomonas sp. and Nitrobacter sp. Regression analysis showed that the concentration of nitrogen in coral tissue was not significantly related to the density of nitrogen fixing bacteria. The result of this research implies that the concentration of nitrogen in the coral tissue may be affected by many other factors.

Keywords: bacteria, coral, fixation, nitrogen, tissue

Please make two column

Introduction **UPCRASE (times new roman, 10 space 1)**

As the development of coastal area increase, the problems related to the coastal ecosystem arises. This includes the coral reef ecosystem which has the most important role in supporting marine biodiversity. Coral also provides micro-biome which shows the relationship among organisms in the coral surface. Various processes occur in the micro-biome, such as sulphur cycling, nitrogen fixing, as well as the it defensive system (Krediet, Ritchie, Paul, & Teplitski, 2013).

Coral is a rocky organism that usually lives in the shallow marine ecosystem ranging from the tropical to sub-tropical area. Ecologically, shallow coral reefs are considered as the huge source of benthic nitrogen fixation (Diaz, 2017). The biodiversity in coral reef ecosystem is supported by the coral associated organisms, such as sulphur recycling bacteria or nitrogen fixing bacteria. All of them existed in the coral ecosystem and provide defense toward various threats.

As a living organism, coral absorbs various minerals to grow. One of the important nutrient for coral is nitrogen. Nitrogen for coral is mainly provided by zooxanthelae which which lives mutualistically with coral polyp (Krediet et al., 2013). Coral polyp is where various ecological processes occur. Various physiological response due to the arising stress in the environment is detected and responded in the polyp (Hohn & Merico, 2012).

Symbiodinium is the algae species which has mutualistic association with coral. However, the micro-biome in coral is just recognized recently. The micro-biome is the system which provide support to the organization of coral ecoystems which involved various microorganisms (Rädecker, Pogoreutz, Woolstra, Wiedenmann, & Wild, 2015). There are many function that the microbiome provide. Among the

functions the microbiome provides, nitrogen cycling is considered as the most important process for coral sustainability.

There are several studies which found that coral polyp also contain nitrogen fixing bacteria. Nitrogen is an important element required by any plant to grow. However, in some cases the availability of nitrogen in the coral reef ecosystem is limited. In this condition, nitrogen fixing is the main source for coral to support its growth (Putnam, Barott, Ainsworth, & Gates, 2017). The nitrogen fixing bacteria are frequently found in coral mucus (Lema, Willis, & Bourne, 2012).

The nitrogen fixing bacteria is commonly called diazotrophs. Nitrogen fixing bacteria are mostly found in a reef building symbiotic coral as the source of nitrogen for coral's growth (Wang et al., 2015). The bacteria favors the nitrogen supply to the coral organism directly through bacterivory, through polyp and zooxanthellae (Diaz, 2017). The species of nitrogen fixing bacteria is varied among coral species and locations. At least there are 23 species of nitrogen fixing bacteria. *Nitrosomonas* and *Nitrobacter* are frequently used as bioremediator for aquaculture. It is mostly functioned as the remediator for ammonia and nitrite (Barik, Ram, Halder, & Vardya, 2018).

Critical concern on the microbiological role on the coral reef ecosystem arises the question whether the associated microorganism also affect the nutrient uptake by coral reef. Least information is known regarding the processes and impact of the nitrogen fixing bacteria to the livelihood of coral community. This research aimed to study the nitrogen concentration in coral and to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in coral.

Methods RESEARCH METHODS

The research was carried out in June – July 2019 in the area of Karimunjawa National Park. The research was carried out by field observation. The research stations were located in Karimunjawa island, Menjangan Kecil island, and Cemara Kecil island, while laboratory analysis was in the *Tropical Marine Biotechnology (TMB), Faculty of Fisheries and Marine Science Diponegoro University Semarang*.

There were three sampling stations occupied in the research. Each station consisted of three sampling spots. The objects of the research were including the concentration of Nitrogen in coral organisms and the density of nitrogen fixing bacteria. Coral samples were obtained from two groups of coral organisms, including the massive and branching corals. Thus, the research design applied 3 x 2 factorial treatment.

Data collection was carried out through field sampling followed

The coral was sampled with the length of 10 cm. The coral samples were stored in a cool box and identified in the laboratory for its density of nitrogen fixing bacteria and the concentration of nitrogen. The bacteria were isolated from the coral and placed in the enrichment media (nutrient agar). The bacteria collected as the object of the research was *Nitrosomonas* sp. and *Nitrobacter* sp.

Data analysis was carried out with ANOVA and regression. *Multivariate* ANOVA was carried out to analyze the difference of nitrogen concentration and the abundance of nitrogen fixing bacteria. Regression was carried out to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in the coral. The regression analysis was carried out through multiple regression to analyze the simultaneous effect of nitrogen fixing bacteria, and curve estimation to analyze the effect of nitrogen fixing bacteria separately. Analysis was carried out by SPSS.

Results

The result of the laboratory analysis showed that there was variation on the concentration of nitrogen in coral samples. Generally, the concentration of nitrogen obtained from massive coral was lower than that in the branching coral. The concentration of nitrogen in Cemara Kecil island was also higher than in Karimunjawa and Menjangan Kecil for respective types of coral. Detailed result of laboratory analysis and the statistical analysis for nitrogen concentration is presented in Table 1.

Table 4. Average concentration of nitrogen in coral (mg/kg) from each sample **(Capital Each Word)**

please format the table according to the template

No.	Location	Types of Coral		Total by Location
		Massive	Branching	
1.	Karimunjawa	61.380 ^{*a}	231.527 ^{‡a}	146.453 ^{‡p}
2.	Menjangan Kecil	73.662 ^{*a}	197.390 ^{‡a}	135.526 ^{‡p}
3.	Cemara Kecil	160.223 ^{‡a}	221.486 ^{‡a}	190.855 ^{‡p}
	Total by Types of Coral	98.422 ^{*x}	216.801 ^{‡y}	

Notation: * = very low

‡ = low

‡ = moderate

Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Table 1 shows that the average concentration of nitrogen in the coral was ranging from very low to moderate. However, moderate nitrogen concentration was obtained from branching coral, while very low concentration was obtained from massive coral. The concentration of nitrogen is considered as very low within the range $< 0.1\%$, low within the range $0.1 - 0.2$ and moderate within the range $0.2 - 0.5$ (Wahyuni, Kusnadi, & Honorita, 2017). Thus, it can be noted that massive coral has a lower accumulation of nitrogen.

Statistical data analysis showed that the nitrogen concentration was affected by the types of coral. Massive coral tended to have lower nitrogen concentration than the branching coral. However, other treatments did not show any significant effect on the nitrogen concentration. However, the difference in nitrogen concentration affected by location, and types of coral altogether was noticeable. Data analysis resulted F value for location was 2.311 ($p = 0.142$), types of coral showed the value of 28.285 ($p = 0.000$), and combination of both 2.009 ($p = 0.251$).

please add description Highest nitrifying bacteria rating in each location

Table 5. Average density of nitrogen fixing bacteria *Nitrosomonas* sp. (cfu/g) from each sample **(Capital Each Word)**

please format the table according to the template

No.	Location	Types of Coral		
		Massive	Branching	Total
1.	Karimunjawa	58.27x10 ^{5(a)}	39.93x10 ^{5(a)}	49.10x10 ^{5(p)}
2.	Menjangan Kecil	11.30x10 ^{6(a)}	15.00x10 ^{6(a)}	13.15x10 ^{6(p)}
3.	Cemara Kecil	11.07x10 ^{6(a)}	64.33x10 ^{5(a)}	87.50x10 ^{5(p)}
	Total	93.98x10 ^{5(x)}	84.76x10 ^{5(x)}	

Notation: Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Statistical analysis showed that there was no significant effect of the source of coral samples and the types of coral to the density of *Nitrosomonas* sp. Data analysis with univariate ANOVA showed that the F value obtained from the location, types of coral and combination of both were 2.847 ($p = 0.097$), 0.107 ($p = 0.749$) and 0.753 ($p = 0.492$) respectively. Thus, it can be suggested that massive coral provided better support to the nitrogen fixing bacteria. The environmental condition of Menjangan Kecil island was also more supportive than the other two islands.

The average density of nitrogen fixing bacteria *Nitrobacter* sp. is the last parameter observed in this research. Assessment on the average density of *Nitrobacter* sp. showed that there was slight difference between the massive and branching corals. However, branching coral had higher *Nitrobacter* sp. density even though the difference was not significant. Noticeable difference was obtained from the bacterial density in Karimunjawa island. The difference was obtained not only from the total average density, but also in the density of respective massive and branching coral types. Table 3 presents the detailed *Nitrobacter* sp. density of the samples obtained from the research location.

Table 6. Average density of nitrogen fixing bacteria *Nitrobacter* sp. (cfu/gr) from each sample (Capital Each Word)

please format the table according to the template

No.	Location	Types of Coral		
		Massive	Branching	Total
1.	Karimunjawa	96.57x10 ^{5(a)}	59.43x10 ^{5(a)}	78.00x10 ^{5(p)}
2.	Menjangan Kecil	15.00x10 ^{6(a)}	19.67x10 ^{6(a)}	17.33x10 ^{6(p)}
3.	Cemara Kecil	15.33x10 ^{6(a)}	16.90x10 ^{6(a)}	16.12x10 ^{6(p)}
	Total	13.33x10 ^{6(x)}	14.17x10 ^{6(x)}	

Notation: Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Table 3 shows that the density of *Nitrobacter* sp. in Karimunjawa National Park was higher than the *Nitrosomonas* sp. However, there was no significant difference in the density of nitrogen fixing bacteria *Nitrobacter* sp. both partially and simultaneously. Statistical analysis with ANOVA showed the F value of coral source, types of coral and combination of both were as much as 1.877 ($p = 0.195$), 0.037 ($p = 0.851$) and 0.313 ($p = 0.737$) respectively.

Among the results from the assessment of nitrogen concentration as well as the density of *Nitrosomonas* sp. and *Nitrobacter* sp., it can be suggested that Karimunjawa always had the lowest value. This could be due to the environmental condition of the island. However, further in-depth research is needed to study the possible causes.

~~The result of statistical tests showed that the effect~~ of nitrogen fixing bacteria showed that the concentration of nitrogen in the coral was not related to the density of *Nitrosomonas* sp. nor *Nitrobacter* sp. Both multiple regression and curve estimation analysis did not show the significance of the relationship. Thus, it can be concluded that the concentration of nitrogen in coral is affected by some other factor, or involves some other factors as moderator or as cooperating factor.

Discussion

The accumulation of nitrogen in massive and branching coral is suggested as the effect of growth rate between the two. Nitrogen is required for coral metabolism. However, the environmental condition significantly affects the concentration of nitrogen. Shaded environment tends to cause the increase of the C/N ratio in coral. Moreover, the concentration of nitrogen is decreased along with the increasing depth (Alamaru, Loya, Brokovich, Yam, & Shemesh, 2009).

The size of coral is related to its capability to tolerate environmental stress (Edmunds & Burgess, 2016). However, each coral species has its defensive method to overcome the environmental stress. The response of coral toward environmental disturbance also involved the metabolic system provided by coral's microstructure (Delgado-Lillo-Nuño, Liñán-Cabello, Reyes-Gómez, & Soriano-Santiago, 2014).

~~Assessment of the abundance of nitrifying bacteria *Nitrosomonas* sp. showed that massive coral had higher average density. The collected data as presented in Table 2 showed that higher density of *Nitrosomonas* sp. in massive coral was obtained from all locations. Furthermore, the density of bacteria~~

from the coral sample obtained from Menjangan Kecil island was the highest, while Karimunjawa provided the lowest density.

The sustainability of coral organisms relies on the nutrient supply. The symbiotic between coral and the symbiodinium is highly reliable to the availability of nitrogen (Rädecker et al., 2015). The significant difference between nitrogen concentration in massive and branching coral resulted in this research can be caused by the absorption rate rather than the nitrification process. Higher nitrogen concentration in the branching corals may be caused by polyp abundance. Branching coral tended to have larger surface due to its three dimensional form, while the massive coral only has the top surface. Thus, since the sample was collected in the same size, the abundance of coral polyp in the branching coral was more than the massive coral. Thus, it affected the nitrogen absorption.

The abundance of nitrogen fixing bacteria indicates its nitrification rate provided by the community (Zulfarina, Rusmana, Mubarik, & Santosa, 2017). It also indicates the potential nitrogen produced from the nitrification process. However, the research found that there was no significant effect of the density of *Nitrosomonas* sp. and *Nitrobacter* sp. to the concentration of nitrogen in coral tissue. The nitrification process occurred in the polyp surface, but it mainly affects the nitrogen availability in the environment. Unfortunately, the high nitrification rate is not followed by higher nitrogen uptake by coral organism.

The difference in the density of nitrogen fixing bacteria is dominantly caused by the environmental condition. One of the affecting factors of nitrifying bacteria is the depth of which the bacteria sampled. The abundance of nitrifying bacteria is higher in the shallow water (Smith, Damashek, Chavez, & Francis, 2016). Unfortunately, measurement on the depth of the sampling location was not carried out.

The bacteria is sensitive to the change of environmental condition, such a temperature (Diaz, 2017). At the research location, Karimunjawa island is the most exploited area. Thus, anthropologic aspects may play a role in changing the coral waters. However, this should be assured by **investigat** the condition of coral reef ecosystem. Any environmental stress may break the symbiosis of coral and algae (zooxanthellae) (Wooldridge, 2013). This may affect the other associated microorganisms in the polyp.

Nitrosomonas and *Nitrobacter* are frequently used as bioremediator in aquaculture. However, the preferences of the two is not much understood. Related to the nitrogen fixing process in Karimunjawa National Park, *Nitrosomonas* and *Nitrobacter* might have an important role in remediating coastal environment. The density of both bacterias may be related to the high input of ammonia since Karimunjawa island has been much exploited for tourism activities. *Nitrosomonas* and *Nitrobacter* mainly act as bioremediator of both pollutant (Barik et al., 2018).

Even though nitrogen fixing bacteria exists in the coral mucus or tissue, but it's suggested that the bacteria are associated with Symbiodinium rather than directly to the coral (Lema et al., 2012). This explains why the *Nitrosomonas* sp. and *Nitrobacter* sp. did not influence the concentration of nitrogen significantly.

The role of nitrogen fixing bacteria is obvious in the nitrogen limited environment (Putnam et al., 2017). However, since the effect of *Nitrosomonas* sp. and *Nitrobacter* sp. in Karimunjawa National Park remains insignificant, it can be concluded that the source of nitrogen is sufficient. Or otherwise, since the coral growth requires nitrogen, the coral can not accumulate the nitrogen it consumed.

Conclusion (UPCRASE)


Based on the analysis result, the concentration of nitrogen is higher in branching coral as the impact of its higher growth rate. However, the concentration of nitrogen in coral tissue is not affected by the abundance of nitrogen fixing bacteria.

ACKNOWLEDGMENT

You should wish to thank those who have supported you and your work, especially an institution that provided the financial support.

References (UPCRASE)

- Alamaru, A., Loya, Y., Brokovich, E., Yam, R., & Shemesh, A. (2009). Carbon and nitrogen utilization in two species of Red Sea corals along a depth gradient: Insights from stable isotope analysis of total organic material and lipids. *Geochimica et Cosmochimica Acta*, *73*, 5333–5342.
- Barik, P., Ram, R., Haldar, C., & Vardia, H. K. (2018). Study on nitrifying bacteria as bioremediator of ammonia in simulated aquaculture system. *Journal of Entomology and Zoology Studies*, *6*, 1200–1206.
- Delgadillo-Nuño, M. A., Liñán-Cabello, M. A., Reyes-Gómez, J., & Soriano-Santiago, O. (2014). Response to pH stress in the reef-building coral *Pocillopora capitata* (Anthozoa: Scleractinia). *Revista de Biología Marina y Oceanografía*, *49*, 449–459.
- Diaz, L. (2017). Response of the coral associated nitrogen fixing bacteria toward elevated water temperature. *Journal of Water Resources and Ocean Science*, *6*, 98–109.
- Edmunds, P. J., & Burgess, S. C. (2016). Size-dependent physiological responses of the branching coral *Pocillopora verrucosa* to elevated temperature and PCO₂. *The Journal of Experimental Biology*, *219*, 3896–3906.
- Hohn, S., & Merico, A. (2012). Modelling coral polyp calcification in relation to ocean acidification. *Biogeosciences*, *9*, 4441–4454.
- Krediet, C. J., Ritchie, K. B., Paul, V. J., & Teplitski, M. (2013). Coral-associated micro-organisms and their roles in promoting coral health and thwarting diseases. *Proceedings of the Royal Society B: Biological Sciences*, *280*, 1755.
- Lema, K. A., Willis, B. L., & Bourne, D. G. (2012). Corals form characteristic associations with symbiotic nitrogen-fixing bacteria. *Applied and Environmental Microbiology*, *78*, 3136–3144.
- Putnam, H. M., Barott, K. L., Ainsworth, T. D., & Gates, R. D. (2017). The vulnerability and resilience of reef-building corals. *Current Biology*, *27*, R528–R540.
- Rädecker, N., Pogoreutz, C., Voolstra, C. R., Wiedenmann, J., & Wild, C. (2015). Nitrogen cycling in corals: the key to understanding holobiont functioning? *Trends in Microbiology*, *23*, 490–497.
- Smith, J. M., Damashek, J., Chavez, F. P., & Francis, C. A. (2016). Factors influencing nitrification rates and the abundance and transcriptional activity of ammonia-oxidizing microorganisms in the dark northeast Pacific Ocean. *Limnology and Oceanography*, *61*, 596–609.
- Wahyuni, T., Kusnadi, H., & Honorita, B. (2017). Status unsur hara karbon organik dan nitrogen tanah sawah tiga Kabupaten di Provinsi Bengkulu. In S. Herlinda (Ed.), *Prosiding Seminar Nasional Lahan Suboptimal* (pp. 978–979). Palembang.
- Wang, X. T., Sigman, D. M., Cohen, A. L., Sinclair, D. J., Sherrell, R. M., Weigand, M. A., ... Ren, H. (2015). Isotopic composition of skeleton-bound organic nitrogen in reef-building symbiotic corals: A new method and proxy evaluation at Bermuda. *Geochimica et Cosmochimica Acta*, *148*, 179–190.
- Wooldridge, S. A. (2013). Breakdown of the coral-algae symbiosis: towards formalising a linkage between warm-water bleaching thresholds and the growth rate of the intracellular zooxanthellae. *Biogeosciences*, *10*, 1647–1658.
- Zulfarina, Rusmana, I., Mubarik, N. R., & Santosa, D. A. (2017). The abundance of nitrogen fixing, nitrifying, denitrifying and ammonifying bacteria in the soil of tropical rainforests and oil palm plantations in Jambi. *Makara Journal of Science*, *21*, 187–194.



[Main Menu \(#editorMenu\)](#)

[Home](https://ejournal.undip.ac.id/index.php/sainstek/index/) /
 [User](https://ejournal.undip.ac.id/index.php/sainstek/user/) /
 [Editor](https://ejournal.undip.ac.id/index.php/sainstek/editor/) /
 [Submissions](https://ejournal.undip.ac.id/index.php/sainstek/submission/) /
 [#31339](https://ejournal.undip.ac.id/index.php/sainstek/submission/31339/) /
 [Review](https://ejournal.undip.ac.id/index.php/sainstek/submissionReview/31339/)

Submission #31339

Summary

[Submission](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/)

[Editors](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/status/)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/status/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/status/)

[Status](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/status/)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/status/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/status/)

[Submission Metadata](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/status/)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/status/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/status/)

Review

[Review Submission](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/)

[Peer Review](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/#peerReview)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/#peerReview\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/#peerReview)

[Editor Decision](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/#peerReview)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/#peerReview\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/#peerReview)

Editing

[Copyediting](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#copyedit)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#copyedit\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#copyedit)

[Scheduling](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#scheduling)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#scheduling\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#scheduling)

[Layout](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#layout)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#layout\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#layout)

[Proofreading](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#proofreading)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#proofreading\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#proofreading)

History

[Event Log](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionHistory/31339/)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionHistory/31339/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionHistory/31339/)

[Email Log](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEmailLog/31339/)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEmailLog/31339/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEmailLog/31339/)

[Submission Notes](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionNotes/31339/)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionNotes/31339/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionNotes/31339/)

#31339 Review

[Summary](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/)

[Review](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/)

[Editing](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/)

[History](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionHistory/31339/)

[References](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionCitations/31339/)

[Metadata](https://ejournal.undip.ac.id/index.php/sainstek/editor/submission/31339/#metadata)

Submission

[Review Submission](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionReview/31339/)

[Authors](https://ejournal.undip.ac.id/index.php/sainstek/user/email?) [Suryanti Suryanti, Supriharyono Supriharyono, Sigit Febrianto, Muslidin Aini](https://ejournal.undip.ac.id/index.php/sainstek/user/email?)

[redirectUrl=https%3A%2F%2Fejournal.undip.ac.id%2Findex.php%2Fsainstek%2Feditor%2FsubmissionReview%2F31339&to%5B%5D=%22Suryanti%20Suryanti%22](https://ejournal.undip.ac.id/index.php/sainstek/user/email?)

[Author Scholar Query](https://ejournal.undip.ac.id/index.php/sainstek/user/email?) <https://scholar.google.com/scholar?q=Suryanti+Suryanti+OR+Supriharyono+Supriharyono+OR+Febrianto+Sigit+OR+Aini+Muslidin>

Title NITROGEN CONCENTRATION IN CORAL REEF: THE CONTRIBUTION OF CORAL FORMS AND NITROGEN FIXING BACTERIA IN KARIMUNJAWA ISLANDS

Section [SAINTEK PERIKANAN Sainstek](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#copyedit)

[SAINTEK PERIKANAN Sainstek](https://ejournal.undip.ac.id/index.php/sainstek/user/email?redirectUrl=https%3A%2F%2Fejournal.undip.ac.id%2Findex.php%2F)

Review [31339-91674-1-SM.pdf](https://ejournal.undip.ac.id/index.php/sainstek/editor/downloadFile/31339/91674/1) 01-07-2020

[Version](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#layout)

[Proofreading](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#proofreading)

[History](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionEditing/31339/#proofreading)

Choose File No file chosen

Peer Review

Round 1

[Select Reviewer](https://ejournal.undip.ac.id/index.php/sainstek/editor/selectReviewer/31339/) [View Regrets, Cancels, Previous Rounds](https://ejournal.undip.ac.id/index.php/sainstek/editor/selectReviewer/31339/)

[\(https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionRegrets/31339/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/submissionRegrets/31339/)

Reviewer A Max Muskananfolo
<http://www.scopus.com/authid/detail.uri?authorId=57053506800>

Review Form FORMULIR REVIEW

Request Underway	Due	Acknowledge
25-05-2020	05-06-2020	2-06-2020
2020	(https://ejournal.undip.ac.id/index.php/sainstek/editor/setDueDate/31339/19859/)	

Recommendation Revisions Required 02-06-2020

Uploaded files [31339-123148-1-RV.docx](https://ejournal.undip.ac.id/index.php/sainstek/editor/downloadFile/31339/123148/1) 02-06-2020 Let author view

Reviewer rating 5 High

Editor Decision

Select decision [Accept Submission](#) [Record Decision](#) [Scholar Title Search](https://scholar.google.com/scholar?q=NITROGEN+CONCENTRATION+IN+CORAL+REEF:+THE+CONTRIBUTION+OF+CORAL+FORMS+AND+NITROGEN+FIXING+BACTERIA++IN+KARIMUNJAWA+ISLANDS)

Decision Accept Submission 02-06-2020

Notify Author [\(https://ejournal.undip.ac.id/index.php/sainstek/editor/emailEditorDecisionComment?articleId=31339/\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/emailEditorDecisionComment?articleId=31339/) Editor/Author Email Record

[\(javascript:openComments\('https://ejournal.undip.ac.id/index.php/sainstek/editor/viewEditorDecisionComments/31339/'\)\)](https://ejournal.undip.ac.id/index.php/sainstek/editor/viewEditorDecisionComments/31339/) No Comments

Review Version [31339-91674-1-SM.pdf](https://ejournal.undip.ac.id/index.php/sainstek/editor/downloadFile/31339/91674/1) 02-06-2020 Sent 01-07-2020

Author Version [31339-91674-1-ED.doc](#)

Editor Version None

Choose File No file chosen



Diponegoro under [Creative Commons Attribution-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-sa/4.0/).

Copyright ©2021 [Universitas Diponegoro](https://www.uns.ac.id/). Powered by [Public Knowledge Project OJS](https://www.pkoj.org/) and [Mason Publishing OJS theme](https://masonpublishing.com/).

Pengiriman Hasil Perbaikan Artikel

The image shows a screenshot of an email client interface. At the top, there is a search bar with the text "Search mail" and a magnifying glass icon. To the right of the search bar are icons for help, settings, and a grid of apps. Below the search bar is a navigation bar with various icons for email actions like back, forward, delete, and search. On the right side of the navigation bar, it says "3 of 1,437" with left and right arrow icons. The main content area shows an email from "Dr. Suryanti FPIK UNDIP" with a profile picture and the text "to me". The email is dated "Wed, Jun 3, 2020, 8:30 PM" and has icons for star, reply, and more options. The body of the email contains the following text:

Yth. Redaksi Sainstek Perikanan

Dengan hormat,
Di bawah ini dilampirkan perbaikan artikel kami sesuai dengan hasil review, mohon untuk dapat diterima. Berkas juga sudah kami upload melalui ojs Sainstek Perikanan. Nuwun

Salam Hormat,
Dr. Suryanti

--
Best Regards,

Dr. Ir. Suryanti, MPi
Department of Aquatic Resources
Faculty of Fisheries and Marine Sciences,
Diponegoro University, Semarang, Indonesia

At the bottom of the email, there is a small rectangular box containing a logo and some text, which appears to be a watermark or a small document preview.

**NITROGEN CONCENTRATION IN CORAL REEF: THE
CONTRIBUTION OF CORAL FORMS AND NITROGEN FIXING
BACTERIA
IN KARIMUNJAWA ISLANDS**

Suryanti Suryanti^{1*}, Supriharyono Supriharyono¹, Sigit Febrianto¹, Musludin Aini²

¹ Departemen of Aquatic Resources, Faculty of Fisheries and Marine Science,
Universitas Diponegoro, Semarang, Indonesia

Jl. Prof. Soedarto, SH Tembalang, Semarang, Central Java, Indonesia 50275

²Research Center, Coastal Environmental and Fisheries,

Jl. RA Kartini no 23, Rakam, Selong, Lombok Timur, NTB. Indonesia . 83619

Email: suryantidr@gmail.com

Received : 10 Mei 2020, Accepted : 18 Juni 2020

ABSTRACT

*Coral is an ecosystem in the coastal area which plays an important role in controlling nutrient cycle in the sea. This research aimed to study the the nitrogen concentration in coral and to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in coral. The research was carried out by field observation, by taking coral reef samples and analyzed in the laboratory for its nitrogen concentration and number of nitrogen fixing bacteria. Data analysis was carried out by ANOVA and regression. The analysis resulted that there was a significant difference of nitrogen concentration in massive and branching corals. However, there is no significant correlation between the sampling location and types of coral on the density of nitrogen fixing bacteria, both *Nitrosomonas sp.* and *Nitrobacter sp.* Regression analysis showed that the concentration of nitrogen in coral tissue was not significantly related to the density of nitrogen fixing bacteria. The result of this research implies that the concentration of nitrogen in the coral tissue may be affected by many other factors.*

Keywords: *bacteria; coral; fixation; nitrogen; tissue*

INTRODUCTION

As the development of coastal area increase, the problems related to the coastal ecosystem arises. This includes the coral reef ecosystem which has the most important role in supporting marine biodiversity. Coral also provides micro-biome which shows the relationship among organisms in the coral surface. Various processes occur in the micro-biome, such as sulphur cycling, nitrogen fixing, as well as the it defensive system (Krediet, Ritchie, Paul, & Teplitski, 2013).

Coral is a rocky organism that lives in the shallow marine ecosystem ranging from the tropical to sub-tropical area. Ecologically, shallow coral reefs are considered as the huge source of benthic nitrogen fixation (Diaz, 2017). The biodiversity in coral reef ecosystem is supported by the coral associated organisms, such as sulphur recycling bacteria or nitrogen fixing bacteria. All of them existed in the coral ecosystem and provide defense toward various threats.

As a living organism, coral absorbs various minerals to grow. One of the important nutrient for coral is nitrogen. Nitrogen for coral is mainly provided by zooxanthelae which which lives mutualistically with coral polyp (Krediet *et al.*, 2013). Coral polyp is where various ecological processes occur. Various physiological response due to the arising stress in the environment is detected and responded in the polyp (Hohn & Merico, 2012).

Symbiodinium is the algae species which has mutualistic association with coral. However, the micro-biome in coral is just recognized recently. The micro-biome is the system which provide support to the organization of coral ecosystems which involved various microorganisms (Rädecker, Pogoreutz, Voolstra, Wiedenmann, & Wild, 2015). Among the functions the microbiome provides, nitrogen cycling is considered as the most important process for coral sustainability.

There are several studies wich found that coral polyp also contain nitrogen fixing bacteria. Nitrogen is an important element required by any plant

to grow. However, in some cases the availability of nitrogen in the coral reef ecosystem is limited. In this condition, nitrogen fixing is the main source for coral to support its growth (Putnam, Barott, Ainsworth, & Gates, 2017). The nitrogen fixing bacteria are frequently found in coral mucus (Lema, Willis, & Bourne, 2012).

The nitrogen fixing bacteria is commonly called diazotrophs. Nitrogen fixing bacteria are mostly found in a reef building symbiotic coral as the source of nitrogen for coral's growth (Wang *et al.*, 2015). The bacteria favors the nitrogen supply to the coral organism directly through bacterivory, through polyp and zooxanthelae (Diaz, 2017). The species of nitrogen fixing bacteria is varied among coral species and locations. At least there are 23 species of nitrogen fixing bacteria. *Nitrosomonas* and *Nitrobacter* are frequently used as bioremediator for aquaculture. It is mostly functioned as the remediator for ammonia and nitrite (Barik, Ram, Haldar, & Vardia, 2018).

Critical concern on the microbiological role on the coral reef ecosystem arises the question whether the associated microorganism also affect the nutrient uptake by coral reef. Least information is known regarding the processes and impact of the nitrogen fixing bacteria to the livelihood of coral community. This research aimed to study the nitrogen concentration in coral and to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in coral.

RESEARCH METHODS

The research was carried out in June – July 2019 in the area of Karimunjawa National Park. The research was carried out by field observation. The research stations were located in Karimunjawa island, Menjangan Kecil island, and Cemara Kecil island, while laboratory analysis was in the *Tropical Marine Biotechnology (TMB), Faculty of Fisheries and Marine Science Diponegoro University Semarang*.

There were three sampling stations occupied in the research. Each station consisted of three sampling spots. The objects of the research were including the concentration of Nitrogen in coral organisms and the density of nitrogen fixing bacteria. Coral samples were obtained from two groups of coral organisms, including the massive and branching corals. Thus, the research design applied 3 x 2 factorial treatment.

Data collection was carried out through field sampling followed by laboratory measurement and identification. The coral was sampled with the length of 10 cm. The coral samples were stored in a cool box and identified in the laboratory for its density of nitrogen fixing bacteria and the concentration of

nitrogen. The bacteria were isolated from the coral and placed in the enrichment media (nutrient agar). The bacteria collected as the object of the research was *Nitrosomonas* sp. and *Nitrobacter* sp.

Data analysis was carried out with ANOVA and regression. Multivariate ANOVA was carried out to analyze the difference of nitrogen concentration and the abundance of nitrogen fixing bacteria. Regression was carried out to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in the coral. The regression analysis was carried out through multiple regression to analyze the simultaneous effect of nitrogen fixing bacteria, and curve estimation to analyze the effect of nitrogen fixing bacteria separately. Analysis was carried out by SPSS.

RESULTS AND DISCUSSION

Data analysis was carried out with ANOVA and regression. Multivariate ANOVA was carried out to analyze the difference of nitrogen concentration and the abundance of nitrogen fixing bacteria. Regression was carried out to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in the coral. The regression analysis was carried out through multiple regression to analyze the simultaneous effect of nitrogen fixing bacteria, and curve estimation to analyze the effect of nitrogen fixing bacteria separately. Analysis was carried out by SPSS.

Table 1 shows that the average concentration of nitrogen in the coral was ranging from very low to moderate. However, moderate nitrogen concentration was obtained from branching coral, while very low concentration was obtained from massive coral. The concentration of nitrogen is considered as very low within the range < 0.1 %, low within the range 0.1 – 0.2 and moderate within the range 0.2 – 0.5 (Wahyuni, Kusnadi, & Honorita, 2017). Thus, it can be noted that massive coral has a lower accumulation of nitrogen.

Table 7. Average Concentration of Nitrogen in Coral (Mg/Kg) from Each Sample

No	Location	Types of Coral		Total by Location
		Massive	Branching	
1.	Karimunjawa	61.380*	231.527 ^{‡a}	146.453 [‡] _p
2.	Menjangan Kecil	73.662*	197.390 ^{‡a}	135.526 [‡] _p
3.	Cemara Kecil	160.223	221.486 ^{‡a}	190.855 [‡] _p
Total by Types of Coral		98.422* _x	216.801 ^{‡y}	

Notation : * = very low; ‡ = low; [‡] = moderate; Cells, column (total by types of coral), and rows (total by

location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Statistical data analysis showed that the nitrogen concentration was affected by the types of coral. Massive coral tended to have lower nitrogen concentration than the branching coral. However, other treatments did not show any significant effect on the nitrogen concentration. However, the difference in nitrogen concentration affected by location, and types of coral altogether was noticeable. Data analysis resulted F value for location was 2.311 ($p = 0.142$), types of coral showed the value of 28.285 ($p = 0.000$), and combination of both 2.009 ($p = 0.251$).

Assessment of the abundance of nitrifying bacteria *Nitrosomonas* sp. showed that massive coral had higher average density. The collected data as presented in Table 2 showed that higher density of *Nitrosomonas* sp. in massive coral was obtained from all locations. Furthermore, the density of bacteria from the coral sample obtained from Menjangan Kecil island was the highest, while Karimunjawa provided the lowest density.

Table 8. Average Density of Nitrogen Fixing Bacteria *Nitrosomonas* Sp. (cfu/g) from Each Sample

No.	Location	Types of Coral		
		Massive	Branching	Total
1.	Karimunjawa	$58.27 \times 10^{5(a)}$	$39.93 \times 10^{5(a)}$	$49.10 \times 10^{5(a)}$
2.	Menjangan Kecil	$11.30 \times 10^{6(a)}$	$15.00 \times 10^{6(a)}$	$13.15 \times 10^{6(a)}$
3.	Cemara Kecil	$11.07 \times 10^{6(a)}$	$64.33 \times 10^{5(a)}$	$87.50 \times 10^{5(a)}$
<i>Total</i>		$93.98 \times 10^{5(x)}$	$84.76 \times 10^{5(x)}$	

Notation: Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Statistical analysis showed that there was no significant effect of the source of coral samples and the types of coral to the density of *Nitrosomonas* sp. Data analysis with univariate ANOVA showed that the F value obtained from the location, types of coral and combination of both were 2.847 ($p = 0.097$), 0.107 ($p = 0.749$) and 0.753 ($p = 0.492$) respectively. Thus, it can be suggested that massive coral provided better support to the nitrogen fixing bacteria. The environmental condition of Menjangan Kecil island was also more supportive than the other two islands.

The average density of nitrogen fixing bacteria *Nitrobacter* sp. is the last parameter observed in this research. Assessment on the average density of *Nitrobacter* sp. showed that there was slight difference between the massive and branching corals. However, branching coral had higher *Nitrobacter* sp. density even though the difference was not significant. Noticeable difference was obtained from the bacterial density in Karimunjawa island. The difference was

obtained not only from the total average density, but also in the density of respective massive and branching coral types. Table 3 presents the detailed *Nitrobacter* sp. density of the samples obtained from the research location.

Table 9. Average Density of Nitrogen Fixing Bacteria *Nitrobacter* sp. (cfu/gr) from Each Sample

No.	Location	Types of Coral		
		Massive	Branching	Total
1.	Karimunjawa	$96.57 \times 10^{5(a)}$	$59.43 \times 10^{5(a)}$	$78.00 \times 10^{5(p)}$
2.	Menjangan Kecil	$15.00 \times 10^{6(a)}$	$19.67 \times 10^{6(a)}$	$17.33 \times 10^{6(p)}$
3.	Cemara Kecil	$15.33 \times 10^{6(a)}$	$16.90 \times 10^{6(a)}$	$16.12 \times 10^{6(p)}$
<i>Total</i>		$13.33 \times 10^{6(x)}$	$14.17 \times 10^{6(x)}$	

Notation: Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Table 3 shows that the density of *Nitrobacter* sp. in Karimunjawa National Park was higher than the *Nitrosomonas* sp. However, there was no significant difference in the density of nitrogen fixing bacteria *Nitrobacter* sp. both partially and simultaneously. Statistical analysis with ANOVA showed the F value of coral source, types of coral and combination of both were as much as 1.877 ($p = 0.195$), 0.037 ($p = 0.851$) and 0.313 ($p = 0.737$) respectively.

Among the results from the assessment of nitrogen concentration as well as the density of *Nitrosomonas* sp. and *Nitrobacter* sp., it can be suggested that Karimunjawa always had the lowest value. This could be due to the environmental condition of the island. However, further indepth research is needed to study the possible causes.

Statistical analysis to assess the influence of nitrogen fixing bacteria showed that the concentration of nitrogen in the coral was not related to the density of *Nitrosomonas* sp. nor *Nitrobacter* sp. Both multiple regression and curve estimation analysis did not show the significance of the relationship. Thus, it can be concluded that the concentration of nitrogen in coral is affected by some other factors, or involves some other factors as moderator or as cooperating factor.

Discussion

The accumulation of nitrogen in massive and branching coral is suggested as the effect of growth rate between the two. Nitrogen is required for coral metabolism. However, the environmental condition significantly affect the concentration of nitrogen. Shaded environment tends to cause the increase of the C/N ratio in coral. Moreover, the concentration of nitrogen is decreased along with the increasing depth (Alamaru, Loya, Brokovich, Yam, & Shemesh, 2009).

The size of coral is related to its capability to tolerate environmental stress (Edmunds & Burgess,

2016). However, each coral species has its defensive method to overcome the environmental stress. The response of coral toward environmental disturbance also involved the metabolic system provided by coral's microstructure (Delgadillo-Nuño, Liñán-Cabello, Reyes-Gómez, & Soriano-Santiago, 2014).

The sustainability of coral organisms relies on the nutrient supply. The symbiotic between coral and the symbiodinium is highly reliable to the availability of nitrogen (Rädecker *et al.*, 2015). The significant difference between nitrogen concentration in massive and branching coral resulted in this research can be caused by the absorption rate rather than the nitrification process. Higher nitrogen concentration in the branching corals may be caused by polyp abundance. Branching coral tended to have larger surface due to its three dimensional form, while the massive coral only has the top surface. Thus, since the sample was collected in the same size, the abundance of coral polyp in the branching coral was more than the massive coral. Thus, it affected the nitrogen absorption.

The abundance of nitrogen fixing bacteria indicates its nitrification rate provided by the community (Zulfarina, Rusmana, Mubarik, & Santosa, 2017). It also indicates the potential nitrogen produced from the nitrification process. However, the research found that there was no significant effect of the density of *Nitrosomonas* sp. and *Nitrobacter* sp. to the concentration of nitrogen in coral tissue. The nitrification process occurred in the polyp surface, but it mainly affects the nitrogen availability in the environment. Unfortunately, the high nitrification rate is not followed by higher nitrogen uptake by coral organism.

The difference in the density of nitrogen fixing bacteria is dominantly caused by the environmental condition. One of the affecting factors of nitrifying bacteria is the depth of which the bacteria sampled. The abundance of nitrifying bacteria is higher in the shallow water (Smith, Damashek, Chavez, & Francis, 2016). Unfortunately, measurement on the depth of the sampling location was not carried out.

The bacteria is sensitive to the change of environmental condition, such a temperature (Diaz, 2017). At the research location, Karimunjawa island is the most exploited area. Thus, anthropologic aspects may play a role in changing the coral waters. However, this should be assured by investigating the condition of coral reef ecosystem. Any environmental stress may break the symbiosis of coral and algae (zooxanthellae) (Wooldridge, 2013). This may affect the other associated microorganisms in the polyp.

Nitrosomonas and *Nitrobacter* are frequently used as bioremediator in aquaculture. However, the

preferences of the two is not much understood. Related to the nitrogen fixing process in Karimunjawa National Park, *Nitrosomonas* and *Nitrobacter* might have an important role in remediating coastal environment. The density of both bacteria may be related to the high input of ammonia since Karimunjawa island has been much exploited for tourism activities. *Nitrosomonas* and *Nitrobacter* mainly act as bioremediator of both pollutant (Barik *et al.*, 2018).

Even though nitrogen fixing bacteria exists in the coral mucus or tissue, but it's suggested that the bacteria are associated with Symbiodinium rather than directly to the coral (Lema *et al.*, 2012). This explains why the *Nitrosomonas* sp. and *Nitrobacter* sp. did not influence the concentration of nitrogen significantly.

The role of nitrogen fixing bacteria is obvious in the nitrogen limited environment (Putnam *et al.*, 2017). However, since the effect of *Nitrosomonas* sp. and *Nitrobacter* sp. in Karimunjawa National Park remains insignificant, it can be concluded that the source of nitrogen is sufficient. Or otherwise, since the coral growth requires nitrogen, the coral can not accumulate the nitrogen it consumed.

CONCLUSION

Based on the analysis result, the concentration of nitrogen is higher in branching coral as the impact of its higher growth rate. However, the concentration of nitrogen in coral tissue is not affected by the abundance of nitrogen fixing bacteria.

ACKNOWLEDGMENTS

We would like to thank: Head of Universitas Diponegoro Research and Community Service Institute for PNBPN research funding No. 474-19 / UN7.P4.3 / PP / 2019, and to the Head of BTNKJ for permission and information

REFERENCES

- Alamaru, A., Loya, Y., Brokovich, E., Yam, R., & Shemesh, A. (2009). Carbon and nitrogen utilization in two species of Red Sea corals along a depth gradient: Insights from stable isotope analysis of total organic material and lipids. *Geochimica et Cosmochimica Acta*, 73, 5333–5342.
- Barik, P., Ram, R., Haldar, C., & Vardia, H. K. (2018). Study on nitrifying bacteria as bioremediator of ammonia in simulated aquaculture system. *Journal of Entomology and Zoology Studies*, 6, 1200–1206.
- Delgadillo-Nuño, M. A., Liñán-Cabello, M. A.,

- Reyes-Gómez, J., & Soriano-Santiago, O. (2014). Response to pH stress in the reef-building coral *Pocillopora capitata* (Anthozoa: Scleractinia). *Revista de Biología Marina y Oceanografía*, *49*, 449–459.
- Diaz, L. (2017). Response of the coral associated nitrogen fixing bacteria toward elevated water temperature. *Journal of Water Resources and Ocean Science*, *6*, 98–109.
- Edmunds, P. J., & Burgess, S. C. (2016). Size-dependent physiological responses of the branching coral *Pocillopora verrucosa* to elevated temperature and PCO₂. *The Journal of Experimental Biology*, *219*, 3896–3906.
- Hohn, S., & Merico, A. (2012). Modelling coral polyp calcification in relation to ocean acidification. *Biogeosciences*, *9*, 4441–4454.
- Krediet, C. J., Ritchie, K. B., Paul, V. J., & Teplitski, M. (2013). Coral-associated micro-organisms and their roles in promoting coral health and thwarting diseases. *Proceedings of the Royal Society B: Biological Sciences*, *280*, 1755.
- Lema, K. A., Willis, B. L., & Bourne, D. G. (2012). Corals form characteristic associations with symbiotic nitrogen-fixing bacteria. *Applied and Environmental Microbiology*, *78*, 3136–3144.
- Putnam, H. M., Barott, K. L., Ainsworth, T. D., & Gates, R. D. (2017). The vulnerability and resilience of reef-building corals. *Current Biology*, *27*, R528–R540.
- Rädecker, N., Pogoreutz, C., Voolstra, C. R., Wiedenmann, J., & Wild, C. (2015). Nitrogen cycling in corals: the key to understanding holobiont functioning? *Trends in Microbiology*, *23*, 490–497.
- Smith, J. M., Damashek, J., Chavez, F. P., & Francis, C. A. (2016). Factors influencing nitrification rates and the abundance and transcriptional activity of ammonia-oxidizing microorganisms in the dark northeast Pacific Ocean. *Limnology and Oceanography*, *61*, 596–609.
- Wahyuni, T., Kusnadi, H., & Honorita, B. (2017). Status unsur hara karbon organik dan nitrogen tanah sawah tiga Kabupaten di Provinsi Bengkulu. In S. Herlinda (Ed.), *Prosiding Seminar Nasional Lahan Suboptimal* (pp. 978–979). Palembang.
- Wang, X. T., Sigman, D. M., Cohen, A. L., Sinclair, D. J., Sherrell, R. M., Weigand, M. A., ... Ren, H. (2015). Isotopic composition of skeleton-bound organic nitrogen in reef-building symbiotic corals: A new method and proxy evaluation at Bermuda. *Geochimica et Cosmochimica Acta*, *148*, 179–190.
- Wooldridge, S. A. (2013). Breakdown of the coral-algae symbiosis: towards formalising a linkage between warm-water bleaching thresholds and the growth rate of the intracellular zooxanthellae. *Biogeosciences*, *10*, 1647–1658.
- Zulfarina, Rusmana, I., Mubarik, N. R., & Santosa, D. A. (2017). The abundance of nitrogen fixing, nitrifying, denitrifying and ammonifying bacteria in the soil of tropical rainforests and oil palm plantations in Jambi. *Makara Journal of Science*, *21*, 187–194.

SAINTEK PERIKANAN
(Indonesian Journal of Fisheries Science and Technology)
Terakreditasi Sinta 2 (SK No. 10/E/KPT/2019)
P-ISSN 1858-4748 E-ISSN 2549-0885
FAKULTAS PERIKANAN DAN ILMU KELAUTAN - UNIVERSITAS DIPONEGORO
Jl. Prof. H. Soedarto SH Tembalang Semarang. 50275 Telp/fak. (024) 76480685
Email : saintekjurnal@gmail.com

SURAT KETERANGAN NASKAH DITERIMA

No : 15/Saintek Pi /VI/2020

Bersama ini, redaksi Sainstek Perikanan memberitahukan bahwa naskah dengan identitas sebagai berikut:

Judul : Nitrogen Concentration In Coral Reef: The Contribution Of Coral Forms And Nitrogen Fixing Bacteria In Karimunjawa Islands

Penulis : Suryanti Suryanti Supriharyono Supriharyono, Sigit Febrianto, Muslidin Aini

Afiliasi / Institusi : Departemen of Aquatic Resources, Faculty of Fisheries and Marine Science, Universitas Diponegoro
Research Center, Coastal Enviromental and Fisheries

Email : suryantidr@gmail.com

Artikel tersebut telah memenuhi kriteria pada publikasi di Sainstek Perikanan untuk diterbitkan pada volume 16, Nomor 2, Agustus 2020 dalam versi cetak dan elektronik.

Demikian surat keterangan ini disampaikan, untuk digunakan sebagaimana mestinya.

Semarang, 15 Juni 2020
Ketua Dewan Redaksi



Prof. Dr. Aristi Dian Purnama Fitri, M.Si
NIP. 19731002 199803 2 001

Copyedit Review Request

Search mail

2 of 1,437

[IJFST] Copyediting Review Request

Saintek jurnal <saintekjurnal@gmail.com>
to suryanti

Mon, Jun 22, 2020, 1:45 PM

Dr. Suryanti :

Your submission "NITROGEN CONCENTRATION IN CORAL REEF: THE CONTRIBUTION OF CORAL FORMS AND NITROGEN FIXING BACTERIA IN KARIMUNJAWA ISLANDS" for Saintek Perikanan : Indonesian Journal of Fisheries Science and Technology has been through the first step of copyediting, and is available for you to review by following these steps:

1. Click on the Submission URL below.
2. Log into the journal and click on the File that appears in Step 1.
3. Open the downloaded submission.
4. Review the text, including copyediting proposals and Author Queries.
5. Make any copyediting changes that would further improve the text.
6. When completed, upload the file in Step 2.
7. Click on METADATA to check indexing information for completeness and accuracy.
8. Send the COMPLETE email to the editor and copyeditor.

Submission URL: <https://ejournal.undip.ac.id/index.php/saintek/author/submissionEditing/31339>
Username: drsuryanti

This is the last opportunity to make substantial copyediting changes to the submission. The proofreading stage, that follows the preparation of the galleys, is restricted to correcting typographical and layout errors.

Vol 16, No 2 (2020): SAINTEK PERIKANAN



Table of Contents

Research Articles

PENGGUNAAN HIGH POWER LED (HPL) PADA PERIKANAN BAGAN APUNG DI SELAT MADURA (The Use Of High Power LED (HPL) Lamp On The Lift Net Fishing In The Madura Strait) PDF 79-85

Hairul Umam, Gondo Puspito, Wazir Mawardi

Views: 835 | Citations: ? | Language: ID | DOI: 10.14710/ijfst.16.2.79-85

Received: 15 Mar 2020; Published: 4 Aug 2020.

NITROGEN CONCENTRATION IN CORAL REEF: THE CONTRIBUTION OF CORAL FORMS AND NITROGEN FIXING BACTERIA IN KARIMUNJAWA ISLANDS PDF 86-89

Suryanti Suryanti, Supriharyono Supriharyono, Sigit Febrianto, Muslidin Aini

Views: 265 | Citations: ? | Language: EN | DOI: 10.14710/ijfst.16.2.86-89

Received: 10 May 2020; Published: 4 Aug 2020.

RESPONS TINGKAH LAKU CUMI-CUMI (*Uroteuthis duvaucelli*, Orbigny 1835) TERHADAP WARNA DAN INTENSITAS CAHAYA YANG PDF

NITROGEN CONCENTRATION IN CORAL REEF: THE CONTRIBUTION OF CORAL FORMS AND NITROGEN FIXING BACTERIA IN KARIMUNJAWA ISLANDS

Suryanti Suryanti^{1*}, Supriharyono Supriharyono¹, Sigit Febrianto¹, Muslidin Aini²

¹Departemen of Aquatic Resources, Faculty of Fisheries and Marine Science, Universitas Diponegoro, Semarang, Indonesia

Jl. Prof. Soedarto, SH Tembalang, Semarang, Central Java, Indonesia 50275

²Research Center, Coastal Environmental and Fisheries, Jl. RA Kartini no 23, Rakam, Selong, Lombok Timur, NTB. Indonesia . 83619

Email: suryantidr@gmail.com

Received : 10 Mei 2020, Accepted : 18 Juni 2020

ABSTRACT

Coral is an ecosystem in the coastal area which plays an important role in controlling nutrient cycle in the sea. This research aimed to study the the nitrogen concentration in coral and to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in coral. The research was carried out by field observation, by taking coral reef samples and analyzed in the laboratory for its nitrogen concentration and number of nitrogen fixing bacteria. Data analysis was carried out by ANOVA and regression. The analysis resulted that there was a significant difference of nitrogen concentration in massive and branching corals. However, there is no significant correlation between the sampling location and types of coral on the density of nitrogen fixing bacteria, both Nitrosomonas sp. and Nitrobacter sp. Regression analysis showed that the concentration of nitrogen in coral tissue was not significantly related to the density of nitrogen fixing bacteria. The result of this research implies that the concentration of nitrogen in the coral tissue may be affected by many other factors.

Keywords: bacteria; coral; fixation; nitrogen; tissue

INTRODUCTION

As the development of coastal area increase, the problems related to the coastal ecosystem arises. This includes the coral reef ecosystem which has the most important role in supporting marine biodiversity. Coral also provides micro-biome which shows the relationship among organisms in the coral surface. Various processes occur in the micro-biome, such as sulphur cycling, nitrogen fixing, as well as the it defensive system (Krediet, Ritchie, Paul, & Teplitski, 2013).

Coral is a rocky organism that lives in the shallow marine ecosystem ranging from the tropical to sub-tropical area. Ecologically, shallow coral reefs are considered as the huge source of benthic nitrogen fixation (Diaz, 2017). The biodiversity in coral reef ecosystem is supported by the coral associated organisms, such as sulphur recycling bacteria or nitrogen fixing bacteria. All of them existed in the coral ecosystem and provide defense toward various threats.

As a living organism, coral absorbs various minerals to grow. One of the important nutrient for coral is nitrogen. Nitrogen for coral is mainly provided by zooxanthelae which lives mutualistically with coral polyp (Krediet *et al.*, 2013). Coral polyp is where various ecological processes occur. Various physiological response due to the arising stress in the environment is detected and responded in the polyp (Hohn & Merico, 2012).

Symbiodinium is the algae species which has mutualistic association with coral. However, the micro-biome in coral is just recognized recently. The micro-biome is the system which provide support to the organization of coral

ecosystems which involved various microorganisms (Rädecker, Pogoreutz, Voolstra, Wiedenmann, & Wild, 2015). Among the functions the microbiome provides, nitrogen cycling is considered as the most important process for coral sustainability.

There are several studies wich found that coral polyp also contain nitrogen fixing bacteria. Nitrogen is an important element required by any plant to grow. However, in some cases the availability of nitrogen in the coral reef ecosystem is limited. In this condition, nitrogen fixing is the main source for coral to support its growth (Putnam, Barott, Ainsworth, & Gates, 2017). The nitrogen fixing bacteria are frequently found in coral mucus (Lema, Willis, & Bourne, 2012).

The nitrogen fixing bacteria is commonly called diazotrophs. Nitrogen fixing bacteria are mostly found in a reef building symbiotic coral as the source of nitrogen for coral's growth (Wang *et al.*, 2015). The bacteria favors the nitrogen supply to the coral organism directly through bacterivory, through polyp and zooxanthelae (Diaz, 2017). The species of nitrogen fixing bacteria is varied among coral species and locations. At least there are 23 species of nitrogen fixing bacteria. Nitrosomonas and nitrobacter are frequently used as bioremediator for aquaculture. It is mostly functioned as the remediator for ammonia and nitrite (Barik, Ram, Haldar, & Vardia, 2018).

Critical concern on the microbiological role on the coral reef ecosystem arises the question wether the associated microorganism also affect the nutrien uptake by coral reef. Least information is known regarding the processes and impact of the nitrogen fixing bacteria to the livelihood of coral

community. This research aimed to study the nitrogen concentration in coral and to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in coral.

RESEARCH METHODS

The research was carried out in June – July 2019 in the area of Karimunjawa National Park. The research was carried out by field observation. The research stations were located in Karimunjawa island, Menjangan Kecil island, and Cemara Kecil island, while laboratory analysis was in the *Tropical Marine Biotechnology (TMB), Faculty of Fisheries and Marine Science Diponegoro University Semarang*.

There were three sampling stations occupied in the research. Each station consisted of three sampling spots. The objects of the research were including the concentration of Nitrogen in coral organisms and the density of nitrogen fixing bacteria. Coral samples were obtained from two groups of coral organisms, including the massive and branching corals. Thus, the research design applied 3 x 2 factorial treatment.

Data collection was carried out through field sampling followed by laboratory measurement and identification. The coral was sampled with the length of 10 cm. The coral samples were stored in a cool box an identified in the laboratory for its density of nitrogen fixing bacteria and the concentration of nitrogen. The bacteria were isolated from the coral and placed in the enrichment media (nutrient agar). The bacteria collected as the object of the research was *Nitrosomonas* sp. and *Nitrobacter* sp.

Data analysis was carried out with ANOVA and regression. Multivariate ANOVA was carried out to analyze the difference of nitrogen concentration and the abundance of nitrogen fixing bacteria. Regression was carried out to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in the coral. The regression analysis was carried out through multiple regression to analyze the simultaneous effect of nitrogen fixing bacteria, and curve estimation to analyze the effect of nitrogen fixing bacteria separately. Analysis was carried out by SPSS.

RESULTS AND DISCUSSION

Data analysis was carried out with ANOVA and regression. Multivariate ANOVA was carried out to analyze the difference of nitrogen concentration and the abundance of nitrogen fixing bacteria. Regression was carried out to analyze the influence of nitrogen fixing bacteria on the concentration of nitrogen in the coral. The regression analysis was carried out through multiple regression to analyze the simultaneous effect of nitrogen fixing bacteria, and curve estimation to analyze the effect of nitrogen fixing bacteria separately. Analysis was carried out by SPSS.

Table 1 shows that the average concentration of nitrogen in the coral was ranging from very low to moderate. However, moderate nitrogen concentration was obtained from branching coral, while very low concentration was obtained from massive coral. The concentration of nitrogen is considered as very low within the range < 0.1 %, low within the range 0.1 – 0.2 and moderate within the range 0,2 – 0.5

(Wahyuni, Kusnadi, & Honorita, 2017). Thus, it can be noted that massive coral has a lower accumulation of nitrogen.

Table 1. Average Concentration of Nitrogen in Coral (Mg/Kg) from Each Sample

No.	Location	Types of Coral		Total by Location
		Massive	Branching	
1.	Karimunjawa	61.380 ^{*a}	231.527 ^{-a}	146.453 ^{‡p}
2.	Menjangan Kecil	73.662 ^{*a}	197.390 ^{‡a}	135.526 ^{‡p}
3.	Cemara Kecil	160.223 ^{‡a}	221.486 ^{□a}	190.855 ^{‡p}
Total by Types of Coral		98.422 ^{*x}	216.801 ^{□y}	

Notation : * = very low; ‡ = low; □ = moderate; Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Statistical data analysis showed that the nitrogen concentration was affected by the types of coral. Massive coral tended to have lower nitrogen concentration than the branching coral. However, other treatments did not show any significant effect on the nitrogen concentration. However, the difference in nitrogen concentration affected by location, and types of coral altogether was noticeable. Data analysis resulted F value for location was 2.311 ($p = 0.142$), types of coral showed the value of 28.285 ($p = 0.000$), and combination of both 2.009 ($p = 0.251$).

Assessment of the abundance of nitrifying bacteria *Nitrosomonas* sp. showed that massive coral had higher average density. The collected data as presented in Table 2 showed that higher density of *Nitrosomonas* sp. in massive coral was obtained from all locations. Furthermore, the density of bacteria from the coral sample obtained from Menjangan Kecil island was the highest, while Karimunjawa provided the lowest density.

Table 2. Average Density of Nitrogen Fixing Bacteria *Nitrosomonas* Sp. (cfu/g) from Each Sample

No.	Location	Types of Coral		
		Massive	Branching	Total
1.	Karimunjawa	58.27x10 ^{5(a)}	39.93x10 ^{5(a)}	49.10x10 ^{5(p)}
2.	Menjangan Kecil	11.30x10 ^{6(a)}	15.00x10 ^{6(a)}	13.15x10 ^{6(p)}
3.	Cemara Kecil	11.07x10 ^{6(a)}	64.33x10 ^{5(a)}	87.50x10 ^{5(p)}
Total		93.98x10 ^{5(x)}	84.76x10 ^{5(x)}	

Notation: Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Statistical analysis showed that there was no significant effect of the source of coral samples and the types of coral to the density of *Nitrosomonas* sp. Data analysis with univariate ANOVA showed that the F value obtained from the location, types of coral and combination of both were 2.847 ($p = 0.097$), 0.107 ($p = 0.749$) and 0.753 ($p = 0.492$) respectively. Thus, it can be suggested that massive coral provided better support to the nitrogen fixing bacteria. The environmental condition of Menjangan Kecil island was also more supportive than the other two islands.

The average density of nitrogen fixing bacteria *Nitrobacter* sp. is the last parameter observed in this research. Assessment on the average density of *Nitrobacter* sp. showed that there was

slight difference between the massive and branching corals. However, branching coral had higher *Nitrobacter* sp. density even though the difference was not significant. Noticeable difference was obtained from the bacterial density in Karimunjawa island. The difference was obtained not only from the total average density, but also in the density of respective massive and branching coral types. Table 3 presents the detailed *Nitrobacter* sp. density of the samples obtained from the research location.

Table 3. Average Density of Nitrogen Fixing Bacteria *Nitrobacter* sp. (cfu/gr) from Each Sample

No.	Location	Types of Coral		
		Massive	Branching	Total
1.	Karimunjawa	96.57x10 ^{5(a)}	59.43x10 ^{5(a)}	78.00x10 ^{5(p)}
2.	Menjangan Kecil	15.00x10 ^{6(a)}	19.67x10 ^{6(a)}	17.33x10 ^{6(p)}
3.	Cemara Kecil	15.33x10 ^{6(a)}	16.90x10 ^{6(a)}	16.12x10 ^{6(p)}
Total		13.33x10 ^{6(x)}	14.17x10 ^{6(x)}	

Notation: Cells, column (total by types of coral), and rows (total by location) with similar letter indicates insignificant difference at $\alpha = 0.05$

Table 3 shows that the density of *Nitrobacter* sp. in Karimunjawa National Park was higher than the *Nitrosomonas* sp. However, there was no significant difference in the density of nitrogen fixing bacteria *Nitrobacter* sp. both partially and simultaneously. Statistical analysis with ANOVA showed the F value of coral source, types of coral and combination of both were as much as 1.877 ($p = 0.195$), 0.037 ($p = 0.851$) and 0.313 ($p = 0.737$) respectively.

Among the results from the assessment of nitrogen concentration as well as the density of *Nitrosomonas* sp. and *Nitrobacter* sp., it can be suggested that Karimunjawa always had the lowest value. This could be due to the environmental condition of the island. However, further indepth research is needed to study the possible causes.

Statistical analysis to assess the influence of nitrogen fixing bacteria showed that the concentration of nitrogen in the coral was not related to the density of *Nitrosomonas* sp. nor *Nitrobacter* sp. Both multiple regression and curve estimation analysis did not show the significance of the relationship. Thus, it can be concluded that the concentration of nitrogen in coral is affected by some other factors, or involves some other factors as moderator or as cooperating factor.

Discussion

The accumulation of nitrogen in massive and branching coral is suggested as the effect of growth rate between the two. Nitrogen is required for coral metabolism. However, the environmental condition significantly affect the concentration of nitrogen. Shaded environment tends to cause the increase of the C/N ratio in coral. Moreover, the concentration of nitrogen is decreased along with the increasing depth (Alamaru, Loya, Brokovich, Yam, & Shemesh, 2009).

The size of coral is related to its capability to tolerate environmental stress (Edmunds & Burgess, 2016). However, each coral species has its defensive method to overcome the environmental stress. The response of coral toward environmental disturbance also involved the metabolic system provided by coral's microstructure (Delgadillo-Nuño, Liñán-Cabello, Reyes-Gómez, & Soriano-Santiago, 2014).

The sustainability of coral organisms relies on the nutrient supply. The symbiotic between coral and the symbiodinium is highly reliable to the availability of nitrogen (Rädecker *et al.*, 2015). The significant difference between nitrogen concentration in massive and branching coral resulted in this research can be caused by the absorption rate rather than the nitrification process. Higher nitrogen concentration in the branching corals may be caused by polyp abundance. Branching coral tended to have larger surface due to its three dimensional form, while the massive coral only has the top surface. Thus, since the sample was collected in the same size, the abundance of coral polyp in the branching coral was more than the massive coral. Thus, it affected the nitrogen absorption.

The abundance of nitrogen fixing bacteria indicates its nitrification rate provided by the community (Zulfarina, Rusmana, Mubarik, & Santosa, 2017). It also indicates the potential nitrogen produced from the nitrification process. However, the research found that there was no significant effect of the density of *Nitrosomonas* sp. and *Nitrobacter* sp. to the concentration of nitrogen in coral tissue. The nitrification process occurred in the polyp surface, but it mainly affects the nitrogen availability in the environment. Unfortunately, the high nitrification rate is not followed by higher nitrogen uptake by coral organism.

The difference in the density of nitrogen fixing bacteria is dominantly caused by the environmental condition. One of the affecting factors of nitrifying bacteria is the depth of which the bacteria sampled. The abundance of nitrifying bacteria is higher in the shallow water (Smith, Damashek, Chavez, & Francis, 2016). Unfortunately, measurement on the depth of the sampling location was not carried out.

The bacteria is sensitive to the change of environmental condition, such a temperature (Diaz, 2017). At the research location, Karimunjawa island is the most exploited area. Thus, anthropologic aspects may play a role in changing the coral waters. However, this should be assured by investigating the condition of coral reef ecosystem. Any environmental stress may break the symbiosis of coral and algae (zooxanthellae) (Wooldridge, 2013). This may affect the other associated microorganisms in the polyp.

Nitrosomonas and *Nitrobacter* are frequently used as bioremediator in aquaculture. However, the preferences of the two is not much understood. Related to the nitrogen fixing process in Karimunjawa National Park, *Nitrosomonas* and *Nitrobacter* might have an important role in remediating coastal environment. The density of both bacterias may be related to the high input of ammonia since Karimunjawa island has been much exploited for tourism activities. *Nitrosomonas* and *Nitrobacter* mainly act as bioremediator of both pollutant (Barik *et al.*, 2018).

Even though nitrogen fixing bacteria exists in the coral mucus or tissue, but it's suggested that the bacteria are associated with Symbiodinium rather than directly to the coral (Lema *et al.*, 2012). This explains why the *Nitrosomonas* sp. and *Nitrobacter* sp. did not influence the concentration of nitrogen significantly.

The role of nitrogen fixing bacteria is obvious in the nitrogen limited environment (Putnam *et al.*, 2017). However, since the effect of *Nitrosomonas* sp. and *Nitrobacter* sp. in Karimunjawa National Park remains insignificant, it can be concluded that the source of nitrogen is sufficient. Or

otherwise, since the coral growth requires nitrogen, the coral can not accumulate the nitrogen it consumed.

CONCLUSION

Based on the analysis result, the concentration of nitrogen is higher in branching coral as the impact of its higher growth rate. However, the concentration of nitrogen in coral tissue is not affected by the abundance of nitrogen fixing bacteria.

ACKNOWLEDGMENTS

We would like to thank: Head of Universitas Diponegoro Research and Community Service Institute for PNPB research funding No. 474-19 / UN7.P4.3 / PP / 2019, and to the Head of BTNKJ for permission and information

REFERENCES

- Alamaru, A., Loya, Y., Brokovich, E., Yam, R., & Shemesh, A. (2009). Carbon and nitrogen utilization in two species of Red Sea corals along a depth gradient: Insights from stable isotope analysis of total organic material and lipids. *Geochimica et Cosmochimica Acta*, *73*, 5333–5342.
- Barik, P., Ram, R., Haldar, C., & Vardia, H. K. (2018). Study on nitrifying bacteria as bioremediator of ammonia in simulated aquaculture system. *Journal of Entomology and Zoology Studies*, *6*, 1200–1206.
- Delgadillo-Nuño, M. A., Liñán-Cabello, M. A., Reyes-Gómez, J., & Soriano-Santiago, O. (2014). Response to pH stress in the reef-building coral *Pocillopora capitata* (Anthozoa: Scleractinia). *Revista de Biología Marina y Oceanografía*, *49*, 449–459.
- Diaz, L. (2017). Response of the coral associated nitrogen fixing bacteria toward elevated water temperature. *Journal of Water Resources and Ocean Science*, *6*, 98–109.
- Edmunds, P. J., & Burgess, S. C. (2016). Size-dependent physiological responses of the branching coral *Pocillopora verrucosa* to elevated temperature and PCO₂. *The Journal of Experimental Biology*, *219*, 3896–3906.
- Hohn, S., & Merico, A. (2012). Modelling coral polyp calcification in relation to ocean acidification. *Biogeosciences*, *9*, 4441–4454.
- Krediet, C. J., Ritchie, K. B., Paul, V. J., & Teplitski, M. (2013). Coral-associated micro-organisms and their roles in promoting coral health and thwarting diseases. *Proceedings of the Royal Society B: Biological Sciences*, *280*, 1755.
- Lema, K. A., Willis, B. L., & Bourne, D. G. (2012). Corals form characteristic associations with symbiotic nitrogen-fixing bacteria. *Applied and Environmental Microbiology*, *78*, 3136–3144.
- Putnam, H. M., Barott, K. L., Ainsworth, T. D., & Gates, R. D. (2017). The vulnerability and resilience of reef-building corals. *Current Biology*, *27*, R528–R540.
- Rädecker, N., Pogoreutz, C., Voolstra, C. R., Wiedenmann, J., & Wild, C. (2015). Nitrogen cycling in corals: the key to understanding holobiont functioning? *Trends in Microbiology*, *23*, 490–497.
- Smith, J. M., Damashek, J., Chavez, F. P., & Francis, C. A. (2016). Factors influencing nitrification rates and the abundance and transcriptional activity of ammonia-oxidizing microorganisms in the dark northeast Pacific Ocean. *Limnology and Oceanography*, *61*, 596–609.
- Wahyuni, T., Kusnadi, H., & Honorita, B. (2017). Status unsur hara karbon organik dan nitrogen tanah sawah tiga Kabupaten di Provinsi Bengkulu. In S. Herlinda (Ed.), *Prosiding Seminar Nasional Lahan Suboptimal* (pp. 978–979). Palembang.
- Wang, X. T., Sigman, D. M., Cohen, A. L., Sinclair, D. J., Sherrell, R. M., Weigand, M. A., ... Ren, H. (2015). Isotopic composition of skeleton-bound organic nitrogen in reef-building symbiotic corals: A new method and proxy evaluation at Bermuda. *Geochimica et Cosmochimica Acta*, *148*, 179–190.
- Wooldridge, S. A. (2013). Breakdown of the coral-algae symbiosis: towards formalising a linkage between warm-water bleaching thresholds and the growth rate of the intracellular zooxanthellae. *Biogeosciences*, *10*, 1647–1658.
- Zulfarina, Rusmana, I., Mubarik, N. R., & Santosa, D. A. (2017). The abundance of nitrogen fixing, nitrifying, denitrifying and ammonifying bacteria in the soil of tropical rainforests and oil palm plantations in Jambi. *Makara Journal of Science*, *21*, 187–194.