

# MAPPING OF SEAGRASS COVER ON CHLOROPHYLL-A, NITRATES AND PHOSPHATE IN THE TELUK AWUR, JEPARA

*by* S. Febrianto , N. Latifah , H. Endarwati M. Zainuri; S. Suryanti

---

**Submission date:** 19-Jun-2020 04:16PM (UTC+0700)

**Submission ID:** 1346460797

**File name:** Mapping\_of\_Seagrass\_Saintek\_2019.pdf (853.11K)

**Word count:** 2258

**Character count:** 12147

## MAPPING OF SEAGRASS COVER ON CHLOROPHYLL-A, NITRATES AND PHOSPHATE IN THE TELUK AWUR, JEPARA

Sigit Febrianto<sup>1</sup>, Nurul Latifah<sup>1</sup>, Hadi Endarwati<sup>2</sup>, Muhammad Zainuri<sup>3</sup>, Suryanti<sup>1</sup>

<sup>1</sup>Aquatic Resources Departement

<sup>2</sup>Marine Science Departement

<sup>3</sup>Oceanography Departement

Faculty of Fisheries and Marine Science Diponegoro University

Jl. Prof. Soedarto SH Tembalang, Semarang, 50275

Email : [febriantosigit40@gmail.com](mailto:febriantosigit40@gmail.com)

Diterima tanggal 20 November 2019, Diterima tanggal 05 Desember 2019

### ABSTRACT

Teluk Awur Beach Jepara is one of the waters in Central Java that has a seagrass ecosystem. Seagrass role is the ability to support life of another biota closely related to the fertility of the waters indicated by the content of chlorophyll-a, nitrate, and phosphate. The purpose of this study is to know the relationship between chlorophyll-a, nitrate, and phosphate to seagrass land cover as well as the variables which have major contributions. The method of study used is descriptive with four station sampling and seagrass with a line transect of 50 m, and a 1x1 m<sup>2</sup> transect Quadrant divided into 16 sub plots measuring 25 x 25 cm<sup>2</sup> a spatial approach and multiple regression statistics. The results showed that there was a positive correlation between seagrass land cover to the concentration of chlorophyll-a and nitrate but less correlated phosphate. The highest chlorophyll-a concentrations of 0.381 mg/m<sup>3</sup>, seagrass land cover 40-60%, seagrass density 314 individual/m<sup>2</sup>, nitrate 1.141 mg/l and phosphate 0.54 mg/l. The lowest chlorophyll-a concentration ranges from 0.32-0.35 mg/m<sup>3</sup>, seagrass cover of 0-10%, seagrass density 3-12 individual/m<sup>2</sup>, nitrate 0.96 mg/l and phosphate 0, 21mg/L. This research shows that there is a strong link between seagrass land cover Concentrations of chlorophyll-a, nitrates and less influential to phosphates.

**Keywords:** chlorophyll-a; nitrate; phosphate; seagrass bed; spatial

### INTRODUCTION

Seagrass is a coastal ecosystem that contributes to the productivity of the water and has the ability to absorb CO<sub>2</sub> which is known as blue carbon and the ecological role of fishery resources, another important role of the ecosystem Seagrass is supporting the life of other biota such as ground nursery and feeding ground (Tuya *et al.*, 2014). Globally, there has been a decrease in the area of one of the seagrass fields in Indonesia (Knudby *et al.*, 2010; Lyons *et al.*, 2013). Some causes of decreased existence and the area of seagrass are natural and anthropogenic factor or combination of the two such as coastal erosion, reclamation, water enrichment due to organic waste (Howari *et al.*, 2009; Yang and Yang, 2009; Barille *et al.*, 2010). The existence of the types and patterns of seagrass fields are caused by some things, among others, seagrass, which is influenced by biological factors and aquatic physics and ecosystem services (Bostrom *et al.*, 2006). Based on this, this research aims to find out the spatial distribution, type and cover of seagrass in the Pantai Awur Bay of Jepara, knowing the difference in the concentration of chlorophyll-a, nitrates and phosphate from the seagrass cover based on density difference from highest to lowest.

### RESEARCH METHODS

The research was conducted on the coast of the Teluk Awur Jepara in October 2018. The station determination method uses purposive sampling with a number of sampling stations by four lines with 44 observation points (Fig 1) which

are based on the Seagrass Watch 2003 guide and a combination of satellite imagery analysis and field survey results.

The materials used in this study are seagrass samples, water samples according to the Seagrass cover as well as the satellite image of Sentinel 2A recording August 2018 with a spatial resolution of 10 m. Satellite imagery Data obtained from Europe Satellite Agency (ESA) and Earth Explorer of the U.S. Geological Survey (USGS). Sampling and seagrass observation are carried out to obtain seagrass, and seagrass with a line transect of 50 m, and a 1x1 m transect Quadrant divided into 16 subplots measuring 25 x 25 cm (Hulopi *et al.*, 2017). The measurement of seagrass refers to the Seagrass watch 2003 with a measurement point distance of 25 m so that in 50 m there are 11 observation points. Water sampling is carried out after known the seagrass density value on each line transect with a very tightly, moderate and rare category that is further filtered by the vacuum pump using *Whatman* paper.

### Laboratory Analysis

Seawater samples that have been distinguished based on seagrass are further analyzed using spectrophotometers by adding acetone, nitrate kits and phosphate kits and storing them in cooling for 24 hours to the next. Readings to know the value of chlorophyll-a, nitrate and phosphate.

### Data analysis

Density and percentage of cover calculated using the formula (Tuwo, 2011) Equation 1

$$K_{ji} = \frac{Ni}{A} \dots\dots\dots 1$$

$K_{ji}$ = Density type -i (ind/m<sup>2</sup>);  $N_i$ = Total number of individuals type-I (ind);  $A$ = Total area of sampling (m<sup>2</sup>)

Seagrass cover type is calculated using the formula (Tuwo,2011) Equation 2

$$PJ = \frac{ai}{A} \dots\dots\dots 2$$

$PJ$ = Seagrass cover type-i (%/m<sup>2</sup>);  $ai$ = Total cover area type-i (%);  $A$ = Total area covered of seagrass (m<sup>2</sup>)

**Statistical analysis**

The statistics used in this study are Principal Component Analysis (PCA), Inverse Distance Weighted (IDW) This is done to know the relationship between factors as well as the contribution of each factor to other factors as well as how spread from the values of these variables spatially, in this case, chlorophyll-a, seagrass cover, nitrate and phosphate (Abdi *et al.*, 2010).



Figure1. Research Location Map

**RESULTS AND DISCUSSION**

There are three types of seagrass found in the waters of Teluk Awur, among others: *Thalasia Hemprichi*, *Cymodocea serrulata*, *Enhalus acoroides*. The density and distribution of seagrass type at the research site are presented in table 1. Based on the results of the analysis of the main components indicate that there is a positive correlation between seagrass land density to the concentration of oxygen solubility of DO phosphate and nitrate but correlated to the concentration of chlorophyll-a and depth. The highest chlorophyll-a concentration of 0.381 mg/m<sup>3</sup>.

Table 1. Seagrass density in Teluk Awur Beach

No	Species	Seagrass density (ind/m <sup>2</sup> )			
		Line 1	line 2	line 3	line 4
1	<i>Thalasia hemprichi</i>	815	465	117	79
2	<i>Cymodocea serrulata</i>	0	63	0	0
3	<i>Enhalus acoroides</i>	0	3	5	12

**Spatial seagrass distribution map, density map, chlorophyll-a map, nitrate and phosphate, comparison graph chlorophyll-A, nitrate-based & phosphate density**

Distribution of phosphate in Teluk Awur beach Jepara scattered unevenly where centered on line 2 and line 3 with a concentration of 0.54 mg/l and increasingly lower when moving away from the mainland (Figure 2). Based on the results of the measurement known that the highest chlorophyll-a concentration is on the line 4 with a value of 0.38 mg/l This is due to the type of seagrass in the form of *Enhalus acoroides* (Lyons *et al.*, 2011).(Figure 4). Seagrass cover in Teluk Awur Beach is scattered uneven between the left side of the port and the right side of the harbor where on the left of the port of seagrass, reached 60% but is derived from the types of *Thalasia Hemprichi* and *Cymodocea serrulata* while on the right side of the port has a low density of less than 30% which is dominated by seagrass type of *Thalasia Hemprichi* and *Enhalus acoroides* so that the closing is low because it has a large morphology (Suryanti *et al.*, 2014; Sakey *et al.* 2015) (Figure 5.)

**Results of PCA statistics of seagrass, chlorophyll-a, nitrate, phosphate and variable water chemistry physics**

Based on the PCA chart and the value of eigenvalue more than 2, the relationship seagrass cover with water quality and nutrient then selected PC1 with a value of 2.56 (28.52%) and PC2 with a value of 2.02 (22.51%) So the total value of both 51.03% is able to explain the relationship that occurred (Coremap-Lipi, 2014). Variable DO has a very close relationship to seagrass density and is close to the central axis while other variables positively related to seagrass density are nitrates, phosphate, salinity, pH and temperature together Affects the existence and density of seagrass place in a location while the variable depth and chlorophyll-a has a negative relation to PC2 indicating the value of chlorophyll-a water and depth has no strong influence on The existence and density of seagrass in a location (McLeod *et al.*, 2011).

Based on Figure 7 it appears that stations 1, 4, 5, 7 are on one quadrant describing the proximity of the station distance and its characteristics where the station representing 4 different lines has the same class of the highest density, While stations 2, 8, 9 represent moderate density and 3.6 stations represent rare stations that are on the negative quadrant of PC1 and PC2 and indeed on this station is not found seagrass only sand substrates.




Figure 2. Map of Phosfat distribution of Teluk Awur beach




Figure 3. Map of nitrate distribution in Teluk Awur beach




Figure 4. Map of chlorophyll-a distribution in Teluk Awur beach




Figure 5. Map of seagrass cover distribution in Teluk Awur beach




Figure. 7. Data retrieval Station distribution




Figure 6. Graphic Principal components analysis and Eigenvalue

## CONCLUSION

The existence of the seagrass ecosystem is important to the coastal resources but there has been a degradation of the area and the condition of seagrass so that. The results of the principal analysis of main components indicate the variables of DO, nitrate, and phosphate become key factors of seagrass place on Teluk Awur Beach with a contribution of 51.3%. While the results of spatial analysis are known that the spread of seagrass scattered unevenly with the highest density inline 1.

## ACKNOWLEDGMENTS

The author thank Diponegoro University through a research grant with skim RPP No. 474-20/UN7. P 4.3/PP/2019. Thanks also to the laboratory PSDIL FPIK UNDIP, UNDIP Livestock Nutrition Laboratory and UNNES Chemical Laboratory which has been helpful in data analysis.

17

## REFERENCES

- Abdi H., Williams L.J., 2010 Principal component analysis. *WIREs Computational Statistics* 2:433-459.
- Barille, L., Robin, M., Harin, N., Bargain, A, Launeau, P., 2010. Increase in seagrass distribution at Bourgneuf Bay (France) detected by spatial remote sensing. *Aqu Bot.* 92, 185-194.
- Bostrom, C., Jackson, E.L, Simenstad, C.A., 2006. Seagrass landscapes and their effects on associated fauna: a review. *Estuar.Coast.Shelf Sci.* 68, 383-403.
- COREMAP-LIPI, 2014. Panduan Monitoring Padang Lamun. Pusat Penelitian Oseanografi LIPI, Jakarta.
- Howari, Fares M., Benjamin, R, Jordan, Naima, Bouhouche, Sandy Wyllie-Echeverria. 2009. Field and Remote-Sensing Assessment of Mangrove Forests and Seagrass Beds in the Northwestern Part of the United Arab Emirates. *Journal of Coastal Research.* 25, 48-56.
- Hulopi, M., Tuahatu, J. W., & Tuhumury, N. C. (2017). Seagrass potency as a blue carbon source in Galala and Tanjung Tiram waters, Ambon Island, Indonesia, *IO(5)*, 1019-1025.
- Knudby, A., Newman, C., Shaghude, Y., Muhando, C., 2010. Simple and effective monitoring of historic changes in nearshore environments using the free archive of Landsat imagery. *International Journal of Applied Earth Observation and Geoinformation* 12, S116S122.
- Lyons, M.B., Phinn, S.R., Roelfsema, C.M., 2011. Integrating Quickbird multi-spectral satellite and field data: mapping bathymetry, seagrass cover, seagrass species and change in Moreton Bay, Australia in 2004 and 2007. *Remote Sensing* 3, 42e64.
- Lyons, Mitchell B., Chris M, Roelfsema., Stuart R, Phinn. 2013. Towards understanding temporal and spatial dynamics of seagrass landscapes using time-series remote sensing. *Estuarine, Coastal and shelf science.* 120, 42-53.
- M.A. Hemminga and C.M. Duarte, *Seagrasses Ecology*, Cambridge University Press, Cambridge, p. 248 - 251 (2000).
- McLeod, E., Chmura, G.L., Bouillon, S., Salm R, Bjork, M., Duarte, C.M., Lovelock, C.E., Schlesinger, W.H., Siliman, B.R., 2011. A blue carbon : toward an improved understanding of the role of vegetated coastal habitats in sequestering CO2. *Front. Ecol.*
- Sakey. W. F., B. T. Wagey dan G. S. Gerung. 2015. Variasi Morfometrik pada Beberapa Lamun di Perairan Semenanjung Minahasa. *Jurnal Pesisir dan Laut Tropis.* Vol 1. No 1.
- Suryanti., Ain, C. & Tismawati, C.N. (2014). Hubungan Kerapatan Lamun (Seagrass) dengan Kelimpahan

- 14 Syngnathidae di Pulau Panggang Kepulauan Seribu. Diponegoro Journal of Maquares, 3(4). 147153
- Tuwo, A. (2011). Pengelolaan Ekowisata Pesisir dan Laut. Brillian Internasional. Sidoarjo.
- 10 Tuya, F., Haroun, R., Espino, F., 2014. Economic assessment of ecosystem services: Monetary value of seagrass meadows for coastal fisheries. Ocean Coast Manag. 96, 181–187.
- 1 Yang, D., Yang, C., 2009. Detection of seagrass distribution changes from 1991 to 2006 in Xincun bay, Hainan, with satellite remote sensing. Sensors 9, 830–844.

# MAPPING OF SEAGRASS COVER ON CHLOROPHYLL-A, NITRATES AND PHOSPHATE IN THE TELUK AWUR, JEPARA

## ORIGINALITY REPORT

22%

SIMILARITY INDEX

%

INTERNET SOURCES

22%

PUBLICATIONS

%

STUDENT PAPERS

## PRIMARY SOURCES

- 1** Bijeesh Kozhikkodan Veetil, Raymond D. Ward, Mariana Do Amaral Camara Lima, Milica Stankovic, Pham Ngoc Hoai, Ngo Xuan Quang. "Opportunities for seagrass research derived from remote sensing: A review of current methods", *Ecological Indicators*, 2020  
Publication **3%**
- 2** Vivi Endar Herawati, Pinandoyo, YS Darmanto, Johannes Hutabarat. " Growth Performance and Nutrient Content of Carp ( ) With the Feeding of Maggot Meal Substitution Cultivated in Different Media ", *IOP Conference Series: Earth and Environmental Science*, 2019  
Publication **2%**
- 3** A Junaidi, C Kartiko. "Design of Pond Water Quality Monitoring System Based on Internet of Things and Pond Fish Market in Real-Time to Support the Industrial Revolution 4.0", *IOP Conference Series: Materials Science and Engineering*, 2020 **2%**

4

Kevan Moffett, William Nardin, Sonia Silvestri, Chen Wang, Stijn Temmerman. "Multiple Stable States and Catastrophic Shifts in Coastal Wetlands: Progress, Challenges, and Opportunities in Validating Theory Using Remote Sensing and Other Methods", Remote Sensing, 2015

Publication

---

2%

5

John A. Burt. "The environmental costs of coastal urbanization in the Arabian Gulf", City, 2014

Publication

---

2%

6

Tiit Kutser, John Hedley, Claudia Giardino, Chris Roelfsema, Vittorio E. Brando. "Remote sensing of shallow waters – A 50 year retrospective and future directions", Remote Sensing of Environment, 2020

Publication

---

1%

7

J. Lucas Pérez-Lloréns, Juan J. Vergara, Irene Olivé, Jesús M. Mercado et al. "Chapter 9 Autochthonous Seagrasses", Springer Science and Business Media LLC, 2014

Publication

---

1%

8

Brenda V.A. Lahiwu, Rose O.S.E. Mantiri, Ferdinand F. Tilaar, Laurentius Th. X. Lalamentik, Ruddy D. Moningkey, Billy T.

1%



Wagey. "Zooplankton Community Structure in Seagrass Ecosystem, Tongkaina Village, Bunaken Darat District, Manado City", JURNAL PERIKANAN DAN KELAUTAN TROPIS, 2019

Publication

---

9

S Suryanti, C Ain, N Latifah. "Mapping of Nitrate, Phospat And Zooxanthelae With Abundance Of Sea Urchins on Massive Coral Reef in Karimunjawa Island", IOP Conference Series: Earth and Environmental Science, 2018

Publication

---

10

"Seagrasses of Australia", Springer Science and Business Media LLC, 2018

Publication

---

11

Hossain, M.S., J.S. Bujang, M.H. Zakaria, and M. Hashim. "The application of remote sensing to seagrass ecosystems: an overview and future research prospects", International Journal of Remote Sensing, 2015.

Publication

---

12

HANNAH M. MURPHY. "Response of fauna in seagrass to habitat edges, patch attributes and hydrodynamics", Austral Ecology, 12/2009

Publication

---

13

Ruiliang Pu, Susan Bell, Cynthia Meyer. "Mapping and assessing seagrass bed changes in Central Florida's west coast using

1%

1%

1%

1%

1%

multitemporal Landsat TM imagery", Estuarine,  
Coastal and Shelf Science, 2014

Publication

---

14

Claudio David Togas, Grace O Tambani, Nurdin Jusuf. "ANALISIS KELAYAKAN USAHA BUDIDAYA IKAN SISTEM KARAMBA JARING TANCAP DI DESA TALIKURANKECAMATAN REMBOKEN KABUPATEN MINAHASA PROVINSI SULAWESI UTARA", AKULTURASI (Jurnal Ilmiah Agrobisnis Perikanan), 2016

Publication

---

1%

15

Radhouan El Zrelli, Lotfi Rabaoui, Rubén H. Roa-Ureta, Nicola Gallai et al. "Economic impact of human-induced shrinkage of Posidonia oceanica meadows on coastal fisheries in the Gabes Gulf (Tunisia, Southern Mediterranean Sea)", Marine Pollution Bulletin, 2020

Publication

---

<1%

16

Rizki Aulia Ansari, Tri Apriadi, Agung Dhamar Syakti. "STOK KARBON LAMUN THALLASIA HEMPRICHII DAN SEDIMEN PULAU BINTAN KEPULAUAN RIAU", Jurnal Ruaya : Jurnal Penelitian dan Kajian Ilmu Perikanan dan Kelautan, 2020

Publication

---

<1%

17

Krzysztof Lamorski, Andrzej Bieganowski, Magdalena Ryżak, Agata Sochan, Cezary

<1%

Sławiński, Wioleta Stelmach. "Assessment of the usefulness of particle size distribution measured by laser diffraction for soil water retention modelling", Journal of Plant Nutrition and Soil Science, 2014

Publication

---

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

# MAPPING OF SEAGRASS COVER ON CHLOROPHYLL-A, NITRATES AND PHOSPHATE IN THE TELUK AWUR, JEPARA

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---