

Effect of Secang Wood (*Ceasalpinia sapan* L.) Extract on Morphology of Spermatozoa, Sperm Count, and Reversible Process in Male Rates

by Suhartono Suhartono

Submission date: 27-Oct-2020 02:28PM (UTC+0700)

Submission ID: 1427922709

File name: 14.pdf (391.35K)

Word count: 2661

Character count: 13957

²
Arum SS, et al. *Belitung Nursing Journal*. 2017 April;3(2):142-147
Accepted: 28 February 2017
<http://belitungraya.org/BRP/index.php/bnj/>

© 2017 The Author(s)

This is an Open Access article distributed under the terms of the [Creative Commons Attribution 4.0 International License](#) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORIGINAL RESEARCH

ISSN: 2477-4073

EFFECT OF SECANG WOOD (*CEASALPINIA SAPPAN L.*) EXTRACT ON MORPHOLOGY OF SPERMATOZOA, SPERM COUNT, AND REVERSIBLE PROCESS IN MALE RATS

Silfia Sekar Arum^{1*}, Syarief Thaufik Hidayat², Nur Khafidhoh², Suryati
Kumorowulan¹, Suhartono³, Ari Suwondo⁴

¹Magister Applied Midwifery, Health Ministry Polytechnic Semarang, Semarang, Indonesia

²RSUP. DR. Kariadi, Semarang, Indonesia

³Universitas Diponegoro, Semarang, Indonesia

⁴Jurusan Kesehatan dan Keselamatan Kerja, Fakultas Kesehatan Masyarakat, Universitas Diponegoro,
Semarang, Indonesia

*Corresponding author:

Silfia Sekar Arum, M.SST

Magister Applied Midwifery, Health Ministry Polytechnic Semarang, Jl. Tirta Agung, Pedalangan,
Banyumanik, Kota Semarang, Jawa Tengah, Indonesia (50268) e-mail: silfiaelly77@gmail.com

¹ABSTRACT

Background: Limited choice of contraception for males is one of the reasons why their participation is low. Secang wood is considered as an alternative contraception that has an influence in the morphology and number of spermatozoa and reversible in nature.

Objective: To examine the effect of extracts of secang wood in the morphology of spermatozoa, sperm count and reversible process in male mice.

Methods: This was a quasi-experimental study with randomized posttest only control group design. There were 32 male rats (*Mus Musculus L*) recruited in this study. Four groups were involved, namely: 1) ethanol group, 2) chloroform fraction, 3) water fraction, and 4) control group. Each group consisted of 8 mice. There were two treatments in this study: 1) each group was given secang extract with dose 50 mg/25 gram of weight, 2) the observation period was 10 days after the treatment. Four mice in each group was dissected to see the morphology and sperm count while the other four mice were allowed to live and maintained until 20 days to see reversible morphology and sperm count.

Results: The extract of secang wood at a dose of 50mg/25gram weight could increase the abnormal sperm morphology and lower sperm count. There was a significant difference between the treatment group and control group with p-value <0.05. The extract was also reversible on the morphology and sperm count.

Conclusion: There was a significant effect of secang wood extract on sperm morphology, the number of spermatozoa, and reversible process. This study provides the insight of scientific information about the effect of the secang extracts on the number and morphology of spermatozoa, and it could be used as a basis for further research in human in the development of natural contraceptive on a reversible man.

Keywords: secang wood, ceasalpinia sappan l, spermatozoa, contraception

³

Belitung Nursing Journal , Volume 3, Issue 2, March-April 2017

142

Editor's Note: This article has been updated on 17 June 2020 in terms of minor changes in the reference format. The update is according to BNJ Policy on article correction.

INTRODUCTION

Reasons for low male participation in family planning program is limited choice of contraception for male, and is more directed to females.^{1,2} Until today, the contraception modes for male are just limited to condom, vasectomy, and hormone injections.² However, ideally, contraception for men should be able to prevent fertilization, safe, fast performance, no side effects, temporary, and no effect on sex and libido.

In Indonesia, one of the things that is under developed is the use of natural medicinal plant as an alternative anti male fertility. One of them is secang wood (*Caesalpinia sappan L*). This plant empirically and in vitro-proof has anti fertility effect on cell donor of human spermatozoa.³

The active compounds of secang wood that can be used as antifertility are flavonoids that can inhibit the enzyme aromatase, namely an enzyme that has a function to catalyze androgen conversion to be estrogen that increases testosterone, so it will inhibit spermatogenesis.⁴ Alkaloid can affect the secretion of hormone reproduction, which is needed for the process of spermatogenesis and lower the quality of spermatozoa.³ Beside flavonoid and alkaloid, another essential compound in the secang wood is aetheric oil that can be used as an alternative infertility, which has function to work on the transportation of sperm, and agglomerate sperm in order to reduce sperm vitality, which consequently affect the sperm not to reach the egg and fertilization can be prevented.⁵

Secang wood is empirically known having many healing properties and is often consumed by community as healthy drink. It also has a powerful antioxidant extracts with antioxidative indexes, which is higher than the commercial antioxidants (BHT and BHA).⁶ Thus, the purpose of this study was to examine the effect of

secang wooden extract on sperm morphology, sperm count and reversible process in male rats (*Mus Musculus L*).

METHODS

Design

This was a quasi-experimental study with randomized posttest only control group design.

Population and Research Subject

Thirty-two male rats (*Mus Musculus L*) recruited in this study with the inclusion criteria: mature and healthy mice, aged 3 months, 20-30 grams weight. There were four groups in this study: 1) ethanol group, 2) chloroform fraction, 3) water fraction, and 4) control group. Each group consisted of 8 mice.

Intervention

There were two treatments in this study: 1) each group was given secang extract with dose 50 mg/25 gram of weight, 2) The observation period was 10 days after the treatment, and 4 mice in each group was dissected to see the morphology and sperm count, while the other 4 mice were allowed to live and maintained until 20 days to see reversible morphology and sperm count.

Instrument

The instrument used was an individual mouse cage, maceration tool, Neubauer hemocytometer micro, microscope, watch glass, surgical equipment, pipette test, pipette leukocytes, petri dishes, cover glass, tray stopwatch, books and recording data.

Data Analysis

Data were analyzed using one-way ANOVA test to find out the mean difference between the treatment groups and then continued with Tukey test to compare the mean of the entire treatments

in order to know the most influence treatment.

Ethical Consideration

This research has been approved by the Health Ministry Polytechnic Semarang with No. 024/KEPK/polytechnic-SMG/EC/2017.

RESULTS

Morphology of Spermatozoa

The morphological changes in spermatozoa after a given treatment for 10 days shown in table 1 indicated that the means

of abnormal sperm morphology in the treatment groups (ethanol extract 13.50, chloroform fraction 11.50, and water fraction 6.75) were higher than sperm morphology in the control group (3.75). The highest average of abnormal sperm morphology was found in the group of ethanol extract (13.50) and the lowest was in the control group (3.75). While the reversible morphological changes after 20 days could be seen that the ethanol extract group (6.00), chloroform fraction (5.25), and the control group (3.75) were lower than the water fraction group (3.00).

Table 1 Morphology of Spermatozoa

Groups	n	Intervention (X±SD %)	Reversible (X±SD %)
Control	4	3.75±0.957	3.75±0.957
Ethanol extract	4	13.50±2.082	6.00±0.816
Chloroform fraction	4	11.50±1.915	5.25±0.957
Water fraction	4	6.75±2.062	3.00±0.816

Table 2 Sperm Count of Male Rats

Groups	n	Intervention (X±SD %)	Reversible (X±SD %)
Control	4	13.25±2.062	11.50±2.246
Ethanol extract	4	8.25±0.957	12.25±1.258
Chloroform fraction	4	7.75±0.957	10.25±1.500
Water fraction	4	8.25±2.500	11.25±3.304

Table 3 One-Way ANOVA of Morphology of Spermatozoa and Its Reversible

	Intervention					Reversible				
	Sum of Squares	df	Mean	F	Sig	Sum of Squares	df	Mean	F	Sig
Between Groups	236.250	3	78.750	23.924	.000	22.500	3	7.500	9.474	.002
Within Group	39.500	12	3.292			9.500	12	.792		
Total	275.750	15				32.000	15			

Table 4 One-Way ANOVA of the Number of Spermatozoa and Its Reversible

	Intervention					Reversible				
	Sum of Squares	df	Mean	F	Sig	Sum of Squares	df	Mean	F	Sig
Between Groups	80.70	3	26.927	8.730	.002	39.688	3	13.229	2.433	.115
Within Group	37.500	12	3.083			65.250	12	5.438		
Total	117.750	15				104.938	15			

Sperm count

Table 2 shows that changes in the number of spermatozoa after given the treatment for 10 days in the treatment groups (ethanol extract 8.25; chloroform fraction 7.75; and water fraction 8.25) were lower than in the control group (13.25). The lowest number of sperm was found in the group of chloroform fraction. However, the reversible number of spermatozoa showed that the means of sperm count in the treatment groups (ethanol extract 12.25; chloroform fraction 10.25; and water fraction 10.25) were increased than the mean in the control group (11.50). The results of Shapiro-Wilk test for all variables showed normal distribution, so further analysis was conducted using one-way ANOVA.

Table 3 shows that morphology of spermatozoa after given the treatment showed p-value = 0.000 and reversible morphology of spermatozoa showed the p-value = 0.002 (<0.05), indicated that there were significant differences in the mean of morphology of spermatozoa and its reversible.

On the other hand, the number of spermatozoa after given the treatment in Table 4 shows p-value = 0.000 (< 0.05), indicated that there was a significant difference in the mean of sperm count. But the reversible of spermatozoa count showed p-value = 0.115 (>0.05), which indicated that there were no significant differences in the mean of reversible sperm count.

Tukey test shows that there were significant differences in the sperm morphology between the control group (0.000) with the other groups (ethanol group 0.001; chloroform 0.014; and water fraction 0.143). Among the treatment groups, ethanol group (0.001) had the most significant differences compared to the other treatment groups (chloroform 0.14; and water fraction 0.14). While in the number of spermatozoa, it also

showed that there were significant differences between the control group (0.008) and the treatment groups (ethanol group 0.977; chloroform 0.008; water fraction 1.000).

The result of the Tukey test on the reversible revealed that there were no significant differences in sperm morphology between the control group (0.017) with the other groups (ethanol group 0.643; chloroform 0.017; and water fraction 0.643). Among the treatment groups, chloroform (0.017) had the most significant differences compared to the other treatment groups (ethanol 0.017; water fraction 0.643).

DISCUSSION

This study found the primary abnormality in the form of a small head, amorphous head, and spiral tail due to a decrease in testosterone levels. Secondary abnormality is seen from the headless spermatozoa and tailless, which is in line with the previous study said that these abnormalities were caused by disruption in the process of sperm maturation in epididymis.⁷ The chemicals contained in secang wood (*Caesalpinia sappan L*) is an alkaloid which can affect spermatogenesis by pressing the secretion of reproductive hormones (FSH and LH) required for the ongoing spermatogenesis and lower the quality of spermatozoa. FSH serves to spur spermatozoa process, namely the formation of spermatogonia into spermatids.⁸ LH serves to stimulate spermatogenesis process, namely the formation of spermatid cells become spermatozoa.⁹

The decline in the number of spermatozoa after given the extract of secang wood was because the active substance contained in the secang wood such as flavonoid compound, as a class of compounds that function as anti-androgenic by inhibiting the enzyme aromatase, the enzyme that serves to

catalyze the immediate conversion of androgens to estrogen that increases the testosterone.¹⁰

Based on the results of this study, it could be said that the effect of secang wood on the morphology and sperm count is reversible. It is because after the treatment was stopped, the morphology and sperm count increased. This is assumed that the active Leydig cells produced testosterone again. These findings were in line with the study conducted by Ermayanti et al¹¹ revealed that there was an increase in testosterone levels back into the blood plasma of mice after secang wood treatment was stopped; and it was also reported that an increase in Leydig cell activity in mice by in vitro after the secang wood treatment stopped.¹⁰

Limitations of the Study

Insignificant difference between the time of observation might be due to the time was too short, which only for 20 days. It was possibly required more time for the recovery, since the sperm had already damaged. It was assumed that the restoration of spermatozoa only occurred with spermatozoa that were able to adapt.

CONCLUSION

The secang wood extract significantly affected the sperm morphology, the number of spermatozoa, and the reversible process. This study provides the insight of scientific information about the effect of the secang extracts on the number and morphology of spermatozoa and could be used as a basis for further research in human in the development of natural contraceptive on a man who is reversible.

Declaration of Conflict of Interest

None declared.

Funding

This study was supported by Health Ministry Polytechnic Semarang, Semarang, Indonesia.

Authorship Contribution

The authors equally contributed in this study.

References

1. Kabagenyi A, Jennings L, Reid A, Nalwadda G, Ntozi J, Atuyambe L. Barriers to male involvement in contraceptive uptake and reproductive health services: A qualitative study of men and women's perceptions in two rural districts in Uganda. *Reproductive Health*. 2014;11(1):21.
2. Page ST, Amory JK, Bremner WJ. Advances in male contraception. *Endocrine Reviews*. 2008;29(4):465-493.
3. Guyton AC. *Human physiology and mechanisms of disease*. Philadelphia: WB Saunders Co; 1992.
4. Alfi NF. *Pengaruh fraksi etanol dan fase air daun gendarussa vulgaris nees pada proses fertilisasi in vitro mencit [Effect of ethanol fraction and water phase of gendarussa vulgaris nees on the process of in-vitro fertilization of mice]*. Surabaya: Universitas Airlangga; 2003.
5. Lim D-K, Choi U, Shin D-H. Antioxidative activity of some solvent extract from caesalpinia sappan L. *Korean Journal of Food Science and Technology*. 1996;28(1):77-82.
6. Alberts B, Johnson A, Lewis J, Raff M, Roberts K, Walter P. *Molecular biology of the cell*. New York: Garland science; 2002.
7. Hardy MP, Sottas CM, Ge R, et al. Trends of reproductive hormones in male rats during psychosocial stress: role of glucocorticoid metabolism in behavioral dominance. *Biology of Reproduction*. 2002;67(6):1750-1755.
8. Rusmiati. *Pengaruh ekstrak kayu secang (caesalpinia sappan L.) terhadap viabilitas spermatozoa mencit jantan (mus musculus) [Effect of caesalpinia sappan L. extract on viability of mus musculus spermatozoa]*. Yogyakarta, Indonesia: Universitas Gadjah Mada; 2005.
9. Zaneveld LJD, Chatterton RT. *Biochemistry of mammalian reproduction*. New York: Wiley; 1982.

10. Priastini R. Tanaman obat alami di Indonesia sebagai alternatif antifertilisasi laki-laki [Natural medicinal plants in Indonesia as an alternative to male antifertilization]. *Jurnal Kedokteran Meditek*. 2014;15(39B).
11. Ermayanti NGA, Suarni NI, Made RAI. Kualitas spermatozoa mencit (mus musculus l.) setelah perlakuan infus kayu amargo (quassia amara linn.) dan pemulihannya [Quality of mus musculus l spermatozoa after given

infusion of quassia amara linn. abd its recovery]. *Jurnal Biologi*. 2010;14(2).

Cite this article as: Arum SS, Hidayat ST, Khafidhoh N, K¹norowulan S, Suhartono, Suwondo A. Effect of secang wood (ceasalpinia sappan l.) extract on morphology of spermatozoa, sperm count, and reversible process in male rats. *Belitung Nursing Journal*. 2017;3(2):142-147. <https://doi.org/10.33546/bnj.78>

Effect of Secang Wood (Ceasalpinia sapan L.) Extract on Morphology of Spermatozoa, Sperm Count, and Reversible Process in Male Rates

ORIGINALITY REPORT

19%

SIMILARITY INDEX

21%

INTERNET SOURCES

8%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1

garuda.ristekbrin.go.id

Internet Source

14%

2

Yuni Sulistiawati, Ari Suwondo, Triana Sri Hardjanti, Ariawan Soejoenoes, M. Choiroel Anwar, Kun Aristiati Susiloretni. "EFFECT OF MORINGA OLEIFERA ON LEVEL OF PROLACTIN AND BREAST MILK PRODUCTION IN POSTPARTUM MOTHERS", Belitung Nursing Journal, 2017

Publication

3%

3

Submitted to Adtalem Global Education

Student Paper

2%

Exclude quotes On

Exclude matches < 2%

Exclude bibliography On