

Length-Weight Relationships and Condition Factor of *Anabas testudineus* in the Semayang Lake, East Kalimantan, Indonesia

by Frida P


Submission date: 26-Jul-2022 09:04PM (UTC+0700)

Submission ID: 1875443567

File name: C5_-_Length-weight_relationships_and_condition_factor.pdf (663.75K)

Word count: 5980

Character count: 28114



Length-weight relationships and condition factor of *Anabas testudineus* in the Semayang Lake, East Kalimantan, Indonesia

^{1,2}Mohammad Mustakim, ¹Sutrisno Anggoro, ¹Frida Purwanti, ¹Haeruddin

¹ Doctoral Program in Coastal Resources Management, Faculty of Fisheries and Marine Science, Diponegoro University, Tembalang, Semarang, Central Java, Indonesia;

² Department Aquatic Resources Management, Faculty of Fisheries and Marine Science, Mulawarman University, East Kalimantan, Indonesia. Corresponding author: M. Mustakim, moh.mustakim@fpik.unmul.ac.id; mustakim.unmul2005@gmail.com

Abstract. The environment of the Semayang Lake is a floodplain lake area that has unique habitat characteristics in the East Kalimantan province, Indonesian. *Anabas testudineus* is one of the fish found in Semayang Lake. The length-weight relationship research have been conducted in Semayang Lake, for one full year (dry and rainy season) in providing basic data in resources management of *A. testudineus* so that this fish species can be harvested optimally and sustainably. Growth pattern analysis of the allometric linear model (LAM) was used to calculate intercept (a) and slope (b). The relative weight (Wr) was determined by the equation $Wr = (W / Ws) \times 100$. The condition factor was determined from the Fulton coefficient (K). The results of the present research suggest that *A. testudineus* has negative allometric growth pattern. The coefficient determination value (R^2) of male individuals at swamp habitat is at range of 76-94% (r value: 0.87-0.97), females ranges 78-98% (r value: 0.889-0.99) and coefficient determination value (R^2) on male in lake habitat ranges 77-93% (r value: 0.877-0.96); female ranges 75-92% (r value: 0.866-0.959). Relative weight (Wr) value obtained from the standard weight and total weight in swamp habitats in males ranged from 100.04 ± 0.72 to 100.93 ± 2.41 , the females ranged from 100.25 ± 1.39 to 101.13 ± 2.60 . In lake habitat the relative weight value in male fish was 100.29 ± 1.57 - 101.06 ± 2.65 , and for females 100.40 ± 2.98 - 101.60 ± 3.07 . The observed weight in swamp habitat was 34.2205 ± 0.397 higher than the average standard weight of 34.012 ± 0.386 . Condition factor of *A. testudineus* in swamp habitat on male fish was 3.102 ± 0.0058 , in female fish 3.141 ± 0.007 , whereas in the lake habitat (Station 2, 3 and 5) condition factor was 3.114 ± 0.007 and 3.143 ± 0.009 for females respectively.

Key Words: growth pattern, relative weight, climbing perch, *A. testudineus*.

Introduction. The Semayang Lake is a floodplain lake area that has unique habitat characteristics in the East Kalimantan province, Indonesia, because on the water bodies there are swamps and lakes habitats. Swamp habitats are characterized by a large coverage of aquatic plants and shallow water (Archibold 1995; Lopez & Kursar 2007; Corlett & Primack 2011). The lake habitat is characterized by relatively broad water surface, high quantity of aquatic plants at the littoral area and has deeper water than the swamp habitat (Pokorný & Björk 2010; Romanowski 2013).

Inundation is the main characteristic of floodplain lake caused by simultaneous hydrodynamics and can be distinguished by the duration and frequency of puddles which are the main cause of ecological differences between swamp and lake habitats. This hydrodynamic occurrence is valid for a one-year cycle, but at the same location even though the flow of water and the height of the inundation from month to month even from year to year shows different traits and characters (Furukawa 1994; Mitsch & Gosselink 1993). This hydrodynamic condition occurs in the Semayang Lake (Noor 2007) reported that puddles length in the surroundings of Semayang Lake have a duration that is almost throughout the year. The main different in this case is the change in the water level, where during the rainy season, that have swamp habitats characteristics is

uniform, otherwise during the dry season the both habitat is separated and connected only by a channel of river.

The connection changes in habitat have implications to the dynamics of fish biology, and similarly, in the case of fish biology (growth patterns and condition factors). *Anabas testudineus* is a resident fish species (Welcomme 1995) and is naturally found in many countries from India to the Wallace line (Kohinoor et al 1991; Tay et al 2006; Mustakim et al 2009), predominantly of lentic habitats (swamps and lakes) and lotic habitats (rivers and canals). *A. testudineus* can thrive in low dissolved oxygen water bodies (Pethiyagoda 1991; Kumary & Raj 2016). This species has a high selling price due to its taste and soft meat texture (Helmizuryani & Muslimin 2016). *A. testudineus* is one of the fish species inhabiting Semayang Lake.

The biological aspect of fish that is one of the instruments in the management of fish resources is the length-weight relationship (Khristenko & Kovska 2017; Tsoumani, et al 2006), focus the length-weight relationship of fish can be used to determine the pattern of fish growth (Mousavi-Sabet et al 2013; Vilizzi et al 2013; Lorenzoni et al 2015), further condition factor that are important derivatives of fish growth (Everhart & Youngs 1981; Tsoumani et al 2006; Froese 2006). Research of length-weight relationship and condition factors of *A. testudineus* have generally been previously conducted in other biotopes (Alam et al 2007; Mustakim et al 2009; Begum & Minar 2012; Kumar et al 2013; Rahman et al 2015; Kumary & Raj 2016). However, length-weight relationship research has not been conducted in Semayang Lake, for a whole year (dry and rainy season), this matter, signify why this research is important, as the preliminary data in resources management of *A. testudineus* so that can be harvested optimally and sustainably.

Material and Method. The research location is located in Semayang Lake, with the selection of research stations based on the characteristics of swamp habitats (stations 1, 4 and 6) and lake habitats (stations 2, 3 and 5) (Figure 1).




Figure 1. Research Location Map (Semayang Lake East Kalimantan Province, Indonesia).

This research has been conducted for one year (February 2017 - January 2018). Samples were collected from the fishermen using non-selective trapping gear category (traditional name is Pengilar and Sawaran). The sample number is 10% of the catch, the sample were then preserved in 10% formalin solution then analyzed in the laboratory for total length and total weight measurements. The total length was measured using a ruler of 1 mm accuracy and the total weight was measured with a digital scale of 0.1 g accuracy (Anderson & Neumann 1996). The total length (TL) was expressed in mm units and total weight (TW) in gram units. Sexes were determined by dissection and observation of the color and gonad shape.

Growth pattern analysis. The Allometric Linear Model (LAM) was applied to calculate a and b parameters through measurement of weight and length changes. The bias correction on the mean weight change of the logarithmic unit was used to predict the weight of length parameters of allometric equations, according to De Robertis & Williams (2008). $W = a TL^b$, where W is the weight of the fish (g), TL is the total length of the fish (mm), a (intercept) and b (slope). The value of a (intercept) obtained from the relationship of total length (mm) and weight (g) was converted by the formula $a' = a10^b$ (Froese 2006). The t_{test} was performed on the value of b to find out whether $b = 3$ (isometric) or $b \neq 3$ (allometric) to examine the value of b , it is necessary to compute the t test with the following hypotheses and formulas. Hypothesis: $H_0: b = 3$ $H_1: b \neq 3$;

$T_s = \frac{\beta_0 - \beta_1}{S\beta_1}$ Decision making on the hypothesis is done by comparing t_{test} with t_{table} at 95% confidence interval. If $t_{test} > t_{table}$, then the decision is to reject H_0 . If $t_{test} < t_{table}$, then the decision is to accept the H_0 (Steel & Torrie 1993).

Condition factor. Relative weight (W_r) and coefficient (K) condition factors were used to evaluate the condition factor of each individual. The relative weight (W_r) was determined by the equation proposed by Anderson & Neumann (1996) as follows: $W_r = (W / W_s) \times 100$, W_r is the relative weight, W weight of each fish, and W_s is the predicted standard weight of the sample which is the same because it is calculated from a combination of length-weight regression over distances between species: $W_s = aL^b$. The coefficient of the Fulton (K) condition Factor was determined by the following formula: $K = WL^{-3} \times 100$ Okgerman (2005), where K is the condition factor, W is the weight (g), L is the total length (mm) and -3 is the coefficient.

Result and Discussion

Length-weight relationship. The total number of *A. testudineus* caught during the study consisted of 1,592 individuals, which of 566 males and 409 females collected in swamp habitats, and 335 males and 282 females collected from lake habitats (Table 1). The average total length and total weight in swamp habitat (Station 1, 4 and 6) of male individuals was 119.8 ± 0.55 mm, 31.72 ± 0.46 g, and 126.6 ± 0.75 mm, 37.69 ± 0.67 g of female specimens. In lake habitat (Station 2, 3 and 5) the average total length and total weight of males was 121.99 ± 0.66 mm and 32.91 ± 0.55 , and 125.81 ± 0.90 and 37.47 ± 0.86 of females. The results of the analysis of the length and total weight relationship month⁻¹ showed different b (slope) values between males and females, in both habitats, swamp (stations 1, 4 and 6) and lake (2, 3 and 5) the value of b (slope) during the study is $b \neq 3$ ($b < 3$; $3 < b$), so that the growth pattern of *A. testudineus* is negative allometric and positive allometric. *A. testudineus* growth patterns in Semayang Lake changed dynamically; this result is consistent with previous studies that the value of b (slope) was depending on physiological and environmental conditions such as temperature, pH, salinity, geographical location and sampling techniques (Jennings et al 2001; Turkmen et al 2002) and also on biological condition development of gonads and food availability including growth phase, season, abdominal fullness, gonadal maturity, gender, range of measures, and disease (Le Cren 1951; Tesch 1971; Neff & Cargnelli 2004; Ecoutin et al 2005; Froese 2006).

Table 1
Length and weight of *Anabas testudineus* (males and females) harvested from Lake Semayang in February 2017 - January 2018

Month	Sexes	N	Swamp			Lake				
			TL (mm)		TW (g)	TL (mm)		TW (g)		
			(Mean±SE)	Range	(Mean±SE)	(Mean±SE)	Range	(Mean±SE)		
Feb-17	M	82	115.2±1.66	87-171	29.23±1.41	11.2-79.4	120.19±2.46	94-173	32.279±1.96	12.9-87.4
	F	45	126.3±2.82	85-193	38.89±2.280	11.2-88.2	122.72±3.33	100-193	35.42±2.63	16.0-83.5
	M	77	117.9±1.38	95-148	30.59±1.28	16.11-58.6	122.16±1.69	100-150	33.02±1.39	16.0-54.4
Mar-17	M	45	120.3±1.95	97-149	34.22±1.89	13.4-60.6	121.41±2.12	100-145	33.54±1.69	16.5-52.1
	F	30	118.2±2.29	94-152	31.00±2.29	13.6-72.4	120.19±1.63	99-140	30.37±1.66	15.6-52.8
Apr-17	M	25	128.9±4.80	83-172	42.51±4.56	10.5-81.4	118.04±3.83	93-154	32.27±4.08	11.2-81.8
	F	53	117.2±2.12	84-151	30.31±1.80	11.7-68.0	118.05±2.07	101-141	32.13±2.22	18.1-55.6
May-17	M	22	116.0±3.84	78-155	30.36±2.45	7.9-81.5	122.00±5.78	97-160	36.35±6.41	16.5-85.4
	F	26	116.5±2.39	94-144	29.43±1.76	16.3-52.3	119.68±2.35	103-145	30.53±2.09	16.6-60.4
Jun-17	M	29	125.3±2.22	102-148	36.27±1.92	20.8-57.2	122.70±2.29	102-144	33.91±2.08	21.8-55.4
	F	29	118.0±2.07	92-142	31.96±1.56	16.0-49.3	117.93±1.68	100-142	30.65±1.30	15.8-51.5
Jul-17	M	34	130.1±2.36	105-158	40.41±2.05	22.5-66.0	125.58±4.25	95-170	37.71±4.09	16.6-92.9
	F	54	116.9±1.45	97-144	28.28±1.04	11.7-48.4	119.11±3.41	99-142	30.63±2.15	18.2-49.4
Aug-17	M	21	123.2±2.18	104-140	31.98±1.89	18.2-47.2	128.00±3.13	116-142	37.25±2.73	28.3-49.2
	F	35	128.3±1.70	115-164	39.82±1.74	25.2-78.0	123.71±2.30	102-143	35.04±1.76	23.2-56.5
Sep-17	M	34	132.0±1.95	105-165	42.90±1.87	21.5-76.4	126.10±2.37	112-152	38.87±2.46	24.6-71.6
	F	48	129.8±1.77	104-154	38.66±1.41	20.9-63.1	128.92±2.46	104-150	38.58±1.80	21.6-55.2
Oct-17	M	39	135.1±1.90	111-161	42.52±1.65	24.9-67.1	132.41±2.49	109-161	41.14±2.64	23.5-81.0
	F	56	118.6±1.41	94-145	30.91±0.96	15.2-45.9	124.73±1.78	102-160	34.91±1.46	23.3-69.9
Nov-17	M	48	124.4±1.97	100-160	35.49±1.71	15.5-67.7	129.22±2.32	102-161	41.91±2.39	16.6-75.1
	F	33	122.1±1.72	102-140	30.99±1.47	15.3-47.3	123.35±2.49	101-150	30.77±2.49	16.6-62.6
Dec-17	M	32	122.6±2.05	100-155	32.65±2.18	13.9-67.6	133.77±2.25	115-154	41.86±2.72	18.4-71.7
	F	43	124.8±1.63	101-143	33.50±1.43	17.8-57.7	128.14±3.00	108-145	38.48±2.63	22.4-58.1
Jan-18	M	35	132.0±1.66	110-152	40.87±1.92	25.0-64.5	126.44±3.08	101-146	37.89±3.07	16.9-63.1
	F	566	119.8±0.55	84-171	31.72±0.46	11.2-79.4	121.99±0.66	94-173	32.91±0.55	12.9-87.4
Total	F	409	126.6±0.75	78-193	37.69±0.67	7.9-88.2	125.81±0.90	93-193	37.47±0.86	11.2-92.9
	A	975	122.6±0.46	78-193	34.22±0.40	7.9-88.2	123.74±0.55	93-193	34.99±0.50	11.2-92.9

Source: Primary Data, 2017-2018

M - Male; F - Female; TL - Total length; TW - Total weight.

Based on the value of b (slope) (Table 2 & Figure 2) in general the growth pattern of *A. testudineus* during the study had a negative allometric growth pattern, where length growth is faster than weight growth. Actually in field, if *A. testudineus* body shape was a combination between rover-predator and bottom rover, that has a slightly elongated body (Moyle & Cech 2004). Reinforced with result of t_{test} analysis concluding that $t_{test} > t_{table}$ (0.05), it means H_0 is rejected ($B \neq 3$). *A. testudineus* growth patterns that had negative allometric patterns were also reported by several studies (Satrawaha & Pilasamorn 2009; Mustakim et al 2009; Kumar et al 2013).

The coefficient determination value (R^2) in males of swamp habitat had a range of 76-94% (r value: 0.87-0.97), and females of 78-98% (r value: 0.889-0.99), and coefficient determination value (R^2) in lake habitat for males was 77-93% (r value: 0.877-0.96), and 75-92% (r value: 0.866-0.959) for females. In terms of the value of r that is near to 1, it can be stated that the relationship of length and weight of *A. testudineus* has a positive correlation, in which each length increase will be followed by weight. The coefficient determination value (R^2) and correlation value (r) of *A. testudineus* near to 1 was also found in several studies (Alam et al 2007; Mustakim et al 2009; Satrawaha & Pilasamorn 2009; Ezzum & Minar 2012; Kumar et al 2013; Rahman et al 2015; Kumary & Raj 2016). The value of the coefficient of determination in general was near to 100%, this signifies that the diversity influenced by other factors (sex, disease, and food of availability) is small and the relationship between the total length and total weight of *A. testudineus* is very tight.




Figure 2. Black circle (male) and back triangle (female). a) The combined length-weight relationship of *A. testudineus* at Swamp habitat. b) The combined length-weight relationship of *A. testudineus* in lake habitat for one year in Semayang Lake East Kalimantan Province, Indonesia.

Relative weight and condition factor. The relative weight (W_r) value obtained from the standard weight and total weight in swamp habitats in males ranged from 100.04 ± 0.72 to 100.93 ± 2.41 , and in females from 100.25 ± 1.39 to 101.13 ± 2.60 . In lake habitat the relative weight value in male individuals was 100.29 ± 1.57 - 101.06 ± 2.65 , and in females 100.40 ± 2.98 - 101.60 ± 3.07 . The average value of relative weight was constantly > 100 , this indicates that the environmental conditions of *A. testudineus* in Semayang Lake which have the character of swamp and lake habitat provide good environment in terms of food availability and predation (Blackwell et al 2000). According to Anderson & Neumann (1996), if the value of weight is below 100, it indicates issue

such as lack of prey or high predation density, and vice versa if it is above 100 it indicates high prey availability or low predator density.

The average value of relative weight >100 is obtained from the comparison between observation weight with standard weight, so that it can be stated that the higher observation weight compared to standard weight can be used as an indicator of the aquatic environment that is good for life, survive and thrive. Based on this understanding, the standard weight data on male and female fish in both swamp and lake habitat measured for 1 year (February 2017 - January 2018) (Table 2, Figure 3 & 4) with relative dynamic value changes. The observed weight in swamp habitat was 34.2205 ± 0.397 g, higher than the average standard weight of 34.012 ± 0.386 g. As well as observed weight in lake habitat was 34.995 ± 0.501 g, higher than the average standard weight of 34.735 ± 0.473 g. The result shows that the observed weight is higher than the predicted weight (standard weight), this indicates that the water condition is healthy enough to support growth.

Table 2
Standard weight, condition factor (Fulton) and relative weight of *Anabas testudineus* (male-female) (in swamp habitat and lake habitat) in Semayang Lake, East Kalimantan Province, Indonesia

Month	Sexes	N	Swamp			N	Lake		
			Standard weight (Mean±SE)	Condition factor (Mean±SE)	Relative weight (Mean±SE)		Standard weight (Mean±SE)	Condition factor (Mean±SE)	Relative weight (Mean±SE)
3									
Feb-17	S	82	29.19±1.48	3.07±0.01	100.04±0.72	43	32.88±1.94	3.12±0.02	100.49±1.54
	F	45	39.29±2.67	3.17±0.09	100.25±1.39	32	35.30±3.05	3.14±0.03	101.46±3.09
Mar-17	M	77	30.35±1.17	3.09±0.01	100.53±1.22	49	32.91±1.40	3.11±0.02	100.60±1.57
	F	45	34.06±1.87	3.15±0.02	100.66±1.74	34	33.34±1.64	3.14±0.02	100.73±2.09
Apr-17	M	30	30.89±2.24	3.09±0.03	100.50±1.83	32	30.00±1.38	2.06±0.03	101.06±2.65
	F	25	42.47±4.75	3.13±0.04	100.66±2.44	24	31.85±3.82	3.03±0.06	101.43±3.61
May-17	M	53	30.15±1.76	3.07±0.02	100.70±1.72	21	31.85±1.95	3.15±0.03	100.90±2.76
	F	22	30.12±3.17	3.05±0.04	100.41±2.02	11	35.72±5.06	3.13±0.05	101.03±4.56
Jun-17	M	26	29.34±1.67	3.10±0.01	100.26±1.53	22	30.29±1.88	3.08±0.03	100.76±2.81
	F	29	36.08±1.76	3.15±0.02	100.29±1.66	23	33.61±1.76	3.14±0.03	100.73±2.61
Jul-17	M	29	31.82±1.49	3.16±0.02	100.54±1.89	27	30.54±1.20	3.13±0.01	100.38±1.74
	F	34	40.05±1.79	3.17±0.02	100.89±2.37	19	37.23±3.63	3.14±0.03	101.11±3.54
Aug-17	M	54	28.07±0.93	3.06±0.02	100.86±1.86	19	30.43±1.99	3.10±0.03	100.67±2.72
	F	21	31.77±1.66	3.08±0.03	100.62±2.48	10	37.11±2.45	3.15±0.03	100.40±2.98
Sep-17	M	35	39.66±2.63	3.20±0.01	100.38±1.55	21	34.92±1.56	3.16±0.02	100.30±1.68
	F	34	42.78±1.82	3.20±0.01	100.33±1.45	20	38.62±2.17	3.21±0.02	100.55±2.56
Oct-17	S	48	38.46±1.28	3.15±0.01	100.52±1.50	24	38.48±1.73	3.16±0.01	100.29±1.57
	F	39	41.30±1.48	3.16±0.01	100.49±1.70	29	40.84±2.35	3.15±0.02	100.47±2.03
Nov-17	M	56	30.74±0.89	3.12±0.02	100.69±1.59	37	34.72±1.29	3.14±0.02	100.55±1.71
	F	48	35.30±1.70	3.13±0.02	100.89±2.02	37	41.27±2.07	3.20±0.03	101.60±3.07
Dec-17	M	33	30.71±1.31	3.06±0.03	100.93±2.41	26	30.49±2.00	3.02±0.03	100.68±2.43
	F	32	32.24±1.91	3.08±0.03	101.13±2.60	25	41.62±2.66	3.14±0.03	100.80±2.62
Jan-18	M	43	33.21±1.20	3.09±0.02	100.83±1.99	14	38.28±2.30	3.217±0.02	100.43±2.60
	F	35	40.58±1.69	3.16±0.02	100.70±2.09	18	37.70±2.88	3.15±0.04	100.54±2.54

Source: Primary Data, 2017-2018.




Figure 3. Standard weight and relative weight of *Anabas testudineus* (males and females) (in swamp and lake habitat respectively) for one year in Semayang Lake East Kalimantan Province, Indonesia.




Figure 4. a) The combined observation weight and standard weight pattern of *Anabas testudineus* in swamp habitat. b) Observed and standard weight pattern of *Anabas testudineus* in lake habitat for one year in Semayang Lake East Kalimantan Province, Indonesia.

Condition factors were calculated to assess general fish health, productivity and physiological conditions of fish populations (Blackwell et al 2000; Richter 2007). This condition factor reflects the characteristics of the body's morphology, lipid content and growth rate (Bister et al 2000; Froese 2006; Stevenson & Woods 2006). The result of the condition factor analysis of *A. testudineus* in swamp habitat in males was 3.102 ± 0.0058 , and in female 3.141 ± 0.007 , whereas in lake habitat (Station 2, 3 and 5) condition factor was 3.114 ± 0.007 for males and 3.143 ± 0.009 for females. Descriptively it can be stated that the condition factor in female individuals is relatively greater than in male individuals

(Figure 5). The difference in the value of this condition factor is due to sexes (Lorenzoni et al 2015). Several other factors that are suspected to be the cause of fluctuation and variation of fish condition factor value are the difference of fish size and age (Enchina & Granado-Lorenzo 1997); during the spawning season the fish do not engage in food activities, but uses the fat reserves of the body for energy supply (Tzikas et al 2007); and parasitic pressure (Neff & Cargnelli 2004).




Figure 5. Combined condition factor of *Anabas testudineus* (males and females) (in swamp and lake habitat respectively) for one year in Semayang Lake East Kalimantan Province, Indonesia.

Conclusions. The general growth pattern of *A. testudineus* is negative allometric. Based on the relative weight value, the observed weight is higher than the standard weight value and factor condition value. Therefore it can be concluded that the Semayang Lake is still providing healthy aquatic habitat for the survival and thrive of *A. testudineus*.

Acknowledgements. The authors would like to thank Ministry of Research, Technology and Higher Education of Indonesia that have provided research grant, Mrs. Ghitari, MSc as the head of the water quality laboratory and Mr. Jailani as head of the laboratory of Aquatic Ecobiology Faculty of Fisheries and Marine Sciences, Mulawarman University which has provided facilities in supporting the present research. We also thank to Syahidul Akbar, Muhammad Reva, Berlian Noor and Veni for the technical support given during the research. Hopefully this research will be useful in managing *A. testudineus* resources so that the potential can be utilized optimally and sustainable.

References

- Anderson R. O., Neumann R. M., 1996 Length, weight and associated structural indices. In: Fisheries techniques. Murphy B. R., Willis D. W. (eds), pp. 447-483, American Fisheries Society, Bethesda.
- Archibold O. W., 1995 Ecology of world vegetation. Springer Netherlands, 510 p.
- Alam M. A., Noor A., Khan M. M. R., Rahman L., 2007 Growth performance and morphological variations of local and Thai climbing perch (*Anabas testudineus*, Bloch). Bangladesh Journal of Fisheries Research 11(2):163-171.
- Begum M., Minar M. H., 2012 Weight-length relationships of koi (*Anabas testudineus*) along with condition factor fed on formulated feed. Trends in Fisheries Research 1(2):1-6.
- Bister T. J., Willis D. W., Brown M. L., Jordan S. M., Neumann R. M., Quist M. C., Guy C. S., 2000 Proposed standard weight (Ws) equations and standard length categories for 18 warm water nongame and riverine fish species. North American Journal of Fisheries Management 20:570-574.
- Blackwell B. G., Brown M. L., Willis D. W., 2000 Relative weight (Wr) status and current use in fisheries assessment and management. Reviews in Fisheries Science 8:1-44.

- Corlett R. T., Primack R. B., 2011 Tropical rain forests: an ecological and biogeographical comparison (2nd edition). Hoboken, NJ: Wiley-Blackwell, 335 p.
- De Robertis A., Williams K., 2008 Weight-length relationship in fisheries studies: the standard allometric model should be applied with caution. *Transaction of the American Fisheries Society* 137(3):707-719.
- Ecoutin J. M., Albaret J. J., Trape S., 2005 Length-wWeight relationships for fish populations of a relatively undisturbed tropical estuary: The Gambia. *Fisheries Research* 72:347-351.
- Encina L., Granado-Lorencio C., 1997 Seasonal changes in condition, nutrition, gonad maturation and energy content in barbel, *Barbus sclateri*, inhabiting a fluctuating river. *Environmental Biology of Fishes* 50:75-84.
- Everhart W. H., Youngs W. D., 1981 Principles of fishery Science. 2nd edition, Comstock Publishing Associates, a division of Cornell University Press, London.
- Froese R., 2006 Cube law, condition factor and weight-length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology* 22:241-253.
- Furukawa H., 1994 Coastal wetland of Indonesia: environmental, subsistence and exploration. Kyoto University Press, Japan, 228 p.
- Helmizuryani, Muslimin B., 2016 Growth performance of mono sex and mixed sex climbing perch (*Anabas testudineus*). *Omni-Akuatika* 12(2):99-103.
- Jennings S., Kaiser M. J., Reynolds J. D., 2001 Marine fishery ecology. Blackwell Sciences, Oxford.
- Khristenko D. S., Kotovska G. O., 2017 Length-weight relationship and condition factors of freshwater bream *Abramis brama* (Linnaeus, 1758) from the Kremenchug Reservoir, Middle Dnieper. *Turkish Journal of Fisheries and Aquatic Sciences* 17:71-80.
- Kohinoor A. H. M., Akhteruzzaman M., Hussain M. G., Shah M. S., 1991 Observation on the induced breeding of koi fish (*Anabas testudineus* Bloch) in Bangladesh. *Bangladesh Journal of Fisheries* 14(1-2):73-77.
- Kumar K., Lalrinsanga P. L., Saho M., Mohanty U. L., Kumar R, Sah A. K., 2013 Length-weight relationship and condition factor of *Anabas testudineus* and *Channa* species under different culture systems. *World Journal of Fish and Marine Sciences* 5(1):74-78.
- Kumary K. S. A., Raj S., 2016 Length-weight relationship and condition of climbing perch *Anabas testudineus* Bloch population in Kuttanad, Kerala. *International Journal of Advanced Research in Biological Sciences* 3(9):21-26.
- Le Cren C. P., 1951 Length-weight relationship and seasonal cycle in gonad weight and condition in the perch (*Perca fluviatilis*). *Journal of Animal Ecology* 20(2):201-219.
- Lopez O. R., Kursar T. A., 2007 Interannual variation in rainfall, drought stress and seedling mortality may mediate monodominance in tropical flooded forests. *Oecologia* 154:35-43.
- Lorenzoni M., Giannetto D., Carosi A., Dolciami R., Ghatti L., Pompei L., 2015 Age, growth and body condition of big-scale sand smelt *Atherina boyeri* Risso, 1810 inhabiting a freshwater environment: Lake Trasimeno (Italy). *Knowledge and Management of Aquatic Ecosystems* 416(9):1-13.
- Mitsch W. J., Gosselink J. G., 1993 Wetland. 2th edition, Van Nostrand Reinhold, New York, 722 p.
- Moyle P. B., Cech Jr. J., 2004 Fishes: an introduction to ichthyology. 5th edition, Prentice-Hall. Inc., New Jersey.
- Mousavi-Sabet H., Abdollahpour S., Salehi-Farsani A., Vatandoust S., Langroudi H. F., Jamalzade H. R., Nasrollahzadeh A., 2013 Length-weight and length-length relationships and condition factor of *Alburnus mossulensis* from the Persian Gulf basin. *AAFL Bioflux* 6(4):297-302.
- Mustakim M., Sunarno D., Affandi R., Kamal M. M., 2009 [Growth of climbing perch (*Anabas testudineus*, Bloch) in different habitats of Lake Melintang Areas, East Kalimantan Province]. *Jurnal Penelitian Perikanan Indonesia (Indonesian Journal of Fisheries Research)* 15(2):113-121. [In Indonesian].

- Neff B. D., Cargnelli L. M., 2004 Relationships between condition factors, parasite load and paternity in bluegill sunfish, *Lepomis macrochirus*. *Environmental Biology of Fishes* 71:297–304.
- Noor M., 2007 Rawa Lebak, Ekologi, Pemanfaatan dan Pengembangannya [The Lebak of swamp, ecology, utilization and development]. PT. Raja Grafindo Persada. Jakarta, 274 p. [In Indonesian].
- Okgerman H., 2005 Seasonal variation of the length weight and condition factor of Rudd (*Scardinius erythrophthalmus* L) in Spanca Lake. *International Journal of Zoological Research* 1(1):6-10.
- Pethiyagoda R., 1991 Freshwater fishes of Sri Lanka. The Wildlife Heritage Trust of Sri Lanka, 362 p.
- Pokorný J., Björk S., 2010 Wetlands: ecology, conservation, and management: restoration of lake, streams, floodplains and bogs in Europa (principles and case studies). Development of aquatic macrophytes in shallow lakes and ponds. Springer Dordrecht Heidelberg London New York, pp. 37–43.
- Rahman A., Talukdar K., Rahman W., Deka P., 2015 Length-Weight relationship and relative condition factor of *Anabas testudineus* (Bloch) of Deepar Beel (wetland) of Assam, India. *International Journal of Applied Research* 1(11):956-958.
- Richter T. J., 2007 Development and evaluation of standard weight equations for bridgelip sucker and largescale suckers. *North American Journal of Fisheries Management* 27:936-939.
- Romanowski N., 2013 Living water: ecology of animal in swamps, rivers, lakes and dams: rebirth of lake (Chapter 9). CSIRO Publishing, Australia, 292 p.
- Satrawaha R., Pilasamorn C., 2009 Length-weight and length-length relationships of fish species from the Chi River, northeastern Thailand. *Journal of Applied Ichthyology* 25:787–788.
- Steel R. G. D., Torrie J. H., 1993 Principles and procedure of statistics. 3th edition, 748 p.
- Stevenson R. D., Woods W. A., 2006 Condition indices for conservation: new uses for evolving tools. *Integrative and Comparative Biology* 46:1169-1190.
- Tay Y. L., Loong A. M., Hiong K. C., Lee S. J. J., Tng Y. Y. M., Wee N. L. J., Lee S. M. L., Wang W. P., Chew S. F., Wilson J. M., Ip Y. K., 2006 Active ammonia transport and excretory nitrogen metabolism in the climbing perch, *Anabas testudineus*, during 4 days of emersion or 10 minutes of forced exercise on land. *Journal of Experimental Biology* 209:4475-4489.
- Tesch F. W., 1971 Age and growth. In: Fish production in fresh waters. W. E. Ricker (ed), pp. 98–130, Blackwell, Oxford.
- Tsoumani M., Liasko R., Moutsaki P., Kagalou I., Leonardos I., 2006 Length-weight relationships of an invasive cyprinid fish (*Carassius gibelio*) from 12 Greek lakes in relation to their trophic states. *Journal of Applied Ichthyology* 22:281–284.
- Turkmen M., Erdogan O., Yildirim A., Akyurt I., 2002 Reproductive tactics, age and growth of *Capoeta capoeta umbra* Heckel 1834 from the Askale Region of the Karsu River, Turkey. *Fisheries Research* 54(3):317-328.
- Tzikas Z., Ambrosiadis I., Soultos N., Georgakis S., 2007 Seasonal size distribution, condition status and muscle yield of mediterranean horse mackerel *Trachurus mediterraneus* from The North Aegean Sea, Greece. *Fisheries Science* 73:453–462.
- Vilizzi L., Copp G. H., Britton J. R., 2013 Age and growth of European barbel *Barbus barbus* (Cyprinidae) in the small, mesotrophic River Lee and relative to other populations in England. *Knowledge and Management of Aquatic Ecosystems* 409(09):1-18.
- Welcomme R. L., 1985 River fisheries. FAO, Fisheries Technical Paper, pp. 262-330.

Received: 14 November 2018. Accepted: 21 February 2019. Published online: 28 February 2019.

Authors:

Mohammad Mustakim, Diponegoro University, Faculty of Fisheries and Marine Science, Doctoral Program in Coastal Resources Management, Indonesia, Central Java, Tembalang, Semarang 50275; Samarinda Mulawaran University, Faculty of Fisheries and Marine Science, Department of Aquatic Resources Management, Indonesia, East Kalimantan, Gunung Tabur Kampus, Gunung Kelua street, e-mail: moh.mustakim@fpik.unmul.ac.id; mustakim.unmul2005@gmail.com.

Sutrisno Anggoro, Diponegoro University, Faculty of Fisheries and Marine Science, Department of Aquatic Resources Management, Indonesia, Semarang 50275, Jl. Prof. H. Soedarto, SH. – Tembalang, e-mail: sutrisnoanggoro52@gmail.com

Frida Purwanti, Diponegoro University, Faculty of Fisheries and Marine Science, Department of Aquatic Resources Management, Indonesia, Semarang 50275, Jl. Prof. H. Soedarto, SH. – Tembalang, e-mail: frpurwanti@gmail.com

Haeruddin, Diponegoro University, Faculty of Fisheries and Marine Science, Department of Aquatic Resources Management, Indonesia, Semarang 50275, Jl. Prof. H. Soedarto, SH. – Tembalang, e-mail: haeruddindaengmile@yahoo.co.id, haeruddindaengmile@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Mustakim M., Anggoro S., Purwanti F., Haeruddin, 2019 Length-weight relationships and condition factor of *Anabas testudineus* in the Semayang Lake, East Kalimantan, Indonesia. *AAFL Bioflux* 12(1):327-337.

Length-Weight Relationships and Condition Factor of *Anabas testudineus* in the Semayang Lake, East Kalimantan, Indonesia

ORIGINALITY REPORT

10%

SIMILARITY INDEX

%

INTERNET SOURCES

10%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Nur Asia Nasution, Rusdi Machrizal. "Bioecological aspect of lamasi (*Barbonymus gonionotus*) in mailil rever labuhanbatu district, indonesia", JP BIO (Jurnal Pendidikan Biologi), 2021 1%
Publication

- 2** Z. A. Muchlisin, M. Musman, M. N. Siti Azizah. "Length-weight relationships and condition factors of two threatened fishes, *Rasbora tawarensis* and *Poropuntius tawarensis*, endemic to Lake Laut Tawar, Aceh Province, Indonesia", Journal of Applied Ichthyology, 2010 1%
Publication

- 3** Mitchell H. Tsai, Michael A. Hall, Melanie S. Cardinal, Max W. Breidenstein, Michael J. Abajian, Richard S. Zubarik. "Changing Anesthesia Block Allocations Improves Endoscopy Suite Efficiency", Journal of Medical Systems, 2019 1%
Publication

4

Orestes Carlos Bello-González, Perla Alonso-EguíaLis, César Arturo Ilizaliturri Hernández, Norman Mercado-Silva. "Variability in mentum deformities of Tanytarsus larvae (Diptera, Chironomidae) in a metal rich basin in Northern Mexico", Water Biology and Security, 2022

Publication

1 %

5

Rima Mohdeb, M. Hichem Kara. " Age, growth and reproduction of the Morocco dentex of the eastern coast of Algeria ", Journal of the Marine Biological Association of the United Kingdom, 2014

Publication

1 %

6

Siddique, Mohammad Abdul Momin, Aziz Arshad, and S.M. Nurul Amin. "Length-weight relationships of the tropical cephalopod Uroteuthis chinensis (Gray, 1849) from Sabah, Malaysia", Zoology and Ecology, 2014.

Publication

<1 %

7

Wayan Kantun, Indra Cahyono, Wayan Suma Arsana. " Biological Aspect of Bullet Tuna (Risso, 1810) in the Makassar Strait, West Sulawesi, Indonesia ", Croatian Journal of Fisheries, 2019

Publication

<1 %

8

Agustu Sholeh Pujokaroni, Hideaki Ichiura. "Preparation of carboxymethyl cellulose

<1 %

(CMC) derived from palm fiber waste: Effect of ozone pretreatment on CMC properties", Research Square Platform LLC, 2022

Publication

9

ŞAHİN, Cemalettin, KASAPOĞLU, Nazlı, GÖZLER, A. Mutlu, KALAYCI, Ferhat, HACIMURTAZAOĞLU, Necati and MUTLU, Cengiz. "Age, growth, and gonadosomatic index (GSI) of Mediterranean horse mackerel (*Trachurus mediterraneus* Steindachner, 1868) in the Eastern Black Sea", TÜBİTAK, 2009.

Publication

<1 %

10

D F Putra, A A Muhammadar, N Muhammad, A Damora, A Waliul, M Z Abidin, N Othman. "Length-weight relationship and condition factor of white shrimp, *Penaeus merguensis* in West Aceh waters, Indonesia", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

11

Dibakar Bhakta, Suhas Prakash Kamble, Wakamban Anand Meetei, Jayesh Kumar Solanki et al. "Relation of fish size with that of its otolith of Pama croaker, *Otolithoides pama* (Hamilton, 1822) from Narmada estuary, India", Research Square Platform LLC, 2022

Publication

<1 %

12

Ecoutin, J.M.. "Length-weight relationships for fish populations of a relatively undisturbed tropical estuary: The Gambia", Fisheries Research, 200505

Publication

<1 %

13

J Tambun, D Bakti, Desrita. " The growth and exploitation rate of yellowstripe scad (*cuvier*, 1833) in the Malacca Strait, Medan Belawan Subdistrict, North Sumatera Province ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

<1 %

14

Azevedo, Juliana de Souza, Jorge Eduardo de Souza Sarkis, Talita Aparecida Oliveira, and João Cristiano Ulrich. "Tissue-specific mercury concentrations in two catfish species from the Brazilian coast", Brazilian Journal of Oceanography, 2012.

Publication

<1 %

15

Boudinar, A. S., L. Chaoui, and M. H. Kara. "Age, growth and reproduction of the sand smelt *Atherina boyeri* Risso, 1810 in Mellah Lagoon (Eastern Algeria)", Journal of Applied Ichthyology, 2016.

Publication

<1 %

16

E Paujiah, D D Solihin, R Affandi, S Sa'adah, I Zulfahmi. " Growth pattern and lenght-weight relationship of , in The Cisadea River, West

<1 %

17

MARGARITA N. LAVIDES, NICHOLAS V. C. POLUNIN, SELINA M. STEAD, DON GEOFF TABARANZA, MIA THERESA COMEROS, JESUS RAY DONGALLO. "Finfish disappearances around Bohol, Philippines inferred from traditional ecological knowledge", Environmental Conservation, 2010

Publication

<1 %

18

Nwabueze I. Igu. "Species Diversity and Structure of an Intact Freshwater Swamp Forest in the Niger Delta", Open Journal of Forestry, 2017

Publication

<1 %

19

Mohamed Hichem Kara, Jean-Pierre Quignard. "Fishes in Lagoons and Estuaries in the Mediterranean 2", Wiley, 2019

Publication

<1 %

20

Fúlvia Cristina Oliveira, Rodrigo Yutaka Dichoff Kasai, Carlos Eurico Fernandes, Wesley Souza da Silva et al. " Probiotic, prebiotic and synbiotics supplementation on growth performance and intestinal histomorphometry larvae ", Journal of Applied Aquaculture, 2020

Publication

<1 %

21 Mirna Dwirastina, Marson. " Length-weight relationships Silver barb () in Mamberamo River, Papua ", E3S Web of Conferences, 2021
Publication <1 %

22 Desrita, I E Susetya, M Suriani, A Rahman. "Biology and growth of Asiatic Hard Clam (Meretrix meretrix) population in Tanjung Balai, North Sumatera", IOP Conference Series: Earth and Environmental Science, 2019
Publication <1 %

23 L. Guardone, E. Ricci, F. Susini, E. Polsinelli, G. Guglielmone, A. Armani. "First detection of Eustrongylides excisus (Nematoda: Dioctophymatidae) in big-scale sand smelt (Atherina boyeri) from the lake Massaciuccoli (Northwest Tuscany, Italy): implications for public health and seafood quality", Food Control, 2021
Publication <1 %

24 R Afdhila, A A Muhammadar, M A Chaliluddin. "Length-weight relationship and condition factors of layang fish (Decapterus macrosoma) that landed at Lampulo Ocean Fishing Port, Banda Aceh", IOP Conference Series: Earth and Environmental Science, 2019
Publication <1 %

25 R Faizah, L Sadiyah. " Some biology aspects of Indian Scad Rupell, 1928) in Pemangkat <1 %

Fisheries Port, West Kalimantan ", IOP
Conference Series: Earth and Environmental
Science, 2020

Publication

26

Y. L. Tay, A. M. Loong, K. C. Hiong, S. J. Lee et al. "Active ammonia transport and excretory nitrogen metabolism in the climbing perch, *Anabas testudineus*, during 4 days of emersion or 10 minutes of forced exercise on land", *Journal of Experimental Biology*, 2006

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On