

Case Report Correlation of Warm and Cold Ischemic Time to Graft Function in Kidney Transplant: A Single Centre Report

by Eriawan Agung Nugroho

Submission date: 02-Mar-2022 09:04AM (UTC+0700)

Submission ID: 1774293810

File name: TOUNJ-12-66.pdf (522.98K)


Word count: 4130

Character count: 22370



The Open Urology & Nephrology Journal

Content list available at: <https://openurologyandnephrologyjournal.com>



CASE REPORT

Correlation of Warm and Cold Ischemic Time to Graft Function in Kidney Transplant: A Single Centre Report

Eriawan A. Nugroho^{1*}, Arif Hidayat² and Ahsanu T. Hidayat²

¹Division of Urology, Dr. Kariadi General Hospital/Medical Faculty Diponegoro University, Semarang, Indonesia

²Department of Surgery, Dr. Kariadi General Hospital/Medical Faculty Diponegoro University, Semarang, Indonesia

Abstract:

Background:

Kidney transplantation has been known as a major advancement of modern medicine which provides high-quality life years to patients with irreversible kidney failure (end-stage renal disease, ESRD) worldwide. The first kidney transplantation in Semarang was carried out at Telogorejo Hospital in 1985. Dr. Kariadi Hospital, has also undertaken kidney transplantation 28 times since January 2014 until September 2018. Kidney ischemic time is one of the most important factors which can affect the outcome of kidney transplantation. Prolonged ischemia of the graft can be related to transplantation's effects. This study aimed to evaluate the influence of warm and cold ischemic time on the outcome of kidney transplantation in Dr. Kariadi Hospital Semarang.

Material and Methods:

This is an observational, cross-sectional study. The data was collected from medical record of all patients who underwent kidney transplantation from January 2014 until December 2018 and the level of creatinine and urine production before and after transplantation was documented. A total of 28 patients were registered and completed the inclusion criteria. The results of the study were tabulated and statistical analysis was performed using SPSS 23.0. The relationship was declared meaningful if $p = 0.05$ was obtained.

Result:

From the analysis of the research that has been done using the Pearson correlation test and hypothesis test Wilcoxon, it was found that there was a relationship between ischemic time and a decrease in creatinine level from $r = -0.4489$ with a value of $p = 0.008$. From the analysis of the research, a relationship between ischemic time and urine production was found after transplantation with a value of $r = -0.562$ and a value of $p = 0.002$.

Conclusion:

The results of this study show a strong correlation between the length of total ischemic time with a decrease in creatinine and urine production which means that the longer ischemic time, the lower the decrease in creatinine levels and the lesser the urine production.

Keywords: Warm ischemic time, Cold ischemic time, Graft function, Kidney transplantation, Urine production, End-Stage Renal Disease.

Article History

Received: May 28, 2019

Revised: September 17, 2019

Accepted: September 20, 2019

1. INTRODUCTION

The number of patients diagnosed with End-Stage Renal Disease (ESRD) in Indonesia is increasing annually, showing similar trends with global prevalence [1, 2]. Kidney transplantation is acknowledged as a major advancement of modern medicine which provides high-quality life years to patients with ESRD worldwide [1, 3]. Patients who have

kidney transplantation live 10 to 15 years longer than patients who undergo dialysis [4].

Since 1985, there have been several renal transplantations performed in Telogorejo Hospital and Dr. Kariadi General Hospital, Semarang. In Dr. Kariadi General Hospital Semarang, kidney transplantation have been done 26 times since January 2014 until September 2018 [5].

The success of kidney transplantation depends on several factors [6, 7]. Prolonged ischemia of the graft can be related to the effect of transplantation. Ischemia can induce a harmful cascade effect that can be worsened with the return of blood

* Address correspondence to this author at the Division of Urology, Dr. Kariadi General Hospital/Medical Faculty Diponegoro University, Semarang, Indonesia; Tel: +088210362784; E-mail: eriawanspu@gmail.com

supply [8]. This injury can produce an inflammatory and immune response that can potentially slow down the kidney function, increase alloimmune reactivation, and develop a progressive pathological change [9, 10].

The aim of this study was to investigate the influence of total ischemic time with the outcome of kidney transplantation in Dr. Kariadi General Hospital Semarang.

2. MATERIALS AND METHODS

This is an observational, cross-sectional study. The data was collected from the medical record of patients who underwent kidney transplantation from January 2014 until December 2018 at Kariadi General Hospital Semarang.

The population in this study are patients with renal failure who had kidney transplantation in Dr. Kariadi General Hospital Semarang. All the patients who underwent kidney transplantation having a medical record from January 2014 until December 2018 were examined for the level of creatinine and urine production before and after transplantation. There were 28 patients who were registered and completed the inclusion criteria.

Descriptive data is presented in the form of tables and box-plots. The normality test was conducted with Shapiro Wilk. Then the data was analyzed using the Wilcoxon test. The results of the study were tabulated and statistical analysis was

performed using SPSS 23.0. The relationship is declared meaningful if $p = 0.05$ is obtained.

In clinical settings, patients with ESRD were actively screened as potential candidates for transplantation. Patients younger than 15 years old, older than 70 years, who were diagnosed with active systemic infection, active malignancy, significant cardiovascular, cerebrovascular or pulmonary disease, and drug/alcohol addiction were considered ineligible to become a recipient. Eligible and compliant patients with active National Health Insurance status were offered renal transplant program. The patient and family members were provided with the list of potential kidney donors. Potential donors who were willing to do the process were screened preliminarily before undergoing further tests to determine their suitability (Table 1).

Table 1. Demographics of kidney transplantations in Semarang (Age and Sex).

Variable	–	Recipient	Donor
Age	Mean (range)	32.5 (15 – 50)	46.8 (24 – 64)
Sex, n (%)	Male	17 (63.0)	14 (51.9)
	Female	10 (37.0)	13 (48.1)

Kidney transplantation and several renal transplantations are performed in Telogorejo Hospital and Dr. Kariadi General Hospital, Semarang. The number of transplants performed increases steadily every year (Table 2).

Table 2. Demographics of kidney transplantations in Semarang (Donor-recipient relationship, Etiology of ESRD, Recipient comorbidities, Recipients Body Mass Index and HLA Matching).

Variable	N	%
Donor-recipient relationship		
Related	20	74.1
Non-related	7	25.9
Etiology of ESRD		
Hypertension	16	59.3
Diabetes mellitus	4	14.8
Autoimmune	1	3.7
Glomerulonephritis	2	7.4
Other	4	14.8
Recipient comorbidities		
Coronary artery disease	3	11.1
Cerebrovascular disease	3	11.1
Pulmonary tuberculosis	2	7.4
Pleural effusion	2	7.4
Ascites	2	7.4
Hepatitis C	2	7.4
Gastric erosion	1	3.7
Depression	1	3.7
Recipients Body Mass Index		
< 18.5 (underweight)	4	14.8
18.5 – 24.9 (normal)	14	51.9
25.0 – 29.9 (overweight)	6	22.2
≥ 30.0 (obese)	3	11.1
HLA Matching		
Full match	20	74.1
4 miss match	1	3.7

(Table 2) contd....

Variable	N	%
3 miss match	1	3.7
2 miss match	2	7.4
1 miss match	2	7.4

HLA: Human Leucocyte Antigen

The criteria for transplant recipients were the age of 15–70 years old, with at least one Human Leukocyte Antigen (HLA) matched with the donor and absence of Donor-Specific Antibody (DSA). Patients in Semarang showed similar demographics with the overall Indonesian cases, where most recipients were younger (33.0 years old) compared to the donors (46.1 years old) [11]. Similarly, hypertension (61.5%) was the major etiology of ESRD. The recipient's comorbidities include cerebrovascular disease, coronary artery disease, history of pulmonary tuberculosis, pleural effusion, ascites, hepatitis C, gastric erosion and depression. The majority of the recipients received kidney from blood-related donors ($n=19$, 73.0%). There were 20 (76.9%) full HLA match, two cases of one and two miss match, one case of four and three miss-match. There were five (19.2%) cases of mortalities, two of them occurred during post-operative recovery period and the rest within 12 months after being discharged from the hospital. The cause of the two “early” mortalities was acute myocardial infarction (Table 3).

Table 3. Demographics of kidney transplantations in Semarang (Year).

Year	No. of Transplant	Mortality
2012	1	0/1
2013	1	0/1
2014	3	2/3
2015	4	1/4
2016	4	2/4
2017	7	0/7
2018	7	2/4

The third death occurred 7 months post-transplant of a 39-year-old male with a history of myocardial infarction and gastric erosion. The cause of death was determined to be heart failure due to ischemic and hypertensive cardiomyopathy. Septic shock was the cause of death for the other two mortalities with one related to bacterial pneumonia and the other with unknown primary site of infection. There were three cases of allograft rejection, one patient with three pre-transplant HLA miss match and the other with two HLA miss-match. Post-transplant immunosuppressive regimen was discontinued and all three patients underwent dialysis again. This is the first report that presents the total number of kidney transplantations performed in Indonesia. The number has risen significantly within the past 5 years, especially in medical centers in Java. However, it is still far below from what is needed or when compared to neighboring Southeast Asian countries like Vietnam, Thailand, and Philippines [12, 13]. Several barriers to transplantation in Indonesia that had been previously elaborated include shortage of specialists, high cost, lack of easy access and information within the community, as well as amongst the medical practitioners.

The situation of kidney transplantation in Semarang was

affected by the events elaborated by Mochtar *et al.* (Level Of Evidence 4). The first recorded kidney transplant in Semarang was in 1977. Afterwards, there were several unrecorded kidney transplantations, mainly performed in Dr. Kariadi General Hospital. The renal transplant system was formally restarted in 2014 and since then, the procedure has been regularly performed in the institution. In-hospital ESRD patients are individually offered the transplant program by the nephrologist team. Although there has been a constant annual increase of kidney transplant performed since 2014, the manual patient recruitment system will eventually limit the center from ever reaching its maximum potential.

The average waiting time for patients in Semarang was 68 months from the start of enrollment until the operation. The waiting time and the total number of transplants can be greatly increased by a coordinated National database online system. We report on 27 cases of living-donor transplantations from January 2012 until July 2018. In contrast, Cipto Mangunkusumo Hospital in Jakarta performed 491 transplants within 6 years. There were five (19.2%) mortalities in Semarang, all within 12 months post-transplant. The mortality rate is similar to Jakarta ($n=28$, 20.28%) as reported by Marbun *et al.* In Semarang, two patients (40.0%) died during post-operative recovery due to acute myocardial infarction, one patient (20.0%) died because of cardiomyopathy, and two patients (40.0%) by sepsis. The main cause of death in Jakarta was sepsis ($n=8$, 40.0%), followed by an unknown cause ($n=7$, 35.0%), pulmonary edema ($n=2$, 10.0%), hepatitis ($n=2$, 10.0%) and stroke ($n=1$, 5.0%). Marcelino *et al.* published the only Indonesian data available for comparison, which was based on Laparoscopic Living-Donor Nephrectomy (LLDN) technique. In terms of procedural time, the average first Warm Ischemic Time (WIT) of open technique in Semarang was longer compared to the LLDN in Jakarta (6.6 minutes vs. 4.3 minutes). Expectedly, the average intraoperative blood loss was also higher in open technique (384 ± 192 mL vs. 194 ± 198 mL). However, no urinary retention, operative wound site infection or severe pain (Visual Analogue Scale ≥ 7) was recorded (Table 4).

Table 4. Demographics of kidney transplantations in Semarang (procedural time).

Procedural Time	Minutes: Seconds
Warm ischemic time I	06:34
Cold ischemic time	08:04
Warm ischemic time II	27:55
Total ischemic time	36:56

Based on the outcome of 17 patients that surpassed the 12-month mark, the 1-year survival rate of a transplant patient in Semarang is 70.6%. Compared to Jakarta, the 1-year survival is 88.5% and 3-year survival is 79.7% [11]. Compared to abroad, the 1-year survival rates for living donor kidney

transplants are 97.2% in the United States, 98.0% in Australia and New Zealand, 95.8% in Europe, and 97.7% in Canada. The limited number of transplants in Semarang contributes to the low survival rates. It also highlights the need for training and expertise. Better survival rate can be achieved with more transplants number as well as reaching the plateau of learning curve within the coming years. The transplant team in Semarang is also in the process of applying LLDN technique to achieve better operative results (Table 5).

Table 5. Demographics of kidney transplantations in Semarang (Complications and outcome).

Intraoperative bleeding, mean (SD) in mL	384 (\pm 192)	
Hospital stay, mean (range) in days	11 (10 – 14)	
Infection, n (%)	2	(7.7)
Cardiovascular, n (%)	3	(11.5)
Total number of deaths, n (%)	5	(19.2)
Re-dialysis, n (%)	3	(11.5)
Re-transplant, n (%)	0	(0.0)

Based on this study, there was an improvement in the total number of kidney transplantations between January 2012 to July 2018 at Dr. Kariadi Hospital Semarang. The main cause of ESRD is hypertension. Most recipients are younger than the donor with majority kidney sources being blood-related donor. This report shows that 1-year survival rate in Semarang is lower than others center. Exclusive hospital-patient communication and medical record system for kidney transplant patients is also being established to ensure a long-term and complete follow-up. The initial goal of our institution is to establish a firm kidney transplant center capable of covering Central Java, aiding the decentralization of kidney transplantation in Indonesia that can be considered to be still in its infancy. There has been a consistent increase in kidney transplantation, however still concentrated within the capital city. Major barriers that have been surmounted include the establishment of National Health Insurance full coverage in 11 appointed hospitals and a National law that permits organ transplantation. As of 2018, the five leading centers that performed the most transplants are Jakarta, Surabaya, Yogyakarta, Semarang and Malang. Currently, there are limited Indonesian data to compare the intra and post-operative results of our center. However, improved experience and expertise are imperative to achieve better and satisfactory results in Semarang. At national level, continuous efforts are being made for the expansion of specialists' training programs and kidney transplantation centers. There is an ongoing yet intangible effort by the Indonesian National Transplantation Society in establishing an organ procurement system for a cadaveric donor, public education, and the ever-needed Transplant National Database.

3. RESULTS

3.1. Creatinine Level

The results of the research conducted on 28 study subjects found that before and after transplantation as many as 28 subjects had decreased creatinine (Fig. 1). From the analysis carried out, we obtained creatinine average before the

transplant to be 7.9 ± 2.53 , mean creatinine value after surgery 5.2 ± 4.4 , and mean creatinine decrease 3.4 ± 1.1 mg/dL. From the analysis of the research using Pearson correlation test and hypothesis test Wilcoxon, it was found that there was a relationship between ischemic time and a decrease in renal postin creatinine from $r = -0.4489$ with a value of $p = 0.008$ (Fig. 2).




Fig. (1). The decreased of creatinin level before and after transplant.

3.2. Urine Production

The results of the study conducted on 28 research subjects found that the average urine production 2.5 hours post anastomosis was 294 ± 87 ml. From the analysis of the research, there is a relationship between Ischemic Time and post-transplant urine production with a value of $r = -0.562$ and a value of $p = 0.002$ (Fig. 3).




Fig. (2). The decrease in creatinine level compared with total ischemic time.




Fig. (3). The decrease in urine production compared with total ischemic time.

4. DISCUSSION

Chronic Kidney Disease (CKD) is damage to the structure or function of the kidney lasting ≥ 3 months, with or without a decrease in Glomerular Filtration Rate (GFR) [11]. The successful kidney transplantation depends on some factors such as ischemic time. [6, 7] Total ischemic time is defined as the time of the donor renal artery interruption or aortic clamp, until the time of release of the clamp on the renal artery in the recipient (in hours). Total ischemic time is a combination of Cold Ischemic Time (CIT) and Warm Ischemic Time (WIT). [12, 13] Prolonged ischemic time is associated with an increase of Delayed Graft Function (DGF) and longer length of stay which consequently increase the cost of transplantation [14, 15].

Reperfusion after CIT activates the complex cascade, which ultimately contributes to apoptotic or necrosis cell death. This phenomenon is also known as Ischemic Reperfusion Injury (IRI) [8]. It is known that injuries caused by ischemia and reperfusion and to some extent also with cooling and rewarming give rise to cascades which ultimately lead to apoptosis, endothelial cell injury, and activation of the innate immune response resulting from organ dysfunction [10].

DGF occurred in 13.5% of recipients with a total ischemic time of 14 hours or longer than 10.9% of those with shorter ischemic time. This effect is greater for those with donors aged 55 years or more. Thus, it appears that the consequences of a long ischemic time are mainly identifiable in the early posttransplant period as DGF and possibly an increased acute rejection rate. The effects of ischemic time on the long-term outcome of renal transplantation in the form of impaired graft function and graft survival are less evident [10, 16].

CONCLUSION

From the results of this study, there is a strong correlation between the length of total ischemic time with a decrease in creatinine levels and urine production which means that the longer ischemic time, the lower the decrease in creatinine levels and consequently the lesser the urine production.

15

ETHICS APPROVAL AND CONSENT TO PARTICIPATE

Not applicable.

HUMAN AND ANIMAL RIGHTS

Not applicable.

CONSENT FOR PUBLICATION

Written informed consent was obtained from all the participants.

AVAILABILITY OF DATA AND MATERIALS

- The article used to support the findings of this study have been deposited in the Ethnicity & Disease, Volume 16, Spring 2006. https://www.google.com/url?sa=t&rc=t=j&q=&esrc=s&source=web&cd=1&cad=rja&uact=8&ved=2ahUKEwi4xafokLPiAhU363MBHQO_BZ0QFjAAegQIBhAC&url=https%3A%2F%2Fwww.ethndis.org%2Fpriorsuparchives%2Fethn-16-2s2-14.pdf&usg=AOvVaw3PK9yA_gCjqlB8hj8469Qt
- The article used to support the findings of this study have been deposited in PLoS One. 2016 Jul 6;11(7):e0158765. doi: 10.1371/journal.pone.0158765.
- The article used to support the findings of this study have been deposited in <https://www.google.com/url?sa=t&rc=t=j&q=&esrc=s&source=web&cd=2&cad=rja&uact=8&ved=2ahUKEwi2uJe2k7PiAhXS7XMBHUnCnEQFjABegQIBBAC&url=https%3A%2F%2Fwww.pernefri.org%2F&usg=AOvVaw183-YUJkiaDeQigB9G0Yhm>
- The article used to support the findings of this study have been deposited in https://www.google.com/url?sa=t&rc=t=j&q=&esrc=s&source=web&cd=1&ved=2ahUKEwisoc7ok7PiAhVh7nMBHeN_DrQQFjAAegQIBBAC&url=http%3A%2F%2Fwww.european-renal-bes-t-practice.org%2Fsites%2Fdefault%2Ffiles%2FBoekje%2520niertransplantatie%2520Indonesian%2520OKDRUK.pdf&usg=AOvVaw0FU86CQLlyk_5F9H-Bs0sy
- The article used to support the findings of this study have been deposited in Asian Jurnal of Urology available online 22 February 2019. doi: <https://doi.org/10.1016/j.ajur.2019.02.003>.
- The article used to support the findings of study have been deposited in Transplant Proc. 2010 Jun;42(5):1413-20. doi: 10.1016/j.transproceed.2010.04.016.
- The article used to support the findings of study have been deposited in Am J Transplant. 2013 May;13(5):1134-48. doi: 10.1111/ajt.12210. Epub 2013 Apr 8.
- The article used to support the findings of study have been deposited in Kidney Int. 2015 Feb;87(2):343-9. doi: 10.1038/ki.2014.304. Epub 2014 Sep 17.
- The article used to support the findings of study have been deposited in Kidney Int. 2015 Feb;87(2):272-5. doi: 10.1038/ki.2014.359.

- The article used to support the findings of study have been deposited in Am J Kidney Dis. 2002 Feb;39(2 Suppl 1):S1-266
- The article used to support the findings of study have been deposited in Transplantation. 2017 Jun;101(6): 1152-1158.
doi: 10.1097/TP.0000000000001351.
- The article used to support the findings of study have been deposited in Transplant Direct. 2018 May; 4(5): e342. Published online 2018 Apr 18.
doi: 10.1097/TXD.0000000000000781.
- The article used to support the findings of study have been deposited in Transplantation. 2019 Feb;103(2): 401-411.
doi: 10.1097/TP.0000000000002309.
- The article used to support the findings of study have been deposited in Transplant Direct. 2017 Jul; 3(7): e177. Published online 2017 Jun 23.
doi:10.1097/TXD.0000000000000680.
- The article used to support the findings of study have been deposited in Curr Opin Organ Transplant. 2013 Apr;18(2):174-8.
doi: 10.1097/MOT.0b013e32835e2a08.

FUNDING

Not applicable.

CONFLICT OF INTEREST

The authors declare no conflict of interest, financial or otherwise.

ACKNOWLEDGEMENTS

The author would like to thank the editorial board and all staff of Renal Transplant Team Dr. Kariadi General Hospital Semarang, for their support to this research.

REFERENCES

- [1] Projosudjadi W. Incidence, prevalence, treatment and cost of end-stage renal disease in Indonesia. *Ethn Dis* 2006; 16(Suppl.2): 14-6.
- [2] Hill NR, Fatoba ST, Oke JL, *et al.* Global prevalence of chronic kidney disease- asystematic review and meta-analysis. *PLoS One* 2016; 11(7):e0158765
[http://dx.doi.org/10.1371/journal.pone.0158765] [PMID: 27383068]
- [3] Perhimpunan Nefrologi Indonesia. Jakarta: Konsensus Transplantasi Ginjal Perhimpunan Nefrologi Indonesia (Pernefri) 2013.
- [4] Dina-Nilasari. Konsensus Evaluasi Donor dan Resipien Transplantasi Ginjal Serta penatalaksanaan Perioperatif. *Medicus* 2015; 28: 45-55.
- [5] Supit T, Nugroho EA, Santosa A, Soedarso MA, Daniswara N, Addin SR. Kidney transplantation in Indonesia: An update. *Asian J Urol* [internet] Elsevier Ltd 2019; 1-7.
[http://dx.doi.org/10.1016/j.ajur.2019.02.003]
- [6] Campbell S, Pilmore H, Gracey D, Mulley W, Russell C, McTaggart S. KHA-CARI guideline: Recipient assessment for transplantation. *Nephrology (Carlton)* 2013; 18(6): 455-62.
[http://dx.doi.org/10.1111/nep.12068] [PMID: 23581832]
- [7] Kayler D. Kidney Transplant Ureteroneocystostomy Techniques and Complications: Review of the Literature 2010; 1413-20.
- [8] Cavaille-Coll M, Bala S, Velidedeoglu E, *et al.* Summary of FDA workshop on ischemia reperfusion injury in kidney transplantation. *Am J Transplant* 2013; 13: 1134-48.
[http://dx.doi.org/10.1111/ajt.12210]
- [9] Debut A, Foucher Y, Trébern-Launay K, *et al.* Each additional hour of cold ischemia time significantly increases the risk of graft failure and mortality following renal transplantation. *Kidney Int* 2015; 87(2): 343-9.
[http://dx.doi.org/10.1038/ki.2014.304] [PMID: 25229341]
- [10] Ponticelli CE. The impact of cold ischemic time on renal transplant outcome. *Kidney Int* 2018; 272-4.
- [11] National Kidney Foundation. K/DOQI clinical practice guidelines for chronic kidney disease: Evaluation, classification and stratification 2002. <http://doi.org/10.1634/theoncologist.2011-S2-45>.
- [12] Wong G, Teixeira-Pinto A, Chapman JR, *et al.* The impact of total ischemic time, donor age and the path-way of donor death on graft outcomes after deceased donor kidney transplantation. *Transplantation* 2017; 101(6): 1152-8.
[http://dx.doi.org/10.1097/TP.0000000000001351] [PMID: 27479168]
- [13] Amanda J. Factors associated with prolonged warm ischemia time among deceased donor kidney. *Transplant Recipients Kidney transplantation* 2018.
- [14] Serrano OK, Vock DM, Chinnakotla S, *et al.* The relationships between cold ischemia time, kidney transplant length of stay, and transplant-related costs. *Transplantation* 2019; 103(2): 401-11.
[http://dx.doi.org/10.1097/TP.0000000000002309] [PMID: 29863580]
- [15] Nugroho EA, Kamar MA, Junita D. The correlation between total ischemic time with length of hospitalization in kidney transplantation: A single centre report. *Int J Sci Eng* 2019; 13(1): 28-3.
- [16] van der Vliet JA, Warlé MC. The need to reduce cold ischemia time in kidney transplantation. *Curr Opin Organ Transplant* 2013; 18(2): 174-8.
[http://dx.doi.org/10.1097/MOT.0b013e32835e2a08] [PMID: 23337956]

© 2019 Nugroho. *et al.*

This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 International Public License (CC-BY 4.0), a copy of which is available at: <https://creativecommons.org/licenses/by/4.0/legalcode>. This license permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Case Report Correlation of Warm and Cold Ischemic Time to Graft Function in Kidney Transplant: A Single Centre Report

ORIGINALITY REPORT

13%

SIMILARITY INDEX

8%

INTERNET SOURCES

9%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1 pubmed.ncbi.nlm.nih.gov 1%
Internet Source

2 repository.ubn.ru.nl 1%
Internet Source

3 www.quintessence-publishing.com 1%
Internet Source

4 www.kidneynews.org 1%
Internet Source

5 Germaine Wong, Armando Teixeira-Pinto, Jeremy R. Chapman, Jonathan C. Craig, Henry Pleass, Stephen McDonald, Wai H. Lim. "The Impact of Total Ischemic Time, Donor Age and the Pathway of Donor Death on Graft Outcomes After Deceased Donor Kidney Transplantation", Transplantation, 2017 1%
Publication

6 Submitted to Mount St. Marys College 1%
Student Paper

"Abstracts", BJU International, 2018

7	Publication	1 %
8	Submitted to Higher Education Commission Pakistan Student Paper	1 %
9	Submitted to Universitas Pendidikan Indonesia Student Paper	1 %
10	"RHEUMATOID ARTHRITIS ? TREATMENT", APLAR Journal of Rheumatology, 8/2006 Publication	<1 %
11	studyres.com Internet Source	<1 %
12	www.nsmc.indoscholar.com Internet Source	<1 %
13	www.arcjournals.org Internet Source	<1 %
14	sjkdt.org Internet Source	<1 %
15	openorthopaedicsjournal.com Internet Source	<1 %
16	Albertus Marcelino, Chaidir Arif Mochtar, Irfan Wahyudi, Agus Rizal Hamid. "Kidney transplantation: A new era of laparoscopic	<1 %

living donor nephrectomy in Indonesia", Asian
Journal of Surgery, 2018

Publication

17

journals.plos.org

Internet Source

<1 %

18

Michael A. J. Moser, Nathan Ginther, Yigang Luo, Gavin Beck et al. "Early experience with hypothermic machine perfusion of living donor kidneys - a retrospective study", Transplant International, 2017

Publication

<1 %

19

www.revistanefrologia.com

Internet Source

<1 %

20

Eriawan Nugroho, Aulia Ahimsa Martawiguna, Ardy Santosa, MOH. Adi Soedarso, Nanda Daniswara. "The Profile of Kidney Transplantation in Dr. Kariadi Hospital Semarang, Indonesia", Transplantation, 2018

Publication

<1 %

21

Kazunari Yoshida. "Factors contributing to long graft survival in non-heart-beating cadaveric renal transplantation in Japan: a single-center study at Kitasato University", Clinical Transplantation, 12/2002

Publication

<1 %

22

Krishnan, Anoushka R, Germaine Wong, Jeremy R Chapman, Patrick T Coates, Graeme

<1 %

R Russ, Henry Pleass, Christine Russell, Bulang He, and Wai H Lim. "Prolonged Ischemic Time, Delayed Graft Function, Graft and Patient Outcomes in Live-Donor Kidney Transplant Recipients", American Journal of Transplantation, 2016.

Publication

23

Mathieu Lesouhaitier, Camille Legeai, Emilie Savoye, Christelle Cantrelle et al. "Impact of donor coronary angiography on kidney transplantation outcomes", Clinical Transplantation, 2018

Publication

24

Tjokorda Gde Dalem Pemayun, Ridho M. Naibaho. "Clinical profile and outcome of diabetic foot ulcer, a view from tertiary care hospital in Semarang, Indonesia", Diabetic Foot & Ankle, 2017

Publication

25

Wolfgang Baar, Kai Kaufmann, Kai Silbach, Bernd Jaenigen et al. "Early Postoperative Use of Diuretics After Kidney Transplantation Showed Increase in Delayed Graft Function", Progress in Transplantation, 2020

Publication

26

"Abstracts", Journal of Hepato-Biliary-Pancreatic Sciences, 2022

Publication

<1 %

<1 %

<1 %

<1 %

27

Chaidir A. Mochtar, Fekhaza Alfarissi, Adhitama A. Soeroto, Agus Rizal A.H. Hamid et al. "Milestones of kidney transplantation in Indonesia", Medical Journal of Indonesia, 2017

Publication

<1 %

28

K. Dziewanowski, M. Myślak, R. Drozd, E. Krzystolik et al. "Factors Influencing Long-Term Survival of Kidney Grafts Transplanted From Deceased Donors—Analysis Based on a Single-Center Experience", Transplantation Proceedings, 2018

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Case Report Correlation of Warm and Cold Ischemic Time to Graft Function in Kidney Transplant: A Single Centre Report

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
