

Fortified Dadih (Fermented Buffalo Milk) with Vitamin D3 Improves Interleukin-6 and Caecum Short Chain Fatty Acids on Diet-Induced Obese Rat

by Yora Nindita

Submission date: 24-Mar-2022 11:04AM (UTC+0700)

Submission ID: 1791528817

File name: FATTY_ACIDS_ON_DIET-INDUCED_OBESE_RAT_-_Azzahra_Mutiara_Ayu.pdf (257.8K)

Word count: 6127

Character count: 31867

Original Article

FORTIFIED DADIH (FERMENTED BUFFALO MILK) WITH VITAMIN D3 IMPROVES INTERLEUKIN-6 AND CAECUM SHORT CHAIN FATTY ACIDS ON DIET-INDUCED OBESE RAT

SARI BEMA RAMDIKA¹, ENDANG MAHATI², ANANG MOHAMAD LEGOWO³, MUFLIHATUL MUNIROH⁴, YORA NINDITA², DIANA NUR AFIFAH^{1*}

¹Department of Nutrition Science, Faculty of Medicine, Diponegoro University, Semarang 50275, Indonesia, ²Department of Pharmacology and Therapy, Faculty of Medicine, Diponegoro University, Semarang 50275, Indonesia, ³Department of Food Technology, Faculty of Animal and Agricultural Sciences, Diponegoro University, Semarang 50275, Indonesia, ⁴Department of Physiology, Faculty of Medicine, Diponegoro University, Semarang 50275, Indonesia
Email: d.nurafifah.dna@fk.undip.ac.id

Received: 24 Jul 2020, Revised and Accepted: 02 Sep 2020

ABSTRACT

Objective: This study aims to determine the effect of fortified dadih with vitamin D3 on IL-6 expression level and the concentration of caecum SCFA in obese rats.

Methods: A total of 30 male *Sprague Dawley* rats were divided into five equal groups: healthy-control-(K-), obese-control-(K+), obese-intervention-(X1, X2, and X3). K(+), X1, X2, and X3 were in obesity conditions, which was induced by a high-fat sucrose diet (HFSD) and K(-) as a healthy-control-group. Furthermore, vitamin D3-fortified dadih at doses of 4 g/200 g-body-weight/d, dadih only at doses of 4 g/200 g-body-weight/d, and vitamin D3 only at 36 IU/200 g-body-weight/d was administered to X1, X2, and X3 groups, respectively.

Results: Treatment using fortified dadih with vitamin D3 showed significantly reduce weight gain ($p < 0.05$) compare to K(+) and X2. In addition, X1 showed a decreased level of Interleukin-6 expression ($p < 0.05$) than K(+), X2, and X3 groups but higher than K(-). Also, it showed the highest total SCFA, acetate, and propionate concentration ($p < 0.05$). However, a moderately negative correlation was discovered between the pair of total SCFA and Interleukin-6 expression, acetate and Interleukin-6 expression, SCFA and body weight, propionate and body weight, butyrate and body weight. On the contrary, a strong positive correlation was observed between the pair of Interleukin-6 expression levels and body weight.

Conclusion: This study shows that fortified dadih with vitamin D3 from fermented foods improve the expression level of Interleukin-6 and increase the production of SCFA. Also, they improve intestinal homeostasis because of the increased SCFA production.

Keywords: Obesity, Dadih, Vitamin D3, IL-6, SCFA

© 2020 The Authors. Published by Innovare Academic Sciences Pvt Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>) DOI: <https://dx.doi.org/10.22159/ijpps.2020v12i11.39209>. Journal homepage: <https://innovareacademics.in/journals/index.php/ijpps>.

41

INTRODUCTION

Obesity is a chronic metabolic disorder characterized by increased fat storage in the body [1, 2]. When energy consumption exceeds expenditure, 60-80% of the excess is stored as body fat [3]. Its development is not only a balance between energy intake and expenditure but also between white and brown adipose tissue in the form of thermogenesis through increased expression of mitochondria Uncoupling Protein 1 (UCP-1) [4]. At the developmental stage, enlargement of adipose tissue caused adipocyte hypertrophy [5], and this increased production of Tumour Necrotic Factor-Alpha (TNF- α), Interleukin-6 (IL-6) and Monocyte Chemotactic Protein-1 (MCP-1) [6].

Dadih is a traditional food from West Sumatera, Indonesia, and it is also known as fermented buffalo milk [7]. Some of the LAB that is usually found in dadih are *Lactobacillus Plantarum*, *Leuconostoc Mesenteroides*, *Streptococcus Faecalis*, *S. lactis* subspecies *lactis*, *S. cremoris*, *L. casei* subspecies *casei*, and *Lactobacillus casei* subspecies *Rhamnosus*, and *Lactococcus* [7]. In addition, as a regulator of obesity, LAB helps to reduce fat absorptions, energy intake, the weight of white fat tissue, the average size of adipocytes, modulates intestinal microbiota, as well as inhibits the differentiation of 3T3-L1 adipocytes, and the growth of pathogenic bacteria in the gastrointestinal tract [8-12]. Pathogenic bacteria within the gut stimulate production and secretion of Lipopolysaccharides (LPS) from intestinal epithelial cells, and they bind to cytokine receptors on hepatocytes and adipocytes to trigger the release of pro-inflammatory cytokines (TNF α , IL6, IL1 β , and MCP1). However, probiotic therapy with LAB reduced the release of LPS from the intestinal epithelial cells, and it decreases the production of proinflammatory cytokines in adipose tissue [13].

Modulated intestinal microbiota with probiotics is a promising approach to improving host health since it is protected from infection and disease as well as produces vitamins and Short Chain Fatty Acids (SCFAs) [14, 15]. In the intestinal lining, it contributes to energy metabolism through the production of SCFAs, which are usually produced from the fermentation of fibers, proteins, and peptides [16]. The end products of this reaction are acetate, propionate, and butyrate. Cell culture studies showed that these products can change the function of adipose tissue by reducing the production of proinflammatory molecules and stimulating adipogenesis. In addition, oral administration of butyrate influences weight control through increased energy expenditure and fat oxidation in obese rats [17].

Vitamin D is an important regulator of immunity, composition and function of intestinal microbial, and metabolic processes in the body [18-20]. The deficiency of vitamin D leads to several diseases such as obesity, diabetes, cardiovascular disease, and metabolic syndrome [21]. Research showed a relationship between its deficiency and obesity since obese individuals tend to have a low 25 hydroxyvitamin D (25(OH)D) serum level [22-24]. Furthermore, the fortification of dadih with vitamin D3 is expected to increase its intake in obese people. The active form of vitamin D, 1,25-dihydroxyvitamin D3 (1,25(OH)2D3) inhibits IL-6 secretion by suppressing phosphorylation of Akt and signaling of Nuclear Factor Kappa B (NF- κ B) [25]. Yogurt is a fermented milk product similar to dadih, and when fortified with vitamin D, it caused weight loss in obese patients with type 2 diabetes mellitus [26].

This study aimed to determine the effect of fortified dadih with vitamin D3 on IL-6 expression level and concentration of caecum SCFA in obese rats.

MATERIALS AND METHODS

Dadiah preparation

100 ml buffalo milk was pasteurized at 72 °C for 15 s [27] and cooled until 30 °C, then vitamin D3 was added at a dose of 900 IU. Pasteurized milk was poured into a clean bamboo tube and then covered with banana leaves (fig. 1). The tube was placed at room temperature for two days. The dadih is stored at -18 °C (frozen) for future uses (fig. 2).



Fig. 1: Fortified dadih during spontaneous fermentation



Fig. 2: Fortified dadih after fermentation

Research design and experimental animals

This research was a true-experiment study using a randomized post-test with control group design only. The animal used was male *Sprague Dawley* rats, aged 8–12 w, weighing 150–200 g. They were acclimatized at the laboratory of the Center for Food and Nutrition Studies at Gajah Mada University, Yogyakarta. In addition, they were placed in individual stainless-steel cages at regulated temperatures (21 °C) and housed in a cleaned and germ-free place. The experimental animals were fed with 20 g/d of the Comfeed II standard-diet (7% fat) during the non-HFSD and intervention period. Also, they received ad-libitum water during the experiment, and animal care was performed following the Guidelines of the Central Laboratory for Food and Nutrition Studies.

Thirty rats were divided into five groups, which were obese-control (K+), obese-intervention (X1, X2, and X3) and healthy-control (K-) groups. The obese-condition was induced by HFSD for 28 d, and after this period, the X1, X2, and X3-groups were treated daily with vitamin D3-fortified dadih at doses of 4 g/200 g-body-weight/d, dadih without fortification at doses of 4 g/200 g-body-weight/d, and vitamin D3 at doses of 36 IU/200 g-body-weight/d, respectively. The blood sample and caecum digesta were obtained at the end of the intervention. Furthermore, IL-6 level and SCFA concentration were analysed by ELISA (EliKine™, China) and gas chromatography,

respectively. This study was approved by the Health Research Ethics Commission (KEPK) of the Faculty of Medicine Diponegoro University, Semarang through ethical clearance No. 129/EC/H/KEPK/FK-UNDIP/X/2019.

Serum IL-6 determination

The determination of serum IL-6 was performed using ELISA-Kit according to the manufacturer's instructions (EliKine™, China). Also, 100 µl of standard and serum of each sample were transferred into well plates, which were sealed with a cover and incubated at room temperature for 2 h. After incubation, the wells were aspirated and washed by filling them with washing buffer (250 µl), repeating the process twice for a total of three washes. 100 µl of diluted Mouse IL-6 detected antibodies within the well and incubated at room temperature for 1 h. Thereafter, the wells were aspirated and washed by filling them with washing buffer (250 µl), repeating the process twice for a total of three washes. Similarly, 100 µl of the working dilution of Streptavidin-HRP was added to each well. In addition, the plate was sealed with a cover and incubated at room temperature for 30 min in the dark, and the wells were aspirated and washed by filling them with washing buffer (250 µl), repeating the process twice for the total of three washes. 100 µl of HRP substrate (TMB) was also added and sealed with a cover. Furthermore, it was incubated at room temperature for 15 min in the dark. Finally, 50 µl of stop solution was added, and the color turned yellow immediately. The sample's absorbance was further read by ELISA Reader at 450 nm.

SCFA determination

The SCFA determination was performed using gas chromatography, and a total of 0.05 g of caecal content was acidified with 0.05 ml of sulphuric acid. Furthermore, SCFA was extracted with 0.6 ml of diethyl ether and centrifuged at 14000 r. p. m. for 30 s. One microliter of the organic phase was injected directly into the capillary column (Nukol) of the gas chromatography, which was combined with a flame ionization detector. The initial temperature was 80 °C while the final was 200 °C. Nitrogen was used as a carrier gas and sample quantification was performed using a calibration curve for acetic, propionic, and butyric acid. A standard curve was performed for the quantitation of acetic, propionic, and butyric acid in the sample [28]. The total SCFA is the amount of acetate, propionate, and butyrate [29].

Chemicals and reagents

Sulphuric acid and diethyl ether were purchased from Sigma-Aldrich Co, St Louis, MO, USA. Vitamin D3 and ELISA Kit were purchased from Wellness Inc., USA, and Abbkine Inc., Wuhan, Hubei, China, respectively.

Statistical analysis

The results were expressed as mean±SD (for normally distributed data), which was otherwise manifested as a median (min-max). The statistical difference was analyzed using One-Way Analysis of Variance (ANOVA) followed by post hoc Bonferroni for normally distributed data. In addition, the Kruskal Wallis test followed by Mann-Whitney-U-test was used (SPSS 16). Spearman's correlative test was used to analyze the correlation between variables, and statistical analyses were performed by a computer. Likewise, the differences and correlations were considered significant at p-value<0.05 as well as 95% confidence intervals, and its strength was determined by r value.

RESULTS AND DISCUSSION

Body weight

The processed data were obtained from 30 *Sprague Dawley*-rats, which were divided into each group consisting of 6. There was an increase in body weight during the HFSD period in groups K (+), X1, X2, and X3 (they had p = 0.001; table 1). Furthermore, the body weight in K (-) increased even though it only received a standard feed. The weight gain of those that received HFSD was higher than K (-) that received a standard feed. However, bodyweight increase during the HFSD period was caused by a higher proportion of fat and

sucrose. Therefore, K (+), X1, X2, and X3 groups have higher weight gain than K (-). The consumption of high-energy foods such as fat and sugar are the main factors that cause obesity [30]. Also, previous studies have suggested that HFSD increased body weight to 51% fat and 24% lean mass due to changes in composition [30].

After intervention of fortified dadih, dadih without fortification, and vitamin D3, K (+) gained more body weight compared to K (-), X1, X2, and X3 groups. A statistical difference was obtained on the pre-post-intervention change (Δ) of rat-BW across the groups (Kruskal Wallis test; $p = 0.001$) followed by a Mann Whitney U-test between two-groups. However, there was no significant difference between X1 and X3 groups compared with K (-) ($p = 0.545$, $p = 0.789$, table 1). In addition, it was reported that fortified dadih with vitamin D3 reduced body weight gain in obese rats since it contained LAB. Several studies have shown that LAB and vitamin D3 can reduce gain in body weight and epididymal fat mass [8, 31]. Similarly, research on the administration of *Lactobacillus Plantarum Strain Ln4* reduced weight gain in mice [32]. Vitamin D has been reported to have a link with obesity as some studies showed that its concentration is low in obese individuals [33, 34]. Furthermore, its deficiency causes a decreased amount of intestinal calcium and phosphorus absorption, which increases PTH levels [35]. This increase has been hypothesized as an independent predictor of obesity. One hypothesis suggested that high physiological levels of PTH will increase calcium influx to adipocytes. In addition, it leads to increased lipogenesis and reduced catecholamine-induced lipolysis as a mechanism to increase fat storage [36].

IL-6 expression level

The post-intervention-IL-6-levels was significantly different between the groups ($p = 0.000$; table 1). Its levels of X1 was significantly different with K (+), X2 and X3 groups ($p = 0.000$, $p = 0.000$, and $p = 0.003$, respectively; table 1), while the mean values were lower compared to X2 and X3. This result showed that fortified dadih with vitamin D3 reduced the production of proinflammatory cytokines compared to those without fortification and vitamin D3.

Obesity, which causes inflammation, is characterized by increased circulation of inflammatory cytokines and adipokines [37]. Various pro-inflammatory and anti-inflammatory factors such as adipokines, leptin, adiponectin, resistin, as well as cytokines and chemokines, such as TNF- α , IL-6, and MCP-1 are produced and released by adipocytes [38].

Lactic Acid (LA) as the main metabolite of LAB reduces the production of inflammatory cytokines. Furthermore, it decreases pro-inflammatory cytokines especially TNF- α and IL-6 by reducing phosphorylation of I κ B α since it inhibits the activation of NF- κ B [39].

As shown in several studies, some LABs similar to dadih may reduce the production of pro-inflammatory cytokines. Research on the administration of *L. curvatus HY7601* and *L. Plantarum KY1032*, which was obtained from the Korean traditional fermentation of cabbage reduced the expression of IL-6 gene in rat epididymal fat [13]. Administration of *Lactobacillus casei strain Shirota* supplements to the experimental animals also reduced its production [40].

An active metabolite of vitamin D, 1,25(OH) $_2$ D inhibits proinflammatory cytokine production (IL-1 β , IL-6, IL-8, IL-12), reduces inflammatory activity in adipocytes and inflammation in visceral fat tissue [34]. In addition, it suppresses the ability of T cell and monocyte proliferation as well as its stimulation and decreases proinflammatory cytokines such as C Reactive Protein (CRP), TNF α , IL 6, IL-1, and IL-8, and increases anti-inflammatory types such as IL-10. This metabolite a ligand for the vitamin D receptor (VDR) [41], which inhibits NF κ B activation and signaling [42]. *In vitro* study also showed a link between the absence of VDR and increased activity of NF- κ B [43]. However, inflammatory signaling was mediated by the NF κ B pathway since it increased the expression of proinflammatory cytokines such as IL-6 and TNF- α [44].

SCFA concentration and ratio

The post-intervention total of SCFA, acetate, and butyrate concentration was significantly different between groups ($p = 0.029$, $p = 0.013$, and $p = 0.022$, respectively; table 1). Their concentrations of X1 were significantly different with K (+) ($p = 0.016$; $p = 0.016$; and $p = 0.016$; table 1, respectively) but not with K (-) group ($p = 0.109$; $p = 0.150$; and $p = 0.262$; table 1, respectively). Similarly, their concentrations were significantly different from the X3 group but not with X2. The mean of total SCFA, acetate, and propionate concentration of X1 was higher compared to other groups. This result showed that fortified dadih with vitamin D3 is better at producing SCFA compared to those without fortification and vitamin D3.

Furthermore, the ratio of acetate, propionate, and butyrate was significantly different between groups ($p = 0.001$, $p = 0.001$, and $p = 0.001$, respectively; table 1). There are several references to this ratio, such as 60:20:20 and 60:25:15 [45, 46]. The difference depends on the type of substrate, diet, and composition of the intestinal microbiota [46, 47].

The production of SCFA by microbial fermentation of dietary fiber was associated with various health benefits, including improvements in body composition, energy homeostasis, lipid profile, and weight loss by stimulating several hormonal and nerve pathways, such as appetite control, increased oxidation, and muscle thermogenesis as well as liver tissue [17]. Its effect on body weight and food intake occurs through G-protein coupled receptors (GPR, GPR41, and GPR 43). GPR41 and GPR 43 are known as a Free Fatty Acid Receptor 3 (FFAR3) and 2 (FFAR2), respectively. FFAR2 is expressed on immune cells and primarily reacts to acetate and propionate, while FFAR3 is mostly present in adipocytes and preferentially binds to propionate and butyrate. Their activation increases the synthesis and signaling of satiety hormones such as GLP-1 and PYY, causing decreased appetite and increased energy expenditure [48].

The correlation between variables

The Spearman test showed that a moderate correlation was obtained between variables. A moderate negative correlation was observed between total SCFA and IL-6-levels ($p = 0.021$; $r = -0.419$; table 2). Furthermore, it was observed between acetate and IL-6-levels ($p = 0.027$; $r = -0.403$; table 2) as well as between body weight and each of total SCFA, propionate, butyrate ($p = 0.020$; $r = -0.423$, $p = 0.037$; $r = -0.383$, $p = 0.008$; $r = -0.473$, respectively; table 2). Also, a very strong positive correlation was observed between body weight and IL-6-levels ($p = 0.000$; $r = 0.802$; table 2).

SCFA prevented the production of LPS from gram-negative bacterial cell membranes, and it has been linked to endotoxemic metabolism, inflammation, insulin resistance, adiposity, and liver fat [49]. Furthermore, SCFA reduces systemic inflammation through molecular signaling pathways by activating G-protein coupled receptor 41/43 and inhibition of the histone deacetylase (HADACs) [50, 51].

The effect of SCFA on body weight and food intake occurs through G-protein coupled receptors (FFAR3 and FFAR2). Their activation increases the synthesis and signaling of satiety hormones such as GLP-1 and PYY, causing decreased appetite and increased energy expenditure [48]. Butyrate prevents obesity by activating β 3 adrenergic receptors in white adipose tissue. Acetate plays a role in regulating appetite in the brain, increasing fat oxidation, improving glucose homeostasis, and inflammatory status [52]. However, its role in the regulation of appetite, fat, and weight gain remains controversial. Besides, a previously conducted study has reported its influence on central appetite regulation and reduced food intake [53]. Other studies showed the opposite effect, causing an increase in ghrelin secretion, hyperphagia, and obesity in rats [54]. These conflicting results may be related to metabolic status and the location of its administration [52]. Similar to these results, acetate does not have a significant correlation with body weight. In humans and mice, propionate is associated with preventing the accumulation of hepatic lipids through the suppression of genes involved in fatty acid synthesis [52].

Table 1: The effect of fortified dadih with vitamin D3 on body weight, IL-6 and SCFA

Groups Marker	K(-)	K(+)	X1	X2	X3	p ¹
Body Weight after HFD administration (gr)						
Pre	181.67±5.2	186.83±5.19	184.5±5.24	181.5±3.72	184.5±8.24	0.468
Post	202.17±5.38 ^{b,c,d,e}	220.33±5.04 ^a	220.5±4.03 ^a	217.5±3.98 ^a	220±7.09 ^a	0.001
Δ	20.5±1.04 ^{b,c,d,e}	33.5±3.88 ^a	36.0±1.67 ^a	36.0±0.63 ^a	35.5±1.37 ^a	0.001
p	0.001	0.001	0.001	0.001	0.001	
Body Weight of intervention (gr)						0.005
Pre	201.0	221.5	220.0	218.0	220.0	0.001
	(196-211) ^{b,c,d,e}	(211-225) ^a	(215-226) ^a	(212-223) ^a	(212-231) ^a	0.001
Post	225.0	256.5	244.0	245.5	243.0	
	(220-234) ^{b,c,d,e}	(248-261) ^{a,c,d,e}	(240-249) ^{a,b}	239-250) ^{a,b}	(237-254) ^{a,b}	
Δ	23.5	35.0	24.5	27.0	24.0	
	(23.0-25.0) ^{b,d}	(35.0-37.0) ^{a,c,d,e}	(23.0-25.0) ^b	(27.0-28.0) ^{a,b}	(23.0-25.0) ^b	
p	0.026	0.023	0.026	0.023	0.024	
IL-6 (pg/ml)						
Post	66.81±6.23 ^{b,c,d,e}	176.53±8.21 ^{a,c,d,e}	85.66±9.08 ^{a,b,d,e}	127.12±1.16 ^{a,b,c,e}	106.6±6.86 ^{a,b,c,d}	0.000
Total SCFA (mmol/kg)						
Post	26.11	15.56	32.07	20.09	22.04	0.029
	(3.23-33.54)	(11.86-32.57) ^{c,d}	(24.12-40.83) ^{b,e}	17.57-40.64) ^b	(16.43-28.32) ^c	
Acetate (mmol/kg)	12.35	7.14	15.78	9.6	8.39	0.013
Post	(1.37-18.15)	(5.73-15.21) ^{c,d}	(11.71-21.02) ^{b,e}	(8.21-19.21) ^b	(6.44-11.11) ^c	
Propionate (mmol/kg)	9.25	6.12	11.11	6.71	8.15	0.093
Post	(1.13-11.66)	(4.21-12.47) ^c	(8.09-14.11) ^b	(5.97-13.70)	(5.82-10.41)	
Butyrate (mmol/kg)	4.24	2.3	5.04	3.78	5.6	0.022
Post	(0.73-5.96)	(1.92-4.89) ^{c,d,e}	(4.32-5.44) ^b	(3.20-7.73) ^b	(4.17-6.80) ^b	
The ratio of Acetate (mmol/kg)	48.60±5.35	46.19±1.71 ^c	49.63±1.63 ^c	47.32±1.03 ^c	39.18±1.34 ^{b,c,d}	0.001
Post						
Ratio of Propionate (mmol/kg)	33.73±2.84 ^b	38.43±1.76 ^{a,c,d}	34.32±1.72 ^b	33.87±0.54 ^b	36.38±1.07	0.001
Post						
The ratio of Butyrate (mmol/kg)	17.63±5.36	15.36±0.67 ^{d,e}	16.06±3.01 ^c	18.79±0.83 ^{b,e}	24.43±1.04 ^{b,c,d}	0.001
Post						

Five groups of rats (n=6 for each group) consists of K(-): control healthy and K(+): control Obese; X1: Fortified Dadih Treatment; X2: Dadih Treatment; X3: Vitamin D3 Treatment; Δ: changes between pre and post value; differences between the groups of rats were analyzed using ANOVA (BW HFD, IL-6, Ratio of Acetate, Propionate, and Butyrate) Kruskal Wallis (BW intervention, Total SCFA, acetate, propionate, butyrate); p: value between pre and post-treatment were analyzed using Paired t-test/Wilcoxon test; alphabetical superscripts showed a significance level of *p<0.05 compared as K(-); ^bp<0.05 as compared to K(+); ^cp<0.05 as compared to X1; ^dp<0.05 as compared to X2; ^ep<0.05 as compared to X3. The data were written as mean±SD for normally distributed data and median (Min-Max) when data were not normally distributed; p¹: value between all of the groups were analyzed using ANOVA if data are normally distributed and Kruskal Wallis if data are not normally distributed.

Table 2: Correlation between SCFA concentration, IL-6 levels, and body weight after fortified dadih with vitamin D3 intervention

	IL-6		Body weight	
	r	p	r	p
Total SCFA	-0.419	0.021*	-0.423	0.020*
Acetate	-0.403	0.027*	-0.359	0.052
Propionate	-0.355	0.054	-0.383	0.037*
Butyrate	-0.286	0.126	-0.473	0.008*
Body Weight	0.802	0.000*	-	-

Thirty-five of rats were analysed, statistical significance calculated by Spearman test *Value are statistically significant at p<0,05

The expansion of adipose tissue in obesity caused high IL-6 in blood circulation [55]. The expression of white adipose tissue and plasma IL-6 correlate with increased body mass, waist circumference, and free fatty acid levels [56]. In addition, increased expression of IL-6 Receptor (IL-6R) and IL-6 in adipose tissue was positively correlated with clinical indicators of obesity, such as body mass index and fat percentage [55].

Considering the results, it is reasonable to conclude that fortified dadih with vitamin D3 significantly reduces IL-6 expression level and increases

the production of SCFA. However, the IL-6 expression level was not significantly different from healthy-control since it was lower than obese-control, dadih without fortification, and vitamin D3 groups. Furthermore, the total SCFA, acetate, and propionate concentration of fortified dadih with vitamin D3 groups were higher compared to others.

CONCLUSION

Fortified dadih with vitamin D3 reduces weight gain, improves the expression level of IL-6, and increases the production of SCFA. Also,

they improve intestinal homeostasis because of the increased SCFA production. The strong correlation between IL-6 and body weight indicated that the improvement of IL-6 related to the reduction of body weight gain.

ACKNOWLEDGMENT

The authors are grateful to The Faculty of Medicine, Diponegoro University

FUNDING

This research was funded by The Faculty of Medicine, Diponegoro University (Grant Number: 1160/UN7.5.4.2.1/PP/PM/2020).

AUTHORS CONTRIBUTIONS

Sari Bema Ramdika: Conceived the study question and the study design, conducted of data collection, data analysis and interpretation, and writing the manuscript.

Endang Mahati: Designed the study, supervised data collection and data analysis, contributed to data interpretation, and review and editing the manuscript.

Anang Mohamad Legowo: Designed the study, contributed to data interpretation, and review and editing the manuscript.

Muflihatul Muniroh: Designed the study, contributed to data interpretation, and review and editing the manuscript.

Yora Nindita: Designed the study, contributed to data interpretation, and review and editing the manuscript.

Diana Nur Afifah: Managed the experimental processes, supervised data collection and data analysis, contributed to data interpretation, and writing the manuscript.

CONFLICT OF INTERESTS

The authors declare no conflict of interest

REFERENCES

- Schwartz MW, Seeley RJ, Zeltser LM, Drewnowski A, Ravussin E, Redman LM, *et al.* Obesity pathogenesis: an endocrine society scientific statement. *Endocr Rev* 2017;38:267–96.
- Gu Z. Tailoring biomaterials for metabolic diseases. *Adv Healthc Mater* 2019;8:1–2.
- Oussaada SM, Galen KA van, Cooiman MI, Kleinendorst L, Hazebroek EJ, Haelst MM van, *et al.* The pathogenesis of obesity. *Metabolism* 2019;92:26–36.
- Gomez Hernandez A, Beneit N, Diaz Castroverde S, Escribano O. Differential role of adipose tissues in obesity and related metabolic and vascular complications. *Int J Endocrinol* 2016;2016:1–15.
- Verboven K, Wouters K, Gaens K, Hansen D, Bijnen M, Wetzels S, *et al.* Abdominal subcutaneous and visceral adipocyte size, lipolysis and inflammation relate to insulin resistance in male obese humans. *Sci Rep* 2018;8:4677–85.
- Chan PC, Hsieh PS. The role of adipocyte hypertrophy and hypoxia in the development of obesity-associated adipose tissue inflammation and insulin resistance. W: Gordeladze JO, red. *Adiposity-Omics and Molecular Understanding*. London: InTech; 2017. p. 127–41.
- Venema K, Surono IS. Microbiota composition of dadih-a traditional fermented buffalo milk of West Sumatra. *Lett Appl Microbiol* 2019;68:234–40.
- Mazloom K, Siddiqi I, Covasa M. Probiotics: how effective are they in the fight against obesity? *Nutrients* 2019;11:258–82.
- Alipin K, Saitir R. The potential of indigenous lactic acid bacteria against *Salmonella* sp; 2016.
- Hong SM, Chung EC, Kim CH. Anti-obesity effect of fermented whey beverage using lactic acid bacteria in diet-induced obese rats. *Korean J Food Sci Anim Resour* 2015;35:653–9.
- Park SY, Cho SA, Lee MK, Lim SD. Effect of *Lactobacillus plantarum* FH185 on the reduction of adipocyte size and gut microbial changes in mice with diet-induced obesity. *Korean J Food Sci Anim Resour* 2015;35:171–8.
- Tsai YT, Cheng PC, Pan TM. Anti-obesity effects of gut microbiota are associated with lactic acid bacteria. *Appl Microbiol Biotechnol* 2014;98:1–10.
- Park DY, Ahn YT, Park SH, Huh CS, Yoo SR, Yu R, *et al.* Supplementation of *Lactobacillus curvatus* HY7601 and *Lactobacillus plantarum* KY1032 in diet-induced obese mice is associated with gut microbial changes and reduction in obesity. *PLoS One* 2013;8:108–35.
- Azad MAK, Sarker M, Li T, Yin J. Probiotic species in the modulation of gut microbiota: an overview. *Biomed Res Int* 2018;2018:1–8.
- Umu OCO, Rudi K, Diep DB. Modulation of the gut microbiota by prebiotic fibres and bacteriocins. *Microb Ecol Health Dis* 2017;28:1–12.
- Baothman OA, Zamzami MA, Taher I, Abubaker J, Abu-Farha M. The role of gut microbiota in the development of obesity and diabetes. *Lipids Health Dis* 2016;15:108–16.
- Lu Y, Fan C, Li P, Lu Y, Chang X, Qi K. Short-chain fatty acids prevent high-fat-diet-induced obesity in mice by regulating G protein-coupled receptors and gut microbiota. *Sci Rep* 2016;6:37589–602.
- Wimalawansa SJ. Associations of vitamin D with insulin resistance, obesity, type 2 diabetes, and metabolic syndrome. *J Steroid Biochem Mol Biol* 2018;175:177–89.
- Dimitrov V, White JH. Vitamin D signaling in intestinal innate immunity and homeostasis. *Mol Cell Endocrinol* 2017;453:68–78.
- Clark A, Mach N. Role of vitamin D in the hygiene hypothesis: the interplay between vitamin D, vitamin D receptors, gut microbiota, and immune response. *Front Immunol* 2016;7:627–39.
- Al-shahwan M, Gacem SA, Shamseddin S, Sammour M. Vitamin D impact on human health and its relation with several diseases. *Int J Appl Pharm* 2018;10:60.
- Fish E, Beverstein G, Olson D, Reinhardt S, Garren M, Gould J. Vitamin D status of morbidly obese bariatric surgery patients. *J Surg Res* 2010;164:198–202.
- Khandalavala BN, Hibma PP, Fang X. Prevalence and persistence of vitamin D deficiency in biliopancreatic diversion patients: a retrospective study. *Obes Surg* 2010;20:881–4.
- Goldner WS, Stoner JA, Thompson J, Taylor K, Larson L, Erickson J, *et al.* Prevalence of vitamin D insufficiency and deficiency in morbidly obese patients: a comparison with non-obese controls. *Obes Surg* 2008;18:145–50.
- Duggan C, Dieu Tapsoba J de, Mason C, Imayama I, Korde L, Wang CY, *et al.* Effect of vitamin D3 supplementation in combination with weight loss on inflammatory biomarkers in postmenopausal women: a randomized controlled trial. *Cancer Prev Res* 2015;8:628–35.
- Shah Bidar S, Neyestani TR, Djazayeri A, Eshraghian MR, Houshiarrad A, Gharavi A, *et al.* Regular consumption of vitamin D-fortified yogurt drink (doogh) improved endothelial biomarkers in subjects with type 2 diabetes: a randomized, double-blind clinical trial. *BMC Med* 2011;9:125–35.
- Ranieri ML, Huck JR, Sonnen M, Barbano DM, Boor KJ. High temperature, short-time pasteurization temperatures inversely affect bacterial numbers during refrigerated storage of pasteurized fluid milk. *J Dairy Sci* 2009;92:4823–32.
- Huazano Garcia AGM. Metabolism of short-chain fatty acids in the colon and faeces of mice after supplementation of diets with agave fructans. W: Valenzuela Baez R. red. *Lipid Metabolism*; 2013.
- Cuesta Zuluaga J de la, Mueller N, Alvarez Quintero R, Velasquez Mejia E, Sierra J, Corrales Agudelo V, *et al.* Higher fecal short-chain fatty acid levels are associated with gut microbiome dysbiosis, obesity, hypertension and cardiometabolic disease risk factors. *Nutrients* 2018;11:51–67.
- Rosas Villegas A, Sanchez Tapia M, Avila Nava A, Ramirez V, Tovar A, Torres N. Differential effect of sucrose and fructose in combination with a high-fat diet on intestinal microbiota and kidney oxidative stress. *Nutrients* 2017;9:393–406.
- Fan Y, Futawaka K, Koyama R, Fukuda Y, Hayashi M, Imamoto M, *et al.* Vitamin D3/VDR resists diet-induced obesity by modulating UCP3 expression in muscles. *J Biomed Sci* 2016;23:56.

32. Lee E, Jung SR, Lee SY, Lee NK, Paik HD, Lim SI. *Lactobacillus plantarum* strain Ln4 attenuates diet-induced obesity, insulin resistance, and changes in hepatic mRNA levels associated with glucose and lipid metabolism. *Nutrients* 2018;10:643–58.
33. Rontoyanni V, Avila J, Kaul S, Wong R, Veeranki S. Association between obesity and serum 25(OH)D concentrations in older Mexican adults. *Nutrients* 2017;9:97.
34. Zakharova I, Klimov L, Kuryaninova V, Nikitina I, Malyavskaya S, Dolbnya S, et al. Vitamin D insufficiency in overweight and obese children and adolescents. *Front Endocrinol* 2019;10:103–16.
35. Hamid Mehmood ZTN, Papandreou D. An updated mini-review of vitamin D and obesity: adipogenesis and inflammation state. *Open Access Maced J Med Sci* 2016;4:526–32.
36. Dura Trave T, Gallinas Victoriano F, Chueca Guindulain MJ, Berrade Zubiri S, Urretavizcaya Martinez M, Ahmed Mohamed L. Assessment of vitamin D status and parathyroid hormone during a combined intervention for the treatment of childhood obesity. *Nutr Diabetes* 2019;9:18–26.
37. Fabersani E, Abejon Mukdsi MC, Ross R, Medina R, Gonzalez S, Gauffin Cano P. Specific strains of lactic acid bacteria differentially modulate the profile of adipokines *in vitro*. *Front Immunol* 2017;8:266–81.
38. Ellulu MS, Patimah I, Khazaai H, Rahmat A, Abed Y. Obesity and inflammation: the linking mechanism and the complications. *Arch Med Sci* 2017;4:851–63.
39. Xu G, Jiang J, Wang M, Li L, Su J, Ren X. Lactic acid reduces LPS-Induced TNF- α and IL-6 mRNA levels through decreasing I κ B α phosphorylation. *J Integr Agric* 2013;12:1073–8.
40. Karimi G, Sabran MR, Jamaluddin R, Parvaneh K, Mohtarrudin N, Ahmad Z, et al. The anti-obesity effects of *Lactobacillus casei* strain shirota versus orlistat on high fat diet-induced obese rats. *Food Nutr Res* 2015;59:29273–82.
41. Bikle DD. Vitamin D metabolism, mechanism of action, and clinical applications. *Chem Biol* 2014;21:319–29.
42. Wöbke TK, Sorg BL, Steinhilber D. Vitamin D in inflammatory diseases. *Front Physiol* 2014;5:244–64.
43. Mousa A, Misso M, Teede H, Scragg R, Courten B de. Effect of vitamin D supplementation on inflammation: protocol for a systematic review. *Br Med J Open* 2016;6:e010804–9.
44. Cortez M, Carmo LS, Rogero MM, Borelli P, Fock RA. A high-fat diet increases IL-1, IL-6, and TNF- α production by increasing NF- κ B and attenuating PPAR- γ expression in bone marrow mesenchymal stem cells. *Inflammation* 2013;36:379–86.
45. Besten G den, Eunen K van, Groen AK, Venema K, Reijngoud DJ, Bakker BM. The role of short-chain fatty acids in the interplay between diet, gut microbiota, and host energy metabolism. *J Lipid Res* 2013;54:2325–40.
46. Abdul Rahim MBH, Chilloux J, Martinez Gili L, Neves AL, Myridakis A, Gooderham N, et al. Diet-induced metabolic changes of the human gut microbiome: importance of short-chain fatty acids, methylamines and indoles. *Acta Diabetol* 2019;56:493–500.
47. Soldavini J, Kaunitz JD. Pathobiology and potential therapeutic value of intestinal short-chain fatty acids in gut inflammation and obesity. *Dig Dis Sci* 2013;58:2756–66.
48. Gabriel FC, Fantuzzi G. The association of short-chain fatty acids and leptin metabolism: a systematic review. *Nutr Res* 2019;72:18–35.
49. Kim KN, Yao Y, Ju SY. Short-chain fatty acids and fecal microbiota abundance in humans with obesity: a systematic review and meta-analysis. *Nutrients* 2019;11:2512–26.
50. Li M, Esch BCAM van, Henricks PAJ, Folkerts G, Garssen J. The anti-inflammatory effects of short-chain fatty acids on lipopolysaccharide-or tumor necrosis factor α -stimulated endothelial cells via activation of GPR41/43 and inhibition of HDACs. *Front Pharmacol* 2018;9:533–45.
51. McLoughlin RF, Berthon BS, Jensen ME, Baines KJ, Wood LG. Short-chain fatty acids, prebiotics, synbiotics, and systemic inflammation: a systematic review and meta-analysis. *Am J Clin Nutr* 2017;106:930–45.
52. Chambers ES, Preston T, Frost G, Morrison DJ. Role of gut microbiota-generated short-chain fatty acids in metabolic and cardiovascular health. *Curr Nutr Rep* 2018;7:198–206.
53. Frost G, Sleeth ML, Sahuri Arisoylu M, Lizarbe B, Cerdan S, Brody L, et al. The short-chain fatty acid acetate reduces appetite via a central homeostatic mechanism. *Nat Commun* 2014;5:3611–22.
54. Perry RJ, Peng L, Barry NA, Cline GW, Zhang D, Cardone RL, et al. Acetate mediates a microbiome-brain- β -cell axis to promote metabolic syndrome. *Nature* 2016;534:213–7.
55. Sindhu S, Thomas R, Shihab P, Sriraman D, Behbehani K, Ahmad R. Obesity is a positive modulator of IL-6R and IL-6 expression in the subcutaneous adipose tissue: significance for metabolic inflammation. *PLoS One* 2015;10:e0133494–511.
56. Makki K, Froguel P, Wolowczuk I. Adipose tissue in obesity-related inflammation and insulin resistance: cells, cytokines, and chemokines. *ISRN Inflammation* 2013;2013:1–12.

Fortified Dadih (Fermented Buffalo Milk) with Vitamin D3 Improves Interleukin-6 and Caecum Short Chain Fatty Acids on Diet-Induced Obese Rat

ORIGINALITY REPORT

23%

SIMILARITY INDEX

16%

INTERNET SOURCES

18%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|---|-----|
| 1 | Fernanda C. Gabriel, Giamila Fantuzzi. "The association of short-chain fatty acids and leptin metabolism: a systematic review", Nutrition Research, 2019
Publication | 1 % |
| 2 | orca.cardiff.ac.uk
Internet Source | 1 % |
| 3 | documents.mx
Internet Source | 1 % |
| 4 | Sasaki, Miyakawa, Watanabe, Nakayama, Lyu, Hama, Shibata. "Mice Microbiota Composition Changes by Inulin Feeding with a Long Fasting Period under a Two-Meals-Per-Day Schedule", Nutrients, 2019
Publication | 1 % |
| 5 | www.nature.com
Internet Source | 1 % |

6	"Sex and Gender Factors Affecting Metabolic Homeostasis, Diabetes and Obesity", Springer Nature, 2017 Publication	1 %
7	Park, Do-Young, Young-Tae Ahn, Se-Hoon Park, Chul-Sung Huh, Sae-Rom Yoo, Rina Yu, Mi-Kyung Sung, Robin A. McGregor, and Myung-Sook Choi. "Supplementation of Lactobacillus curvatus KY1032 in Diet-Induced Obese Mice Is Associated with Gut Microbial Changes and Reduction in Obesity", PLoS ONE, 2013. Publication	1 %
8	Submitted to Higher Education Commission Pakistan Student Paper	1 %
9	www.ajtmh.org Internet Source	1 %
10	Submitted to Queen Margaret University College, Edinburgh Student Paper	1 %
11	pubmed.ncbi.nlm.nih.gov Internet Source	<1 %
12	Kyu Nam Kim, Yao Yao, Sang Yhun Ju. "Short Chain Fatty Acids and Fecal Microbiota Abundance in Humans with Obesity: A	<1 %

Systematic Review and Meta-Analysis", Nutrients, 2019

Publication

13

Submitted to University of Wales Institute,
Cardiff

Student Paper

<1 %

14

Submitted to October University for Modern
Sciences and Arts (MSA)

Student Paper

<1 %

15

res.mdpi.com

Internet Source

<1 %

16

v3r.esp.org

Internet Source

<1 %

17

Ding, Cherlyn, John P. H. Wilding, and Chen
Bing. "1,25-dihydroxyvitamin D3 Protects
against Macrophage-Induced Activation of
NFkB and MAPK Signalling and Chemokine
Release in Human Adipocytes", PLoS ONE,
2013.

Publication

<1 %

18

Sardar Sindhu, Reeby Thomas, Puthiyaveetil
Shihab, Devarajan Sriraman, Kazem
Behbehani, Rasheed Ahmad. "Obesity Is a
Positive Modulator of IL-6R and IL-6
Expression in the Subcutaneous Adipose
Tissue: Significance for Metabolic
Inflammation", PLOS ONE, 2015

Publication

<1 %

19	USMAN PATO, MUHAMMAD ALI, ABDUL K. PARLINDUNGAN. "Taurocholate Deconjugation and Cholesterol Binding by Indigenous Dadih Lactic Acid Bacteria", HAYATI Journal of Biosciences, 2005 Publication	<1 %
20	www.biorxiv.org Internet Source	<1 %
21	www.pravara.com Internet Source	<1 %
22	www.veterinaryworld.org Internet Source	<1 %
23	Edward S. Chambers, Tom Preston, Gary Frost, Douglas J. Morrison. "Role of Gut Microbiota-Generated Short-Chain Fatty Acids in Metabolic and Cardiovascular Health", Current Nutrition Reports, 2018 Publication	<1 %
24	jeccr.biomedcentral.com Internet Source	<1 %
25	www.pubfacts.com Internet Source	<1 %
26	WWW.MDPI.COM Internet Source	<1 %
27	vitamindwiki.com Internet Source	<1 %

28

Aya Mousa, Marie Misso, Helena Teede, Robert Scragg, Barbora de Courten. "Effect of vitamin D supplementation on inflammation: protocol for a systematic review", BMJ Open, 2016

Publication

<1 %

29

Emanuel E. Canfora, Christina M. van der Beek, Johan W. E. Jocken, Gijs H. Goossens et al. "Colonic infusions of short-chain fatty acid mixtures promote energy metabolism in overweight/obese men: a randomized crossover trial", Scientific Reports, 2017

Publication

<1 %

30

Yueh-Ting Tsai, Po-Ching Cheng, Tzu-Ming Pan. "Anti-obesity effects of gut microbiota are associated with lactic acid bacteria", Applied Microbiology and Biotechnology, 2013

Publication

<1 %

31

balimedicaljournal.org

Internet Source

<1 %

32

José Antonio De Jesús Batún-Garrido, Marisol Salas-Magaña, Isela Esther Juárez-Rojop. "Association between leptin and IL-6 concentrations with cardiovascular risk in patients with rheumatoid arthritis", Clinical Rheumatology, 2017

Publication

<1 %

33	Nicole Mohajer, Chrislyn Y. Du, Christian Checkcinco, Bruce Blumberg. "Obesogens: How They Are Identified and Molecular Mechanisms Underlying Their Action", Frontiers in Endocrinology, 2021 Publication	<1 %
34	Wwww.dovepress.com Internet Source	<1 %
35	dokumen.pub Internet Source	<1 %
36	subjectguides.york.ac.uk Internet Source	<1 %
37	www.karger.com Internet Source	<1 %
38	www.spandidos-publications.com Internet Source	<1 %
39	www.wjgnet.com Internet Source	<1 %
40	Submitted to Arcadia University Student Paper	<1 %
41	Lee, Y.S.. "Effects of a Citrus depressa Hayata (shiikuwasa) extract on obesity in high-fat diet-induced obese mice", Phytomedicine, 20110615 Publication	<1 %

42	Lisa Dannenberg, Dorothee Zikeli, Marcel Benkhoff, Samantha Ahlbrecht, Malte Kelm, Bodo Levkau, Amin Polzin. "Targeting the human microbiome and its metabolite TMAO in cardiovascular prevention and therapy", Pharmacology & Therapeutics, 2020 Publication	<1 %
43	academic.oup.com Internet Source	<1 %
44	doiserbia.nb.rs Internet Source	<1 %
45	lipidworld.biomedcentral.com Internet Source	<1 %
46	annalsofintensivecare.springeropen.com Internet Source	<1 %
47	www.newswise.com Internet Source	<1 %
48	Bingwei Ma, Yao Chen, Chunjun Sheng, Peng Yang, Xingchun Wang, Shen Qu. "Low 25-hydroxyvitamin D is associated with arterial stiffness in Chinese with Type 2 diabetes mellitus", European Journal of Clinical Nutrition, 2021 Publication	<1 %
49	ephysician.ir Internet Source	<1 %

- | | | |
|-------|--|------|
| 50 | wjgnet.com
Internet Source | <1 % |
| <hr/> | | |
| 51 | www.plosone.org
Internet Source | <1 % |
| <hr/> | | |
| 52 | D. Sklan, M. Shelly, B. Makovsky, A. Geyra, E. Klipper, A. Friedman. "The effect of chronic feeding of diacetoxyscirpenol and T-2 toxin on performance, health, small intestinal physiology and antibody production in turkey poults", British Poultry Science, 2003
Publication | <1 % |
| <hr/> | | |
| 53 | George Mastorakos. "The Hypothalamic-Pituitary-Adrenal Axis in the Neuroendocrine Regulation of Food Intake and Obesity: The Role of Corticotropin Releasing Hormone", Nutritional Neuroscience, 10/1/2004
Publication | <1 % |
| <hr/> | | |
| 54 | Mia J Coleman, Luis M Espino, Hernan Lebensohn, Marija V Zimkute et al. "Individuals with Metabolic Syndrome show altered fecal lipidomic profiles with no signs of intestinal inflammation or increased intestinal permeability: a pilot study", Cold Spring Harbor Laboratory, 2022
Publication | <1 % |
| <hr/> | | |
| 55 | Teodoro Durá-Travé, Fidel Gallinas-Victoriano, Diego Mauricio Peñafiel-Freire, María | <1 % |

Urretavizcaya-Martinez et al.

"Hypovitaminosis D and Cardiometabolic Risk Factors in Adolescents with Severe Obesity",
Children, 2020

Publication

56	Www.lifeextension.com Internet Source	<1 %
57	Www.mdpi.com Internet Source	<1 %
58	bmccardiovascdisord.biomedcentral.com Internet Source	<1 %
59	docksci.com Internet Source	<1 %
60	tel.archives-ouvertes.fr Internet Source	<1 %
61	worldwidescience.org Internet Source	<1 %
62	wrap.warwick.ac.uk Internet Source	<1 %
63	www.advrehab.org Internet Source	<1 %
64	www.clonerresources.com Internet Source	<1 %
65	www.mdpi.com Internet Source	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Fortified Dadih (Fermented Buffalo Milk) with Vitamin D3 Improves Interleukin-6 and Caecum Short Chain Fatty Acids on Diet-Induced Obese Rat

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6