

**LEMBAR**  
**HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW**  
**KARYA ILMIAH : JURNAL ILMIAH**

Judul Artikel Ilmiah : The Effect of Tempeh Gembus Substitution on Protein Content, Calcium, Protein Digestibility and Organoleptic Quality of Meatballs

Penulis Artikel Ilmiah : Audrey Mutiara Setya Arini, Diana Nur Afifah, **Fillah Fithra Dieny**

Status Pengusul : Penulis Anggota

Identitas Jurnal Ilmiah :  
 a. Nama Jurnal : Current Research in Nutrition and Food Science  
 b. ISSN : 2347-467X  
 c. Nomor/Volume/Hal : 3/7/828-841  
 d. Edisi (bulan/tahun) : 2019  
 e. Penerbit : Enviro Research Publishers Bhopal India  
 f. Jumlah halaman : 14  
 g. DOI artikel (Jika ada) : <http://dx.doi.org/10.12944/CRNFSJ.7.3.22>  
 h. Alamat web Jurnal : <https://www.foodandnutritionjournal.org/>  
 i. Terindeks di : Google Scholar, Scopus  
 j. Turnitin : [https://doc-pak.undip.ac.id/2335/1/Turnitin\\_C4\\_Fillah.FD.pdf](https://doc-pak.undip.ac.id/2335/1/Turnitin_C4_Fillah.FD.pdf)

Kategori Publikasi Jurnal Ilmiah : ☒ Jurnal Ilmiah Internasional  
 (beri ✓ pada kategori yang tepat) ☐ Jurnal Ilmiah Nasional Terakreditasi  
☐ Jurnal Ilmiah Nasional tidak Terakreditasi

**I. Hasil Penilaian Peer Review :**

Komponen Yang Dinilai		Nilai Maksimal Karya Ilmiah (isikan di kolom yang sesuai)			Nilai Akhir Yang Diperoleh
		Internasional	Nasional Terakreditasi	Nasional tidak Terakreditasi	
		40			
a.	Kelengkapan dan Kesesuaian unsur isi artikel (10%)	4.00			4.00
b.	Ruang lingkup dan kedalaman pembahasan (30%)	12.00			11.00
c.	Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12.00			11.00
d.	Kelengkapan unsur dan kualitas penerbit (30%)	12.00			11.00
Nilai Total = (100%)		40.00			37.00
Nilai pengusul $(40\% \times 37)/2 =$					7.40
KOMENTAR/ULASAN PEER REVIEW					
Kelengkapan dan Kesesuaian Unsur		: Unsur lengkap dan terlah diuraikan pada tiap bagiannya			
Ruang Lingkup dan Kedalaman Pembahasan		: Penelitian tentang pengaruh tempe gembus memiliki hasil yang signifikan telah dibahas dengan mendalam dan didukung jurnal terkini			
Kecukupan & Kemutakhiran Data & Metodologi		: Metode yang digunakan tepat dan diuraikan dengan lengkap			
Kelengkapan Unsur dan Kualitas Penerbit		: Jurnal diterbitkan oleh Enviro Research Publisher Bhopal India. Jurnal terindeks SCOPUS Q3 dengan nilai SJR 0,22			

Semarang,  
 Penilai 1



dr. Achmad Zulfa Juniarto, M.Si.Med., Sp.And (K), M.M.R., Ph.D.  
 NIP 19700608 199702 1 001  
 Unit kerja : Fakultas Kedokteran  
 Bidang Ilmu : Ilmu Kedokteran  
 Jabatan/Pangkat : Lektor Kepala / Penata Tk I, III/d

**LEMBAR**  
**HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW**  
**KARYA ILMIAH : JURNAL ILMIAH**

Judul Artikel Ilmiah : The Effect of Tempeh Gembus Substitution on Protein Content, Calcium, Protein Digestibility and Organoleptic Quality of Meatballs

Penulis Artikel Ilmiah : Audrey Mutiara Setya Arini, Diana Nur Afifah, **Fillah Fithra Dieny**

Status Pengusul : Penulis Anggota

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : Current Research in Nutrition and Food Science
- b. ISSN : 2347-467X
- c. Nomor/Volume/Ha : 3/7/828-841
- d. Edisi (bulan/tahun) : 2019
- e. Penerbit : Enviro Research Publishers Bhopal India
- f. Jumlah halaman : 14
- g. DOI artikel (Jika ada) : <http://dx.doi.org/10.12944/CRNFSJ.7.3.22>
- h. Alamat web Jurnal : <https://www.foodandnutritionjournal.org/>
- i. Terindeks di : Google Scholar, Scopus
- j. Turnitin : [https://doc-pak.undip.ac.id/2335/1/Turnitin\\_C4\\_Fillah.FD.pdf](https://doc-pak.undip.ac.id/2335/1/Turnitin_C4_Fillah.FD.pdf)

Kategori Publikasi Jurnal Ilmiah : ☒ Jurnal Ilmiah Internasional  
 (beri ✓ pada kategori yang tepat) ☐ Jurnal Ilmiah Nasional Terakreditasi  
☐ Jurnal Ilmiah Nasional tidak Terakreditasi

**I. Hasil Penilaian Peer Review :**

Komponen Yang Dinilai	Nilai Maksimal Karya Ilmiah (isikan di kolom yang sesuai)			Nilai Akhir Yang Diperoleh
	Internasional	Nasional Terakreditasi	Nasional tidak Terakreditasi	
	40			
a. Kelengkapan dan Kesesuaian unsur isi artikel (10%)	4.00			3.00
b. Ruang lingkup dan kedalaman pembahasan (30%)	12.00			11.00
c. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12.00			11.00
d. Kelengkapan unsur dan kualitas penerbit (30%)	12.00			10.00
Nilai Total = (100%)	40.00			35.00

Nilai pengusul (40% x 35)/2 = 7.00

**KOMENTAR/ULASAN PEER REVIEW**

Kelengkapan dan Kesesuaian Unsur	: Jurnal internasional bereputasi terindeks di SCOPUS (Q3), unsur lengkap dari abstract, introduction, methods, discussion, acknowledgement, dan references.
Ruang Lingkup dan Kedalaman Pembahasan	: Ruang lingkup penelitian pangan fungsional dengan memodifikasi bakso daging dengan substitusi tempe gembus. Pembahasan mendalam dengan sebagian besar referensi adalah pustaka primer.
Kecukupan & Kemutakhiran Data & Metodologi	: Penelitian eksperimental di laboratorium hingga uji organoleptik yang umum digunakan.
Kelengkapan Unsur dan Kualitas Penerbit	: Terbitan lengkap memuat vol, no, tahun dan daftar isi serta terindeks di SCOPUS (Q3).

Semarang,  
 Penilai 2



Dr. Diana Nur Afifah, S.TP., M.Si.  
 NIP 198007312008012011

Unit kerja : Fakultas Kedokteran

Bidang Ilmu : Ilmu Gizi

Jabatan/Pangkat : Lektor Kepala / Penata Tk I, III/d

ISSN: 2347-467X

Volume 9 Number 3 Decembert 2021



## Current Research in Nutrition and Food Science



Published by:  
Enviro Research Publishers  
[www.foodandnutritionjournal.org](http://www.foodandnutritionjournal.org)



## Editorial Board

[Home](#) » [About](#) » [Editorial Board](#)

### Chief Editor



**Dr. Shih-Min Hsia**

Distinguished Professor,  
School of Nutrition and Health Sciences,  
Taipei Medical University,  
Taiwan.  
Email: [bryanhhsia@tmu.edu.tw](mailto:bryanhhsia@tmu.edu.tw)  
Scopus ID: 13409857600

### Co-Editor in Chief



**Dr. Ravi Shukla, Ph.D**

Associate Professor,  
STEM- School of Science,  
RMIT University,  
Australia.  
Email: [ravi.shukla@rmit.edu.au](mailto:ravi.shukla@rmit.edu.au)  
Scopus ID: 9437484300

### Managing Editors

**Dr. Neha Sanwalka**  
Founder & Director Nutri Canvas,  
Mumbai, MH, India.  
Email: [neha.sanwalka@gmail.com](mailto:neha.sanwalka@gmail.com)  
Scopus ID: 22954784500

**Dr. Adele Papetti**  
Assistant Professor  
Department of Drug Sciences  
Universita degli Studi di Pavia,  
Pavia, Italy.  
Email: [adele.papetti@unipv.it](mailto:adele.papetti@unipv.it)  
Scopus ID: 6603497119

**Prof. Y. Kourkoutas**  
Associate Professor,  
Department of Molecular Biology & Genetics  
Democritus University of Thrace  
Alexandroupolis, Greece.  
Email: [ikourkou@mbg.duth.gr](mailto:ikourkou@mbg.duth.gr)  
Scopus ID: 55919995700

**Dr. Amalia Tsiami**  
Associate Professor,  
London School of Hospitality and Tourism  
University of West London,  
London, United Kingdom.  
Email: [Amalia.Tsiami@uwl.ac.uk](mailto:Amalia.Tsiami@uwl.ac.uk)  
Scopus ID: 6506464729

### Associate Editors

**Prof. R. Jeewon**  
Associate Professor

**Prof. Reema F. Tayyem**  
Department of Nutrition and Food Technology,

### Scopus Journal Metrics

**CiteScore 2020: 1.3**

**CiteScore Details**

**SJR 2020**



This journal is a member of, and subscribes to  
the principles of, the Committee on  
Publication Ethics (COPE)

### Journal is Indexed in:

**CABELLS**

SCHOLARLY ANALYTICS

**Cabells Whitelist**

Follow us on:



[Submit Manuscript Online](#)

[Join our team of Reviewers](#)

[Journal is indexed in...](#)

## Most Popular Articles

Low Alkaline Phosphatase (ALP) In Adult  
Population an...

(174,636)

Colostrum - its Composition, Benefits as  
a...

(121,924)

Importance of Exclusive Breastfeeding  
and Complementary...

(97,557)

Processing and Nutritional Composition of  
Rice Bran

(40,767)

Processing and Nutritive Value of Mango  
Seed Kernel Flour

(38,318)

Dept. of Health Sciences,  
The University of Mauritius,  
Moka, Mauritius.  
Email: r.jeewon@uom.ac.mu  
Scopus ID: 6602641191

Faculty of Agriculture,  
The University of Jordan,  
Amman, Jordan.  
Email: r\_tayyem@yahoo.com  
Scopus ID: 22959129200

### Statistical Editors

**Dr. Ravindra Boojhawon**  
Associate Professor  
The University of Mauritius,  
Moka, Mauritius.  
Email: r.boojhawon@uom.ac.mu  
Scopus ID: 6507224879

### Editorial cum Advisory Board

**Dr. Vikas Kumar**  
Assistant Food Technologist,  
Department of Food Science and  
Technology, Punjab Agricultural University,  
Ludhiana, Punjab, India.  
Email: vkchoprafst@rediffmail.com  
Scopus ID: 57193417700

**Dr. Dan-Cristian Vodnar**  
Faculty of Food Science and Technology,  
University of Agricultural Science and  
Veterinary Medicine,  
Cluj-Napoca, Romania.  
Email: dan.vodnar@usamvcluj.ro  
Scopus ID: 35604846100

**Prof. Shwu-Huey Sherry Yang**  
School of Nutrition and Health Sciences  
Taipei Medical University,  
Taipei, Taiwan.  
Email: sherry@tmu.edu.tw  
Scopus ID: 8676122700

**Prof. Sitthipong Nalinanon**  
Associate Dean  
Faculty of Agro-Industry, King Mongkut's  
Institute of Technology Ladkrabang  
Ladkrabang,  
Bangkok, Thailand.  
Email: sitthipong.na@kmitl.ac.th  
Scopus ID: 16245779000

**Dr. Maria Papageorgiou**  
Associate Professor  
Department of Food Technology  
Alexander Technological Educational  
Institute of Thessaloniki (ATEITH)  
Thessaloniki, Greece.  
Email: mariapapage@food.teithe.gr  
Scopus ID: 8645438400

**Dr. Pierluigi Plastina**  
Department of Pharmacy,  
Health and Nutrition,  
University of Calabria,  
Cosenza, Italy.  
Email: pierluigi.plastina@unical.it  
Scopus ID: 8360032300

**Dr. Miguel Herrero**  
Institute of Food Science Research (CIAL-  
CSIC), Laboratory of Foodomics,  
Madrid, Spain.  
Email: m.herrero@csic.es  
Scopus ID: 7102420113

**Prof. Giovanni Leone Zabot**  
Food Engineer  
Federal University of Santa Maria (UFSM)  
Santa Maria, Brazil.  
Email: giovani.zabot@gmail.com  
Scopus ID: 26639885600

**Dr. Ardiansyah**  
Associate Professor and Head  
Department of Food Technology,  
Universitas Bakrie,  
Jakarta, Indonesia.  
Email: ardiansyah@bakrie.ac.id  
Scopus ID: 12798333700

**Prof. Aly El Sheikh**  
Visiting Professor, School of Nutrition  
Sciences, University of Ottawa, Canada;  
Full Professor at College of Bioscience and  
Bioengineering, Jiangxi Agricultural  
University, China.  
Email: elsheikha\_aly@yahoo.com  
Scopus ID: 35174115900

**Prof. Ana Laura Isabel De la Garza  
Hernández**  
School of Public Health and Nutrition,  
Autonomous University of Nuevo Leon,  
Nuevo Leon, Mexico.  
Email: ana.dlgarzah@uanl.mx  
Scopus ID: 57195297780

**Prof. Victor J. Temple**  
School of Medicine and Health Sciences,  
University of Papua New Guinea,  
Port Moresby,  
Papua New Guinea.  
Email: templevj@upng.ac.pg  
Scopus ID: 57194389852

**Prof. Dr. Dariusz Dziki**  
Professor  
Department of Thermal Technology and  
Food Process Engineering,  
Faculty of Production Engineering,  
University of Life Sciences in Lublin, Lublin,

**Dr. Keshab Chandra Mondal**  
Professor  
Department of Microbiology,  
Vidyasagar University Midnapur,  
West Bengal, India.



Poland.  
Email: [dariusz.dziki@up.lublin.pl](mailto:dariusz.dziki@up.lublin.pl)  
Scopus ID: 24314765900

**Dr. Mattia Di Nunzio**  
Interdepartmental Centre for Agri-Food  
Industrial Research, Alma Mater Studiorum  
Universita di Bologna, Bologna, Italy.  
Email: [mattia.dinunzio@unibo.it](mailto:mattia.dinunzio@unibo.it)  
Scopus ID: 23093741900

**Dr. Pasquale Russo**  
Department of Science of Agriculture,  
Food and Environment, University of  
Foggia, Foggia, Italy.  
Email: [pasquale.russo@unifg.it](mailto:pasquale.russo@unifg.it)  
Scopus ID: 35726052700

**Prof. Suhad Maatoug Bahijri**  
Professor  
Clinical Biochemistry Department,  
Faculty of Medicine, King Abdulaziz  
University,  
Jeddah, Saudi Arabia.  
Email: [sbahijri@gmail.com](mailto:sbahijri@gmail.com)  
Scopus ID: 6603404358

**Dr. Nurul Huda**  
Associate Professor  
Faculty of Food Science and Nutrition,  
Universiti Malaysia Sabah,  
Sabah, Malaysia.  
Email: [drnurulhuda@yahoo.com](mailto:drnurulhuda@yahoo.com)  
Scopus ID: 6701695514

**Prof. Dr. Harun AKSU**  
Faculty of Veterinary Faculty  
Food Hygiene and Technology  
Department, Istanbul University,  
Avcılar-Istanbul, Turkey.  
Email: [h.aksu@istanbul.edu.tr](mailto:h.aksu@istanbul.edu.tr)  
Scopus ID: 56235721300

**Dr. Norazmir Md Nor**  
Senior Lecturer  
Faculty of Health Sciences, Universiti  
Teknologi,  
Selangor, Malaysia.  
Email: [azmir2790@gmail.com](mailto:azmir2790@gmail.com)  
Scopus ID: 54417702600

**Dr. Prasad S. Variyar**  
Professor, Homi Bhabha National Institute  
Scientific Officer (H) & Head,  
Food Flavor and Aroma Chemistry Section  
Food Technology Division  
Bhabha Atomic Research centre  
Mumbai, Maharashtra, India.  
Email: [prasadpsv@rediffmail.com](mailto:prasadpsv@rediffmail.com)  
Scopus ID: 6603637229

**Prof Quazi Syed Zahir Uddin**  
Director Research & Development  
Associate Dean (Global Health)  
Professor, Community Medicine  
J N Medical College, Datta Meghe Institute  
of Medical Sciences

Email: [kcmondal@mail.vidyasagar.ac.in](mailto:kcmondal@mail.vidyasagar.ac.in)  
Scopus ID: 7004865977

**Dr. Krešimir Mastanjević**  
Assistant Professor  
Faculty of Food Technology,  
University of Osijek,  
Osijek, Croatia.  
Email: [kmastanj@ptfos.hr](mailto:kmastanj@ptfos.hr)  
Scopus ID: 26655948300

**Prof. Gabriel O. Adegoke**  
Department of Food Technology,  
University of Ibadan, Ibadan.  
Nigeria.  
Email: [goadegoke@hotmail.com](mailto:goadegoke@hotmail.com)  
Scopus ID: 7003766493

**Prof. Jiwan S. Sidhu**  
Director,  
Department of Food Science and Nutrition,  
College of Life Sciences,  
Kuwait University,  
Safat, Kuwait.  
Email: [dr.jiwan.sidhu@gmail.com](mailto:dr.jiwan.sidhu@gmail.com)  
Scopus ID: 7004614004

**Dr Jose Manuel Lorenzo Rodriguez**  
Head of Research  
Department of Food Science and  
Technology, Centro Tecnológico de la Carne  
de Galicia, Parque Tecnológico de Galicia,  
Spain.  
Email: [jmlorenzo@ceteca.net](mailto:jmlorenzo@ceteca.net)  
Scopus ID: 57195195602

**Prof. Chunpeng Wan**  
Jiangxi Key Laboratory for Postharvest  
Technology and Nondestructive Testing of  
Fruits & Vegetables,  
Jiangxi Agricultural University  
China  
Email: [chunpengwan@jxau.edu.cn](mailto:chunpengwan@jxau.edu.cn)  
Scopus ID: 36731790300

**B.N. Dar**  
UGC Research Awardee  
Assistant Professor  
Department of Food Technology  
Islamic University of Science & Technology,  
Jammu & Kashmir, India.  
Email: [darnabi@gmail.com](mailto:darnabi@gmail.com)  
Scopus ID: 56068490400

**Dr. Nikhil Kumar Mahnot**  
Assistant Professor  
Department of Food Technology  
Rajiv Gandhi University (A central  
University)  
Itanagar, India.  
Email: [nikhil.mahnot@gmail.com](mailto:nikhil.mahnot@gmail.com)  
Scopus ID: 56254346400

**Dr K. V Sunooj**  
Assistant professor and President AFST(I)  
Department of Food Science & Technology,  
Pondicherry University  
Union territory of Pondicherry,  
India.

Wardha, Maharashtra, India.  
Email: [zahirquazi@gmail.com](mailto:zahirquazi@gmail.com)  
Scopus ID: 36337182700

**Dr. Jeyan A. Moses**  
Assistant Professor and In-charge  
Computational Modeling and Nanoscale  
Processing Unit  
Indian Institute of Food Processing  
Technology (IIFPT)  
Ministry of Food Processing Industries |  
Government of India  
Thanjavur, Tamil Nadu, India.  
Email: [moses.ja@iifpt.edu.in](mailto:moses.ja@iifpt.edu.in)  
Scopus ID: 22981136400

Email: [sunoojpufst786@gmail.com](mailto:sunoojpufst786@gmail.com)  
Scopus ID: 25822810100

**Dr. Netsanet Fentahun**  
Associate Professor of Public Health  
Nutrition  
Nutrition and Qualitative Research  
Consultant  
Founder and Executive Director for Ethio-  
Star Consulting PLC  
Department of Nutrition and Dietetics  
School of Public Health  
College of Medicine and Health Sciences  
Bahir Dar University -Ethiopia.  
Email: [netsanet\\_fentahun@yahoo.com](mailto:netsanet_fentahun@yahoo.com)  
Scopus ID: 56394665400

## Links

[About](#)

[Archives](#)

[Article Processing Charges](#)

[Coming Issue](#)

[Contact](#)

[Current Issue](#)

[Order Print Issue](#)

[Social Media](#)

[Submission](#)

## Contact Us

Your Name (required)

Your Email (required)

Your Message

**C 5 D Y**

Type the above text in box below (Case sensitive)

Complte the quiz  
2+2=?

**Send**

Follow us on:



## License



This work is licensed under a Creative Commons Attribution 4.0 International License.

## Recent Articles

Synbiotic (*L. plantarum* Dad-13 and Fructo-oligosaccharide) Powder on Gut Microbiota (*L. plantarum*, *Bifidobacterium* and *Enterobacteriaceae*) on Stunting Children in Yogyakarta, Indonesia

Spirulina (*Arthrospira* spp) as a Complementary COVID-19 Response Option: Early Evidence of Promise

Assessment of Fast and Junk Food Consumption and Addiction among Saudi Population. National Descriptive Study 2020.


## Volume 7, Number 3 // Volume 7, Number 3

[Home](#) » [Volume 7, Number 3](#)

**25 Dec**

Volume 7, Number 3 | Comments Off

### Study on the Effect of Prawn (*Machrobrachium rosenbergii*) Chitosan Coating on Peeled Shallot (*Allium ascalonicum*)



Introduction Shallot (*Allium ascalonicum*) is an important culinary component in Indian cuisine and it is typically cultivated thrice a year. Intended use of Shallot is because of its flavor enhancing and taste adding feature. It contains high nutrients also serve as an appetizer, food digestive agent and immunity booster. Shallots are added in commercial and [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | Comments Off

### The Impact of Front-of-Package Traffic Light (Foptl) in the Senior High School Students' Nutrition Labels Comprehension



Introduction The number of children affected by the obesity epidemic has increased in the last few years.<sup>1</sup> In 2013, the rates of obesity among Indonesian teenagers aged 13-15 and 16-18 years old were 2.5% and 1.6%, respectively.<sup>2</sup> It has long been observed that overweight teenagers will continue to carry increased weight when they enter adulthood. [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | Comments Off

### Prevalence Rate, Antibiotic Resistance and Biotyping of Thermotolerant *Campylobacter* Isolated from Poultry Products Vended in Wasit Markets



Scopus Journal Metrics

CiteScore 2020: 1.3

CiteScore Details

**SJR 2020**



JM14653

This journal is a member of, and subscribes to the principles of, the Committee on Publication Ethics (COPE)

Journal is Indexed in:

**CABELLS**

SCHOLARLY ANALYTICS

Cabells Whitelist

Follow us on:



[Submit Manuscript Online](#)

[Join our team of Reviewers](#)

[Journal is indexed in...](#)

## Most Popular Articles

Low Alkaline Phosphatase (ALP) In Adult Population an...

(174,636)

Colostrum - its Composition, Benefits as a...

(121,924)

Importance of Exclusive Breastfeeding and Complementary...

(97,557)

Processing and Nutritional Composition of Rice Bran

(40,767)

Processing and Nutritive Value of Mango Seed Kernel Flour

(38,318)





Introduction *Campylobacter* is considered as a principal cause of gastroenteritis in humans responsible for approximately 166 million diarrheal cases and 37,600 deaths per year globally.<sup>1</sup> In addition to gastrointestinal infections, campylobacteriosis is the best predisposing factor of Guillain-Barré syndrome, a serious demyelinating neuropathy, which roughly occurs in 3/10 000 cases of *Campylobacter* infections.<sup>2</sup> Thermotolerant *Campylobacter* [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Quality Changes of Chilled Green Mussel (*Perna Viridis*) Pre-Treated with Organic Acids and Sodium Tripolyphosphate



Introduction Green mussel (*Perna viridis*) is a commercially significant commodity in the Philippines.<sup>1</sup> Its latest production had amounted to 18,800 metric tons and considered as top-six among commercially aquaculture species in our country. The major mussel producing provinces include Cavite, Capiz, Samar, Bataan, and Negros Occidental.<sup>2</sup> The species is favored as part of the human [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Preservation of the Polyphenolic Content and Antioxidant Properties of *Rubus Glaucus* Benth



Introduction The Andes berry (*Rubus glaucus* Benth) is from America high tropical zone; it was growth in Colombia, Ecuador, Panamá, Guatemala, Honduras, México and Salvador. This fruit plays an important role in the human diet due to its high content of antioxidants, nutrients, and proteins, all of which are key to a balanced diet.<sup>1</sup> It [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Toxic Effects of Food Colorants Erythrosine and Tartrazine on Zebrafish Embryo Development



**Introduction** Food colorants are natural or synthetic substances used to artificially color and enhance the appearance of food products. Natural food dyes though are safe to use but, do not provide a bright Thus, artificial food colorants (AFCs) such as erythrosine which provide bright color are popular in food industry.<sup>1</sup> Most of the AFCs are [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Study of Factors Affecting Exclusive Breastfeeding and Early Introduction of Complementary Food to Infants in the Aqaba Region of Jordan



**Introduction** Good nutrition in early life is essential to lifelong health; therefore, receiving exclusive breastfeeding (EBF) is very important to protect against diseases and prevent morbidity and mortality of infants and young children, including malnutrition and obesity.<sup>1</sup> For this reason, countries should have solutions to reach the target of 50% EBF through the first 6 [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Mep3s – The Nutri Cookie for Children with Renal Diseases



**Introduction** Adequate nutrition is important for growth and development in children. In general, children are fussy eaters. They prefer and consume more of convenient foods which are mostly processed and ready to eat. The products prepared commercially are tasty but lack almost all important nutrients, thus meeting their daily nutritional requirements becomes difficult. The scenario [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Development and Nutritional Evaluation of Multigrain Gluten Free Cookies and Pasta Products





Introduction Celiac disease is considered as chronic gut disorder that can happen in genetically predisposed people where the consumption of gluten in diet can damage the lining of the small intestine. It is encouraged by a protein known as “gluten” existing in cereals such as wheat, barley, rye etc. It disturbs approximately one per cent [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## The Effect of Tempeh Gembus Substitution on Protein Content, Calcium, Protein Digestibility and Organoleptic Quality of Meatballs



Introduction Osteoporosis is a bone metabolic disease characterized by a decrease in bone mass due to reduced bone matrix and minerals accompanied by microarchitecture damage from bone tissue, with a consequent decrease in bone strength, resulting in a tendency for bones to break easily.<sup>1</sup> Osteoporosis is a disease that is classified as a silent disease [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Identification of Single Nucleotide Polymorphisms in Selenium-Glutathione Peroxidase (GPX1) Gene



Introduction GPX1 was the first GPX to be discovered<sup>1</sup> and is also known as classical GPX, cytosolic GPX or GPX1. It has a wide distribution in animal cells and is present in the cytosol.<sup>1,2,3</sup> It is reported that high levels of GPX present in different tissues such as RBC, placenta, lung, liver and kidney which [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Whey/Broken Chickpea Extract for Manufacture of Probiotic Frozen Yoghurt



Introduction The trend towards the use of dairy by-products such as whey, skim milk, butter milk and permeate and some residues of agricultural crops such as broken chickpea seeds, wheat germ, wheat bran, rice bran and etc, as non-dairy by-products in the manufacture of functional dairy foods is increasing worldwide. Whey is one of the [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Waste Pomelo (Citrus Maxima) Peels – A Natural Source of Antioxidant and Its Utilization in Peanut Oil for Suppressing the Development of Rancidity



Introduction The oxidative rancidity of fats and oils in food is one of the serious problems of the modern society. Oxidative rancidity is responsible for off-flavor and odor, decreasing storage life, altering texture, nutritional quality also decreased due to the formation of secondary compounds which are potentially toxic.<sup>1</sup> The need of addition of antioxidants is [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Soft Drink Consumption and its Influence on BMI and Academic Performance among Selected School Children in Dhaka City of Bangladesh



Introduction Soft drink consumption is increasing among the children and has become a controversial public health concern. Soft drink is a nonalcoholic drink, especially one that is carbonated<sup>1</sup> consisting of a unique chemical blend of sugar or artificial sweetening agents, caffeine, colour, flavour and preservatives in water, injected with carbon-dioxide gas. All of these ingredients [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Technological Quality of New Croatian Barley Varieties Intended for Whisky Malt Production



Introduction Barley is one of the earliest cultivated cereals and besides the use as human and animal feed it was utilised as a source of fermentable extract for the production of beer and spirits. There are over 300.000 barley varieties that are generally categorized as two-row and six-row.<sup>1</sup> Today's barley selection has been significantly advanced [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Ethanol Extracts of Ficus carica Fruit and Leaf Normalize High Serum Lipid Profile, TNF- $\alpha$ , and MDA due to High Fat Diet in Sprague Dawley Rat



Introduction A diet called Western pattern diet (WPD) is characterized by a high intake of red meat, processed meat, pre-packed foods, butter, fried foods, potatoes, high-fructose corn syrup, and high-fat dairy products.<sup>1</sup> This dietary pattern is widespread in some Asian countries including Indonesia and it can result in dyslipidemia.<sup>2</sup> Dyslipidemia is characterized by abnormal levels [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Physico-Chemical, Textural and Sensory Characteristics of Wheat Flour Biscuits Supplemented with Different Levels of Whey Protein Concentrate



Introduction Bakery products have become very popular worldwide because of their unique taste and easy availability at a reasonable cost. Biscuits are one of the most popular bakery products due to their ready to eat nature, easy handling for convenience, and possess long shelf life.<sup>1</sup> In addition to that; most of the bakery products can [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)



## Effect of Intrinsic and Extrinsic Factors on the Storage Stability of Sardine Oil



**Introduction** The major sources of dietary lipids are edible oils, which include both vegetable and fish oils. Edible oils supply calories, essential fats and helps in absorbing fat-soluble vitamins. Type of lipid consumed plays a major role in the health of an individual. Oil sardines are one of the richest and cheapest sources of n-3 [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | Comments Off

## Congruence with Health by Divergence in Uses of Water: An Ayurvedic Notion



**Introduction** Water has been deemed as the pedestal of life since ages and considered as a single most imperative nutrient of our body. It has a preeminent role in subsistence and decadence of any living being. It is improbable to live more than 8-21 days without food and water,<sup>1</sup> although in starvation, the stored fuels [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | Comments Off

## Is Soft Drink Consumption Linked to Higher Body Mass Index and Energy Intake Among Adults in Mauritius?



**Introduction** The addition of sugar is not nutritionally necessary in an individual's diet as most of our food consist of natural sugars that provide us with sufficient energy. Nevertheless, for the last few decades, there has been an increase in the consumption of sugary products. Soft drinks (SD) have become an ultimate liquid to quench [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | Comments Off

## Modified Atmospheric Packaging (MAP) of Trichosanthes Dioica (Parwal) Sweet and Effect of Storage Temperature on the Physicochemical, Microbial and Sensory Characteristics



**Introduction** From time immemorial ethnic food such as traditional sweets has played a significant role in our society and religion. Parwal or pointed gourd (*Trichosanthes dioica*) sweet is a type of indigenous dairy product prepared by filling of khoa in previously boiled parwal in sugar syrup.<sup>1-3</sup> Parwal is a member of cucurbitaceae family and perennial, [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## The Spectrum of Disordered Eating Attitudes Among Female University Students: A Cross-Sectional Study



**Introduction** Eating disorders (ED) rank high among the most prevalent psychiatric problems, adolescent girls are experiencing today, where they develop incorrect attitudes to their body weight and shape, as well as entertain distorted perceptions of body shape.<sup>1-3</sup> The core characteristics of EDs include feeding and eating disturbances and a distorted understanding of body image, causing [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Biochemical Composition and Sensory Evaluation of Desert Date Flowers (*Balanites aegyptiaca* Del) Infusion



**Introduction** *Balanites aegyptiaca* Del can reach a height of 6 to 9 m, it is ramous and thorny tree, also known as desert date in English.<sup>1</sup> It belongs to the family of Zygophyllaceae. The tree is one of the most common but neglected wild plant species of the dry land areas. The tree grown in [...]

[Read more...](#)

## Proportion and Associated Factors of Vitamin B12 Deficiency – A Retrospective Laboratory-Based Study



Introduction Vitamin B12 (B12) is a water-soluble vitamin that belongs to the set of “corrinoids” which contains a corrin macrocycle in the structure.<sup>1,2</sup> B12 acts as a cofactor in one-carbon metabolism through methylation and molecular rearrangement.<sup>3</sup> It is a vital micronutrient essential for the optimum hemopoetic, neurologic and cardio-metabolic function.<sup>4, 5</sup> Deficiency of B12 and [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Quantitative Risk Assessment of Acrylamide in Indonesian Deep Fried Fritters as Street Food Products



Introduction Toxicological studies have shown that acrylamide is neurotoxicant, reproductive toxicant, genotoxic and carcinogen in animals.<sup>1</sup> In fact, it has been categorized as “probable carcinogen to human” (category 2A) by International Agency for Research on Cancer (IARC) in 1994. Although the toxicity of acrylamide has been acknowledged long ago, it was not until the finding [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Storage Stability of Spray Dried Nigella Sativa (Ranunculaceae Family) Instant Beverage Powder: Effect of Carrier Agents on the Physicochemical, Phenolic Compounds and Antioxidant Properties



Introduction Spices serve many functions and one of the major functions of spices is its use in the food industry to infuse flavour and increased the aroma, texture and colour of food. Spices possess antimicrobial, antioxidant and protective functions for human health.<sup>1</sup> They contain various compounds such as protein, fibre, vitamins, minerals and volatile compounds. [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Copper Deficiency Anemia in Older Patients with Swallowing Disorders: A Case Report



**Introduction** It is not uncommon for swallowing function to deteriorate due to causes such as old age, dementia, and cerebrovascular disease sequelae.<sup>1,2</sup> Changing the consistency of food making them safer to swallow, such as introducing easy-to-swallow food, termed dysphagia diet, are recommended for such patients, in order to prevent aspiration.<sup>3,4</sup> Dysphagia diet is a general [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Application of Edible and Biodegradable Starch-Based Films in Food Packaging: A Systematic Review and Meta-Analysis



**Introduction** Today adequate and adequate food supplies have become one of the most important concerns of the authorities and community elites, in order to overcome many efforts in various areas, such as increasing production, increasing the amount of time consuming, improving Storage and storage practices, and developing methods for protecting them against destructive factors such [...]

[Read more...](#)

**25 Dec**

Volume 7, Number 3 | [Comments Off](#)

## Determinants of Household Food Insecurity and Its Association with Child Malnutrition in Sub-Saharan Africa: A Review of the Literature



Introduction Household food insecurity is increasingly known to be a global health problem, especially in Sub-Saharan Africa.<sup>1</sup> The Sustainable Development Goals (SDGs) have emphasised the urgent need to tackle food insecurity in achieving human rights by the year 2030.<sup>2</sup> The first goal is to end ‘poverty in all its forms everywhere’, and the second is [...]

[Read more...](#)

25 Dec

Volume 7, Number 3 | Comments Off

## Food for Thought: The Digital Disruption and the Future of Food Production



The world population will increase to more than 9 billion people by 2050.<sup>1-2</sup> In this context, changes might occur at several points of the food supply and value chains, where the sustainable production of raw materials and commodities, waste reduction, adaptation to climate change, and food security are top of the list.<sup>1</sup> The use of [...]

[Read more...](#)

### Links

[About](#)

[Archives](#)

[Article Processing Charges](#)

[Coming Issue](#)

[Contact](#)

[Current Issue](#)

[Order Print Issue](#)

[Social Media](#)

[Submission](#)

### Contact Us

Your Name (required)

Your Email (required)

Your Message

### LCDS

Type the above text in box below (Case sensitive)

Complete the quiz

3+3=?

**Send**

Follow us on:



### License



This work is licensed under a Creative Commons Attribution 4.0 International License.

### Recent Articles

Synbiotic (*L. plantarum* Dad-13 and Fructo-oligosaccharide) Powder on Gut Microbiota (*L. plantarum*, *Bifidobacterium* and *Enterobacteriaceae*) on Stunting Children in Yogyakarta, Indonesia

Spirulina (*Arthrospira* spp) as a Complementary COVID-19 Response Option: Early Evidence of Promise

Assessment of Fast and Junk Food Consumption and Addiction among Saudi Population. National Descriptive Study 2020.







## Source details

### Current Research in Nutrition and Food Science

Open Access ⓘ

Scopus coverage years: from 2013 to 2019

Publisher: Enviro Research Publishers

ISSN: 2347-467X E-ISSN: 2322-0007

Subject area: [Agricultural and Biological Sciences: Food Science](#) [Medicine: Medicine \(miscellaneous\)](#)

[View all documents >](#)

[Set document alert](#)

[Save to source list](#)

CiteScore 2018

0.78



SJR 2018

0.230



SNIP 2018

0.397



[CiteScore](#) [CiteScore rank & trend](#) [CiteScore presets](#) [Scopus content coverage](#)

CiteScore 2018 ▾

Calculated using data from 30 April, 2019

CiteScore rank ⓘ

$$0.78 = \frac{\text{Citation Count 2018}}{\text{Documents 2015 - 2017}^*} = \frac{102 \text{ Citations} >}{130 \text{ Documents} >}$$

\*CiteScore includes all available document types

[View CiteScore methodology >](#)

[CiteScore FAQ >](#)

Category	Rank	Percentile
Agricultural and Biological Sciences	#179/272	34th
Food Science		
Medicine		
Medicine (miscellaneous)	#141/205	31st

CiteScoreTracker 2019 ⓘ

Last updated on 08 January, 2020  
Updated monthly

$$0.82 = \frac{\text{Citation Count 2019}}{\text{Documents 2016 - 2018}} = \frac{157 \text{ Citations to date} >}{192 \text{ Documents to date} >}$$

[View CiteScore trends >](#)

[Add CiteScore to your site](#)

Metrics displaying this icon are compiled according to Snowball Metrics ↗, a collaboration between industry and academia.

#### About Scopus

[What is Scopus](#)  
[Content coverage](#)  
[Scopus blog](#)  
[Scopus API](#)  
[Privacy matters](#)

#### Language

[日本語に切り替える](#)  
[切换到简体中文](#)  
[切换到繁體中文](#)  
[Русский язык](#)

#### Customer Service

[Help](#)  
[Contact us](#)

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.



# Document details

< Back to results | 1 of 4 Next >

↗ Export ↕ Download 🖨 Print ✉ E-mail 💾 Save to PDF ☆ Add to List More... >

View at Publisher

Current Research in Nutrition and Food Science [Open Access](#)  
Volume 7, Issue 3, December 2019, Pages 828-841

## The effect of tempeh gembus substitution on protein content, calcium, protein digestibility and organoleptic quality of meatballs (Article) [\(Open Access\)](#)

Arini, A.M.S., Afifah, D.N. ✉, Dieny, F.F. 🔍

Department of Nutrition Science, Diponegoro University, Jl. Prof. H. Soedarto, SH, Tembalang, Semarang, 50275, Indonesia

### Abstract

⌵ View references (46)

Osteopenia is a sign of osteoporosis that usually occurs in adolescent aged 18-24 years. One of the efforts to manage osteopenia is by giving a high calcium diet. Meatballs substituted by tempeh gembus can be used as an alternative to local food-based snacks as source of calcium. To analyze the effect of the substitution of tempeh gembus on the protein content, calcium, digestibility of the protein, and the acceptability of meatballs. The study was conducted in two stages, there are preliminary study and main study, and using randomized single factor experimental, tempeh gembus levels (25%, 50%, 75%, 100%) and control (0% tempeh gembus). The analysis of the content of protein was conducted using Kjeldahl method, calcium contents using Uv-Vis spectrophotometer, and protein digestibility using in-vitro method. Data was analyzed with mann-whitney. Organoleptic tests were carried out by hedonic testing of 30 semi-trained panelists. Substitution of tempeh gembus had significant effect on the protein contents and protein digestibility, but not significant effect on the calcium contents. Formulation with 25% tempeh gembus substitution was found to be a best formulation of meatballs containing protein content of 8.03%, calcium contents of 351.19 mg/100 g, and protein digestibility of 53.22%. Meatball with 25% tempeh gembus with 78 mg per serving is sufficient for nutrition label reference in general category, 13% energy; protein by 11%; calcium by 25%. © 2019 The Author(s). Published by Enviro Research Publishers.

### SciVal Topic Prominence ⓘ

Topic: Enzymes | Peptide Hydrolases | Fibrinolytic enzymes

Prominence percentile: 79.987 ⓘ

### Author keywords

Meatballs Nutrient Content Substitution Tempeh Gembus

### Funding details

Funding sponsor	Funding number	Acronym
	474-71/UN7.P4.3/ PP/2018	

### Funding text #1

The authors are thankful to the Research Institute of Diponegoro University which facilitated and funded this research.

Metrics ⓘ View all metrics >



### PlumX Metrics ⌵

Usage, Captures, Mentions, Social Media and Citations beyond Scopus.

Cited by 0 documents

Inform me when this document is cited in Scopus:

Set citation alert >

Set citation feed >

### Related documents

The characteristics of Kerupuk Gembus

Affah, D.N. , Nugrahani, G. , Hastuti, V.N.  
(2019) *IOP Conference Series: Earth and Environmental Science*

Antioxidant activity and soluble protein content of tempeh gembus hydrolysate

Agustina, R.K. , Dieny, F.F. , Rustanti, N.  
(2018) *Hiroshima Journal of Medical Sciences*

The effect of tempeh gembus variations to serum levels of high sensitivity c-reactive protein (Hscrp) and serum levels of fibrinogen of sprague dawley rats with aterogenic diet

Dewi, P.K. , Afifah, D.N. , Rustanti, N.  
(2018) *Romanian Journal of Diabetes, Nutrition and Metabolic Diseases*

View all related documents based on references

ISSN: 2347467X

Source Type: Journal

Original language: English



DOI: 10.12944/CRNFSJ.7.3.22

Document Type: Article

Publisher: Enviro Research Publishers

## References (46)

[View in search results format >](#)

☐ All   [Export](#)    [Print](#)    [E-mail](#)    [Save to PDF](#)   [Create bibliography](#)

- ☐ 1   Oleson, C.V.  
**Osteoporosis rehabilitation: A practical approach**  
  
(2017) *Osteoporosis Rehabilitation: A Practical Approach*, pp. 1-498.  
<https://www.springer.com/gp/book/9783319450827>  
ISBN: 978-331945084-1; 978-331945082-7  
doi: 10.1007/978-3-319-45084-1  
  
[View at Publisher](#)
- ☐ 2   Kementerian Kesehatan, R.I.  
(2015) *Data Dan Kondisi Penyakit Osteoporosis Di Indonesia*. Cited 3 times.  
Jakarta
- ☐ 3   Nafilah, Fitranti, D.Y.  
Hubungan Indeks Massa Tubuh (IMT), Persen Lemak Tubuh, Asupan Zat Gizi, dan Aktivitas Fisik dengan Kepadatan Tulang pada Remaja Putri  
(2014) *Nutrition College*, 3 (4), pp. 680-688.
- ☐ 4   Faizah, L.N., Fitranti, D.Y.  
Hubungan Asupan Protein, Fosfor, dan Kalsium dengan Kepadatan Tulang pada Wanita Dewasa Awal  
(2015) *Nutrition College*, 4 (2), pp. 335-341.
- ☐ 5   Fikawati, S., Syafiq, A., Puspasari, P.  
Faktor-Faktor yang Berhubungan dengan Asupan Kalsium pada Remaja di Kota Bandung  
(2005) *Jurnal Kedokteran Trisakti Universa Medicina*, 24 (1), pp. 24-34.
- ☐ 6   Mahan, L.K., Escott-Stump, S.  
(2008) *Krause's Food & Nutrition Therapy*. Cited 731 times.  
12th ed. Canada: Saunders Elsevier
- ☐ 7   Rosen, H.N.  
(2015) *Patient Education: Calcium and Vitamin D for Bone Health (Beyond the Basics)*  
MD, Available from  
<http://www.uptodate.com/contents/calciumand-vitamin-d-for-bone-health-beyondthe-basics>



- 
- ☐ 8 Damardjati, D., Marwoto, D.K., Swastika, D., Arsyad, Hilman, Y.  
(2005) *Prospek Dan Arah Pengembangan Agribisnis Kedelai*. Cited 8 times.  
Jakarta
- 
- ☐ 9 Deshinta, R.W.  
(2016) *Pengaruh Substitusi Tepung Ampas Tahu Terhadap Tingkat Kekerasan Dan Daya Terima Bakso* [Skripsi]. Surakarta: Universitas Muhammadiyah Surakarta
- 
- ☐ 10 Damanik, R.N.S., Pratiwi, D.Y.W., Widyastuti, N., Rustanti, N., Anjani, G., Afifah, D.N.  
**Nutritional Composition Changes during Tempeh Gembus Processing** ([Open Access](#))  
  
(2018) *IOP Conference Series: Earth and Environmental Science*, 116 (1), art. no. 012026. Cited 4 times.  
<http://www.iop.org/EJ/volume/1755-1315>  
doi: 10.1088/1755-1315/116/1/012026  
  
[View at Publisher](#)
- 
- ☐ 11 Murdiati, Sardjono, Amaliah, A.  
Perubahan Komposisi Kimia Tempe Gembus yang Dibuat dari Bahan Dasar Ampas Tahu dan Bekatul  
(2002) *Agri-tech*, 20 (2), pp. 106-110.
- 
- ☐ 12 Afifah, D.N., Nugrahani, G., Hastuti, V.N., Arifan, F.  
**The characteristics of Kerupuk Gembus** ([Open Access](#))  
  
(2019) *IOP Conference Series: Earth and Environmental Science*, 292 (1), art. no. 012055.  
<https://iopscience.iop.org/journal/1755-1315>  
doi: 10.1088/1755-1315/292/1/012055  
  
[View at Publisher](#)
- 
- ☐ 13 Noviana, A., Dieny, F.F., Rustanti, N., Anjani, G., Afifah, D.N.  
**Antimicrobial activity of tempeh gembus hydrolyzate** ([Open Access](#))  
  
(2018) *IOP Conference Series: Earth and Environmental Science*, 116 (1), art. no. 012044. Cited 3 times.  
<http://www.iop.org/EJ/volume/1755-1315>  
doi: 10.1088/1755-1315/116/1/012044  
  
[View at Publisher](#)
- 
- ☐ 14 Kurniasari, R., Sulchan, M., Afifah, D.N., Anjani, G., Rustanti, N.  
**Influence variation of tempe gembus (an Indonesian fermented food) on homocysteine and malondialdehyde of rats fed an atherogenic diet** ([Open Access](#))  
  
(2017) *Romanian Journal of Diabetes, Nutrition and Metabolic Diseases*, 24 (3), pp. 203-211. Cited 4 times.  
<https://www.degruyter.com/view/j/rjdnmd>  
doi: 10.1515/rjdnmd-2017-0026  
  
[View at Publisher](#)
-

- ☐ 15 Affah, D.N., Sulchan, M., Syah, D., Yanti, Suhartono, M.T., Kim, J.H.  
Purification and characterization of a fibrinolytic enzyme from *Bacillus pumilus* 2.g isolated from gembus, an Indonesian fermented food  
(2014) *Preventive Nutrition and Food Science*, 19 (3), pp. 213-219. Cited 20 times.  
<http://pdf.medrang.co.kr/paper/pdf/PNF/2014/019/PNF019-03-11.pdf>  
doi: 10.3746/pnf.2014.19.3.213  
[View at Publisher](#)
- 
- ☐ 16 Affah, D.N., Sulchan, M., Syah, D., Yanti, Suhartono, M.T.  
Isolation and identification of fibrinolytic protease-producing microorganisms from Red Oncom and Gembus, Indonesian fermented soybean cakes  
(2014) *Malaysian Journal of Microbiology*, 10 (4), pp. 273-279. Cited 6 times.  
<http://web.usm.my/mjm/issues/vol10no4/Research%207.pdf>
- 
- ☐ 17 Affah, D.N., Rustanti, N., Anjani, G., Syah, D., Yanti, Suhartono, M.T.  
Proteomics study of extracellular fibrinolytic proteases from *Bacillus licheniformis* RO3 and *Bacillus pumilus* 2.g isolated from Indonesian fermented food ([Open Access](#))  
(2017) *IOP Conference Series: Earth and Environmental Science*, 55 (1), art. no. 012025. Cited 3 times.  
<http://www.iop.org/EJ/volume/1755-1315>  
doi: 10.1088/1755-1315/55/1/012025  
[View at Publisher](#)
- 
- ☐ 18 Dewi, P.K., Affah, D.N., Rustanti, N., Sulchan, M., Anjani, G.  
The effect of tempeh gembus variations to serum levels of high sensitivity c-reactive protein (HsCRP) and serum levels of fibrinogen of sprague dawley rats with atherogenic diet ([Open Access](#))  
(2018) *Romanian Journal of Diabetes, Nutrition and Metabolic Diseases*, 25 (1), pp. 91-97. Cited 2 times.  
<https://www.degruyter.com/view/j/rjdnmd>  
doi: 10.2478/rjdnmd-2018-0010  
[View at Publisher](#)
- 
- ☐ 19 Affah, D.N., Nabilah, N., Supraba, G.T., Pratiwi, S.N., Nuryanto, Sulchan, M.  
The Effect of Tempeh gembus, an Indonesian Fermented Food, on Lipid Profiles in Women with Hyperlipidemia  
(2018) *Current Nutrition & Food Science*, 14 (1), pp. 1-9. Cited 2 times.
- 
- ☐ 20 Agustina, R.K., Dieny, F.F., Rustanti, N., Anjani, G., Affah, D.N.  
Antioxidant activity and soluble protein content of tempeh gembus hydrolysate  
(2018) *Hiroshima Journal of Medical Sciences*, 67 (Special issue), pp. 1-7. Cited 2 times.  
<http://home.hiroshima-u.ac.jp/medpress/Welcome.html>
- 
- ☐ 21 Sulchan, M., Endang, N.W.  
Nilai Gizi dan Komposisi Asam Amino Tempe Gembus serta Pengaruhnya terhadap Pertumbuhan Tikus  
(2007) *Maj Kedokt*, 57 (3), pp. 80-85. Cited 4 times.
-

□ 22 Bastian, F., Ishak, Tawali, Bilang, M.  
(2013) *Daya Terima Dan Kandungan Zat Gizi Formula Tepung Tempe Dengan Penambahan Semi Refined Carrageenan (SRC) Dan Bubuk Kakao*, p. 2.  
Aplikasi Teknologi Pangan

---

□ 23 Alakali, J.S., Kucha, C.T., Rabi, I.A.  
Effect of drying temperature on the nutritional quality of Moringa oleifera leaves  
(2015) *African J Food Sci*, 9 (7), pp. 395-399. Cited 7 times.

---

□ 24 *Standar Nasional Indonesia: Bakso Daging*, pp. 2-3.  
Jakarta: BSN; 014

---

□ 25 Purwanto, A., Ali, A., Herawati, N.  
Kajian Mutu Gizi Bakso Berbasis Daging Sapi dan Jamur Merang (*Volvariella volvacea*)  
(2015) *Teknologi Hasil Pertanian*, 14 (2), pp. 1-8. Cited 2 times.

---

□ 26 Sayudi, S., Herawati, N., Ali, A.  
Potensi Biji Lamtoro Gung dan Biji Kedelai sebagai Bahan Baku Pembuatan Tempe Komplementasi  
(2015) *Jom Faperta*, 1 (2), pp. 1-9.

---

□ 27 Affah, D.N., Rahma, A., Nuryandari, S.S., Alvice, L., Hartono, P.I., Kurniawati, D.M., Wijayanti, H.S., (...), Purwanti, R.  
Nutrition Content, Protein Quality, and Antioxidant Activity of Various Tempeh Gembus Preparations  
(2019) *Journal of Food and Nutrition Research*, 7 (8), pp. 605-612.

---

□ 28 Sumiati, T.  
(2008) *Pengaruh Pengolahan Terhadap Mutu Cerna Protein Ikan Mujair (Tilapia Mossambica)*  
[Skripsi]. Bogor: Institut Pertanian Bogor

---

□ 29 Takarina, H.L.  
(2013) *Kandungan Kalsium Dan Karbohidrat Bakso Daging Sapi Dengan Penambahan Jamur Tiram (Pleurotus Sp)*  
[Skripsi]. Surakarta: Universitas Muhammadiyah Surakarta

---

□ 30 Gropper, S.S., Smith, J.L., Groff, J.L.  
(2009) *Advanced Nutrition and Human Metabolism*, p. 431. Cited 510 times.  
5th ed. USA: Cengage Learning

---

□ 31 Gandjar, I.G., Rahman, A.  
Kimia Farmasi Analisis  
(2007) *Yogyakarta: Pustaka Pelajar*, 46374, pp. 240-262.

---

- 32 Muchtadi, T.R., Sugiyono, Ayustaningwarno, F.  
(2010) *Ilmu Pengetahuan Bahan Pangan*. Cited 15 times.  
Bandung: Alfabeta
- 
- 33 Soemarno  
(2007) *Rancangan Teknologi Proses Pengolahan Tapioka Dan Produk-Produknya*. Cited 2 times.  
Malang: Universitas Brawijaya
- 
- 34 Wibowo, S.  
(2008) *Budidaya Bawang Merah, Bawang Putih, Bawang Bombay*  
Jakarta: Penebar Swadaya
- 
- 35 (2010) *National Nutrient Database for Standard Reference of Raw Garlic [Internet]*  
Available from  
<https://ndb.nal.usda.gov/ndb/foods/show/301861>
- 
- 36 (2018) *National Nutrient Database for Standard Reference of Raw Pepper [Internet]*  
Available from  
<https://ndb.nal.usda.gov/ndb/foods/show/302197>
- 
- 37 (2018) *National Nutrient Database for Standard Reference of Raw Baking Soda [Internet]*  
Available from  
<https://ndb.nal.usda.gov/ndb/foods/show/45016339>
- 
- 38 Saputra, D.  
Penentuan Daya Cerna Protein In Vitro Ikan Bawal (*Colossoma macropomum*) pada Umur Panen Berbeda  
(2014) *Food Technology*, 5 (2), pp. 1127-1133.
- 
- 39 Hanifah, Kusharto, C.M.  
(2017) *Daya Cerna Protein Serta Kandungan Mineral (Kalsium Dan Fosfor) Pada Snack-Bar Substitusi Tepung Pisang (Musa Paradisiaca) Dengan Tepung Pury Ulat Sutera (Bombyx Mori) Dan Tepung Lele (Clarias Gariepinus)*  
[Skripsi]. Bogor: Institut Pertanian Bogor
- 
- 40 Budiman, Wulandari, Z., Suryati, T.  
Suplementasi Tepung Putih Telur untuk Memperbaiki Nilai Nutrisi Snack Ekstrusi Berbahan Grits Jagung  
(2009) *Media Peternak*, 32 (3), pp. 179-184.
- 
- 41 Negara, J.  
Aspek Mikrobiologis serta Sensori (Rasa, Warna, Tekstur, Aroma) pada Dua Bentuk Penyajian Keju yang Berbeda  
(2016) *Ilmu Produksi Dan Teknologi Hasil Peternakan*, 4 (2), pp. 286-290.
-

□ 42 Firahmi, N., Dharmawati, S., Aldrin, M.  
(2015) *Sifat Fisik Dan Organoleptik Bakso Yang Dibuat Dari Daging Sapi Dengan Lama Pelayuan Berbeda*  
Sains dan Teknologi

□ 43 Hatta, M.  
Pengaruh Level dan Waktu Penambahan Fosfat (Sodium Tripolifosfat/STTP) terhadap Kualitas Bakso  
(2011) *Jurnal Agrisistem*, 7, pp. 87-95.

□ 44 Soeparno  
(2009) *Ilmu Dan Teknologi Daging*. Cited 70 times.  
Yogyakarta: Gadjah Mada University Press

□ 45 Murni, M.  
The Effect of Addition Tempeh Flour to The Quality and the Taste of Chicken Nugget  
(2014) *Berita Litbang Industri*, 3 (2), pp. 117-123. Cited 2 times.

□ 46 (2016) *Acuan Label Gizi*  
Jakarta

🔍 Afifah, D.N.; Department of Nutrition Science, Diponegoro University, Jl. Prof. H. Soedarto, SH, Tembalang,  
Semarang, Indonesia; email:diananurafifah@live.undip.ac.id

© Copyright 2020 Elsevier B.V., All rights reserved.

< Back to results | 1 of 4 Next >

^ Top of page

## About Scopus

What is Scopus  
Content coverage  
Scopus blog  
Scopus API  
Privacy matters

## Language

日本語に切り替える  
切换到简体中文  
切换到繁體中文  
Русский язык

## Customer Service

Help  
Contact us

ELSEVIER

Terms and conditions ↗ Privacy policy ↗

Copyright © Elsevier B.V. ↗. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

RELX





## Current Research in Nutrition and Food Science

Current Research in Nutrition and Food Science Journal is an open access, peer reviewed international journal covering all aspects of Nutrition and Food Science.



Print ISSN: 2347-467X  
Online ISSN: 2322-0007

Frequency: Triannual ( April, August, December)

DOI: Prefix 10.12944

Chief Editor: Prof. Min-Hsiung Pan

CRNFSJ is a RoMEO green Journal

Journal Information.

## Recent Articles

### A Cross Sectional Study on Analysis of Human milk as per Prakriti (Psychosomatic Constitution).

Deeksha Singh<sup>1</sup> , B. M. Singh<sup>2</sup> , Sangeeta Gehlot<sup>1</sup> , Vandana Verma<sup>1</sup> 

[Read full article...](#)

### The Study of Sorghum (*Sorghum bicolor* L.), Mung Bean (*Vigna radiata*) and Sago (*Metroxylon sagu*) Noodles: Formulation and Physical Characterization

Sri Budi Wahjuningsih<sup>1\*</sup>, Sudjatinah<sup>1</sup>, Mita Nurul Azkia<sup>1</sup>, Dian Anggraeni<sup>2</sup>

## Scopus Journal Metrics

CiteScore 2018: 0.78

CiteScore Details



Journal Archived in Portico

Archived In



Follow us on:



[Submit Manuscript Online](#)



[Join our team of Reviewers](#)

[Read full article...](#)

## Effects of Frozen Storage on Viability of Probiotics and Antioxidant Capacities of Synbiotic Riceberry and Sesame-Riceberry Milk Ice Creams

Varongsiri Kemsawasd<sup>1</sup>, Pittaya Chaikham<sup>2\*</sup>[Read full article...](#)

## Sensory Evaluation of Noodles Substituted by Sweet Potato Flour and Rice Bran

Retno Dwi Astuti, Wahyudi David , Ardiansyah\* [Read full article...](#)

## Lactoferrin Association with Maternal Nutritional Status and Lactation Stages

Mohammad Zen Rahfiludin , Dina Rahayuning Pangestuti\* [Read full article...](#)

Journal is indexed in...

## Most Popular Articles

Colostrum - its Composition, Benefits as a...

(82,268)

Importance of Exclusive Breastfeeding and Complementary...

(54,022)

Low Alkaline Phosphatase (ALP) In Adult Population an...

(49,306)

Processing and Nutritive Value of Mango Seed Kernel Flour

(28,511)

Processing and Nutritional Composition of Rice Bran

(26,579)

### Links

[About](#)[Archives](#)[Article Processing Charges](#)[Coming Issue](#)[Contact](#)[Current Issue](#)[Order Print Issue](#)[Submission](#)[Conflict of Interest](#)[Editorial Policy](#)

### Contact Us

Your Name (required)

Your Email (required)

Your Message

YVAU

Type the above text in box below (Case sensitive)

Compte the quiz

2+2=?

Send

Follow us on:



### License



This work is licensed under a Creative Commons Attribution 4.0 International License.

### Recent Articles

A Cross Sectional Study on Analysis of Human milk as per Prakriti (Psychosomatic Constitution).

The Study of Sorghum (*Sorghum bicolor* L.), Mung Bean (*Vigna radiata*) and Sago (*Metroxylon sagu*) Noodles: Formulation and Physical Characterization



Effects of Frozen Storage on Viability of Probiotics and Antioxidant Capacities of Synbiotic Riceberry and Sesame-Riceberry Milk Ice Creams

Copyright © 2019 - This Site - All rights reserved. Conforms to W3C Standard XHTML & CSS



Home » The Effect of Tempeh Gembus Substitution on Protein Content, Calcium, Protein Digestibility and Organoleptic Quality of Meatballs

## The Effect of Tempeh Gembus Substitution on Protein Content, Calcium, Protein Digestibility and Organoleptic Quality of Meatballs

Audrey Mutiara Setya Arini, Diana Nur Afifah\* , Fillah Fithra Dieny 

Department of Nutrition Science, Faculty of Medicine, Diponegoro university, Semarang, Indonesia, 50275

Corresponding Author Email: diananurafifah@live.undip.ac.id

DOI : <http://dx.doi.org/10.12944/CRNFSJ.7.3.22>

### Article Publishing History

Received: 27/07/2019

Accepted: 19/11/2019

Published Online: 20/11/2019

Plagiarism Check: Yes

Reviewed by: Dr. Perna Gupta   India

Second Review by: Dr. Abhijeet B. Muley   Spain

Final Approval by: Dr. Dan Cristian Vodnar

### Article Metrics

 Views: 213

 PDF Downloads: 44

### Scopus Journal Metrics

CiteScore 2018: 0.78

CiteScore Details

**SJR 2018**



Journal Archived in Portico

Archived In



[Submit Manuscript Online](#)

[Join our team of Reviewers](#)

[Journal is indexed in...](#)

### Abstract:

Osteopenia is a sign of osteoporosis that usually occurs in adolescent aged 18-24 years. One of the efforts to manage osteopenia is by giving a high calcium diet. Meatballs substituted by tempeh gembus can be used as an alternative to local food-based snacks as source of calcium. To analyze the effect of the substitution of tempeh gembus on the protein content, calcium, digestibility of the protein, and the acceptability of meatballs. The study was conducted in two stages, there are preliminary study and main study, and using randomized single factor experimental, tempeh gembus levels (25%, 50%, 75%, 100%) and control (0% tempeh gembus). The analysis of the content of protein was conducted using Kjeldahl method, calcium contents using Uv-Vis spectrophotometer, and protein digestibility using in-vitro method. Data was analyzed with mann-whitney. Organoleptic tests were carried out by hedonic testing of 30 semi-trained panelists. Substitution of tempeh gembus had significant effect on the protein contents and protein digestibility, but not significant effect on the calcium contents. Formulation with 25% tempeh gembus substitution was found to be a best formulation of meatballs containing protein content of 8.03%, calcium contents of 351.19 mg / 100 g, and protein digestibility of 53.22%. Meatball with 25% tempeh gembus with 78 mg per serving is sufficient for nutrition label reference in general category, 13% energy; protein by 11%; calcium by 25%.

### Keywords:

Meatballs; Nutrient Content; Substitution; Tempeh Gembus



Download this article as:

### Copy the following to cite this article:

Arini A. M. S, Afifah D. N, Dieny F. F. The Effect of Tempeh Gembus Substitution on Protein Content, Calcium, Protein Digestibility and Organoleptic Quality of Meatballs. Curr Res Nutr Food Sci 2019; 7(3). doi : <http://dx.doi.org/10.12944/CRNFSJ.7.3.22>

### Copy the following to cite this URL:

## Most Popular Articles

Colostrum - its Composition, Benefits as a...

(82,268)

Importance of Exclusive Breastfeeding and Complementary...

(54,022)

Low Alkaline Phosphatase (ALP) In Adult Population an...

(49,306)

Processing and Nutritive Value of Mango Seed Kernel Flour

(28,511)

Processing and Nutritional Composition of Rice Bran

(26,579)

Arini A. M. S, Afifah D. N, Dieny F. F. The Effect of Tempeh Gembus Substitution on Protein Content, Calcium, Protein Digestibility and Organoleptic Quality of Meatballs. Curr Res Nutr Food Sci 2019; 7(3). <https://bit.ly/2s1iVSR>

## Introduction

Osteoporosis is a bone metabolic disease characterized by a decrease in bone mass due to reduced bone matrix and minerals accompanied by microarchitecture damage from bone tissue, with a consequent decrease in bone strength, resulting in a tendency for bones to break easily.<sup>1</sup> Osteoporosis is a disease that is classified as a silent disease because it does not show specific symptoms.<sup>2</sup> Osteopenia is a sign of osteoporosis that begins with low bone density, if it lasts for a long time it can lead to decreased bone strength resulting in osteoporosis.<sup>3</sup>

A lifestyle that occurs in adolescent aged 18-24 years such as smoking, consumption of different types of beverages, not exercising, not drinking milk, sedentary lifestyle, consumption of fast food which is generally high in protein and inappropriate calcium intake are the factors lead to an increase the risk of osteoporosis at this time.<sup>4</sup> Research in high school adolescents showed that 76.2% of adolescents are in a situation of lack of calcium consuming category. The average calcium intake is only 559.05 mg/day or 55.9% of the Daily Value (DV).<sup>5</sup>

Calcium intake recommended for women who have not yet menopause is around 1000-1200 mg /day.<sup>6</sup> The main sources of calcium found in foods are milk and other dairy products such as cheese or yogurt, green vegetables such as broccoli, fortified juices with calcium, and soybeans and soy products.<sup>7</sup> Indonesia has a variety of soy products such as soybean milk, tofu, tempeh, soy sauce, tauco, and *tempeh gembus* which can be an alternative affordable source of calcium.<sup>8</sup>

Meatballs are a type of food that is shaped like a ball made from meat and flour. In general, meatballs in the community vary in name according to the types of basic ingredients such as fish meatballs, chicken meatballs, and beef meatballs or original meatball.<sup>9</sup>

*Tempeh gembus* is one of the functional food products that is widely eaten by Indonesians. *Tempeh gembus* are usually made from tofu residue from soybeans which are commonly used as animal feed. When compared with other soybean products, such as tofu and *tempeh*, *tempeh gembus* has relatively a little nutrient content. This is because *tempeh gembus* is the residual of tofu processing. Protein loss in tofu processing is mainly due to boiling process and unused residue from tofu.<sup>10</sup>

In 2002, Murdiati reported that tofu residue contains 22.28% protein, 5.87% fat, and 71.83% carbohydrates.<sup>11</sup> Another study showed that an innovative snack product like *kerupuk* made from tapioca flour combined with *tempeh gembus* still contains high fiber ranged between 38.1 to 67%.<sup>12</sup> *Tempeh gembus* contains of essential fatty acids, such as linoleic acid 21.51%, linolenic acid 1.82% and unsaturated fatty acids 16.72%.<sup>13</sup> *Tempeh gembus* (dry weight per 100 g) also had some important nutrition contents, such as protein 4.07 g, carbohydrate 14.25 g, fiber 4.69 g, calcium 159.98 mg and iron 0.48 mg.<sup>14</sup> *Tempeh gembus* had health effects such as proteolytic, fibrinolytic and anti-inflammation effect that might be able to act as antithrombotic.<sup>15-19</sup> Based on the previous study, *tempeh gembus* also known had antimicrobial<sup>13</sup> (against *S. aureus*, *B. subtilis*, and *S. mutans*) and antioxidant activities.<sup>20</sup> Raw *tempeh gembus* contains fiber 3.93 g, calcium 143 mg, phosphorus 50 mg, and iron 0.40 mg per 100 g.<sup>21</sup> The content of these nutrients will also be easier to digest due to fermentation. The digestive enzymes produced by tempeh mold make protein, fat, and carbohydrates in tempeh easier to digest.<sup>22</sup> Protein digestibility values of tofu residue are also quite high even though it is still lower than tofu. Protein digestibility on soybean yield was 78.7% for tofu residue, 92.7% for tofu and 65.3% for boiled soybeans.<sup>11</sup>

Analysis of nutrient content in the form of calcium and protein contents was carried out to determine the nutritional content of *tempeh gembus* meatballs. Protein digestibility test was carried out to determine the quality of protein digestibility in *tempeh gembus* meatballs. The acceptance level test was conducted to test the acceptance of *tempeh gembus* meatball products among teenagers. Hence, the objective of the present work was found the best formulation of tempeh gembus meatballs and analyze the effect of the substitution of *tempeh gembus* on the protein content, calcium, digestibility of the protein, and the acceptability of *tempeh gembus* meatballs.

## Materials and Methods

Materials of this study sample used *tempeh gembus* which is obtained from the manufacturer located in Semarang Indah Cluster. *Tempeh gembus* used was from the local soybean processing in the Grobogan area.

This research was an experimental study with a complete randomized one-factor design. The factor in this research is the comparison of the composition of tempeh gembus with beef in units of percent. In this study, 4 levels of treatment were carried out with a percentage of 25%, 50%, 75%, 100% and 1 control with a percentage of 0%. Each group was repeated five times. All samples will be analyzed for protein contents was conducted using *Kjeldahl* method, calcium contents using spectrophotometer Uv-Vis, protein digestibility using *in-vitro* method, and organoleptic tests on one sample from each treatment without repetition using hedonic tests (level of preference) on semi-trained panelists. Determination of the formulation of adding *tempeh gembus* is through preliminary research. Salt, garlic, shallots, and pepper were used as flavoring agents. Egg white and baking soda were used as adhesives and dough developers.

#### Preliminary Treatment Samples

The main study used a completely randomized design of 1 factor, namely the substitution of *tempeh gembus* with 4 levels (25%, 50%, 75%, 100%) and 1 control (0% *tempeh gembus*) with 5 repetitions to obtain 25 experimental units for analysis of nutrient content and the level of acceptance.

The meatballs were formulated by weighing the ingredients according to the level of treatment. The *tempeh gembus* steamed and smoothed using a mortar. *Tempeh gembus* mixed with all mashed ingredients and ground meat. After that, shaping and boiling process is done in boiling water until the meatballs appear on the surface of the water  $\pm$  15 min.

Table 1: Formulation of Treatment in Research

Material	Amount of Material				
	0%	25%	50%	75%	100%
Minced beef	100 g	75 g	50 g	25 g	-
<i>Tempeh Gembus</i>	-	25 g	50 g	75 g	100 g
Tapioca flour	20 g	20 g	20 g	20 g	20 g
Pepper	2,5 g	2,5 g	2,5 g	2,5 g	2,5 g
Egg	20 g	20 g	20 g	20 g	20 g
Garlic	2 g	2 g	2 g	2 g	2 g
Red onion	2 g	2 g	2 g	2 g	2 g
Salt	1 g	1 g	1 g	1 g	1 g
Baking soda	1 g	1 g	1 g	1 g	1 g

#### Protein Content Analysis

The refined samples were weighed as much as 0.2 g and dissolved into the Kjeldahl flask. As much as 0.7 g of nitrogen catalyst which includes 250 g of  $\text{Na}_2\text{SO}_4$  + 5 g of  $\text{CuSO}_4$  + 0.7 g of Selenium /  $\text{TiO}_2$  after that added 4 ml of concentrated  $\text{H}_2\text{SO}_4$  is inserted to the flask through the flask wall. The solution was destroyed in the fume hood until the color of the solution turned clear green, then added 10 ml of aquadest. The solution was distilled by adding 20 ml of NaOH -  $\text{TiO}$  (NaOH 40% +  $\text{Na}_2\text{S}_2\text{O}_3$  5%) then the distillate was accommodated in Erlenmeyer which has been filled with 4%  $\text{H}_3\text{BO}_3$  and the indicator Mr-BCG up to 60 ml where the color changes from red to blue. The distillate solution is tested using a standard 0.02 N HCl solution until it reaches the end point of the titration, which changes color from blue to pink and records the volume of the titration.

#### Calcium Contents Analysis

The samples were weighed as much as 500 mg. 25 mL of concentrated HCL added and heated it for 15 minutes. The sample was filtered in to a 50 mL measuring flask and distilled water to the limit.

The sample solution that was made was taken as much as 1 mL and neutralized with 40% NaOH. After neutral, the volume is sufficient to 10 mL. The neutral solution was taken as much as 1 mL and put it in a 25 mL measuring flask. After that, 1 mL of murexide solution and 2 mL of NaOH were added, then filled with distilled water to 25 mL. The solution is shaken until it is homogeneous and inserted into the cuvette to be read on the length of a wave of 534.6  $\mu\text{m}$ .

### Protein Digestibility Analysis

The samples weighed as much as 5 g were put into Erlenmeyer then added 20 ml of pH 2 whaffole buffer as much as 20 ml and 1% pepsin enzyme as much as 2 ml. The solution was incubated at 40°C for an hour. The solution was filtered or centrifuged and then 5 ml of 5% TCA was added. The solution was left to stand for an hour then 5 ml of filtrate was taken to analyze the protein content.

### Acceptance Level Analysis

The organoleptic tests using hedonic tests (level of preference) on semi-trained panelists as many as 30 people from Nutrition Science students of Diponegoro University in the semester 4 and 6 grades with 5 preference scales, namely 1 = Very dislike, 2 = Dislike, 3 = Neutral, 4 = Like, 5 = Really Like. In this organoleptic test, there are 5 formulations in the form of 1 control formulation and 4 substitution formulations.

### Statistical Analysis

All results of the experiments are expressed in median and mean  $\pm$  standard deviation. Data were analyzed by using statistical software. Kruskal-Wallis and Friedman were used to test the mean differences and the statistical significance differences between mean values was established at  $p < 0.05$ . Mann-Whitney and Wilcoxon were used to post-hoc test.

### Results

**Table 2: Results of Protein Content Analysis of *Tempeh Gembus* Meatballs**

Addition of <i>Tempeh Gembus</i> Levels	n	Median (maximum-minimum)	Mean $\pm$ SD (%)	p*
0%	5	11.27 (11.16-11.61)	11.30 $\pm$ 0.18 <sup>a</sup>	< 0.05
25%	5	8.04 (8.02-8.04)	8.03 $\pm$ 0.01 <sup>b</sup>	
50%	5	7.25 (7.19- 7.34)	7.26 $\pm$ 0.06 <sup>c</sup>	
75%	5	4.45 (4.40- 4.50)	4.45 $\pm$ 0.04 <sup>d</sup>	
100%	5	3.25 (3.08- 3.37)	3.23 $\pm$ 0.14 <sup>e</sup>	

\*Kruskal-Wallis Test, Mann-Whitney Post-hoc Test; \*\* Different superscript letters (a, b, c) show significant differences ( $p < 0.05$ )

### Protein Contents

The results of the analysis of protein contents of *tempeh gembus* meatballs showed the difference between the addition of *tempeh gembus* in meatballs and protein contents ( $p = 0.00$ ). Protein contents of all treatment groups are significantly different from controls and different between treatment groups. The highest protein content in the control class of *tempeh gembus* meatballs were 11.21%. The more additions of tempeh to the *tempeh gembus* meatballs, the protein content decreases significantly.

**Table 3: Results of Calcium Analysis of *Tempeh Gembus* Meatballs**

Addition of <i>Tempeh Gembus</i> Levels	n	Median (maximum-minimum)	Mean $\pm$ SD (mg/100g)	p*
0%	5	350.68 (343.35- 357.74)	351.44 $\pm$ 6.21	< 0.05
25%	5	351.45 (343.35- 356.84)	351.19 $\pm$ 5.36	
50%	5	354.40 (332.42- 356.20)	350.49 $\pm$ 10.13	
75%	5	354.53 (350.16- 356.20)	353.94 $\pm$ 2.29	
100%	5	355.30 (351.32-356.07)	354.53 $\pm$ 1.94	

\* Kruskal-Wallis Test

### Calcium Contents

The results of the analysis of calcium content in *tempeh gembus* meatballs showed no difference between the addition of *tempeh gembus* in meatballs on calcium content ( $p = 0.944$ ). Calcium contents across treatment groups did not differ significantly from controls. The highest calcium content is found in 100% *tempeh gembus* meatballs for 354.53 mg/100g.



Table 4: Results of Analysis of Protein Digestibility of *Tempeh Gembus* Meatballs

Addition of <i>Tempeh Gembus</i> Levels	n	Median (maximum-minimum)	Mean $\pm$ SD (%)	p*
0%	5	83.71 (75.28- 89.21)	82.17 $\pm$ 5.37 <sup>a</sup>	< 0.05
25%	5	54.06 (45.14- 60.15)	53.22 $\pm$ 5.52 <sup>b</sup>	
50%	5	64.10 (47.82- 64.93)	58.44 $\pm$ 8.52 <sup>b</sup>	
75%	5	56.22 (52.98- 75.92)	61.27 $\pm$ 10.13 <sup>b</sup>	
100%	5	5 87.11 (71.15- 91.05)	83.96 $\pm$ 8.19 <sup>a</sup>	

\*Kruskal-Wallis Test, Mann-Whitney Post-hoc Test; \*\*Different superscript letters (a, b, c) show significant differences (p <0.05)

#### Protein Digestibility

The results of the analysis of protein digestibility of *tempeh gembus* meatballs showed the difference between adding *tempeh gembus* in meatballs and protein digestibility (p = 0.00). There is no significant difference between the treatment of 0% and 100%, 25% and 50%, 25%, and also 75%, and 50% and 75%. The highest protein digestibility in 100% *tempeh gembus* meatballs was 83.97%.

Table 5: Results of Analysis of Acceptance Levels on the Color of Meatballs with *Tempeh Gembus* Substitution

Addition of <i>Tempeh Gembus</i> Levels	n	Median (maximum-minimum)	Mean $\pm$ SD	p*
0%	30	4.00 (2.00 - 5.00)	3.63 $\pm$ 1.03 <sup>a,b</sup>	< 0.05
25%	30	4.00 (2.00 - 5.00)	3.80 $\pm$ 1.06 <sup>a</sup>	
50%	30	3.00 (2.00 - 5.00)	3.16 $\pm$ 0.87 <sup>b</sup>	
75%	30	3.00 (2.00- 5.00)	3.26 $\pm$ 0.78 <sup>b</sup>	
100%	30	2.00 (2.00- 5.00)	2.73 $\pm$ 0.98 <sup>c</sup>	

\*Friedman Test, Wilcoxon Post-hoc Test; \*\* Different superscript letters (a, b, c) show significant differences (p <0.05)

#### Acceptance Level Test

##### Color

Based on the statistical analysis with the Friedman test showing the addition of *tempeh gembus* in meatballs showed a significant increase in the color of meatballs (p = 0.00). The color of *tempeh gembus* meatballs with the substitution of 25% *tempeh gembus* had the highest level of acceptance of 3.80 (likes), while meatballs with 100% *tempeh gembus* content had the lowest level of acceptance of color of 2.73 (neutral).

The results of further test analysis showed that there were no significant differences in the treatment of 0% and 25%, 0% and 50%, 0% and 75%, 50%, and 75%.

Table 6: Results of Analysis of Acceptance Levels on the Taste of Meatballs with *Tempeh Gembus* Substitution

Addition of <i>Tempeh Gembus</i> Levels	n	Median (maximum-minimum)	Mean $\pm$ SD	p*
0%	30	3.00 (2.00- 5.00)	3.23 $\pm$ 1.04	< 0.05
25%	30	3.00 (2.00- 5.00)	2.80 $\pm$ 0.80	
50%	30	3.00 (2.00- 5.00)	2.76 $\pm$ 0.89	
75%	30	3.00 (2.00- 5.00)	3.13 $\pm$ 1.22	
100%	30	2.00 (2.00- 5.00)	2.76 $\pm$ 0.97	

\*Freidman Test

### Taste

Based on statistical analysis with the Friedman test, the addition of *tempeh gembus* in meatballs did not significantly increase or decrease the meatball taste ( $p = 0.24$ ). There was no significant difference in taste in the addition of *tempeh gembus* to the meatballs.

**Table 7: Results of Analysis of Acceptance Levels on the Flavor of Meatballs with *Tempeh Gembus* Substitution**

Addition of <i>Tempeh Gembus</i> Levels	n	Median (maximum-minimum)	Mean $\pm$ SD	p*
0%	30	3.00 (2.00- 5.00)	3.23 $\pm$ 1.04	< 0.05
25%	30	3.00 (2.00- 5.00)	2.80 $\pm$ 0.80	
50%	30	3.00 (2.00- 5.00)	2.76 $\pm$ 0.89	
75%	30	3.00 (2.00- 5.00)	3.13 $\pm$ 1.22	
100%	30	2.00 (2.00- 5.00)	2.76 $\pm$ 0.97	

\*Friedman Test, Wilcoxon Post-hoc Test; \*\* Different superscript letters (a, b, c) show significant differences ( $p < 0.05$ )

### Flavor

The analysis of the level of acceptance of meatball flavor with the addition of *tempeh gembus* showed significant results. There is a difference between the flavor of meatballs without the addition of *tempeh gembus* and with the addition of *tempeh gembus*.

The addition of *tempeh gembus* in meatballs significantly increased the level of acceptance of meatball flavor ( $p = 0.04$ ). The most preferred meatballs flavor is meatballs with the addition of *tempeh gembus* by 25%, while meatballs with 50% addition have the lowest level of acceptance of the flavor which is equal to 3.03 (neutral).

The results of further test analysis showed that there were significant differences in the treatment of 0% and 50%.

**Table 8: Results of Analysis of Acceptance Levels on the Texture of Meatballs with *Tempeh Gembus* Substitution**

Addition of <i>Tempeh Gembus</i> Levels	n	Median (maximum-minimum)	Mean $\pm$ SD	p*
0%	30	3.00 (2.00- 5.00)	2.93 $\pm$ 0.94 <sup>a,b</sup>	< 0.05
25%	30	3.00 (2.00- 5.00)	3.16 $\pm$ 0.79 <sup>a</sup>	
50%	30	3.00 (2.00- 5.00)	2.60 $\pm$ 0.89 <sup>b,d</sup>	
75%	30	3.00 (2.00- 5.00)	2.00 $\pm$ 0.87 <sup>c</sup>	
100%	30	2.00 (2.00- 5.00)	2.46 $\pm$ 1.13 <sup>d</sup>	

\*Friedman Test, Wilcoxon Post-hoc Test; \*\* Different superscript letters (a, b, c) show significant differences ( $p < 0.05$ )

### Texture

The addition of *tempeh gembus* in meatballs gives a difference to the texture of meatballs ( $p = 0.00$ ). The texture of meatballs with the addition of 25% *tempeh gembus* has the highest level of texture acceptance which is 3.16 (neutral). Whereas meatballs with the addition of 75% *tempeh gembus* have a low texture acceptance which is only 2.00 (dislike).

The results of further test analysis showed that there were no significant differences in the treatment of 0% and 25%, 0% and 50%, and 50% and 100%.

### Discussion

#### Protein Contents

Addition of *tempeh gembus* in meatballs significantly reduced protein contents. In meatballs added with *tempeh gembus*, the protein content decreased due to the number of beef used and heat treatment during cooking. Meanwhile, previous study reported that the increase in temperature corresponded to the increase in protein denaturation and significant decrease in

protein content as observed in the present study.<sup>23</sup> Protein contents in meatballs with *tempeh gembus* content of 25%, 50%, 75%, and 100%, differ significantly from the control. Meatballs with *tempeh gembus* content of 0% or controls have the highest protein content (11.21%), while meatballs without meat additives or 100% *tempeh gembus* have the lowest protein content (3.24%). However, when compared with meatball quality standards according to the National Standard of Indonesia, the meatball protein content with the formulation of 0% *tempeh gembus* or control has met these standards which is minimum 11% protein.<sup>24</sup>

Meatball protein contents are influenced by protein content from raw materials. The increasing use of beef, it will show a tendency to increase contents of meatball protein produced. This is because beef has a higher protein content than *tempeh gembus*. The protein content of beef reaches 18.80 g per 100 g of ingredients,<sup>25</sup> Meanwhile, the protein content of *tempeh gembus* is only 6.7 g per 100 g of ingredients.<sup>10</sup>

In the results of the study, there was a decrease in protein contents in *tempeh gembus* meatballs when compared with the previous theory. This can be affected due to processing that occurs when making *tempeh gembus* and when making *tempeh gembus* meatballs. Making *tempeh gembus* through the fermentation process wherein the process is used *Rhizopus sp.* strain. The nitrogen compounds contained in the *tempeh gembus* protein are utilized by the mold of *Rhizopus sp.* for its growth.<sup>26</sup>

The steaming factor in *tempeh gembus* before making meatballs also affects the decrease of protein in meatballs. This was proven based on the research that had been done before, the *tempeh gembus* with steaming treatment had decreased in protein content compared to the *tempeh gembus* without any treatment.<sup>27</sup> Protein will be coagulated at a temperature of 100°C and the water contained in the food will come out. This is because the protein loses its binding power to water so that the increase in steaming temperature will cause the protein to be dissolved and denatured.<sup>28</sup> In addition, the decrease in protein content in meatballs is also alleged to be the result of the addition of *tempeh gembus*, where the *tempeh gembus* contains a lot of fiber. Fiber can bind to water. Water can dissolve protein. The presence of fiber on meatballs due to the addition of *tempeh gembus* causes water to be unable to bind proteins perfectly because water binds to fiber so that a lot of protein is dissolved in the process of cooking meatballs.<sup>29</sup>

#### Calcium Contents

Calcium is the micronutrient needed by the body and the most abundant minerals in the body, which is 1.5-2% of an adult's body weight, or around 1,000-1,400 g per 70 kg of body weight.<sup>30</sup> Calcium analysis in this study used the Uv-Vis spectrophotometric method. This method is used for quantitative analysis of metal elements in trace amounts and ultra-trace amounts. This method gives the total amount of metal elements in a sample and does not depend on the molecular form of the metal in the sample.<sup>31</sup>

Calcium contents in meatballs do not increase or decrease significantly. Calcium contents in the addition of *tempeh gembus* at 100% level with the highest calcium content of 354.53 mg/100 g of *tempeh gembus* meatballs. This result shows that the calcium content of the *tempeh gembus* meatballs is almost the same as beef meatballs. However, this is not in accordance with the theory of previous research. In a study conducted by Sulchan *et al.*, calcium contents contained in *tempeh gembus* were 143 mg /100g.<sup>21</sup> Whereas from the research conducted by Ruth *et al.*, that is equal to 232.09 mg/100g.<sup>10</sup>

Increased calcium contents are alleged due to the addition of ingredients found in *tempeh gembus* meatballs. In the process of making meatballs, added some ingredients such as 20% egg white, 20% tapioca flour, 2% onion, 2% garlic, 2.5% pepper, and 1% baking soda. Calcium content in egg white is 8 mg/100g<sup>32</sup>, tapioca flour 20 mg /100mg<sup>33</sup>, onion 36 mg /100 mg<sup>34</sup>, garlic 181 mg /100mg<sup>35</sup>, pepper 18 mg /100g<sup>36</sup>, baking soda 40 mg /100g.<sup>37</sup> Calcium content contained in the composer material of *tempeh gembus* can have an effect on the increase of calcium in *tempeh gembus* meatballs.

#### Protein Digestibility

Protein digestibility is the ability of proteins to be hydrolyzed into amino acids by digestive enzymes; which if the protein digestibility is high means that proteins can be hydrolyzed properly into amino acids so that the number of amino acids that can be absorbed and used by the body is high, while low protein digestibility means proteins are difficult to hydrolyze into amino acids so that the amount of amino acids that can be absorbed and is used by a low body because most of it will be removed by the body together with feces.<sup>38</sup>

Addition of *tempeh gembus* in meatballs had significant effect on digestibility of meatball protein ( $p = 0.00$ ). The highest protein digestibility is found in meatballs with 100% *tempeh gembus* content of 83.97%. This result shows a high number. Protein digestibility is high if the digestive power is equal to or greater than 80%.<sup>39</sup> While the lowest digestibility of the protein is found in meatballs with the *tempeh gembus* content of 25% and 50%. This is not in accordance with previous research. In the previous study, it was stated that the *tempeh gembus* with the steaming process had a protein digestibility of 48.68%.<sup>27</sup> This was alleged because, at the time of making the *tempeh gembus* meatballs, meatballs underwent a boiling stage. The boiling stage can make composed proteins become denatured. Denaturation is the breaking of bonds in molecules so that protein molecules will tend to be easily digested by digestive enzymes.<sup>28</sup> Protein that has been denatured will be easily digested.<sup>38</sup>

The digestibility of *tempeh gembus* meatball protein in the treatment of 25%, 50%, and 75%, decreased in value compared to meatballs with a treatment *tempeh gembus* of 0% and 100%. This is alleged due to the formation of cross-linking proteins. Cross-linking that occurs causes proteases such as trypsin to find difficulty in breaking or breaking peptide bonds in cross-linked proteins. Lysine will easily interact with the aliphatic carbon group  $[-(CH_2)_4-NH_3]$  so that the availability of lysine becomes less useful.<sup>40</sup>

### Acceptance Level

#### Color

In organoleptic tests, at first, the product is assessed visually by looking at the colors they have. Color is the first sensory that can be seen directly by panelists. Determination of the quality of food generally depends on the color it has. Color that does not deviate from the color that should be, will give the panelists a distinctive impression.<sup>41</sup>

The addition of *tempeh gembus* to the meatball had a significant effect on the level of acceptance of meatball color ( $p = 0.00$ ). The average rating value of the panelist's hedonic test on meatball color ranges from 2.73 to 3.80 (neutral-like). Panelists generally like the color of meatballs from all treatments caused by the color of meatballs in this study the same as the color of beef meatballs on the market. However, on meatballs with a content of 100% *tempeh gembus* without the addition of meat, the resulting color is slightly paler compared to meatballs with a mixture of meat.

The color of grey brownish meatballs comes from the process of heating or boiling meatballs dough. During heating, the color of the meat will gradually change from pink to paler. The color change is a result of the amount of myoglobin pigment being oxidized to metmyoglobin and protein polymerization.<sup>42</sup> In making meatballs, meat should be used in the pre-rigormortis phase. This is because the use of meat in this phase affects the color of the meatballs produced. The use of post-rigor beef will produce a whiter meatball color when compared to pre-rigor meat.<sup>43</sup>

#### Taste

Taste is a determining factor for consumer acceptance of food products. Formulations of spices, composer and meat conditions for making meatballs greatly affect the taste of the meatballs produced.<sup>42</sup> The amount of *tempeh gembus* added to the meatballs did not affect the level of acceptance of the *tempeh gembus* meatballs ( $p = 0.24$ ). The average value of the panelist assessment of the hedonic test for meatball taste ranges from 2.76 to 3.23 (neutral). The 0% treatment of *tempeh gembus* or 100% beef meatballs received the highest score of 3.23 which indicates that the panelists prefer meatballs with a content of 100% beef because it is more savory.

The savory typical value of meatballs is obtained from glutamic acid contained in beef.<sup>25</sup> Glutamic acid contained in beef reaches 14.4 g/100g of ingredients. Whereas, glutamic acid contained in the *tempeh gembus* only 0.29%/100g *tempeh gembus* wet weight.<sup>21</sup> Besides glutamic acid, meatballs from pre-rigor meat also have a better taste. This is because pre-rigor meat has a water binding capacity and high pH which increases tenderness and juicy in meat.<sup>44</sup>

#### Flavor

The assessment of the flavor of food is an evaluation of the sense of smell. Addition of *tempeh gembus* to the meatball had a significant effect on the level of acceptance of meatball flavor ( $p = 0.04$ ). The average value of the hedonic test by the panelists on meatball flavor ranges from 3.03 to 3.70 (neutral-like). The highest value (most preferred) for the flavor assessment was obtained from the chopped *tempeh gembus* meatballs with the *tempeh gembus* formulations of 0%. Spices such as garlic can enhance and modify the flavor. Spices are material that is

intentionally added to improve consistency, nutritional value, taste, control acidity, and basicity, and to strengthen the shape and appearance of products.<sup>42</sup> Spices are also useful for increasing taste, as an antioxidant that can reduce rancidity, and as an anti-microbial that can extend the shelf life of meatballs. In addition, based on research on soybean *tempeh*, the fat contained in soybean *tempeh* is resistant to the rancidity process which is influenced by the production of natural antioxidants by mold *tempeh*.<sup>45</sup>

### Texture

Texture assessment is done to find out whether the surface of the *tempeh gembus* is elastic. Elasticity is the ability of food to return to its original form after being pressured<sup>33</sup>. The addition of *tempeh gembus* to the meatballs has a significant effect on the level of acceptance of meatball texture ( $p = 0.00$ ). The average value of the assessment level of the hedonic test by the panelists on the texture of meatballs ranges from 2.00 to 3.16 (dislike-neutral). The highest value (most preferred) for the texture assessment was obtained from the *tempeh gembus* meatballs with the *tempeh gembus* formulations of 25%. Whereas, the smallest value (disliked and highly disliked) was obtained on the *tempeh gembus* meatballs with the *tempeh gembus* formulations of 100% and 75%. Factors that influence the tenderness value are connective tissue and marbling fat contained in the product, as well as the temperature which has an influence on the binding capacity of water by meat protein, cooking shrinkage, pH and meat juice content.<sup>44</sup>

The elasticity of 25% meatballs is alleged due to the high binding capacity of water from meat. Beef has 66% moisture content per 100g of ingredients. Whereas, the steaming *tempeh gembus* has a water content of 65.22%.<sup>27</sup> The binding capacity of water can be defined as the ability of the meat to maintain its water content during external treatments such as cutting, heating, grinding, and processing. Protein contents also become an influencer, because the higher the protein content, the higher the water bound, thus the meatballs will be more elastic.<sup>42</sup>

### Determination of Selected Formula<sup>39</sup>

The best products are selected by giving weighting to the test results on *tempeh gembus* meatballs. The first thing to do is to give weight to the results of the hedonic test conducted by Mansyur (2017). Weighting is given to four indicators which are scored corresponding to their interests according to the panelists. The taste and flavor are given the highest weighting, which is 40% because taste and flavor are the most important indicators in determining the first acceptance of a product.<sup>43</sup> Furthermore, the color and texture are given a weight of 10%.

Table 9: Hedonic Test Attribute Weighting Results

Indicators	0%	25%	50%	75%	100%
Color	1.48	1.4	1.21	1.39	1.37
Taste	1.29	1.12	1.21	1.25	1.11
Flavor	1.48	1.4	1.21	1.39	1.37
Texture	0.29	0.32	0.26	0.2	0.25
Over all	4.54	4.24	3.79	4.23	4.10

The first rank is obtained by a hedonic test with a control treatment or 0%, while the second rank is obtained by a hedonic test with a treatment of 25%. Because in this study the researchers sought the best substitution formulation, the researchers took the second rank with 25% treatment.

The next step after weighting the results of the hedonic test is weighting of each formula according to the results of the test of protein content, calcium content, and protein digestibility. The formula that has the highest value of the test done will get the highest score.

Table 10: Selected Formulation Determination Scores

Formula	Hedonic Test (40%)	Score	Protein Content (20%)	Score	Calcium Content (20%)	Score	Protein Digestibility (20%)	Score	Score (100%)	Scoring Result
0%	4	1.6	5	1	3	0.6	4	0.8	4	
25%	5	2	4	0.8	2	0.4	1	0.2	3.4	Selected Formula
50%	1	0.4	3	0.6	1	0.2	2	0.4	1.6	
75%	3	1.2	2	0.4	4	0.8	3	0.6	3	

100%	2	0.8	1	0.2	5	1	5	1	3	
------	---	-----	---	-----	---	---	---	---	---	--

The formulation with the highest value is the treatment of 0% and 25%. However, because the researchers wanted to find the best treatment for the substitution of *tempeh gembus*, the treatment of meatballs with the substitution of 25% *tempeh gembus* can be used as an alternative in making local food-based snacks. Meatballs formulation of 25% has a protein content of 8.03%, calcium content 351.19 mg/100g, and protein digestibility of 53.22%.

The results of the hedonic quality assessment show that meatball formulation of 25% has characteristics such as gray which is favored by panelists (3.80), taste that is considered neutral by panelists (2.80), meatball flavor preferred by panelists (3.50), and a little compact texture that is considered neutral by panelists (3.17).

Furthermore, the results of the analysis of protein and calcium content from the formulation were 25% compared to the Nutritional Label Reference (NLR). NLR for the general category according to PKBPOM Number 9 of 2016 concerning Reference to Nutritional Label Reference is 60 g protein and 1100 mg for calcium.<sup>46</sup> Below is a comparison of the results of the analysis of % NLR

Table 11: Energy Content, Protein, Calcium per Serving Dose

Components	Analysis Result	% NLR (%)
Protein (g/100g)	8.03	13.4
Calcium (mg/100g)	351.19	31.9

#### Analysis of the Contribution of Nutritional Content to DV and NRL

The serving dose of the *tempeh gembus* meatball products is determined based on the number of the daily value (DV) in adolescents 16-18 years. The serving dose is determined to assume the meatballs are consumed as a snack with a minimum percentage of 10%.

Table 12: Contribution of Nutritional Content to DV and NRL

Components	Nutritional Content (per serving 78 g)	% DV in Adolescents 16-18 Years		% NLR
		Boys 16-18 Years	Girls 16-18 Years	
Energy (kkal)	270	10%	13%	13%
Protein (g)	6.3	10%	11%	11%
Calcium (mg)	273.9	23%	23%	25%

From the results above, it can be seen that the fulfillment of the DV and NLR from 25% *tempeh gembus* meatballs is only about 10-20% of the daily requirement. Therefore, other additional foods are needed to meet the needs of calcium and protein for body needs in a day.

#### Conclusion

The interaction of substitution of *tempeh gembus* with meat content in *tempeh gembus* meatballs has an effect on protein contents and protein digestibility but had no effect on increasing calcium meatball contents for body needs in a day.

Based on an analysis of nutrient content, protein digestibility, and organoleptic tests formulation with 25% *tempeh gembus* substitution was found to be a best formulation of meatballs containing protein content of 8.03%, calcium contents of 351.19 mg / 100 g, and protein digestibility of 53.22%. It is therefore important that advances in development of *tempeh gembus* combination that has significant value in increased the calcium content in addition to the increase the protein content, protein digestibility and acceptability of *tempeh gembus* meatballs.

#### Suggestions

Need to do further research to increase the level of preference for the taste of *tempeh gembus* meatballs.

#### Acknowledgment

The authors are thankful to the Research Institute of Diponegoro University which facilitated and funded this research.

## Funding

The author(s) received financial support for the research, authorship, and/or publication of this article from Research Institute of Diponegoro University with number of grant: 474-71/UN7.P4.3/PP/2018.

## Conflict of Interest

The authors do not have any conflict of interest.

## References

- Oleson C.V., Morina A. B. Osteoporosis Rehabilitation: A Practical Approach. *Springer*; 2017. 5 p. <https://doi.org/10.1007/978-3-319-45084-1>
- Kementerian Kesehatan RI. Data dan Kondisi Penyakit Osteoporosis di Indonesia. Jakarta; 2015.
- Nafilah, Fitranti D. Y. Hubungan Indeks Massa Tubuh (IMT), Persen Lemak Tubuh, Asupan Zat Gizi, dan Aktivitas Fisik dengan Kepadatan Tulang pada Remaja Putri. *Nutrition College*. 2014;3(4):680-688.
- Faizah L. N., Fitranti D. Y. Hubungan Asupan Protein, Fosfor, dan Kalsium dengan Kepadatan Tulang pada Wanita Dewasa Awal. *Nutrition College*. 2015;4(2):335-341.
- Fikawati S., Syafiq A., Puspasari P. Faktor-Faktor yang Berhubungan dengan Asupan Kalsium pada Remaja di Kota Bandung. *Jurnal Kedokteran Trisakti Universa Medicina*. 2005;24(1):24-34.
- Mahan L. Kathleen., and Sylvia Escott-Stump. Krause's Food & Nutrition Therapy. 12th ed. Canada: *Saunders Elsevier*; 2008.
- Harold N Rosen, MD. Patient education: Calcium and Vitamin D for Bone Health (Beyond the Basics). 2015. Available from: <http://www.uptodate.com/contents/calcium-and-vitamin-d-for-bone-health-beyond-the-basics>
- Damardjati D., Marwoto D.K., Swastika D., Arsyad, Hilman Y. Prospek dan Arah Pengembangan Agribisnis Kedelai. Jakarta; 2005.
- Rahmawati Widi Deshinta. Pengaruh Substitusi Tepung Ampas Tahu terhadap Tingkat Kekerasan dan Daya Terima Bakso. [Skripsi]. Surakarta: Universitas Muhammadiyah Surakarta; 2016.
- Damanik R. N. S., Pratiwi D. Y. W., Widyastuti N, Rustanti N, Anjani G, Afifah D. N. Nutritional Composition Changes during *Tempeh Gembus* IOP Conf Ser Earth Environ Sci. 2018;116(1). doi:10.1088/1755-1315/116/1/012026
- Murdiati, A., Sardjono, Amaliah. Perubahan Komposisi Kimia Tempe Gembus yang Dibuat dari Bahan Dasar Ampas Tahu dan Bekatul. *Agritech*. 2002;20(2):106-110.
- Afifah D. N., Nugrahani G., Hastuti V. N., Arifan F. The Characteristics of Kerupuk Gembus The Characteristics of Kerupuk Gembus. *IOP Conf Ser Earth Environ Sci*. 2019;292(1). doi:10.1088/1755-1315/292/1/012055
- Noviana A., Dieny F. F., Rustanti N., Anjani G., Afifah D. N. Antimicrobial Activity of *Tempeh Gembus* IOP Conf Ser Earth Environ Sci. 2018;116(1). doi:10.1088/1755-1315/116/1/012044
- Kurniasari R, Sulchan M, Afifah DN, Anjani G, Rustanti N. Influence Variation of Tempe Gembus (an Indonesian Fermented Food) on Homocysteine and Malondialdehyde of Rats fed an Atherogenic Diet. *Rom J Diabetes, Nutr Metab Dis*. 2017;24(3):203-211. doi:10.1515/rjdnmd-2017-0026
- Afifah D. N., Sulchan M., Syah D., Yanti, Suhartono M. T., Kim J. H. Purification and Characterization of A Fibrinolytic Enzyme from *Bacillus pumilus* 2.g isolated from gembus, an Indonesian fermented food. *Prev Nutr Food Sci*. 2014;19(3):213-219. doi:10.3746/pnf.2014.19.3.213
- Afifah D. N., Sulchan M., Syah D., Yanti, Suhartono M. T. Isolation and Identification of Fibrinolytic Protease-Producing Microorganisms from Red Oncom and Gembus, Indonesian Fermented Soybean Cakes. *Malaysian Journal of Microbiology*. 2014;10(4):273-279.
- Afifah D. N., Rustanti N., Anjani G., Syah D., Yanti, Suhartono M. T. Proteomics Study of Extracellular Fibrinolytic Proteases from *Bacillus licheniformis* RO3 and *Bacillus pumilus* 2.g isolated from Indonesian Fermented Food. *IOP Conf Ser Earth Environ Sci*. 2017;55:1-10.
- Dewi P. K., Afifah D. N., Rustanti N., Sulchan M., Anjani G. The Effect of *Tempeh Gembus* Variations to Serum Levels of High Sensitivity C-Reactive (hsCRP) and Serum Levels of Fibrinogen of Sprague Dawley Rats with Atherogenic Diet. *Rom J Diabetes, Nutr Metab Dis*. 2018;25(1):91-97. doi:2478/rjdnmd-2018-0010.
- Afifah D. N., Nabilah N., Supraba G. T., Pratiwi S. N., Nuryanto., Sulchan M. The Effect of *Tempeh Gembus*, an Indonesian Fermented Food, on Lipid Profiles in Women with Hyperlipidemia. *Current Nutrition & Food Science*. 2018;14(1):1-9. doi:10.2174/1573401314666180807112549
- Agustina R. K., Dieny F. F., Rustanti N., Anjani G., Afifah D. N. Antioxidant activity and soluble protein content of *tempeh gembus* *Hiroshima J Med Sci*. 2018;67:1-7.
- Sulchan M., N.W, Endang. Nilai Gizi dan Komposisi Asam Amino Tempe Gembus serta Pengaruhnya terhadap Pertumbuhan Tikus. *Maj Kedokt*. 2007;57(3):80-85.



22. Bastian F., Ishak, Tawali, Bilang M. Daya Terima dan Kandungan Zat Gizi Formula Tepung Tempe dengan Penambahan Semi Refined Carrageenan (SRC) dan Bubuk Kakao. *Aplikasi Teknologi Pangan*. 2013;2.
23. Alakali JS, Kucha CT, Rabiou IA. Effect of drying temperature on the nutritional quality of Moringa oleifera leaves. *African J Food Sci*. 2015;9(7):395-9.
24. Badan Standar Nasional. Standar Nasional Indonesia: Bakso Daging. Jakarta: BSN;2014. 2-3 p.
25. Purwanto A., Ali A., Herawati N. Kajian Mutu Gizi Bakso Berbasis Daging Sapi dan Jamur Merang (*Volvariella volvaceae*). *Teknologi Hasil Pertanian*. 2015;14(2):1-8.
26. Sayudi S., Herawati N., Ali A. Potensi Biji Lamtoro Gung dan Biji Kedelai sebagai Bahan Baku Pembuatan Tempe Komplementasi. *Jom Faperta*. 2015;1(2):1-9.
27. Afifah D.N., Rahma A., Nuryandari S.S., Alvice L., Hartono P.I., Kurniawati D.M., Wijayanti H.S., Fitranti D.Y., Purwanti R. Nutrition Content, Protein Quality, and Antioxidant Activity of Various Tempeh Gembus Preparations. *Journal of Food and Nutrition Research*. 2019; 7(8):605-612. doi:10.12691/jfnr-7-8-8.
28. Sumiati T. Pengaruh Pengolahan terhadap Mutu Cerna Protein Ikan Mujair (*Tilapia mossambica*). [Skripsi]. Bogor: Institut Pertanian Bogor; 2008.
29. Takarina H. L. Kandungan Kalsium dan Karbohidrat Bakso Daging Sapi dengan Penambahan Jamur Tiram (*Pleurotus* sp). [Skripsi]. Surakarta: Universitas Muhammadiyah Surakarta; 2013.
30. Gropper S. S., Smith J. L., Groff J. L. Advanced Nutrition and Human Metabolism. 5th ed. USA: Cengage Learning; 2009. 431 p.
31. Gandjar I. G., Rahman A. Kimia Farmasi Analisis. Yogyakarta: Pustaka Pelajar; 2007. p. 240-62, 463-74.
32. Muchtadi T. R., Sugiyono, Ayustaningwarno F. Ilmu Pengetahuan Bahan Pangan. Bandung: Alfabeta; 2010.
33. Rancangan Teknologi Proses Pengolahan Tapioka dan Produk-Produknya. Malang: Universitas Brawijaya; 2007.
34. Wibowo S. Budidaya Bawang Merah, Bawang Putih, Bawang Bombay. Jakarta: Penebar Swadaya; 2008.
35. National Nutrient Database for Standard Reference of Raw Garlic [Internet]. 2010. Available from: <https://ndb.nal.usda.gov/ndb/foods/show/301861>
36. National Nutrient Database for Standard Reference of Raw Pepper [Internet]. 2018. Available from: <https://ndb.nal.usda.gov/ndb/foods/show/302197>
37. National Nutrient Database for Standard Reference of Raw Baking Soda [Internet]. 2018. Available from: <https://ndb.nal.usda.gov/ndb/foods/show/45016339>
38. Saputra D. Penentuan Daya Cerna Protein In Vitro Ikan Bawal (*Colossoma macropomum*) pada Umur Panen Berbeda. *Food Technology*. 2014;5(2):1127-1133.
39. , Kusharto C. M. Daya Cerna Protein Serta Kandungan Mineral (Kalsium dan Fosfor) pada Snack-Bar Substitusi Tepung Pisang (*Musa paradisiaca*) dengan Tepung Pury Ulat Sutera (*Bombyx mori*) dan Tepung Lele (*Clarias gariepinus*). [Skripsi]. Bogor: Institut Pertanian Bogor; 2017.
40. , Wulandari Z., Suryati T. Suplementasi Tepung Putih Telur untuk Memperbaiki Nilai Nutrisi Snack Ekstrusi Berbahan Grits Jagung. *Media Peternak*. 2009;32(3):179-184.
41. Negara J. Aspek Mikrobiologis serta Sensori (Rasa, Warna, Tekstur, Aroma) pada Dua Bentuk Penyajian Keju yang Berbeda. *Ilmu Produksi dan Teknologi Hasil Peternakan*. 2016;4(2):286-290.
42. Firahmi N., Dharmawati S., Aldrin M. Sifat Fisik dan Organoleptik Bakso yang Dibuat dari Daging Sapi dengan Lama Pelayuan Berbeda. *Sains dan Teknologi*. 2015.
43. Hatta M. Pengaruh Level dan Waktu Penambahan Fosfat (Sodium Tripolifosfat/ STTP) terhadap Kualitas Bakso. *Jurnal Agrisistem*. 2011;7:87-95.
44. Ilmu dan Teknologi Daging. Yogyakarta: Gadjah Mada University Press; 2009.
45. Murni M. The Effect of Addition Tempeh Flour to The Quality and the Taste of Chicken Nugget. *Berita Litbang Industri*. 2014;3(2):117-23.
46. PKBPOM. Acuan Label Gizi. Jakarta; 2016.



This work is licensed under a Creative Commons Attribution 4.0 International License.

## Links

About

Archives

## Contact Us

Your Name (required)

## License



This work is licensed under a Creative

[Article Processing Charges](#)[Coming Issue](#)[Contact](#)[Current Issue](#)[Order Print Issue](#)[Submission](#)[Conflict of Interest](#)[Editorial Policy](#)

Your Email (required)

Your Message

**Y F 4 B**

Type the above text in box below (Case sensitive)

Complete the quiz

2+2=?

**Send**

Follow us on:



Commons Attribution 4.0 International License.

## Recent Articles

[A Cross Sectional Study on Analysis of Human milk as per Prakriti \(Psychosomatic Constitution\).](#)[The Study of Sorghum \(Sorghum bicolor L.\), Mung Bean \(Vigna radiata\) and Sago \(Metroxylon sagu\) Noodles: Formulation and Physical Characterization](#)[Effects of Frozen Storage on Viability of Probiotics and Antioxidant Capacities of Synbiotic Riceberry and Sesame-Riceberry Milk Ice Creams](#)