

# The Agar Production, Pigment and Nutrient Content in Gracilaria sp. Grown in Two Habitats with Varying Salinity and Nutrient Levels

*by Yudiati Et Al.*

---

**Submission date:** 04-May-2022 11:39AM (UTC+0700)

**Submission ID:** 1827829753

**File name:** Lamp.\_V\_C-4.pdf (298.86K)

**Word count:** 5238

**Character count:** 28235

# The Agar Production, Pigment and Nutrient Content in *Gracilaria* sp. Grown in Two Habitats with Varying Salinity and Nutrient Levels

Ervia Yudiati<sup>1,2,\*</sup>, Annisa Afifah Nugroho<sup>1</sup>, Sri Sedjati<sup>1</sup>, Zaenal Arifin<sup>3</sup>, Ali Ridlo<sup>1</sup>

<sup>1</sup>Department of Marine Science, Faculty of Fisheries and Marine Science, Diponegoro University, Jl. Prof. Soedarto, SH Tembalang, Semarang, 50275; <sup>2</sup>Laboratory of Marine Micro/Macroalgae Technology, Department of Marine Science, Faculty of Fisheries and Marine Science, Diponegoro University, Jl. Prof. Soedarto, SH Tembalang, Semarang, Indonesia 50275; <sup>3</sup>Brackishwater Aquaculture Development Centre, Jl. Cik Lanang, Jepara, Central Java, Indonesia

Received: March 22, 2020; Revised: Oct 10, 2020; Accepted: Nov 12, 2020

## Abstract

*Gracilaria* sp. is rich in agar, pigment, carbohydrate and mineral contents. This study aims to determine the content of agar, chlorophyll a, carotenoids and nutrient content (protein, carbohydrate, total lipid, ash, water) in *Gracilaria* sp. grown in different habitats with high and low salinity. Samples were brought from high salinity (44±0.33) ppt from the reservoir habitat and low salinity (26.98±0.15 ppt) from biofilter's shrimp waste pond habitat. Salinity and water quality parameters were observed in three different subsites. Agar was extracted by alkali methods. Agar spectra were compared to standard agarose and characterized by FT-IR analysis. Results showed that agar content (25.79 ± 0.28%), chlorophyll a (25.79 ± 0.28 mg/g), carotenoid (5.90 ± 0.07 µmol/g) on low salinity was significantly higher (P<0.05) than high salinity. In high salinity, the agar (6.6±0.34%), chlorophyll a (3.62±0.15, carotenoids (0.71±0.1 µmol/g) contents was lower. The protein level of low salinity (14.4 ± 0.70%) was also significantly higher compared to high salinity (8.15±0.25%), respectively. The total lipid, carbohydrate, water, and ash content were similar. FT-IR analysis spectra show the presence of 3,6-anhydro-L-galactose. These can be useful data concerning optimum salinity in *Gracilaria* sp. culture.

**Keywords:** Salinity, *Gracilaria* sp., Agar, Pigment, Nutrient content

## 1. Introduction

Naturally, *Gracilaria* sp. is a euryhaline macroalga (Kumar *et al.*, 2010), has grown all over the coast with high economic value due to its nutrient content (Du *et al.*, 2016; Hernandez, 2017). *Gracilaria/Gracilariopsis* have been mostly cultivated in Asia, especially Indonesia and China. The production is about 98% of global production (FAO, 2016; Kim *et al.*, 2017). Asian people consume seaweed daily (Cikos *et al.*, 2018). Seaweed empirically improves health and able to reduce the chronic disease incidence such as cancer, cardiovascular and heart diseases (Rioux *et al.*, 2017; Xu *et al.*, 2017). *Gracilaria* sp. serves as an excellent source of polysaccharide, i.e. agar (Xu *et al.*, 2017) also rich in pigmented antioxidants such as chlorophyll and carotenoids (Stengel *et al.*, 2015; Asih *et al.*, 2019). Polysaccharides from seaweed are frequently related to pharmacological activities (Hamed *et al.*, 2015) such as anticoagulant, antioxidant (Yudiati *et al.*, 2018a; 2018b), antitumor, and immunomodulatory of shrimp (*Litopenaeus vannamei*) (Yudiati *et al.*, 2016, 2019) as well as Zebrafish (Yudiati *et al.*, 2020)

Common salinity at sea in marine waters is around 35 ppt. Precipitation or freshwater influxes, as in reservoir and mangrove areas, may lead to salinity variation from 10 to 70 (Graham & Wilcox, 2000; Kumar *et al.*, 2010). Some

former researchers have reported the response of estuarine macroalgae to the abiotic factors such as salinity light, pH, temperature, nutrient load (Kumar *et al.*, 2010; Choi *et al.*, 2010) associated to agar yield (Israel *et al.*, 1999) and photosynthetic performance (Phooprong *et al.*, 2007). Later studies have also discussed the possible effects of environmental stresses on seaweed extracts according to its kinetic parameters (Deyab, 2016).

Salinity has been displayed to the reason of osmotic (Kumar *et al.*, 2010) as well as turgor pressure regulation (Pereira *et al.*, 2017). Salinity will improve the upregulation and accumulation of the essential enzymes (Odat, 2018). The seaweed tolerance to high salinity is supported by the internal and external osmotic capacities and on elasticity of the cell wall (Wu *et al.*, 2018). When the cell is located in hypertonic solution, water will flow quickly out of the cell, and turgor pressure will be decreased affecting plasmolysis, which is commonly permanent. If the cell is located in hypotonic solution, water will go into the cell, causing an enlarge of cell volume. Salinity in plants stimulates the ROS generation which produces cellular oxidative damage when overproduced in large amounts (Luis *et al.*, 2018). These mechanisms play a role in combating the accumulation of reactive oxygen species (ROS) by a diverse set of enzymes such the superoxide dismutase (SOD), which dismutase the O<sub>2</sub>-radicals to H<sub>2</sub>O<sub>2</sub> (Luis *et al.*

\* Corresponding author e-mail: eyudiati@gmail.com.

*al.*, 2018). *Gracilaria* sp. is abundant in chlorophyll a and carotenoids. The pigment important role is to neutralize the free radicals (Sedjati *et al.*, 2020). In some periods of time, these mechanisms influence the agar, pigment as well as protein and mineral content caused by salinity stress. The objective of this study is to determine the agar, pigment (chlorophyll a and carotenoids) and nutrient content (water, protein, total lipid, ash, carbohydrate) from low (reservoir) and high (biofilter habitats) salinity of *Gracilaria* sp.

## 2. Materials and Methods

### 2.1. Water Quality Parameters

Measurements of water quality parameters such as salinity (1) (Go Refractometer), pH (pH meter RoHS), dissolved oxygen and water temperature (Water Quality Checker "Amstast") nitrate and phosphate content was administered. Assessment on nitrate content (SNI-06-2480-1991) was applied spectrophotometrically (Shimadzu 1900) while the phosphate content was done by Badan Standarisasi Nasional methods (SNI-06-2412-1991).

### 2.2. Site Location and Sample Preparation

Sample was collected from low salinity/LS (reservoir) and shrimp (*Litopenaeus vannamei*) pond waste (biofilter) (18) which represents high salinity/HS of Center of Brackishwater and Aquaculture Development Center, Jepara, Central Java, Indonesia (Figure 1). Coordinates of sampling sites in HS was S 06°35'12.74" E 110°39'10.32" while in LS was S 06°35'05.64" E 110°38'49.03". The salinity of the reservoir from this study tends to be mild-low salinity. In contrast, biofilter shallow waters illustrate the different condition. The water supply of biofilter came from shrimp (*L. vannamei*) pond waste from uneaten feed and shrimp's fecal. Due to the static and unchanged water supply, biofilter media was relatively high in salinity.

The water and *Gracilaria* sp. sample was taken at three different subsites from two different habitats at 800 cm in depth. Seaweed was taken randomly. Soon as arrived (10) at the laboratory, samples were rinsed with tap water to clean up from debris and salt. This was followed by dried up samples indoor with room temperature. Dried *Gracilaria* sp. was cut off to 0.5 cm and then stored in a clean and dry pack and sealed with aluminium foil



**Figure 1.** Site location of *Gracilaria* sp. samples (reservoir and shrimp pond biofilter).

### 2.3. Pigment Extraction

Pigment extraction was administered by single extraction. Ethyl acetate was used (1:10) to extract the dried seaweed for 24 hr maceration at room temperature (Hidayati *et al.*, 2019). Whatman no. 41 was used to sieve the extract. The extract was then concentrated with a rotary evaporator (Bucchi) and stored (3) a refrigerator and ready to use. Percentage of yield was calculated using the formula:

$$\text{Yield (\%)} = \frac{\text{weight of extract (g)}}{\text{weight of seaweed}} \times 100\%$$

### 2.4. Determination on Chlorophyll a and Carotenoids Contents

Chlorophyll a and carotenoids (Dere, 1993; Harborne, 1984) content were done spectrophotometrically (Shimadzu 1800). Five g of extract was diluted with acetone p.a. (1:1) The absorbance of samples was recorded (31) 645 nm, 663 nm dan 470 nm wavelength. The chlorophyll a and carotenoids content was determined based on the (1) formulation

$$\text{Chlorophyll a } \mu\text{g/g sample (Ca)} = 12.21 \times A_{663} - 2.81 \times A_{646}$$

$$\text{Carotenoids } \mu\text{mol/g sample (Cx+c)} =$$

$$\frac{A_{470} + 0.114 \times A_{663} - 0.638 \times A_{646}}{V} \times 1000$$

$$112.5 \times 0.1 \times 10$$

### 2.5. Agar extraction

Agar extraction methods were basically done by Jayasinghe *et al.*, (2016) with some modifications. Fifty grams of dried seaweed was extracted for 1.5 hr in hot water (85°C) added with 750 mL NaOH 5%. The extract was then rinsed with continuous tap water for discoloration. This was followed by aquadest 750 mL and homogenized. Acetic acid (CH<sub>3</sub>COOH) was then added, stirred and boiled at ± 90-95°C for two hrs. The agar was then sieved, dropped with KCl 6 g, homogenized and poured into the container. Finally, 18 hrs later, the gel was performed and yield (%) was counted.

### 2.6. Nutrient Content

(23) The proximate analysis was referred to as the Association of Official Analytical Chemists (AOAC, 2005) and conducted to 5 g samples.

#### 2.6.1. Water content

Sample was put into the oven (105°C) for three hrs. Dried samples were then placed into desiccator until the weight was constant.

$$\text{Water content (\%)} = \frac{a-b}{c} \times 100\%$$

a = cup + dried sample  
b = cup without sample  
c = initial sample weight

#### 2.6.2. Protein Analysis

Protein analysis was done by Kjeldahl methods, by located samples into Kjeldahl vial and destructed using 20 mL of hot sulfuric acid. The process was continued until samples were colorless and clear. This was then diluted and distilled with 10 mL of NaOH 10%. The distillate was then put into 25 mL of 3% H<sub>3</sub>BO<sub>3</sub> and titered with HCl standard using methyl red as an indicator. The volume of titrants was used to calculate the percentage of total

nitrogen. The protein level was counted by multiplied the total nitrogen and correction factor.

Total nitrogen % = mL HCl x NHCl/sample weight x 14008 x f

Total protein % = total nitrogen x 6.25

denoted : f = correction factor (6.25).

#### 2.6.3. Total Lipid Content

Total lipid analysis was done by soxhlet methods. Dried samples were wrapped with cotton wool. Sample was then placed into soxhlet extractor, using diethyl ether as solvent. Reflux was conducted by the samples until done. Vial was then put into the oven (105°C) and weight. The percentage of total lipid was determined by this formula:

Total lipid (%) = Lipid weight/Sample weight x 100%

#### 2.6.4. Ash Content

Sample was weighed and burned at the top of bunsen until smoke produced and put in the muffle furnace at 500 - 600°C until turned to ash formation. Cup was cooled down and weighed. The ash content was counted using formula:

Ash content (%) = ash weight/sample weight x 100%

#### 2.6.5. Total Carbohydrate

Total carbohydrate was counted simply by this equation:

Total carbohydrate (%) = 100% - (% ash + % water + % protein + % total lipid).

#### 2.6.6. Characterization of Agar

Agar characterization was determined by Fourier Transform Infrared (FT-IR) spectroscopy (Thermo Nicolet 380 FTIR, Germany). Samples were mixed with KBr pellets (10% w/w). Similar to alginate, the pellet was recorded at 4000-500 cm<sup>-1</sup> (Yudiati and Isnansetyo, 2017).

### 3. Statistical Data Analysis

All data surveys were analyzed non parametrically (Mann Whitney test) using SPSS version 20.0 computer software. The laboratory data were analyzed using One Way ANOVA (Analysis of variance) using the same software, with a 95% level of significance.

## 4. Results

#### 4.1. Water Quality Parameters


Salinity, nitrate and phosphate contents and others water quality parameter LS and HS with three subsites are shown in Table 1. There was a differences in salinity, nitrate and phosphate contents, while other parameters were similar.

**Table 1.** Water quality parameters of *Gracilaria* sp. grown in low salinity (reservoir) and high salinity (pond waste biofilter)

Parameter	LS	HS
Temperature (°C)	29.74 ± 0.15 <sup>a</sup>	29.70 ± 0.03 <sup>a</sup>
Salinity (ppt)	26.89 ± 3.15 <sup>a</sup>	44.00 ± 0.33 <sup>b</sup>
pH	8.0 ± 0 <sup>a</sup>	8.3 ± 0 <sup>a</sup>
Dissolve Oxygen (ppm)	3.55 ± 0.72 <sup>a</sup>	3.45 ± 0.05 <sup>a</sup>
Nitrate (mg/L)	0.43 ± 0.56 <sup>b</sup>	1.77 ± 1.44 <sup>a</sup>
Phosphate (mg/L)	0.78 ± 0.26 <sup>b</sup>	0.29 ± 0.19 <sup>a</sup>

#### 4.2. Yield of pigment extraction, chlorophyll a and carotenoids content

Yield, chlorophyll a and carotenoids content in reservoir and shrimp pond biofilter is shown in Figure 1. Yield were similar, chlorophyll a and carotenoids of *Gracillaria* sp. sample in reservoir was higher.




**Figure 2.** Yield, Chlorophyll and Carotenoids content of *Gracillaria* sp. grown in high and low salinity. Different superscript indicates a significantly difference (p<0.05).

#### 4.3. Agar Percentage and Characterization

Percentage of agar in *Gracillaria* sp. samples from LS and HS is shown at Table 2. Agar characterization of sample and standard agarose (Merck, USA) is depicted in Figure 3a, b and Tab 22. Concentration of agar sample from LS habitat was significantly different compared to HS (p<0.05). On the other hand, pairwise comparison spectra of agar either from low or high salinity were fit to the standard.

**Table 2.** Percentage agar yield of *Gracillaria* sp. samples from low salinity (reservoir) and high salinity (biofilter)

Habitat	Agar (% w/w)
LS	25.79 ± 0.28
HS	6.60 ± 0.34




**Figure 3.** FT-IR spectra of agar from *Gracilaria* sp. from low salinity, LS (a) and high salinity, HS (b)

**Table 3.** The vibration signal of Standard, Low Salinity (LS) and High Salinity (HS) of Agar ( 1500- 400 cm<sup>-1</sup>)

Location	Type of Bonding	Wave Number (cm <sup>-1</sup> )	
Standar Agarose	O-H	930.65 bending	
	C-H	1378.56 deformation	
		891.50 bending	
		772.41 bending	
	C-O	1159.50 stretching	
		1073.15 stretching	
	LS	O-H	932.72 bending
		C-H	1419.95 bending
			891.27 bending
			662.67 bending
C-O	1156.23 stretching		
	1074.22 stretching		
HS	O-H	933.27 bending	
	C-H	1417.73 bending	
		1277.30 stretching	
		659.39 bending	
	C-O	1074.93 stretching	

4.4. Nutrient Content

Percentage of protein, carbohydrate, total lipid, water and ash of *Gracilaria* sp. samples from LS and HS presented in Figure 4. Protein content of samples from low salinity is significantly different from the samples from high salinity (p<0.05). On the other hand, other concentrations were similar.



**Figure 4.** Protein, carbohydrate, lipid, water and ash contents of *Gracilaria* sp. grown in LS and HS

5. Discussion

5.1. Pigment Yield, Chlorophyll a and Carotenoids content of *Gracilaria* sp. from LS and HS

In terms of pigment extraction yield, the extraction of *Gracilaria* sp. from LS and HS resulted in a similar yield (± 2,8%) (Fig. 2). This value was possibly derived from other source of pigments based on water extraction from phycobiliprotein groups. Kumar *et al.* (2010) reported that decline concentration of chlorophyll a and carotenoids was in accordance with higher allophycocyanin (APC) and phycoerythrin (PE) content in hyper-salinity (45-50 ppt). Additionally, the similar researchers stated that the increment approximately from 52% and up to 70% from initial contents. This finding was also in agreement with Pereira *et al.* (2017) that high salinity reducing the chlorophyll contents and depigmentation of apices.

As shown in similar figure, photosynthetic pigments i.e. chlorophyll a and carotenoids content of *Gracilaria* sp. samples from the LS (26.98 ppt) was significantly higher than HS (44 ppt) (p<0.05). Chlorophyll is a compound with free electrons structure from nitrogen atoms (Sedjati *et al.*, 2020). Meanwhile, Martinez *et al.*, 2010 stated that carotenoids neutralize free radicals in three ways which is electron transfer, addition of radical species and the hydrogen abstraction. This indicated that low salinity was more tolerable. So, the physiology was relatively normal. Moreover, the phenomenon of oxidative stress, osmotic and turgor pressure regulation might not appear. In addition, chlorophyll a content from the Indonesian North Java coast in this study (14.81 µg/g) is distinctly higher compared to the study from Thondi Coast, India (8.96 µg/g) (Rosemary *et al.*, 2019). In addition, visually, the colour green intensity of pigment from low salinity was much brighter and darker.

In general, visually, *Gracilaria* sp. thallus in this present study from the LS were thicker and bigger that might be indicated the better growth. Pereira *et al.* (2017), Wong and Chang (2000) reported similar results. Moreover, *Gracilaria tenuistipitata* from Songkhla Lagoon in Thailand showed the best growth rates at the salinity of 25 psu (Bunsom and Prathep., 2012). The growth of macroalgae in the marine ecosystem is often decreased in hypersaline water. This is due to the cumulative enzyme effects such as reduction of



turgor pressure, this then leads to restrict division of cells and at last, affected the growth (Lee and Liu, 1999).

Nitrate content at HS in this study was higher than LS (Table 1). This is probably caused by instant efflux nutrient from the shrimp pond. Pond's feed waste and shrimp faecal contribute the nitrate content. Nutrient factor highly fluctuates on photosynthesis especially nitrate and phosphate (Ismail and Osman, 2016). Research by Wu *et al.*, (2018) reported that the nitrate and phosphorus uptake were higher at lower salinities (less than 20 psu) than higher salinity conditions (6 to 20 psu). Study from Choi *et al.* (2010) exhibits that nitrate and phosphate uptake of macroalgae were greater in certain levels of salinity (20 and 25 ppt). These reports demonstrate similar suggestion to this study. Moreover, the water supply in LS from this study was more diverse, either from upside regimes from the sea as well as freshwater influxes from the river. Table 1 noted that phosphate content in LS was higher. The role of phosphate pathway is by transferring the high adenosine triphosphate (ATP) energy and other high energy compounds. This occurs in respiration and photosynthetic process (Ismail and Osman, 2016). These researchers' results were similar to our previous data, that photosynthetic pigments (chlorophyll a and carotenoids) of *Gracilaria* sp. in LS was higher when compared to the HS habitat (data is not shown).

#### 5.2. Agar Percentage and Characterization

Data from Table 2 pointed that agar percentage of *Gracilaria* sp. in LS was significantly higher than HS. This is probably due to a relationship between salinity and the agar content. Research by Sasikumar *et al.* (1999) stated that a high salinity of *G. verrucosa* in summer (43.8‰) was found to be negatively correlated with agar yield. The best agar production of *Gracilaria tenuistipitata* ( $24.8 \pm 3.0$  %DW) in laboratory was found at 25 psu (Bunsom and Prathep., 2012). Rocha *et al.* (2018) stated that, often, yield of agar of *G. tikvahiae* has been clearly connected with salinity and adversely with nitrogen content, similarly reported to *G. gracil* species (Martin *et al.*, 2013). Nitrogen concentration in LS, in fact, is lower than HS (Table 1). Less concentration of nitrogen, synthesis of protein declines in favor of polysaccharide synthesis. However, other factors such as nutrient availability, environmental and geographic factors, seasonal variations can influence the yield, chemistry and biosynthesis of agar (Lahaye and Rochas, 1993). Moreover, Lee *et al.* (2017) also stated that it is not easy to investigate the effects of a single factor on the yield of agar extracted from seaweeds grown in the natural habitat.

Based on spectra of FT-IR analyses, (Figure 2) shows that two *Gracilaria* sp. samples are fit to the standard agarose (Merck) and positively fingerprinted at a specific agar (1500-400  $\text{cm}^{-1}$ ) with galactose bond (around 1070  $\text{cm}^{-1}$ ). The vibration signal is slightly different, but overall those are agar characterizations. Agarose has a basic repeating unit of 1,3-linked  $\beta$ -D-galactopyranose and 1,4-linked 3,6-anhydro- $\alpha$ -L-galactopyranose. Based on figure 3 above, there is a 3,6 anhydrogalactose unit at a wave number of 928-933  $\text{cm}^{-1}$  (Hii *et al.*, 2015). Observed bands at 930  $\text{cm}^{-1}$  indicates O-H bending. The hydroxyl (O-H) unit appears at 3400  $\text{cm}^{-1}$  wavenumber. Alkene ( $\text{CH}_3$  or  $\text{CH}_2$ ) associated to metoxil appeared at 2900  $\text{cm}^{-1}$ .

Aldehyde group (-CHO) markedly appears at 1600  $\text{cm}^{-1}$  (Pereira *et al.*, 2009).

Based on Figure 4, it can be seen that protein content in *Gracilaria* sp. samples from LS was significantly higher compared to HS ( $p < 0.05$ ). Protein percentage is related to the nitrate content. In high salinity, nitrate compound has not taken up, easily (Choi *et al.*, 2010). However, in special cases, nutrient uptake can be occasionally imbalanced during unfavorable periods of situations (Trimmer *et al.*, 2000).

Results from Table 1 show that phosphate compound from the LS habitat was higher, and this may force the production of protein in the algal cell. Energy from photosynthesis will be used for amino acid biosynthesis that comes from the surrounding water. ATP is energy synthesized by photosynthesis, and these surely need phosphate. The reduction of protein synthesis triggered the decrease of the protein content and this was then affected the other cell components such as chlorophyll and other pigments (Ismail and Osman, 2016). This phenomenon was performed in *Gracilaria* sp. from HS habitat. Compared to other research, protein content in this study (14.40%) was higher than *G. changii* from Sarawak, Malaysia (12.57%) (Chan and Matanjun, 2016). Even though, this protein content was lower compared to the study on *G. corticata* by Rosemary *et al.* (2019).

Carbohydrate percentages form *Gracilaria* sp. samples grown in LS and HS were similar ( $p > 0.05$ ) (Figure 4). Carbohydrate content in this study was higher than *G. changii* (Chan and Matanjun, 2016) and *G. edulis* (Rosemary *et al.*, 2019). Total lipid, water and ash content of *Gracilaria* sp. in both samples was not significantly different ( $p > 0.05$ ). Generally, total lipid from all macroalgae, including *Gracilaria* sp. in this research, was relatively low at the range of 0.9-40% (Khairy and El Shafay, 2013). Seaweed is rich in minerals. The high ash content indicates high mineral content. In accordance to our finding, the results from Wu *et al.* (2018) reported that nutrient uptake, tissue nutrient contents were affected by salinity and the ideal salinity was around 20 psu. This information could be beneficial to define optimal salinity in *Gracilaria* sp. culture systems to get the maximal agar yield and pigment quality, protein and mineral content. This initiates better economic incomes for agarophyte farming.

#### 6. Conclusion

Agar, chlorophyll a, carotenoids, protein and mineral contents of *Gracilaria* sp. grown from low salinity habitat (reservoir) were higher compared to high salinity habitat (biofilter shrimp ponds). Other compounds such as the total lipid, carbohydrate, water, and ash content from high and low salinity were similar. In this study, it can be concluded that salinity affected *Gracilaria* sp. nutrient content. Similar to the standard agar, FT-IR spectra from different salinity show the existence of 3,6-anhydro-L-galactose. Our findings can be useful for considering the application of *Gracilaria* sp. culture in the future.

## References

- AOAC. 2005. Official Methods of Analysis of the Association of Official Analytical Chemists. Benjamin Franklin Station, Washington D.C.
- Asih T, Khayuridlo M, Noor R and Muhfahroyin. 2019. Biodiversity and Potential Use of Macro Algae in Pesisir Barat Lampung. *Biosaintifika*, **11**(1): 100-107.
- Badan Standarisasi Nasional. 1991. SNI 06-2412-1991. Metode Pengambilan Contoh Uji Kualitas Air.
- Badan Standarisasi Nasional. 1991. SNI 06-2480-1991. Cara Uji Nitrat Secara Spektrofotometri.
- Bunsom C and Prathep A. 2012. Effects of salinity, light intensity and sediment on growth, pigments, agar production and reproduction in *Gracilaria tenuistipitata* from Songkhla Lagoon in Thailand. *Phycological Research*, **60**: 169-178
- Chan PT, and Matanjun P. 2016. Chemical Composition and Physicochemical Properties of Tropical Red Sea Weed, *Gracilaria changii*. *Food Chem*, **(16)**: 1-37. Cikos AM, Jokic S, Subaric D and Jerkovic I. 2018. Overview on the application of modern methods for the extraction of bioactive compounds from marine macroalgae. *Mar Drugs*, **16**(10): 348-368.
- Dere S, Gunes T and Sivaci R. 1998. Spectrophotometric determination of chlorophyll-a, b, and total carotenoid content of some algae species using different solvent. *J of Botany*, **22**:13-17.
- Dayay MA. 2016. Inhibition activity of Seaweed extract for mild carbon steel corrosion in saline formation water. *Desalination*, **385**: 60-67.
- Du Q, Guiqi B, Yunxiang M and Zhenghong S. 2016. The complete chloroplast genome of *Gracilariopsis lemaneiformis* (Rhodophyta) gives new insight into the evolution of family Gracilariaceae. *J Phycol*, **52** (3): 441-450
- FAO. 2016. *The State of World Fisheries and Aquaculture 2016 – Contributing to food security and nutrition for all*. Rome.
- Graham LE and Wilcox LW. 2000. Algae. Prentice Hall, New York. (640pp.)
- Hamed I, Özogul F, Özogul Y and Regenstein JM. 2015. Marine bioactive compounds and their health benefits: A review. *Compr Rev Food Sci Food Saf*, **14**: 446-465
- Harborne JB. 1984. **Phytochemical Methods: A Guide to Modern Techniques of Plant Analysis**. second ed. Chapman and Hall, London.
- Hernandez A. 2017. The Effect of Salinity on the Growth of the Red Alga, *Gracilaria epihippisor*. Marine Science Department University of Hawai'i at Hilo.
- Hidayati JR, Yudiati E, Pringgienis D, Anrif Z and Oktaviyanti DT. 2019. *Sargassum* sp. Extract Macerated In Different Solvents Polarity. *Jurnal Kelautan Tropis*, **22** (1): 73 – 80.
- Hii SL, Lim JY, Ong WT and Wong CL. 2015. Agar From Malaysian Red Seaweed as Potential Material for Synthesis of Bioplastic Film. *Journal of Engineering Science and Technology Special Issue on SOMCHE*, : 1 – 15.
- Ismail MM and Osman MEH. 2016. Seasonal fluctuation of photosynthetic pigments of most common red seaweeds species collected from Abu Qir, Alexandria, Egypt. *Revista de Biologia Marina y Oceanografia*, **51**(3): 515-525.
- Israel A, Martinez-Goss M and Friedlander M. 1999. Effect of salinity and pH on growth and agar yield of *Gracilaria tenuistipitata* var. *liui* in laboratory and outdoor cultivation. *J Appl Phycol*, **11**: 543-549.
- Jayasinghe P, Pahalawattarachchi S and Ranaweera KKDS. 2016. Effect of Extraction Methods on the Yield and Physicochemical Properties of Polysaccharides Extracted from Seaweed Available in Sri Lanka. *Poultry, Fisheries and Wildlife Sciences*, **4**(1): 1 - 6.
- Khairy HM and El-Shafay SM. 2013. Seasonal Variations In The Biochemical Composition of Some Common Seaweed Species From The Coast of Abu Qir Bay, Alexandria, Egypt. *Oceanologia*, **55**(2):435-452.
- Kim JK, Yarish C, Hwang EK, Park M and Kim Y. 2017. Seaweed aquaculture: cultivation technologies, challenges and its ecosystem services. *ALGAE*, **32**(1): 1-13.
- Kumar M, Kumari P, Gupta PV, Reddy CRK and Jha B. 2010. Biochemical responses of red alga *Gracilaria corticata* (Gracilariales, Rhodophyta) to salinity induced oxidative stress. *J Exp Mar Bio Ecol*, **391**: 27-34
- Lahaye M. and Rochas C. 1991. Chemical-structure and physicochemical properties of agar. *Hydrobiologia*, **221**: 137-148
- Lee, TM and Liu, CH. 1999. Correlation of decreased calcium contents with proline accumulation in the marine green macroalga *Ulva fasciata* exposed to elevated NaCl contents in seawater. *J Exp Bot*, **50**:1855-1862.
- Lee W-K, Lim Y-Y, Leow AT-C, Namasivayam P, Abdullah JO and Ho C-L. 2017. Factors affecting yield and gelling properties of agar. *J Appl Phycol*, **29**:1527-1540 DOI 10.1007/s10811-016-1009-y
- Luis A, Corpas FJ, López-Huertas E and Palma JM. 2018. **Plant Superoxide Dismutases: Function Under Abiotic Stress Conditions**. *Antioxidants and Antioxidant Enzymes in Higher Plants*. Springer, pp1-26.
- Martin LA, Rodriguez MC, Matulewicz MC, Fissore EN, Gerschenson LN and Leonardi PI. 2013. Seasonal variation in agar composition and properties from *Gracilaria gracilis* (Gracilariales, Rhodophyta) of the Patagonian coast of Argentina. *Phycological Research*, **61**(3): 163-171
- Odat N. 2018. Differential Gene Expression of Durum Wheat (*Triticum turgidum* L. var. durum) in Relation to Genotypic Variation under NaCl Salinity Stress. *Jordan Journal of Biological Science*, **11**(5): 591-595
- Phooprang S, Ogawa H and Hayashizaki K. 2007. Photosynthetic and respiratory responses of *Gracilaria salicornia* (C. Agardh) Dawson (Gracilariales, Rhodophyta) from Thailand and Japan. *J Appl Phycol*, **19**: 795-801.
- Pereira L, Amando AM, Critchley AT, Velde FVD and Ribeiro PJA. 2009. Identification of Selected Seaweed Polysaccharides (Phycocolloids) by Vibrational Spectroscopy (FTIR-ATR and FT-Raman). *Food Hydrocol*, **300**: (1-7).
- Pereira, DT, Simioni C, Filipin EP, Bouvie F, Ramlov F, Maraschin M., Bouzon ZL and Schmidt EC. 2017. Effects of salinity on the physiology of the red macroalga, *Acanthophora spicifera* (Rhodophyta, Ceramiales). *Acta Bot Brasilica*, **31**(4):555-565
- Rioux LE, Beaulieu L and Turgeon SL. 2017. Seaweeds: A traditional ingredients for new gastronomic sensation. *Food Hydrocoll*, **68**: 255-265.
- Rocha CMR, Sousa AMM, Kim JJ, Magalhães JMCS, Yarish C and Gonçalves MP. 2018. Characterization of agar from *Gracilaria tikvahiae* cultivated for nutrient bioextraction in open water farms. *Food Hydrocoll*, doi: <https://doi.org/10.1016/j.foodhyd.2018.10.048>.
- Rosemary T, Anulkumar A, Paramasivam S, Mondragon A and Miranda JM. 2019. Biochemical, Micronutrient and Physicochemical Properties of the Dried Red Seaweeds *Gracilaria edulis* and *Gracilaria corticata*. *Molecules*, **24** (2225): 1-5.

- Sasikumar C, Rao VNR and Rengasamy R. 1999. The effect of environmental factors on the qualitative and quantitative characteristics of agar from the marine red alga *Gracilaria verrucosa* (Gracilariales, Rhodophyta). *Indian J Mar Sci*, **28**:270–273
- Sedjati S, Pringgenies D and Fajri M. 2020. Determination of the Pigment Content and Antioxidant Activity of the Marine Microalga *Tetraselmis suecica*. *Jordan Journal of Biological Sciences*, **13**(1): 55-58
- Stengel DB and Connan S. 2015. Marine algae: a source of biomass for biotechnological applications, in: D.B. Stengel, S. Connan (Eds.), *Natural Products From Marine Algae - Methods and Protocols*, Humana Press, New York, NY, 2015, pp. 1–37, [https://doi.org/10.1007/978-1-4939-2684-8\\_1](https://doi.org/10.1007/978-1-4939-2684-8_1).
- Trimmer M, Nedwell DB, Sivyer DB and Malcolm SJ. 2000. Seasonal organic mineralisation and denitrification in intertidal sediments and their relationship to the abundance of *Enteromorpha* sp. and *Ulva* sp. *Mar Ecol Prog Ser*, **203**:67-80.
- Wong SL and Chang J. 2000. Salinity and light effects on growth, photosynthesis, and respiration of *Grateloupia filicina* (Rhodophyta). *Aquaculture*, **182**:387–395
- Wu H, Shin SK, Jang S, Yarish C and Kim JK. 2018. Growth and Nutrient Bioextraction of *Gracilaria chorda*, *G. vermiculophylla*, *Ulva prolifera*, and *U. compressa* Under Hypo- and Hyper-Osmotic Conditions. *ALGAE*, **33**(4): 329-340.
- Xu SY, Xuesong H and Kit-Leong C. 2017. Recent Advances in Marine Algae Polysaccharides: Isolation, Structure, and Activities. A review. *Mar Drugs*, **15** (388) : 1-16
- Xu SY, Huang X and Cheong KL. 2017. Recent Advances in Marine Algae Polysaccharides: Isolation, Structure, and Activities. *Mar Drugs*, **15** (388): 1-16
- Yudiati E and Isnansetyo A. 2017. Characterizing the Three Different Alginate Type of *Sargassum siliquosum*. *ILMU KELAUTAN: Indonesian Journal of Marine Sciences*, **22**(1):7-14
- Yudiati E, Isnansetyo A, Ayuningtyas, Handayani CR, Murwantoko and Triyanto. 2016. Innate immune-stimulating and immune genes up-regulating activities of three types of alginate from *Sargassum siliquosum* in Pacific white shrimp, *Litopenaeus vannamei*. *Fish & Shellfish Immunol*, **54**: 46-53.
- Yudiati E, Santosa GW, Tontowi MR, Sedjati S, Supriyantini E and Khakimah M. 2018a. Optimization of alginate alkaline extraction technology from *Sargassum polycystum* and its antioxidant properties. *IOP Conf. Ser.: Earth Environ. Sci.* **139**. 012052. doi :10.1088/1755-1315/139/1/012052
- Yudiati E, Pringgenies D, Djunaedi A, Arifin Z and Sudaryono A. 2018b. Free radical Scavenging of Low Molecular Weight Sodium Alginate (LMWSA) from *Sargassum polycystum* Produced by Thermal Treatment. *Aquac Ind*, **19** (1): 21-27.
- Yudiati E, Isnansetyo A, Murwantoko, Triyanto and Handayani CR. 2019. Alginate from *Sargassum siliquosum* Simultaneously Stimulates Innate Immunity, Upregulates Immune Genes, and Enhances Resistance of Pacific White Shrimp (*Litopenaeus vannamei*) Against White Spot Syndrome Virus (WSSV). *Marine Biotechnology* **21**(4):503-514. <https://doi.org/10.1007/s10126-019-09898-7>
- Yudiati, E, Rustadi, Ginzel FI, Hidayati JR., Rizfa MS, Azhar N, Djarod MSR, Heriyati E, and Alghazeer R. 2020. Oral Administration of Alginate Oligosaccharide from *Padina* sp. Enhances Tolerance of Oxygen Exposure Stress in Zebrafish (*Danio rerio*). *ILMU KELAUTAN: Indonesian Journal of Marine Sciences* **25**(1):7-14.



# The Agar Production, Pigment and Nutrient Content in Gracilaria sp. Grown in Two Habitats with Varying Salinity and Nutrient Levels

## ORIGINALITY REPORT

15%

SIMILARITY INDEX

11%

INTERNET SOURCES

11%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://ejournal.undip.ac.id">ejournal.undip.ac.id</a> Internet Source	4%
2	<a href="http://www.e-algae.org">www.e-algae.org</a> Internet Source	2%
3	<a href="http://link.springer.com">link.springer.com</a> Internet Source	2%
4	Submitted to University of East London Student Paper	1%
5	Submitted to University of Strathclyde Student Paper	<1%
6	<a href="http://pdfs.semanticscholar.org">pdfs.semanticscholar.org</a> Internet Source	<1%
7	Kumar, M.. "Biochemical responses of red alga <i>Gracilaria corticata</i> (Gracilariales, Rhodophyta) to salinity induced oxidative stress", <i>Journal of Experimental Marine Biology and Ecology</i> , 20100815 Publication	<1%

8

Jonathan A. Hillman. "A homotopy fibration theorem in dimension four", *Topology and its Applications*, 1989

Publication

<1 %

9

Wei-Kang Lee, Yi-Yi Lim, Adam Thean-Chor Leow, Parameswari Namasivayam, Janna Ong Abdullah, Chai-Ling Ho. "Factors affecting yield and gelling properties of agar", *Journal of Applied Phycology*, 2016

Publication

<1 %

10

[ejournal2.undip.ac.id](http://ejournal2.undip.ac.id)

Internet Source

<1 %

11

Sang Sook Kim, Han Sub Kwak, Mi Jeong Kim. "The effect of various salinity levels on metabolomic profiles, antioxidant capacities and sensory attributes of doenjang, a fermented soybean paste", *Food Chemistry*, 2020

Publication

<1 %

12

Vivek Kumar. "Effect of inland saline water ionic profiles on growth, chemical composition and agar characteristics of *Gracilaria cliftonii* (Withell, Miller and Kraft 1994) under laboratory conditions", *Aquaculture International*, 12/24/2009

Publication

<1 %

13 Xiao Zhang, Jude Juventus Aweya, Zong-Xun Huang, Zhuo-Ying Kang et al. "In vitro fermentation of Gracilaria lemaneiformis sulfated polysaccharides and its agaro-oligosaccharides by human fecal inocula and its impact on microbiota", Carbohydrate Polymers, 2020  
Publication

---

14 [digilib.library.usp.ac.fj](http://digilib.library.usp.ac.fj)  
Internet Source

---

15 [www.mdpi.com](http://www.mdpi.com)  
Internet Source

---

16 "Sixteenth International Seaweed Symposium", Springer Science and Business Media LLC, 1999  
Publication

---

17 C. B. Martins, O. Ferreira, T. Rosado, E. Gallardo, S. Silvestre, L. M. A. Santos. "Eustigmatophyte strains with potential interest in cancer prevention and treatment: partial chemical characterization and evaluation of cytotoxic and antioxidant activity", Biotechnology Letters, 2021  
Publication

---

18 Ervia Yudiati, D P Wijayanti, N Azhar, A I Chairunnisa, S Sedjati, Z Arifin. "Alginate oligosaccharide/polysaccharide and lactic acid

bacteria (*Lactobacillus bulgaricus* FNCC-0041 & *Streptococcus thermophilus* FNCC-0040) as immunostimulants against pathogenic *Vibrio* spp. using *Artemia* bio model", IOP Conference Series: Earth and Environmental Science, 2021

Publication

---

19

Ping Zhong, Zhongneng Xu, Wei Chen, Kai Peng et al. "Molecular characterization of a c-Jun NH2-terminal kinase (JNK)-interacting protein 4 (Lv-JIP4) in *Litopenaeus vannamei* and its potential role in the Lv-p38MAPK pathway in response to low temperature", *Aquaculture Reports*, 2021

<1 %

Publication

---

20

Qikun Xing, Guiqi Bi, Min Cao, Arnaud Belcour, Méziane Aite, Zhaolan Mo, Yunxiang Mao. "Comparative transcriptome analysis provides insights into the response of to the fluctuating salinity conditions ", *Journal of Phycology*, 2021

<1 %

Publication

---

21

Xiang Li, Achen Wang, Wenjie Wan, Xuesong Luo, Liuxia Zheng, Guangwen He, Daqing Huang, Wenli Chen, Qiaoyun Huang. "High Salinity Inhibits Soil Bacterial Community Mediating Nitrogen Cycling", *Applied and Environmental Microbiology*, 2021

<1 %

Publication

---

22	<a href="https://docsdrive.com">docsdrive.com</a> Internet Source	<1 %
23	<a href="https://eprints.ucm.es">eprints.ucm.es</a> Internet Source	<1 %
24	<a href="https://repositorium.sdum.uminho.pt">repositorium.sdum.uminho.pt</a> Internet Source	<1 %
25	<a href="http://www.scilit.net">www.scilit.net</a> Internet Source	<1 %
26	"Antioxidants and Antioxidant Enzymes in Higher Plants", Springer Science and Business Media LLC, 2018 Publication	<1 %
27	"Bioactive Molecules in Food", Springer Science and Business Media LLC, 2019 Publication	<1 %
28	Cristina M.R. Rocha, Ana M.M. Sousa, Jang K. Kim, Júlia M.C.S. Magalhães, Charles Yarish, Maria do Pilar Gonçalves. "Characterization of agar from Gracilaria tikvahiae cultivated for nutrient bioextraction in open water farms", Food Hydrocolloids, 2019 Publication	<1 %
29	Ervia Yudiati, Alim Isnansetyo, Murwantoko, Triyanto, Christina Retna Handayani. "Alginate from Sargassum siliquosum Simultaneously Stimulates Innate Immunity, Upregulates	<1 %



Immune Genes, and Enhances Resistance of Pacific White Shrimp (*Litopenaeus vannamei*) Against White Spot Syndrome Virus (WSSV)", *Marine Biotechnology*, 2019

Publication

30

[researchonline.jcu.edu.au](http://researchonline.jcu.edu.au)

Internet Source

<1 %

31

Palas Samanta, Sojin Jang, Sookkyung Shin, Jang Kyun Kim. "Effects of pH on growth and biochemical responses in *Agarophyton vermiculophyllum* under different temperature conditions", *Journal of Applied Phycology*, 2019

Publication

<1 %

32

Wei-Kang Lee, Yi-Yi Lim, Chai-Ling Ho. "pH affects growth, physiology and agar properties of agarophyte *Gracilaria changii* (Rhodophyta) under low light intensity from Morib, Malaysia", *Regional Studies in Marine Science*, 2019

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On