

# Correlation between Nutric Score and Adequacy of Energy and Protein Intake with Duration of Mechanical Ventilation in the Intensive Care Unit Dr. Kariadi Hospital, Semarang-Indonesia

*by* Enny Probosari

---

**Submission date:** 01-Nov-2022 09:57AM (UTC+0700)

**Submission ID:** 1941049469

**File name:** Lampiran\_C1\_-\_kiky\_nafilia.PDF (283.65K)

**Word count:** 3558

**Character count:** 18739

## Correlation between nutric score and adequacy of energy and protein intake with duration of mechanical ventilation in the intensive care unit Dr. Kariadi Hospital, Semarang-Indonesia



Ika Sutrisnawati<sup>1</sup>, Darmono<sup>2</sup>, Etisa Adi Murbawani<sup>2\*</sup>, Niken Puruhita<sup>2</sup>, Enny Probosari<sup>2</sup>

### ABSTRACT

**Background:** Malnutrition in the ICU patients increases morbidity, mortality, cost, and mechanical ventilation (MV) duration. Nutric score has been validated as a screening tool in critically ill patients to identify patients at risk of malnutrition. The pathophysiology of malnutrition in critically ill is caused by severity of disease and inflammation resulting in hypercatabolism. Delivering protein-energy intake will reduce hypercatabolism and maintain lean body mass that affects outcome. The study aims to analyze the correlation between nutric score and adequacy of energy and protein intake with MV duration.

**Methods:** This cross sectional study enrolled 65 subjects  $\geq 18$  years with mechanical ventilation in the ICU of Dr. Kariadi Hospital Semarang during May to July 2020. The nutric score was determined within 24 - 48 hours. Statistical analysis used correlation test.

**Results:** The mean nutric score was  $2.3 \pm 1.49$ , duration of MV was  $3.9 \pm 4.44$ . There was significant correlation between nutric score and duration of MV ( $r = 0.685$ ;  $p < 0.001$ ). The mean adequacy of energy intake was  $1448.7 \pm 235.87$  and protein intake was  $57.9 \pm 11.94$ . There was significant negative correlation between adequacy of energy intake and MV duration ( $r = -0.246$ ;  $p = 0.048$ ). There was significant negative correlation between protein adequacy intake and MV duration ( $r = -0.34$ ;  $p = 0.006$ ).

**Conclusion:** Nutric score was correlated with duration of MV in the ICU patients. Adequacy of energy and protein intake had significant correlation with duration of MV.

**Keywords:** critically ill, duration of MV, malnutrition, nutric score, nutritional screening.

**Cite This Article:** Sutrisnawati, I., Darmono., Murbawani, E.A., Puruhita, N., Probosari, E. 2021. Correlation between nutric score and adequacy of energy and protein intake with duration of mechanical ventilation in the intensive care unit Dr. Kariadi Hospital, Semarang-Indonesia. *Bali Medical Journal* 10(1): 392-396. DOI: 10.15562/bmj.v10i1.2196

<sup>1</sup>Resident of Clinical Nutrition, Faculty of Medicine, Universitas Diponegoro - Dr. Kariadi Hospital, Semarang, Indonesia.

<sup>2</sup>Clinical Nutrition Department, Faculty of Medicine, Universitas Diponegoro - Dr. Kariadi Hospital, Semarang, Indonesia.

\*Corresponding author:  
Etisa Adi Murbawani;  
Clinical Nutrition Department, Faculty of Medicine, Universitas Diponegoro - Dr. Kariadi Hospital, Semarang, Indonesia;  
etisatitits@gmail.com

Received: 2021-01-19

Accepted: 2021-04-20

Published: 2021-04-30

### INTRODUCTION

The duration of mechanical ventilation (MV) is one of the outcome parameters used to evaluate care quality in the intensive care unit (ICU). Malnutrition in critically ill patients increases morbidity, nosocomial infections, cost, prolonged MV and mortality. The prevalence of malnutrition in intensive care ranged from 38% to 78%. It is very important to make the diagnosis of malnutrition in ICU patients by using validated screening tools.<sup>1,2</sup> Nutritional screening is crucial in ICU patients therefore nutritional screening should be done to determine patients at risk of malnutrition. It should describe severity of disease and inflammation

which caused malnutrition.<sup>2-4</sup>

Critically ill patients associated with hypercatabolism caused by inflammation that will worsen nutritional status. The catabolic process will decrease muscle mass which causes several complications such as increased morbidity, infection, multi-organ dysfunction, failure of weaning, length of stay, and increased mortality. Nutritional therapy is needed to prevent catabolism due to metabolic stress and oxidative cell injury modulating the immune response.<sup>4,5</sup> Energy and protein achievement is crucial need for ICU patients. Energy delivery prevents catabolic state, gluconeogenesis, and glycaemic control. The benefits of protein to achieve the synthesis of acute-phase

response proteins related to immune response and muscle protein synthesis.<sup>6</sup>

Nutric score was found by Heyland et al.<sup>7</sup> in England as a screening tool in the ICU by considering that previous screening tools such as NRS (Nutritional Risk Score) 2002 and SGA (Subjective Global Assessment) met the difficulties such as energy-protein intake history, functional status before treatment, gastro intestinal symptoms before ICU admission, and changes in body weight. This difficulty is due to the majority of patients using mechanical ventilation (MV) and sedation, difficult to collect recall of energy-protein intake. It is an objective and quantitative parameter and supports the role of nutrition in patients at

risk of malnutrition in critically ill patients who benefit from nutritional therapy interventions.<sup>4-7</sup>

## METHOD

This cross-sectional study was conducted on 65 subjects admitted to the ICU Dr. Kariadi Hospital Semarang during May to July 2020. The inclusion criteria were mechanically ventilated patients aged  $\geq 18$  years who were admitted to the ICU with the criteria for medical and surgical patients using MV. The Nutric score of subjects who met the inclusion criteria was assessed within 24 - 48 hours admitted in ICU. The variables of the nutric score are age, APACHE score (Acute Physiology Health Disease Classification System II), SOFA score (Sequential Organ Failure Assessment), level of acute inflammation (IL-6) with Human IL-6 ELISA Kit, comorbid, days of hospitalization prior to ICU. Patients died in ICU were excluded. Patients were classified for high-risk ICU patients with nutric score  $\geq 6$  and low nutritional scores (0-5).

Nutritional adequacy was recorded while the patient was in the ICU. Energy and protein target requirements are based on ASPEN guidelines (American Society Parenteral and Enteral Nutrition) with energy 25-30 kcal/kg/day and protein 1.2-2 g/kg/day. The Adequacy of energy and protein was calculated at the last hospitalization in the ICU.

Data analysis was performed using Spearman correlation test to analyze the correlation between the nutric score and adequacy energy and protein intake with duration of MV. All values considered significant if  $p < 0.05$ .

## RESULTS

Most of the subjects were 39 females (60%). The subjects' mean age was  $44.7 \pm 13.73$  years old with 27 subjects were  $\geq 50$  years old (41.5%). The subjects' mean BMI was  $22.14 \text{ kg/m}^2$  with 25% were underweight and 24.6% were obese (Table 1).

There were 63 (96.9%) subjects with low nutritional score (nutric score 0-5) and 2 subjects were at high risk of malnutrition (nutric score  $\geq 6$ ) were 2 subjects (3.1%). Most of the subjects, 45 subjects with a percentage of 75.8%, had an APACHE II

**Table 1. Demography Characteristics and Body Mass Index (BMI)**

Variable	n	%	Mean $\pm$ SD	Median (min-max)
Sex				
Male	26	40		
Female	39	60		
Age (years)			$44.7 \pm 13.73$	46 (18 - 72)
<50	38	58.5		
50-<75	27	41.5		
BMI ( $\text{kg/m}^2$ )			$22.1 \pm 4.78$	21,5 (13.84 - 41.4)
< 18,5	16	24.6		
18.5 - 22.9	26	40.0		
23.0 - 24.9	7	10.8		
$\geq 25$	16	24.6		

**Table 2. Characteristic data of nutric score and duration MV**

Variable	n	%	Mean $\pm$ SD	Median (min-max)
Age (years)			$44.7 \pm 13.73$	46 (18 - 72)
< 50	38	58.5		
50 - < 75	27	41.5		
APACHE II			$10.6 \pm 6.07$	9 (4 - 25)
<15	48	73.8		
15-<20	9	13.8		
20-28	8	12.3		
SOFA			$4.7 \pm 2.64$	
<6	48	73.8		
6- <10	13	20.0		
$\geq 10$	4	6.2		
Comorbid			$0.8 \pm 0.87$	1 (0-2)
0-1	46	70.8		
$\geq 2$	19	29.2		
Days of hospitalization before admission ICU (day)			$6.1 \pm 4.99$	4(0-17)
0-<1	4	6.2		
$\geq 1$	61	93.8		
IL-6 (pg/ml)	65	100	$3.01 \pm 4.14$	1,37 (0.39 - 16.54)
0-<400	0	0		
$\geq 400$			$2.35 \pm 1.49$	2 (1-6)
Nutric				
low (0-5)	63	96.9		
high ( $\geq 6$ )	2	3.1		
Duration of MV (day)			$3.9 \pm 4.44$	2 (1-17)
1-<7	50	76.9		
$\geq 7$	15	23.1		

Abbreviation; sequential organ failure assessment (SOFA); acute physiology and chronic health evaluation II (APACHE II).

score  $\leq 15$ . There were 9 subjects or 9 % subjects who got APACHE score 15 - <20. The mean APACHE II score was 10.65.

The mean SOFA score was <6. The mean duration of MV was  $3.9 \pm 4.44$  days. There were 50 subjects (76.9%) with duration

**Table 3. Characteristic data-target energi- protein dan Adequacy of energy-protein intake**

Variable	Mean± SD	Median (min-max)
Energy target requirement (kcal)	1537.4 ± 89.98	1530 (1140 – 1980)
Protein target requirement (gram)	63.6 ± 10.43	65 (42 – 86)
Adequacy of energy intake (kcal)	1448.7 ± 235.87	1500 (980 – 1980)
Adequacy of protein intake (gram)	57.9 ± 11.94	57 (34-86)

Abbreviation; kilo calories (kcal)

**Table 4. Correlation between nutric score and adequacy energy protein intake and mechanical ventilation**

Variable	VM	
	p-value	r
Nutric score	<0.001	0.685
Adequacy of energy intake (kcal)	0.048	-0.246
Adequacy of protein intake (gram)	0.006	-0.340

Abbreviation; coefficient correlation (r)

of MV <7 and 15 subjects (15%) with duration of MV ≥ 7 (Table 2).

Table 3 showed the mean energy target requirement of this study was 1537 ± 89.89 kcal. The mean protein prescription was 63.6 ± 10.43 grams. The mean adequate protein intake was 57.9 ± 11.94 grams. Table 3 described that most of the samples had received nutrition intervention close related >80% was based on guideline ASPEN 2016 in critically ill patients (Table 3).

This study used the Spearman non-parametric correlation test. Table 4 showed that was significant positive correlation. There was significant correlation between nutric score and duration of MV (r=0.685; p<0.001). There was significant negative correlation between adequacy of energy intake and MV duration (r = -0.246; p = 0.048). There was significant negative correlation between protein adequacy intake and MV duration (r=-0.34; p=0.006) (Table 4).

## DISCUSSION

All patients who were treated for 48 hours in the ICU will be high risk of malnutrition. Nutric score is a screening tool to assess the risk of malnutrition that has been

validated explicitly for ICU patients based on age, disease severity in the form of APACHE score, SOFA score, number of comorbid, length of stay until admission to ICU and IL-6. [8-9] APACHE II and SOFA were used to predict the mortality rate of patients who were admitted to the ICU. Several studies suggested that the APACHE II score to be a predictor of weaning outcome.<sup>10-11</sup>

This study's results were different from previous studies conducted by Rahman et al.<sup>4</sup> and Mendess et al.<sup>3</sup> The difference in previous studies was to describe the correlation between modified nutric score (mnutric) and mortality rate. Rahman et al. described the correlation between adequacy energy and protein intake and 28-day mortality rate. Kalaisevan et al.<sup>8</sup> described the correlation between nutric score and the duration of MV while in this study we analyzed the correlation between the nutric score and the adequacy protein-energy intake and the duration of MV.<sup>3,8</sup>

Table 2 shows that 96.9% subjects of the nutric score were ≤ 6 which meant that it had a low risk of malnutrition. The prolonged MV will have a high risk of malnutrition. Mostly APACHE results are ≤ 15 and SOFA results are < 6. This result presumes that IL-6 concentration

was low. No one subjects concentrations were ≥ 400 pg/dl and influences the result of nutric score.<sup>7,10</sup>

Table 3 showed a significant negative correlation, which meant that each energy and protein supply would reduce MV duration. In this study that achievement of energy intake during in ICU had achieved >80%. This study revealed that the patient received nutritional interventions according to the target energy needs. The energy optimization strategy in the early phase would reduce hyperglycemia and insulin resistance. In some cases, excessive energy intake (overfeeding) was avoided to reduce morbidity of infection, duration of MV use, and length of stay.<sup>11,12</sup> In our study, we did not find an overfeeding subject.

Based on a previous study by Heyland et al.<sup>7</sup>, patients with a nutritional score ≥6 received greater nutritional interventions that affected outcome. Patients who are at high risk of malnutrition will most benefit from early nutritional therapy. Not all ICU patients are the same risk for malnutrition and will benefit equally from nutritional therapy.<sup>7,9,12</sup>

In this study, the mean protein target requirement was found 63 grams. The adequacy protein intake is 57 grams with > 80% of the protein target requirement. Achievement of protein intake of ≥ 80% of the protein target was associated with decreased mortality. Delivering of energy and protein intake in this study was in accordance with the guidelines for providing nutritional therapy in critical patients based on the 2016 ASPEN that most patients had received nutritional interventions according to protein target requirement.<sup>10</sup> This research was different from the study of Rahman et al.<sup>9</sup>, which suggested the correlation between nutritional therapy's adequacy with the 28-day mortality rate and the mnutric score with the mortality rate.

The critical illness causes loss of muscle mass in the first few weeks. Loss of muscle mass decreases respiratory function and thereby prolonged MV. Loss of muscle mass is associated with increased mortality after discharge from hospital. Based on this theory, protein intake must be optimal in critical patients.<sup>12-13</sup>

Recent studies in critical illness have shown protein administration is more closely associated with positive outcomes than energy prescription. A prospective observational study in patients with MV demonstrated protein attainment of 1.3 g/kg bodyweight protein attainment and was associated with a 50% reduction in 28-day mortality. Adequacy protein intake is required to prevent disease-related catabolism or injury in critically ill patients, support protein synthesis and resolve anabolic resistance. There is much controversy regarding optimal energy and protein intake during the initial phase of ICU admission. A recent study by Koekkoek and Looijaard et al. found that giving protein intake of  $\geq 80\%$  of the described value for survival for 60 days and reducing MV duration. Delivering of high protein was controversial in recent studies.<sup>12,14,15</sup>

Adequacy nutritional therapy in the initial phase is only beneficial for patients at high risk of malnutrition, whereas patients at low risk have a good outcome even though they do not receive adequate nutritional therapy for several days.<sup>6</sup> In our study, all subjects had duration of MV  $\leq 7$  days. This positive correlation might occur because most of the study sample had a low risk of malnutrition (63%) and 3.1% had a high risk of malnutrition.

Delivering optimal energy protein intake in critically ill patients was still controversial. Several randomized studies suggest that giving underfeeding during the first week in the ICU provides a better outcome effect. Another study based on large-scale observational data showed that an intake greater than 2/3 of the target requirement was associated with reduced mortality. In a multicentre study conducted by Wei et al.<sup>16</sup> with a cohort study, patients with VM who were given adequate caloric intake in the first week were associated with a better 6-month survival rate and accelerated the recovery phase within 3 months.

The achievement of energy and protein intake depends on hemodynamic instability, vasopressor drugs, hematemesis, and gastrointestinal intolerance disorders, which evaluated gastric residue volume (GRV). Nutric score assessment was carried out 24-48

hours of admission to the ICU so that it only described the physiological condition at the first time of admission but the following day during treatment in the ICU when the patient experienced a decrease in hemodynamics, it was not reidentified so that this condition would affect the adequacy of energy and protein intake to reach the nutrition target via the enteral route was a challenge because early EN was often hindered by delayed gastric emptying or gastrointestinal dysfunction.<sup>17,18</sup>

The limitation of this study was subjected only a few person has nutric score  $\geq 6$ . Therefore, it needs more subjects whom nutric score  $\geq 6$  to see the adequacy of protein energy intake intake between patients at high risk and low-risk malnutrition.

## CONCLUSION

Nutric score was correlated with duration of MV in the ICU patients. Adequacy of energy and protein intake had significant correlation with duration of MV.

## ETHICAL APPROVAL

This study was approved by Ethical Study Committee of Kariadi Hospital with ethical clearance reference number 453/EC/KEPK-RSDK/2020.

## CONFLICTS OF INTEREST

There were no other conflicts of interest in this study.

## FUNDING

No specific funding was provided for this article.

## AUTHOR CONTRIBUTION

Ika Sutrisnawati and Etisa Adi Murbani responsible for project administration, writing the original draft, and data gathering. Darmono, Etisa Adi Murbawani, Niken Puruhita, Enny Probosari for supervision and guiding the main idea.

## ACKNOWLEDGMENTS

This work was supported by Department of Clinical Nutrition, Faculty of Medicine, Universitas Diponegoro Kariadi Hospital Semarang, Indonesia.

## REFERENCES

- Awad A, Bader-El-Den M, McNicholas J. Patient length of stay and mortality prediction: A survey. *Health Serv Manage Res.* 2017;30(2):105-20.
- Lew CCH, Yandell R, Fraser RJL, Chua AP, Chong MFF, Miller M. Association Between Malnutrition and Clinical Outcomes in the Intensive Care Unit: A Systematic Review. *JPEN J Parenter Enteral Nutr.* 2017;41(5):744-58.
- Mendes R, Policarpo S, Fortuna P, Alves M, Virella D, Heyland DK. Nutritional risk assessment and cultural validation of the modified NUTRIC score in critically ill patients-A multicenter prospective cohort study. *Journal of Critical Care.* 2017;37:45-9.
- Rahman A, Hasan RM, Agarwala R, Martin C, Day AG, Heyland DK. Identifying critically-ill patients who will benefit most from nutritional therapy: Further validation of the "modified NUTRIC" nutritional risk assessment tool. *Clinical Nutrition.* 2016;35(1):158-62.
- Canales C, Elsayes A, Yeh DD, Belcher D, Nakayama A, Mc Carthy CM, et al. Nutrition Risk in Critically Ill Versus the Nutritional Risk Screening 2002: Are They Comparable for Assessing Risk of Malnutrition in Critically Ill Patients? *Journal of Parenteral and Enteral Nutrition.* 2019;43(1):81-7.
- Preiser JC, van Zanten AR, Berger MM, Biolo G, Casaer MP, Doig GS, Griffiths RD, Heyland DK, Hiesmayr M, Iapichino G, Laviano A, Pichard C, Singer P, Van den Berghe G, Wernerman J, Wischmeyer P, Vincent JL. Metabolic and nutritional support of critically ill patients: consensus and controversies. *Crit Care.* 2015;19(1):35. doi: 10.1186/s13054-015-0737-8.
- Heyland DK, Dhaliwal R, Jiang X, Day AG. Identifying critically ill patients who benefit the from nutrition therapy: the development and initial validation of a novel risk assessment tool. *Crit Care.* 2011;15(6):R268.
- Kalaiselvan M, Renuka MK, Arunkumar AS. Use of nutrition risk in critically ill (NUTRIC) score to assess nutritional risk in mechanically ventilated patients: A prospective observational study. *Indian Journal of Critical Care Medicine.* 2017;21(5):253-6.
- Rahman A, Hasan RM, Agarwala R, Martin C, Day AG, Heyland DK. Identifying critically-ill patients who will benefit most from nutritional therapy: Further validation of the "modified

- NUTRIC” nutritional risk assessment tool. *Clinical Nutrition*. 2016;35(1):158–62.
10. Jarmula AR, Hombach R, Krzych LJ. APACHE II score predicts mortality in patients required prolonged ventilation in weaning center. *Anesthesiology Intensive Therapy*. 2016;48(4):215–219.
  11. Chung W, Sheu C, Hung J, Hsu T, Yang S, Tsai J. Novel mechanical ventilator weaning predictive model. *The Kaohsiung Journal of Medical Sciences*. 2020;36(10):841–9.
  12. McClave SA, Taylor BE, Martindale RG, Warren MM, Johnson DR, Braunschweig C, et al. Guidelines for the Provision and Assessment of Nutrition Support Therapy in the Adult Critically Ill Patient: Society of Critical Care Medicine (SCCM) and American Society for Parenteral and Enteral Nutrition (A.S.P.E.N.). *JPEN J Parenter Enteral Nutr*. 2016;40(2):159–211.
  13. Sharma K, Mogensen KM, Robinson MK. Pathophysiology of Critical Illness and Role of Nutrition. *Nutrition in Clinical Practice*. 2019;34(1):12–22.
  14. Singer P, Blaser AR, Berger MM, Alhazzani W, Calder PC, Casaer MP, et al. ESPEN guideline on clinical nutrition in the intensive care unit. *Clinical Nutrition*. 2019;38(1):48–79.
  15. Looijaard WGPM, Denneman N, Broens B, Girbes ARJ, Weijs PJM, Oudemans-van Straaten HM. Achieving protein targets without energy overfeeding in critically ill patients: A prospective feasibility study. *Clinical Nutrition*. 2019;38(6):2623–31.
  16. Wei X, Day AG, Ouellette-Kuntz H, Heyland DK. The Association Between Nutritional Adequacy and Long-Term Outcomes in Critically Ill Patients Requiring Prolonged Mechanical Ventilation: A Multicenter Cohort Study. *Critical Care Medicine*. 2015;43(8):1569–79.
  17. Kim H, Stotts NA, Froelicher ES, Engler MM, Porter C. Why patients in critical care do not receive adequate enteral nutrition? A review of the literature. *Journal of Critical Care*. 2012;27(6):702–13.
  18. Flordelis Lasiera JL, Pérez-Vela JL, Montejo González JC. Enteral nutrition in the hemodynamically unstable critically ill patient. *Medicina Intensiva (English Edition)*. 2015;39(1):40–8.



This work is licensed under a Creative Commons Attribution

# Correlation between Nutric Score and Adequacy of Energy and Protein Intake with Duration of Mechanical Ventilation in the Intensive Care Unit Dr. Kariadi Hospital, Semarang-Indonesia

## ORIGINALITY REPORT

13%

SIMILARITY INDEX

8%

INTERNET SOURCES

11%

PUBLICATIONS

1%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="https://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a> Internet Source	1%
2	<a href="https://rbti.org.br">rbti.org.br</a> Internet Source	1%
3	W.G.P.M. Looijaard, N. Denneman, B. Broens, A.R.J. Girbes, P.J.M. Weijs, H.M. Oudemans-van Straaten. "Achieving protein targets without energy overfeeding in critically ill patients: A prospective feasibility study", <i>Clinical Nutrition</i> , 2018 Publication	1%
4	<a href="https://www.mdpi.com">www.mdpi.com</a> Internet Source	1%
5	"Surgical Critical Care Therapy", Springer Science and Business Media LLC, 2018 Publication	1%
6	<a href="https://www.worldwidejournals.com">www.worldwidejournals.com</a> Internet Source	1%

7	<a href="http://www.unboundmedicine.com">www.unboundmedicine.com</a> Internet Source	1 %
8	"Surgical Intensive Care Medicine", Springer Nature, 2016 Publication	<1 %
9	Na Wang, Ping Wang, Wen Li, Li Jiang, Meiping Wang, Bo Zhu, Xiuming Xi. "Prognostic significance of malnutrition risk in elderly patients with acute kidney injury in the intensive care unit", BMC Nephrology, 2022 Publication	<1 %
10	Júlia Lima, Igor Eckert, Maria Cristina Gonzalez, Flávia Moraes Silva. "Prognostic value of phase angle and bioelectrical impedance vector in critically ill patients: A meta-analysis of observational studies", Clinical Nutrition, 2022 Publication	<1 %
11	Zeinab Javid, Niyaz Mohammad Zadeh Honarvar, Majid Khadem-Rezaiyan, Daren K. Heyland et al. "Translation and adaptation of the modified NUTRIC score for critically ill patients", Clinical Nutrition ESPEN, 2021 Publication	<1 %
12	<a href="http://jcancer.org">jcancer.org</a> Internet Source	<1 %



13

Yin-qiu Zhang, Yan Liang, Chun Pan, Yuanyuan Li, Yang Liu, Li-Hua Ren, Wei Xu, Rui-Hua Shi, Yadong Feng. "Emergency digital cholangioscope-assisted one-stage bedside endoscopic stone removal combined with biliary drainage for severe acute cholangitis in intensive care unit: experience from a tertiary Chinese academic hospital", Research Square Platform LLC, 2022

Publication

<1 %

14

[www.ijccm.org](http://www.ijccm.org)

Internet Source

<1 %

15

Mariane Kubiszewski Coruja, Yasmini Cobalchini, Cristiane Wentzel, Jaqueline da Silva Fink. "Nutrition Risk Screening in Intensive Care Units: Agreement Between NUTRIC and NRS 2002 Tools", Nutrition in Clinical Practice, 2019

Publication

<1 %

16

Matteo Luigi Giuseppe Leoni, Elisa Moschini, Maurizio Beretta, Marco Zanello, Massimo Nolli. "The modified NUTRIC score (mNUTRIC) is associated with increased 28-day mortality in critically ill COVID-19 patients: internal validation of a prediction model", Clinical Nutrition ESPEN, 2022

Publication

<1 %

17

Shanti Balasubramanian, Dena H. Tran, Monica Serra, Elizabeth A. Parker et al. "Assessing Calorie and Protein Recommendations for Survivors of Critical Illness Weaning from Prolonged Mechanical Ventilation – Can We Find a Proper Balance?", Clinical Nutrition ESPEN, 2021

Publication

<1 %

18

[jcn.nutrition.or.kr](http://jcn.nutrition.or.kr)

Internet Source

<1 %

19

[oak.ulsan.ac.kr](http://oak.ulsan.ac.kr)

Internet Source

<1 %

20

Camila F. A. Silva, Simone G. de Vasconcelos, Thales A. da Silva, Flávia M. Silva. "Permissive or Trophic Enteral Nutrition and Full Enteral Nutrition Had Similar Effects on Clinical Outcomes in Intensive Care: A Systematic Review of Randomized Clinical Trials", Nutrition in Clinical Practice, 2018

Publication

<1 %

21

Lee-anne S. Chapple, Matthew J. Summers, Luke M. Weinel, Kylie Lange, Woo Han Yang, Adam M. Deane, Marianne J. Chapman. "Muscle size, strength, and physical function in response to augmented calorie delivery: A TARGET sub-study", Journal of Critical Care, 2022

Publication

<1 %

- 22 Tiffany M. N. Otero, D. Dante Yeh, Ednan K. Bajwa, Ruben J. Azocar, Andrea L. Tsai, Donna M. Belcher, Sadeq A. Quraishi. "Elevated Red Cell Distribution Width Is Associated With Decreased Ventilator-Free Days in Critically Ill Patients", *Journal of Intensive Care Medicine*, 2016  
Publication <1 %
- 
- 23 Zheng Yii Lee, Mohd Yusof Barakatun-Nisak, Ibrahim Noor Airini, Daren K. Heyland. "Enhanced Protein-Energy Provision via the Enteral Route in Critically Ill Patients (PEP uP Protocol)", *Nutrition in Clinical Practice*, 2015  
Publication <1 %
- 
- 24 [annalsofintensivecare.springeropen.com](http://annalsofintensivecare.springeropen.com)  
Internet Source <1 %
- 
- 25 [assets.researchsquare.com](http://assets.researchsquare.com)  
Internet Source <1 %
- 
- 26 [www.atsjournals.org](http://www.atsjournals.org)  
Internet Source <1 %
- 
- 27 [www.ijrc.in](http://www.ijrc.in)  
Internet Source <1 %
- 
- 28 [www.tandfonline.com](http://www.tandfonline.com)  
Internet Source <1 %
- 
- 29 Bhuvana Krishna, Shiva K Narayan, Kiran K Gudivada. "Assessment of Nutritional Status <1 %

## in the Critically Ill", Indian Journal of Critical Care Medicine, 2020

Publication

---

30

"Annual Update in Intensive Care and Emergency Medicine 2018", Springer Science and Business Media LLC, 2018

Publication

---

<1 %

31

"Pattern Recognition. ICPR International Workshops and Challenges", Springer Science and Business Media LLC, 2021

Publication

---

<1 %

32

Manoela Lima Oliveira, Daren Keith Heyland, Flávia Moraes Silva, Estela Iraci Rabito et al. "Complementarity of modified NUTRIC score with or without C-reactive protein and subjective global assessment in predicting mortality in critically ill patients", Revista Brasileira de Terapia Intensiva, 2019

Publication

---

<1 %

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On

# Correlation between Nutric Score and Adequacy of Energy and Protein Intake with Duration of Mechanical Ventilation in the Intensive Care Unit Dr. Kariadi Hospital, Semarang-Indonesia

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---