

Identification of agents causing vibriosis in *Litopenaeus vannamei* shrimp culture in Kendal, Central Java, Indonesia and application of microalgae *Dunaliella salina* and *Tetraselmis chui* as bio-control

by Ita Widowati


Submission date: 28-Jul-2022 10:17AM (UTC+0700)

Submission ID: 1876042348

File name: C-4-IW-Ident-AAAL.pdf (106.11K)

Word count: 3564

Character count: 19877



Identification of agents causing vibriosis in *Litopenaeus vannamei* shrimp culture in Kendal, Central Java, Indonesia and application of microalgae *Dunaliella salina* and *Tetraselmis chui* as bio-control agents against vibriosis

¹Ita Widowati, ²Muhammad Zainuri, ³Hemin P. Kusumaningrum, ⁴Yusri Maesaroh, ⁵Yann Hardivillier, ⁵Vincent Leignel, ⁶Nathalie Bourgougnon, ⁵Jean-Luc Mouget

¹ Department of Marine Science, Faculty of Fisheries and Marine Science, Diponegoro University, Tembalang, Semarang, Indonesia; ² Department of Oceanography, Faculty of Fisheries and Marine Science, Diponegoro University, Tembalang, Semarang, Indonesia; ³ Department of Biology, Faculty of Sciences and Mathematic, Diponegoro University, Tembalang, Semarang, Indonesia; ⁴ Marine Science MSc Program, Faculty of Fisheries and Marine Science, Diponegoro University, Tembalang, Semarang, Indonesia; ⁵ Laboratoire du Mer Molécule et Santé (MMS), Faculté des Sciences et Techniques, Le Mans Université, France; ⁶ Laboratoire de Biotechnologie et Chimie Marines (LBCM) Université de Bretagne-Sud, Vannes, France.

Corresponding author: I. Widowati, ita_jusup@yahoo.co.id

Abstract. Aquaculture of Vannamei shrimp at Kendal, Central Java, Indonesia has been using an intensive system, which could provoke the degradation of water quality in the pond. One of the emerging effects due to such degradation becoming threat for shrimp aquaculture is the loss caused by vibriosis disease. This study aimed to identify the presence of vibriosis-causing bacteria in shrimp pond at Kaliwungu, Kendal, Central Java and use the microalgae *Dunaliella salina* and *Tetraselmis chui* as bio control agents against Vibriosis. The Vibriosis-causing bacteria were isolated from the hepatopancreas and tails of ten shrimps showing clinical signs of vibriosis infection, cultured on the Thiosulfate Citrate Bile Salt Sucrose media. Out of 20 isolates, three isolates were selected for further gene sequence identification of 16rRNA using the Repetitive Sequence-based Polymerase Chain Reaction (rep-PCR) and were analyzed with BLAST. Universal primer 27F (5'AGAGTTTGATCMTGGCTCAG-3') and 1492R (5'TACGGTTAACCTTGTTACGACTT-3') were selected as the primers to identify the isolated bacteria. Results demonstrated that the three isolated bacteria were positive as vibriosis-causing agent in shrimp, which were U1H1 AND U4H1, identified as *Vibrio alginolyticus*, and U1E1 as *V. harveyi*. The use of microalgae as biocontrols was performed. Forty-five shrimps infected by vibrio reared during 21 days and feed with *D. salina* and *T. chui* showed a decreased of bacteria amount, which were counted by Total Plate Count method. The result indicated that the microalgae was capable to produce an antibacterial compounds against vibrio.

Key Words: aquaculture, disease, pond water quality, bacteriosis, clinical symptom.

Introduction. Community trend towards the need of fisheries products is constantly increasing, thus, the aquaculture activities are highly developed, and one of them is the Vannamei shrimp culture. This species is a fresh water commodity. The production of Vannamei in Central Java reached 411.729 tons in 2014. From 2010 to 2014, the production of this commodity increased with 20.49% (LAKIP-Directory General of Aquaculture Fisheries, 2014). The augmentation of Vannamei production at Central Java has been generated by the application of intensive systems characterized by spreading high quantity of breed stocks and the usage of 100% of artificial feeds. The application of intensive systems decreases the water quality in ponds making the Vannamei shrimp

susceptible to diseases caused by parasites, virus and bacteria. The mortality of shrimp due to vibriosis infection might reach 80 to 100% in the larval, post larval, juvenile, and adult stages within 1 to 3 days. Case of mass mortality in shrimp aquaculture occurring across every region in Indonesia in the 1900s, including Central Java, had made great loss for most of the companies.

Vibrio sp. has been widely reported to be one of the main pathogenic bacteria causing high mortality rate in shrimp aquaculture (Haldar et al 2011; Heenatigala & Fernando 2016; Solidum et al 2016). The presence of this pathogenic bacteria itself is a part of natural microflora contributing to 60% of total bacteria population as mentioned by Heenatigala & Fernando (2016). As opportunistic organism, the infection of *Vibriosis* to the cultivated organisms occurs when the environmental condition degrades affecting the health condition of the cultivated organism (Kumaran & Citarasu 2016). Bacterial disease outbreak provoked by *Vibriosis* has become constraint on the sustainable production of shrimp (Manilal et al 2010). In order to suppress the outbreak, marine secondary metabolites have been mentioned as promising resources to further develop, especially from microalgae. It has been reported that the extracts of microalgae could prevent the bacterial infections (Cadiz et al 2016; Dash et al 2017).

Despite of the intensive shrimp farming activities in order to fulfill markets demand, infectious disease caused by bacteria remains the main problem that still happens present days. As a consequence, it is crucial to carry out a study that evaluates and identifies the vibriosis-causing bacteria as an update and also serves as basic information prior to further treatment. It is meant to apply the suitable and appropriate treatment to overcome the infection. The objectives of this study were to identify the agents causing vibriosis in the cultivated *Litopenaeus vannamei* and to evaluate the efficacy of microalgae extracts from *Dunaliella salina* and *Tetraselmis chui* in suppressing the vibriosis infection.

Material and Method

Collection of infected Vannamei shrimp. The sampling location of Vannamei shrimp, infected with vibriosis, was collected from the Panggang Ayom village, Kaliwungu district, Kendal, Central Java, Indonesia.

L. vannamei samples had an average weight of 4.30 g and length of 8 cm. Samples was chosen having morphological signs of vibriosis infection (Prayitno & Lachtford 1995). The research was conducted in October 2016 – March 2017. Sample was analyzed at Integrated Laboratory, Diponegoro University, Indonesia.

Isolation of vibriosis-causing bacteria. Tryptone Soya Agar (TSA) as universal agar media and Thiosulfate Citrate Bile Salt (TCBS) (Merck, Jerman) as vibriosis-specific agar media for bacterial culture were selected. Bacteria were isolated from the infected shrimp using Streaking method (Haldar et al 2011).

DNA extraction, sequencing and identification of isolated bacteria. DNA extraction was conducted by using chelex method from Walsh et al (2013) as performed in Susilowati et al (2015). Selected colonies were inoculated in 50-100 µL ddH₂O and 1 mL of 0.5% saponin in PBS 1x (saved overnight). The mixture was centrifuged (12,000 RPM, 10 min). Supernatant was discarded. Then 100 µL ddH₂O and 50 µL of 20% chelex 100 (shake up chelex solution and ensure that some of the crystals make it into sample) were added to a final solution and the solution was boiled for 10 min and vortex once after 5 min. The mixture was centrifuged (12,000 RPM, 10 min) and stored at -20°C. The DNA concentrations were quantified and qualified by using NanoDrop 2000 spectrophotometer (Thermo Scientific). The concentration of 1 µL DNA sample was determined by using the NanoDrop 2000 spectrophotometer (Thermo Scientific). The 260/280 and 260/230 nm ratios was calculated by the NanoDrop spectrophotometer and used to evaluate the DNA purity and also the concentration of DNA.

DNA extracts for 16S rRNA genes sequences were amplified by PCR using universal primers 27F (5'AGAGTTTGATCMTGGCTCAG-3') and 1492 R (5'TACGGTTAACCTTGTTACGACTT-

3'). The PCR mixture consisted GoTaq®Green Master Mix Promega (25 µL), primer 27F (0.5-5 µL), primer 1492R (0.5-5 µL), DNA extract (1-5 µL), and Nuclease-Free Water (50 µL). The PCR reaction was performed in a MJ Mini Personal Thermal Cycler (BIO RAD) using cycling conditions consisting of an initial denaturation at 95°C for 3 min followed by 30 cycles of denaturation at 95°C for 1 min, annealing at 55°C for 1 min, and extension at 72°C for 1 min. A final extension was performed at 72°C for 7 min (Lee et al 2006). The PCR products were analyzed by agarose 1% gel electrophoresis and the result was showed by using UVIDoc HD5 (UVITEC Cambridge).

The isolated bacteria were identified by means of homology at <https://blast.ncbi.nlm.nih.gov/Blast.cgi>. It is an analysis using BLAST based on the database from the GenBank of the National Center for Biotechnology Information (NCBI).

Extraction of microalgae. Dry biomass of microalgae *T. chui* (10 g) and *D. salina* (9 g) were dissolved in 100 mL methanol and was left for 5 x 24 h. Methanolic extract of *T. chui* was passed in ultrasound for 15 min and was evaporated to dryness.

Preparation of feeds. Dry extracts of *T. chui* and *D. salina* (2 g) were blended with 10 g of commercial pellet and were left to dry in the drying machine for 2 x 24 h. Commercial pellet without microalgae extracts served as negative control. The commercial pellets used in this study contain 30% of protein, 5% of lipid, 4% of fiber and with 12% moisture.

The use of microalgae as bio-control (in vivo assay). Forty-five vibriosis infected *L. vannamei* samples were taken based on the previous biomolecular test, used for the *in vivo* assay. Shrimp were reared in aerated aquarium with a density of five individuals in 5 L of water. The water quality was maintained at 26-28°C and salinity 25-30 ppt. Shrimp were then acclimatized for three days before the *in vivo* assay.

Shrimps were placed in three different tanks. Each tank represented different treatments related to the type of feeds given. Treated shrimps were fed three times a day, i.e. at 08:00, 12:00 and 17:00 WIB (local time). All experiments were performed in three replicates.

Treatment I: commercial pellet mixed with *T. chui* extract of 200 ppm against *L. vannamei* infected with vibriosis.

Treatment II: commercial pellet mixed with *D. salina* extract of 200 ppm against *L. vannamei* infected with vibriosis.

Treatment III: commercial pellet without addition of microalgae extract to *L. vannamei* shrimp infected with vibriosis.

Microbial enumeration from the hepatopancreas of shrimp. The observed parameters were amount of bacteria in the hepatopancreas of treated and control shrimps counted using the Total Plate Count (TPC) method (Benson 2001). Hepatopancreas of treated and control shrimp were dissolved in 10 mL sterile seawater. The solutions were stirred for 15 min and were diluted three times (10^{-1} , 10^{-2} and 10^{-3}). The treatment was carried out for 21 days and the bacteria amount was counted at the 7th, 14th, and 21st days.

Results

Morphological signs of the infected shrimp. The clinical signs of the infected shrimps were 'luminescent pleopoda', melanocyst on the shrimp's body, reddish tail, and brownish-red hepatopancreas.

Bacterial isolates. Based on the isolation of vibriosis-causing bacteria, there were 13 isolates that was morphologically characterized based on their colonial edge form, characteristic and color. Out of 13 bacterial isolates, three potential bacteria were selected and results are presented in Table 1.

Table 1

Morphological characteristic sand color of 13 bacterial isolates

No	Colonial edge form	Colonial characteristic	Colonial color	Total	Isolates code
1	Jagged	Convex	Green	1	U1H1
2	Smooth	Convex	Yellow	7	U1H2,U1E1,U2H1,U2H1, U3H2, U4H1, U5H1,
3	Smooth	Convex	Green	2	U1E2, U3H1
4	Smooth	Convex	Milk white	1	U7E1
5	Jagged	Convex	Milk white	1	U8E1
6	Jagged	Convex	Yellow	1	U8E2

Bacterial identification. Based on the biomolecular analysis, there were two bacterial isolates that were classified as same species. This analysis managed to identify the presence of *V. alginolyticus* and *V. harveyi* as two bacteria strain being responsible for the vibriosis infection on the treated shrimps. Biomolecular analysis is presented in Table 2.

Table 2

Biochemical analysis of three selected bacterial isolates from Vannamei shrimp

No	Code	Universal Primer		Species	Homology
		27F	1429R		
1	U1H1	1.27F	1.1429R	<i>V. alginolyticus</i> strain ATCC 17749	97%
2	U1E1	2.27F	2.1429R	<i>V. harveyi</i> strain NCIMB 1280	96%
3	U4H1	3.27F	3.1429R	<i>V. alginolyticus</i> strain ATCC 17749	97%

In vivo assay. Forty-five shrimps infected by vibrio reared during 21 days and feed with *D. salina* and *T. chui* showed a decreased of bacteria amount (Figure 1).




Figure 1. Number of bacteria observed from infected *Litopenaeus vannamei* during the 21 days experiment.

Discussion. Clinical signs observed on the infected *L. vannamei* collected from Kaliwungu, Kendal were in accordance as infection signs in Vannamei caused by vibrio's

as reported by Lina et al (2007) and Huang et al (2013). Another features remarked correlates to the behavior of shrimp are passive movement, slow response to feed and fluorescent swimming legs. Prayitno & Lachtford (1995) stated that *V. harveyi* is one of causative agent that also known as luminescent bacteria. Those clinical signs became the reference in isolating the vibriosis-causing bacteria from the shrimp.

Bacterial isolates demonstrated that there were 13 isolates with diverse morphology. The selected isolate, which was U1H1, had a jagged colonial edge, convex with green color. Meanwhile, the U1E1 AND U4H1 were characterized with smooth colonial edge, convex with yellow color. These three isolates showed similar characteristic as *Vibrio* bacteria. Sarjito et al (2015) revealed that morphological features of bacteria with smooth colonial edge, convex with yellow color tested biochemically were identical with those of *V. harveyi*.

DNA amplification results from the rep-PCR exhibited three isolates with similar band profiles as seen on marker 800 bp. Rep-PCR method has been proven as an efficient method to identify bacterial species and the genetic relationship (Lina et al 2007). DNA sequencing of isolate U1H1 and U4H1 had identical nucleotide with *V. alginolyticus* strain ATCC 17749 with homological level 97%. Isolate U1E1 showed to have similar nucleotide with *V. harveyi* strain NCIMB 1280 with homological level 96%.

Presence and abundance of *V. alginolyticus* and *V. harveyi* are highly correlated with environmental conditions. Asplund (2013) stated that the abundance of bacteria *Vibrio sp.* usually relates to temperature and salinity and increases in stable environmental condition. The abundance of *V. alginolyticus* and *V. harveyi* counted was 21 (isolate U1H1) and 4 (isolate U4H1) colony/plate and 11 (isolate U1E1) colony/plate. Environmental condition in the pond when the sampling was performed in October-November was fluctuated due to the rainy season. In this season, the temperature and salinity in the pond tended to change in short period; thus, it affected the abundance of bacterial colony, either in the pond or on the *L. vannamei* body.

The result of the in vivo assay shown that the vibrio infected shrimps feed with mixed *T. chui* extract and *D. salina* reared for 21 days shown a decreased of the number of bacterial infection as shown on Figure 1. Widowati et al (2017) observed that *Dunaliella sp.*, *T. chui* showed antioxidant potential which could be considered for future applications in aquaculture. The result of this study indicated that the microalgae are capable to produce antibacterial compounds against vibrio.

Conclusions. The present study showed that vibriosis-causing bacteria isolated from *L. vannamei*, which showed clinical signs of vibriosis infection, was identified as *V. alginolyticus*, and as *V. harveyi*. The shrimps infected by vibrio and feed with *D. salina* and *T. chui* reared during 21 days showed a decreased of bacteria amount. The result indicated that the microalgae were capable to produce antibacterial compounds against vibrio. Based on the present findings, it could be inferred that the secondary metabolites of *T. chui* and *D. salina* may be an excellent source in order to develop potent formulations for sustainable shrimp farming.

Acknowledgements. The authors wish to thank the Directorate of Research and Community Service, Directorate General of Research Strengthening and Development, Indonesian Ministry of Research Technology and Higher Education, for funding this research through the Implementation Agreement Assignment of Research Collaboration and International Publication 2017.

References

- Asplund M. E., 2013 Ecological aspect of marine vibrio bacteria. PhD Thesis in Marine Ecology, University of Gothenburg, Kristineberg, 48 p.
- Benson H. J., 2001 Microbiological applications: Laboratory manual in general microbiology. Short version, 8th edition, The McGraw-Hill Companies.
- Cadiz R. E., Ferdinand R., Traifalgar M., Sanares R. C., Grace K., Andrino-Felarca S., Corre V. L., Corre V. L., 2016 Comparative efficacies of tilapia green water and

- biofloc technology (BFT) in suppressing population growth of green *Vibrios* and *Vibrio parahaemolyticus* in the intensive tank culture of *Penaeus vannamei*. *AACL Bioflux* 9(2):195-203.
- Dash P., Avunje S., Tandel R. S., Sandeep K. P., Panigrahi, A. 2017 Biocontrol of Luminous Vibriosis in Shrimp Aquaculture: A Review of Current Approaches and Future Perspectives. *Reviews in Fisheries Science & Aquaculture* 25:245-255.
- Haldar S., Chatterjee S., Sugimoto N., Das S., Chowdhury N., Hinenoya A., Asakura M., Yamasaki, S. 2011. Identification of *Vibrio campbellii* isolated from diseased farm-shrimps from south India and establishment of its pathogenic potential in an *Artemia* model. *Microbiology* 157(1):179-188.
- Heenatigala P. P. M., Fernando M. U. L., 2016 Occurance of bacteria species responsible for vibriosis in shrimp pond culture systems in Sri Lanka and assesment of the suitable control measures. *Sri Lanka Journal of Aquatic Science* 21:1-17.
- Huang H., Liu X., Xiang J., Wang P., 2013 Immune response of *Litopenaeus vannamei* after infection with *Vibrio harveyi*. *Aquaculture* 406-407:115-120.
- Kumaran T., Citarasu T., 2016 Isolation and characterization of *Vibrio* species from shrimp and *Artemia* culture and evaluation of the potential virulence factor. *Intellectual Property Rights* 4:1-5.
- Lee Y. K., Jung H. J., Lee H. K., 2006 Marine bacteria associated with the Korean brown alga, *Undaria pinnatifida*. *Journal of Microbiology (Seoul, Korea)* 44:694-698.
- Lina N., Prayitno S. B., Sarjito, 2007 Study of the causes of bacteria on the shrimp vibriosis by biomolecular. *Journal of Coastal Resources Management* (3):1-22.
- Manilal A., Sujith S., Selvin J., Shakir C., Gandhimathi R., Kiran G. S., 2010 Virulence of *Vibrios* isolated from diseased black tiger shrimp, *Penaeus monodon*, Fabricius. *Journal of the World Aquaculture Society* 41:332-343.
- Prayitno S. B., Latchford J. W., 1995 Experimental infections of crustacean with luminous bacteria related to photobacterium and vibrio, effect of salinity and on infectiosity. *Aquaculture* (132):105-112.
- Sarjito M., Apriliani D., Afriani A. H., Condro H., 2015 Agensia Penyebab Vibriosis Pada Udang Vaname (*Litopenaeus vannamei*) yang Dibudidayakan Secara Intensif di Kendal. *Jurnal Kelautan Tropis* 16(3):159-196.
- Solidum N. S., Sanares R. C., Andriano-Felarca K. G. S., Corre Jr. V. L., 2016 Immune responses and resistance to vibriosis of juvenile Pacific whiteleg shrimp *Penaeus vannamei* fed with high dose mannan oligosakarida and B-glucan. *AACL Bioflux* 9:239-249.
- Susilowati R., Sabdono A., Widowati I., 2015 Isolation and Characterization of Bacteria Associated with Brown Algae *Sargassum* spp. from Panjang Island and their Antibacterial Activities. *Procedia Environmental Sciences* 23:240-246.
- Walsh P. S., Metzger D. A., Higuchi R., 2013 Chelex 100 as a medium for simple extraction of DNA for PCR-based typing from forensic material. *BioTechniques* 54(3):134-139.
- Widowati I., Zainuri M., Kusumaningrum H. P., Susilowati R., Hardivillier Y., Leignel V., Bourgougnon N., Mouget J.-L., 2017 Antioxidant activity of three microalgae *Dunaliella salina*, *Tetraselmis chuii* and *Isochrysis galbana* clone Tahiti. *IOP Conference Series: Earth and Environmental Science* 55:1-7.
- *** <https://blast.ncbi.nlm.nih.gov/Blast.cgi>. Access on [21.02.2017: 22.49 WIB]
- *** LAKIP, 2014 Kementerian Kelautan dan Perikanan. Jakarta, 327 p.

Received: 22 November 2017. Accepted: 28 January 2018. Published online: 06 February 2018.

Authors:

Ita Widowati, Diponegoro University, Faculty of Fisheries and Marine Science, Department of Marine Science, Indonesia, Jalan Prof. Soedarto SH, Tembalang, Semarang 50275, e-mail: ita.widowati@undip.ac.id
ita_jusup@yahoo.co.id

Muhammad Zainuri, Diponegoro University, Faculty of Fisheries and Marine Science, Department of Oceanography, Indonesia, Jalan Prof. Soedarto SH, Tembalang, Semarang 50275, e-mail: muhammad.zainuri@yahoo.co.id

Hemin Pancasakti Kusumaningrum, Diponegoro University, Faculty of Sciences and Mathematic, Department of Biology, Indonesia, Jl. Prof Soedarto SH, Tembalang, Semarang 50275, e-mail: herminsakti@gmail.com

Yusri Maesaroh, Diponegoro University, Faculty of Fisheries and Marine Science, Marine Science MSc Program, Indonesia, Jalan Prof Soedarto SH Tembalang, Semarang 50275, e-mail: yusrimaesaroh@gmail.com

Yann Hardivillier, Le Mans Université, Faculté des Sciences et Techniques, Laboratoire du Mer Molécule et Santé (MMS), France, EA 2160 FR CNRS 3473 IUML, Avenue Olivier Messiaen, 72085 Le Mans cedex 9, e-mail: yann.hardivillier@univ-lemans.fr

Vincent Leignel, Le Mans Université, Faculté des Sciences et Techniques, Laboratoire du Mer Molécule et Santé (MMS), France, EA 2160 FR CNRS 3473 IUML, Avenue Olivier Messiaen, 72085 Le Mans cedex 9, e-mail: vincent.leignel@univ-lemans.fr

Nathalie Bourgougnon, Université de Bretagne-Sud, Laboratoire de Biotechnologie et Chimie Marines (LBCM), France, 56017 Vannes, Campus de Tohannic, e-mail: nathalie.bourgougnon@univ-ubs.fr

Jean-Luc Mouget, Le Mans Université, Faculté des Sciences et Techniques, Laboratoire du Mer Molécule et Santé (MMS), France, EA 2160 FR CNRS 3473 IUML, Avenue Olivier Messiaen, 72085 Le Mans cedex 9, e-mail: jean-luc.mouget@univ-lemans.fr

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Widowati I., Zainuri M., Kusumaningrum H. P., Maesaroh Y., Hardivillier Y., Leignel V., Bourgougnon N., Mouget J.-L., 2018 Identification of agents causing vibriosis in *Litopenaeus vannamei* shrimp culture in Kendal, Central Java, Indonesia and application of microalgae *Dunaliella salina* and *Tetraselmis chui* as bio-control agents against vibriosis. *AAFL Bioflux* 11(1):101-107.

Identification of agents causing vibriosis in Litopenaeus vannamei shrimp culture in Kendal, Central Java, Indonesia and application of microalgae Dunaliella salina and Tetraselmis chui as bio-control

ORIGINALITY REPORT

14%

SIMILARITY INDEX

%

INTERNET SOURCES

14%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1 **Edi Setiyono, Marcelinus Alfasisurya Setya Adhiwibawa, Renny Indrawati, Monika Nur Utami Prihastyanti et al. " An Indonesian Marine Bacterium, , Produces Antimicrobial Prodiginine Pigments ", ACS Omega, 2020** 7%

Publication
- 2 **Thomas v. O. Hansen, Mette K. Simonsen, Finn C. Nielsen, Yrsa Andersen Hundrup. "Collection of Blood, Saliva, and Buccal Cell Samples in a Pilot Study on the Danish Nurse Cohort: Comparison of the Response Rate and Quality of Genomic DNA", Cancer Epidemiology, Biomarkers & Prevention, 2007** 1%

Publication
- 3 **Suwarni Suwarni, Joeharnani Tresnati, Sharifuddin Bin Andy Omar, Ambo Tuwo. "Some Reproductive Biology Studies of Rabbitfish Siganus canaliculatus (Park, 1797) from the Southern Coastal Waters of** 1%

Jenepono, South Sulawesi, Indonesia",
Biosciences Biotechnology Research Asia,
2019

Publication

4

Irman Hermadi, Andi Muhammad Chaerul Hafidz, Auzi Asfarian, Yani Nurhadryani, Nadya Farchana Fidaroina. "Development of Back-end of a Rural Participation Based Knowledge Management System of Smallholder Palm Plantation", 2020 International Conference on Computer Science and Its Application in Agriculture (ICOSICA), 2020

Publication

1 %

5

Norhan E. Saleh, Fady R. Michael, Mohamed M. Toutou. "Evaluation of garlic and onion powder as phyto-additives in the diet of sea bass (*Dicentrarchus labrax*)", The Egyptian Journal of Aquatic Research, 2015

Publication

1 %

6

I. KHAN. "MORPHOLOGICAL, BIOCHEMICAL AND MOLECULAR CHARACTERIZATION OF CULTURABLE EPILITHIC AND ENDOLITHIC BACTERIA FROM ROCKS OF AYUBIA (MURREE), LOWER HIMALAYA, PAKISTAN", Applied Ecology and Environmental Research, 2018

Publication

1 %

7

Crépin, Alexandre, Corinne Barbey, Amélie Beury-Cirou, Valérie Hélias, Laure Taupin, Sylvie Reverchon, William Nasser, Denis Faure, Alain Dufour, Nicole Orange, Marc Feuilloley, Karin Heurlier, Jean-François Burini, and Xavier Latour. "Quorum Sensing Signaling Molecules Produced by Reference and Emerging Soft-Rot Bacteria (*Dickeya* and *Pectobacterium* spp.)", PLoS ONE, 2012.

Publication

1 %

8

Manilal, Aseer, Sugathan Sujith, Balu Sabarathnam, George Seghal Kiran, Joseph Selvin, Chippu Shakir, and Aaron Premnath LIPTON. "Bioactivity of the red algae *Asparagopsis taxiformis* collected from the Southwestern coast of India", Brazilian Journal of Oceanography, 2010.

Publication

<1 %

9

Fernando J. Sutili, Letícia T. Gressler. "Antimicrobial agents", Elsevier BV, 2021

Publication

<1 %

10

Linda A. Ndiana, Gianvito Lanave, Aya A. K. Zarea, Costantina Desario et al. "Molecular characterization of carnivore protoparvovirus 1 circulating in domestic carnivores in Egypt", Frontiers in Veterinary Science, 2022

Publication

<1 %

11

Miriam Esther Garcia - Mendoza, Jorge Cáceres - Martínez, Rebeca Vásquez - Yeomans, Roberto Cruz - Flores. "

Bacteriological water quality of recirculating aquatic systems for maintenance of yellowtail amberjack ", Journal of the World Aquaculture Society, 2019

Publication

<1 %

12

S. Haldar. "Identification of *Vibrio campbellii* isolated from diseased farm-shrimps from south India and establishment of its pathogenic potential in an *Artemia* model", Microbiology, 01/01/2011

Publication

<1 %

13

Silkina, Alla, Alexandra Bazes, Jean-Luc Mouget, and Nathalie Bourgougnon. "Comparative efficiency of macroalgal extracts and booster biocides as antifouling agents to control growth of three diatom species", Marine Pollution Bulletin, 2012.

Publication

<1 %

14

Umi Salmah Binti Ahmed Sajali, Nathan L. Atkinson, Andrew P. Desbois, David C. Little, Francis J. Murray, Andrew P. Shinn. "Prophylactic properties of biofloc- or Nile tilapia-conditioned water against *Vibrio parahaemolyticus* infection of whiteleg shrimp (*Penaeus vannamei*)", Aquaculture, 2019

<1 %

Publication

Exclude quotes On

Exclude bibliography On

Exclude matches Off

Identification of agents causing vibriosis in *Litopenaeus vannamei* shrimp culture in Kendal, Central Java, Indonesia and application of microalgae *Dunaliella salina* and *Tetraselmis chui* as bio-control

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7
