

Redescription of *Stichopus* *monotuberculatus* (Echinodermata, Holothuroidea, Stichopodidae) of Parang island, Karimunjawa Archipelago, Central Java, Indonesia

by Widianingsih Widianingsih

Submission date: 11-Jan-2023 10:06AM (UTC+0700)

Submission ID: 1991037554

File name: ang_island,_Karimunjawa_Archipelago,_Central_Java,_Indonesia.pdf (317.89K)

Word count: 3255

Character count: 17924

Redescription of *Stichopus monotuberculatus* (Echinodermata, Holothuroidea, Stichopodidae) of Parang Island, Karimunjawa Archipelago, Central Java, Indonesia

Widianingsih Widianingsih¹, Retno Hartati¹, Hadi Endrawati¹,
Robertus T. Mahendrajaya¹ and Agoes Soegianto^{2*}

6

¹Department of Marine Science, Faculty of Fisheries and Marine Science, Diponegoro University,
Jl. Prof. Dr. Soedharto, SH Kampus Tembalang, Semarang, Indonesia

²Department of Biology, Faculty of Sciences and Technology, Universitas Airlangga Surabaya,
Indonesia

(Received 22 August, 2020; Accepted 30 October, 2020)

ABSTRACT

Stichopus monotuberculatus (Quoy & Gaimard, 1833) belongs to family Stichopodidae, has high commercial value in fisheries trade world. Most of the sea cucumbers belong to family Stichopodidae have long been used for nutritional food and medicine. The objective of the study was to redescribe this species by its phenotype characteristics through morphological keys and review of their ossicles as well as genotype by DNA mapping. The samples were collected from Parang Island and preserved in alcohol 96%. Commercial bleach were used to remove tissue and take the ossicles, then they were examined under light microscope. For genotype characteristic (DNA mapping) extraction DNEasy of tissue (qiagen) was used for extraction of DNA. Primer CO1 was used to amplify the genome. Samples showed that morphologically they were grey-green colour with numerous small dark patches ventrally and grey green to orange-brown with dark green to black patches dorsally with square thick integument/body wall. Calcareous ring radial pieces had a posterior notch and four short anterior points whereas the interradial pieces had a long anterior tooth. The ossicles showed numerous tables-shaped ossicles in the anterior and dorsal tissue of the bodywall, but norossete-shape in dorsal body wall which is the characteristic of *S. monotuberculatus*. DNA sequencing of nine samples showed that all samples had got 93-99% similarity with *Stichopus monotuberculatus* haplotype 4, 5, 9, and 13. This result confirmed the identification through morphology and ossicles characters. It is approved the presence of *S. monotuberculatus* (Quoy and Gaimard, 1833) in Parang Island, Karimunjawa Archipelago, Jepara.

Key words: *Stichopus monotuberculatus*, Phenotype, Genotype, Parang Island.

Introduction

Stichopus monotuberculatus (Quoy and Gaimard, 1833) belongs to Class Holothuroidea (Echinodermata), the marine benthic invertebrates which has

high economical value. It is inhabit at coral reef, seagrass and deep seas ecosystem across the ocean. Ecologically, it has important role in benthic ecosystem through its bioturbation activity in benthic ecosystem (Hartati *et al.*, 2009). Study on Genus

Stichopus from Karimunjawa has been conducted by Hartati *et al.* (2009), Purwati *et al.* (2010), and Widianingsih *et al.* (2013). Widianingsih *et al.* (2015) has tried to examine three Stichopodid samples, but has not been concluded the name of the species. Setyastuti and Purwati (2015) added list of Purwati *et al.* (2010) with *Stichopus cf. monotuberculatus*.

The external morphology, internal organs, and spicules/ossicles have been widely used to identify sea cucumber. These characteristics may indicate a high level of similarity for particular families, including family Stichopodidae (Wirawati and Purwati, 2016). The molecular methods have been used to identify various marine organisms such as sea cucumbers (Byrne *et al.*, 2010; Amin *et al.* 2016, Madduppa *et al.*, 2017; Patantis *et al.*, 2019). Therefore, this research was aimed to redescribe *Stichopus monotuberculatus* (Family Stichopodidae) from Parang Island, Karimunjawa Archipelago, Central of Java, Indonesia through its phenotype (morphology and ossicles characteristic) and genotype characteristic with barcoding technique.

Materials and Method

Morphological examination and ossicles observation

Fresh sample of sea cucumbers were collected from intertidal zone of Parang Island, Karimunjawa Archipelago and were examined morphologically for its overall shape, body wall, the position of mouth and anus, tentacle shape, tube feet, rete mirabile, pedicels, respiratory tree, gonad (if any) based on Samyn *et al.*, (2006).

Ossicles were extracted from tissues of the mid part of dorsal body, papilla tips, tentacles, tube feet, and cloaca wall. A small piece (about 2 × 2 mm) of tissue is cut and placed on a clean glass slide. A few drops of weak commercial bleach were added using a pipette to dissolve the tissue. Tissues were left to dissolve for about 5 minutes. The ossicles were then washed several times with deionized water and finally with absolute alcohol. The spicules were then observed under a light microscope. The terms used to describe different types of ossicles followed Massin (1999) and Samyn *et al.* (2006).

DNA extraction, PCR, and sequencing

Total genomic DNA was extracted from approximately 10–20 mg of tissue from the tube feet using

⁵ DNeasy Blood and Tissue Kit (QIAGEN GmbH, Hilden, Germany) following the manufacturer's specification. Sections of the cytochrome c oxidase subunit I (COI) and the small subunit 16S ribosomal DNA (16S rDNA) genes were amplified using the primers COle-F and COle-R (Arndt *et al.*, 1996) and 16Ss-F (5'GTAGCATAATCACTTGCTCTTAAA-3') and ¹⁵ 16Ss-R (5'TTATTTAGAAGATAGAAGCTGACCT-3'), respectively. PCR amplification was conducted following Byrne *et al.* (2010).

PCR reaction with Kapa (2x Kapa Taq Extra Hotstart Ready Mix Dye (2mM MgCl₂ at 1x) consisted of one cycle initial denaturation at 95 °C for 3 × 60 s followed by 40 cycles of denaturation at 95 °C for 30s, annealing Tm at -5°C for 30 s, extension at 72 °C for 80 s, and final extension at 72 °C for 1 min. All nucleotide sequences were determined using BigDye Terminator Kit ver. 3.1 and ABI 3730 Genetic Analyzer (Life Technologies, California, USA).

The PCR products were visualized to check DNA quality on the Mufid-EXU (Submarine electrophoresis system) specification with duration of 30 minutes and 100 volts. The positive PCR products were sent to a sequencing facility and loaded into an ABI 3130xl automated sequencer (Applied Biosystems). Sequences editing and alignment will be conducted using MEGA 5 (Tamura *et al.* 2011; Lucas *et al.*, 2012). Sample identification was performed using the Basic Local Assignment Search Tool (BLAST; Altschul *et al.* 1990).

For phylogenetic analysis, one or more reference sequences from GenBank sequence database with the highest maximum identity to each amplicon sequence were downloaded. In the phylogenetic tree, *Portunus pelagicus* (accession number KP976342) was used as an out-group comparison (Madduppa *et al.* 2017). A neighbor-joining phylogenetic tree using Kimura-2 parameter models was reconstructed with 1000 replicates of bootstrap value. The genetic distance within and between species was investigated.

Results and Discussion

Family Stichopodidae belonged to Ordo Aspidochirotida Class Holothuroidea (Echinodermata) in which consisted mostly the commercially trade sea cucumber in Indonesia and high commercial value due to its medicinal and edible properties (Maryam *et al.*, 2012). One of Stichopodidae species found in Karimunjawa Archipelago was *Stichopus*

monotuberculatus. It was widely distributed in the Indo-Pacific Ocean (Massin, 1996) as well as in Indonesia and had synonym name of *Holothuria monotuberculata* Quoy and Gaimard, 1833 (Massin, 1996).

Phenotype (morphology and ossicles characteristics)

Nine fresh samples of *S. monotuberculatus* were observed their morphological characters. It is usually medium to large-sized sea cucumbers, the length of the samples was in the range of 100-320 mm. Their body colour were grey-green with numerous small dark patches ventrally and grey green to orange-brown with dark green to black patches dorsally. During preservation, the colour was changed became light green with light brown patches. According to Massin (1996) this species appeared to exhibit a wide colour range from green/bluish-green to orange-brown. His living specimens colour were pale brown to rust coloured dorsally and ventrally. Dorsally, there were numerous white and deep brown dots. The white dots often clumped together forming a whitish surface.

As other member of Family Stichopodidae, it was usually square in cross section, and distinctly flattened below (Purwati *et al.*, 2010). Ventral side of the body of the samples was flattened with swollen dorsal side. Mouth was in anterior ventral position surrounded by 20large tentacles in which encircled by a large papilla as found by Massin (1996) in *S. monotuberculatus* samples. Anus position was terminal and there was no an anal tooth or papillae. Large conical papillae were distributed in bivium. There was a distinct fringe of 8-10 larger papillae laterally. This character was similar found by Maryam *et al.* (2012) in Gulf of Persian. In the samples of Massin (1996), the numerous papillae were also found in

the dorsal with a white apex. There was one row of 9-10 very large papillae on each side of the body, posterior ones being the largest.

The sample's tube feet in the present studies were large and cylindrical shape, their colour were yellowish brown. According to Massin (1996), the tube feet of *S. monotuberculatus* were well developed and abundant along the three ventral ambulacra (trivium). They were located on 3-6 rows along the lateral ambulacra and on 8-14 rows along the media nambulacrum. The tube feet were translucent, ending in a large deep brown sucker.

Examination of the calcareous ring and ossicles found that the calcareous ring of the samples had large radial pieces and narrow interradial pieces. The radial pieces had a posterior notch and four short anterior points whereas the interradial pieces had a long anterior tooth. This character also found in Massin's sample (1996) of *S. monotuberculatus*. After calcareous ring, there was one large polian vesicle and one contorted stone canal, embedded in the dorsal mesentery.

There were numerous tables-shaped ossicles in the anterior tissue of the bodywall (Fig. 1A) but very few the C- (Fig. 1H), Rosette- (Fig.1D), and plate-shape rods (Fig. 1E). It was also very numerous in the posterior tissue (Fig. 1B) but no rosette-shape. The modified-rods also were found in this part of the body (Fig.1J). In the tube feet there are reduced tables and spiny rods with an enlarged central process, very often perforated.

In the ventral tissue of the body wall, there were only tables (Fig. 1G, N) which are very small. At the base of the dorsal papillae, the ossicles are the same as in the body wall whereas at the apex there were large perforated plates (Fig. 1F), smooth or spiny or modified rods (Fig. 1J), a few tables and C-shaped rods (Fig. 1H). In the dorsal tissue of the body wall,

Table 1. COI gene mtDNA sequence blast analyses result of sea cucumbers samples from Parang Island, Karimunjawa Archipelago, Jepara

No.	Sample ID	Blast sequence	ID GenBank	Length bp	Similarity %
1	S1-OK-F1	<i>Stichopus monotuberculatus</i> haplotype 9	KC424504.1	667	99
2	S1-OK-F	<i>Stichopus monotuberculatus</i> haplotype 4	KC424499.1	667	99
3	S1-OK-R1	<i>Stichopus monotuberculatus</i> haplotype 9	KC424504.1	667	99
4	S1-OK-R	<i>Stichopus monotuberculatus</i> haplotype 4	KC424499.1	667	99
5	S3-OK-F2	<i>Stichopus monotuberculatus</i> haplotype 5	KC424500.1	667	99
6	S3-OK-F3	<i>Stichopus monotuberculatus</i> haplotype 13	KC424508.1	674	99
7	S3-OK-R1	<i>Stichopus monotuberculatus</i> haplotype 13	KC424508.1	674	99
8	S3-OK-R2	<i>Stichopus monotuberculatus</i> haplotype 13	KC424508.1	674	99
9	S4-OK-F 1	<i>Stichopus monotuberculatus</i> haplotype 13	KC424508.1	674	93

there were only few tables (Fig. 1C) and C-shape rods (Fig. 1H), but the rosette (Fig. 1I) were numerous as found by Cherbonnier (1988). The tables were small (39–61 mm in diameter) with a squarish disc perforated by 4 large central holes and 4–10 small peripheral holes. The four short pillars are united by one cross beam and end in a crown of spines which is often as wide as the disc. The rosettes are sparse and irregular (Fig. 1J). The C-shaped rods are long and smooth or with a few excrescences (Fig. 1H).




Fig. 1. Ossicles of *Stichopus monotuberculatus* (Quoy & Gaimard, 1833) from Parang Island, Karimunjawa Archipelago, Jepara, Indonesia

A : tables of anterior, B : tables of posterior, C : tables of dorsal body wall, D: rosette of anterior, E: plate-shape anterior, F: plate of dorsal, G: tables of ventral body wall, H: plate of ventral body wall, H: C-shape rods of body wall, I: rosette shape of dorsal body wall, J: rods of body wall, K: plate of posterior, L: S-rod shape of dorsal, N: plate of ventral body wall, M: rod shape, N: table of ventral body wall.

The character of ossicles found in the present study is similar to that Massin (1996).

The result of present study revealed that the general aspect of the specimen, morphology and of the remaining ossicles of the body wall and the tentacles matched with *S. monotuberculatus*.

Genotype Character with DNA barcoding

Nine samples of sea cucumber from Parang Island, Karimunjawa Archipelago which had been morphology and ossicles examined as *S. monotuberculatus* were then analyzed for their molecular identification. COI gene mtDNA of all sea cucumbers samples have been successfully amplified using universal primers of the COI gene. Amplification results obtained DNA fragments with a length of about 674–667 bp. Blast analysis of this sea cucumber COI gene sequence showed that all sample of sea cucumber had similarity with *Stichopus monotuberculatus* H haplotype 4, 5, 9 and 13 with acc No. KC424499.1, KC424500.1, KC424504.1 and KC424508.1 NCBI Genbank at 93–99% (Table 2). These results confirmed their identification through morphology and ossicles characters.

The molecular methods have been widely used to identify various marine organisms and their processed products, including sea cucumbers, because of their specificity. The result of molecular identification (COI gene mtDNA and Blast analysis) of samples of sea cucumber with morphological features of grey-green with numerous small dark patches ventrally and grey green to orange-brown with dark green to black patches dorsally with square thick integument/body wall and had same spicules characteristics showed that they had similarities with *Stichopus monotuberculatus* at 93–99%. This method has been successfully identified one species of sea cucumber from Surabaya (Amin et al. 2016), seven species of sea cucumbers from Kepulauan Seribu (Madduppa et al., 2017), and 25 beche-de-mere from Boalemo, Pesawaran, Surabaya, and West Lombok (Patantis et al., 2019).

As other member of Family Stichopodidae, because of high demand and good economy value, *S. monotuberculatus* was very prone to overexploitation. Therefore, it was need urgent management intention (Purcell et al. 2013) and effort to conservation effort. The idea of sea cucumber ranching (releasing cultured/wild juvenile/young sea cucumber into unenclosed coastal environments for harvest at a larger size in 'put and take' opera-

tion) (Bartley and Bell, 2008) has been initiated by Hartati *et al.*, (2021 Inpress) with lower value sea cucumber of *Holothuria atra*, it might be applied for *S. monotuberculatus* since it has been demonstrated that this species can be reared in captivity, thus sea ranching or farming of this species would provide an alternative to fisheries (Hu *et al.* 2010; Chen *et al.*, 2015).

Conclusion

Nine samples have been successfully identified through morphology and ossicles examination as *Stichopus monotuberculatus*. Based on genetic DNA barcoding analysis, all samples had 99% similarity with *Stichopus monotuberculatus* haplotype 4, 5 and 9 and lower (93%) similarity with *Stichopus monotuberculatus* haplotype 13. These results confirmed the existence of species in Parang Island, Karimunjawa Archipelago, Jepara.

Acknowledgments

The Directorate General of Higher Education, Ministry of National Education and Culture and Rector Diponegoro University are gratefully acknowledged for funding this research work through Fundamental Research Scheme of 2015 by contract No. 140-05/UN7.5.1/PG/2015.

References

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. 1990. Basic local alignment search tool. *Journal of Molecular Biology*. 215(3):403-410
- Amin, M.H.F.A., Pidada, I.B.R., Sugiharto, Widyatmoko, J.N. and Irawan, B., 2016, March. Sea cucumber species identification of family Caudinidae from Surabaya based on morphological and mitochondrial DNA evidence. *AIP Conference Proceedings*, 1:030001. DOI: 10.1063/1.4943311.
- Arndt A, Marquez C, Lambert P, Smith M.J. 1996. Molecular phylogeny of Eastern Pacific sea cucumbers (Echinodermata: Holothuroidea) based on mitochondrial DNA sequence. *Molecular Phylogenetics and Evolution* 6(3): 425–437.
- Bartley, D.M. and Bell J.D. 2008. Restocking, Stock Enhancement, and Sea Ranching: Arenas of Progress. *Reviews in Fisheries Science*, 16(1-3):357-365. DOI: 10.1080/10641260701678058
- Byrne, M., Rowe, F. and Uthicke, S. 2010 Molecular taxonomy, phylogeny and evolution in the family Stichopodidae (Aspidochirotida: Holothuroidea) based on COI and 16S mitochondrial DNA. *Molecular Phylogenetics and Evolution*. 56(3) : 1068–1081.
- Chen, Y., Hu, C. and Ren, C. 2015. Application of wet waste from shrimp (*Litopenaeus vannamei*) with or without sea mud to feeding sea cucumber (*Stichopus monotuberculatus*). *J. Ocean Univ. China*, 14 : 114–120. <https://doi.org/10.1007/s11802-015-2348-z>
- Cherbonnier, G., 1988. Echinoderms: Holothurides. *Faune de Madagascar*. 70 : 1-292.
- Hartati, R., P. Purwati and Widianingsih, 2010. Field Guide Timun Laut di Kep. Karimunjawa. Undip. Press. 46 pp.
- Hartati R, Ambariyanto, Zainuri M, and Widianingsih. 2021. Sea ranching of *Holothuria atra*: effects of stocking density and stocking time. *BIOTROPIA* Inpress.
- Hu, C.Q., Xu, Y. H., Wen, J., Zhang, L. P., Fan, S. G. and Su, T. 2010. Larval development and juvenile growth of the sea cucumber *Stichopus* sp. (Curry fish). *Aquaculture*. 300(1-4): 73–79.
- Madduppa, H., Taurusman, A.A., Subhan, B., Anggraini, N.P., Fadillah, R. and Tarman, K. 2017. DNA barcoding reveals vulnerable and not evaluated species of sea cucumbers (Holothuroidea and Stichopodidae) from Kepulauan Seribu reefs, Indonesia. *Biodiversitas*, 18:893-898. DOI: 10.13057/biodiv/d180305
- Maryam, E., Majid, A., Houshang, B.A. and Abdolrasoul, D. 2012. First Report of a Sea cucumber, *Stichopus cf. monotuberculatus* (Quoy & Gaimard, 1833), from Hengam Island (Persian Gulf, Iran). *European Journal of Experimental Biology*. 2 (3) : 547-550.
- Massin, C. 1996. Result of the Rumphius Biohistorical Expedition to Ambon (1990) Part 4. The Holothuroidea (Echinodermata) collected at Ambon during the Rumphius Biohistorical Expedition. *Zool. Verh. Leiden*. 307 : 1-53.
- Massin, C. 1999. Reef dwelling Holothuroidea (Echinodermata) of the Spermonde Archipelago (South-West Sulawesi) Indonesia. *Zool. Verh. Leiden*. 329 : 1-143.
- Patantis, G., Dewi, A.S., Fawzya, Y.N. and Nursid, M. 2019. Identification of Beche-de-mers from Indonesia by molecular approach. *Biodiversitas*. 20 : 537-543. DOI: 10.13057/biodiv/d200233
- Purcell S.W., Mercier A., Conand C., Hamel J.-F., Toral-Granda V.M., Lovatelli A. and Uthicke S. 2013. Sea cucumber fisheries: Global analysis of stocks, management measures and drivers of overfishing. *Fish and Fisheries*. 14(1) : 34–59.
- Purwati, P., Hartati, R. and Widianingsih, 2010. Eighteen Sea Cucumber Species Fished in Karimunjawa Island, Java sea. *Marine Research in Indonesia*. 35 (2) : 23 – 30.
- Samyn, Y.D., Vandenspiegel and Massin, C. 2006. A new Indo-West Pasific Species of Actinopyga

- (Holothuroidea: Aspidochirotida: Holothuriidae). *Zootaxa*. 1138 : 53-68.
- Samyn, Y., Van Den Spiegel, D. and Massin, C. 2006. Taxonomie des holothuries des Comores. Cooperation Belge. 130 pages
- Setyastuti, S. and Purwati, P. 2015. Species list of Indonesian trepan. *SPC Beche-de-mer Information Bulletin* #35: 19-25.
- Tamura, K., Peterson, D., Peterson, N., Stecher, G., Nei, M. and Kumar, S. 2011. MEGA5: Molecular evolutionary genetics analysis using maximum likelihood, evolutionary distance, and maximum Parsimony methods. *Molecular Biology and Evolution*. 28(10) : 2731-2739.
- Widianingsih, Hartati, R. and Endrawati, H. 2015. The study of ossicles Family Stichopodidae (sea cucumber) from Karimunjawa Island, Central of Java. *1st International Symposium on Marine and Fisheries Research. August 7th 2015*. Univeritas Gadjahmada, Yogyakarta.
- Wirawati, I. and Purwati, P. 2016. Filogeni Timun Laut (Holothuroidea: Stichopodidae) Berdasarkan Karakteristik Morfologis. *Oseanologi and Limnologi di Indonesia* 1(2) : 1-14. (In Indonesian).

Redescription of *Stichopus monotuberculatus* (Echinodermata, Holothuroidea, Stichopodidae) of Parang island, Karimunjawa Archipelago, Central Java, Indonesia

ORIGINALITY REPORT



PRIMARY SOURCES

- | | | |
|---|--|-----|
| 1 | CLAUDE MASSIN. "Taxonomy of the heavily exploited Indo-Pacific sandfish complex (Echinodermata: Holothuriidae)", <i>Zoological Journal of the Linnean Society</i> , 01/2009
Publication | 1 % |
| 2 | Ji Hye Choi, Atef Omar, Min-Seop Kim, Jae-Ho Jung. "Diophys pauciciliata n. sp. (Ciliophora, Euplotida), a new marine spirotrich ciliate with highly static morphometrics", <i>European Journal of Protistology</i> , 2022
Publication | 1 % |
| 3 | Maria Byrne, Frank Rowe, Sven Uthicke. "Molecular taxonomy, phylogeny and evolution in the family Stichopodidae (Aspidochirotida: Holothuroidea) based on COI and 16S mitochondrial DNA", <i>Molecular Phylogenetics and Evolution</i> , 2010
Publication | 1 % |

- 4 Akito Ogawa, Antonina Kremenetskaia, Shimpei F. Hiruta, Yasutoki Shibata et al. "Rehabilitation of two deep-sea holothurian species in genus *Pannychia* from the northwest Pacific Ocean", Deep Sea Research Part II: Topical Studies in Oceanography, 2022
Publication 1 %
- 5 Keiichi Kakui, Toru Katoh, Shimpei F. Hiruta, Norio Kobayashi, Hiroshi Kajihara. "Molecular Systematics of Tanaidacea (Crustacea: Peracarida) Based on 18S Sequence Data, with an Amendment of Suborder/Superfamily-Level Classification", Zoological Science, 2011
Publication 1 %
- 6 R Hartati, M Zainuri, A Ambariyanto, W Widianiingsih, Mustagpirin, F Panji. "Ranching of sea cucumber in Indonesia : A study case of *Holothuria atra*", IOP Conference Series: Earth and Environmental Science, 2021
Publication 1 %
- 7 Devin M. Bartley, Johann D. Bell. "Restocking, Stock Enhancement, and Sea Ranching: Arenas of Progress", Reviews in Fisheries Science, 2008
Publication 1 %
- 8 Hannes Löwe, Marleen Beentjes, Katharina Pflüger-Grau, Andreas Kremling. "From gas to sugar: Trehalose production in from CO and

hydrogen gas ", Cold Spring Harbor
Laboratory, 2020

Publication

-
- 9 Keiichi Kakui. "Descriptions of two new species of *Rhizorhina* Hansen, 1892 (Copepoda: Siphonostomatoidea: Nicothoidae) parasitic on tanaidacean crustaceans, with a note on their phylogenetic position", *Systematic Parasitology*, 2016 1 %
Publication
-
- 10 Shengping Zhong, Longyan Zhao, Guoqiang Huang, Lianghua Huang, Yonghong Liu. "Characterization and phylogenetic analysis of the complete mitogenome of sea cucumber (Massin, Zulfigar, Hwai & Boss, 2002) (Aspidochirotida: Stichopodidae)", *Mitochondrial DNA Part B*, 2022 <1 %
Publication
-
- 11 SiGang Fan, ChaoQun Hu, Jing Wen, LvPing Zhang. "Characterization of mitochondrial genome of sea cucumber *Stichopus horrens*: A novel gene arrangement in Holothuroidea", *Science China Life Sciences*, 2011 <1 %
Publication
-
- 12 Ying Zhang, Qingsong Yang, Juan Ling, Yanying Zhang et al. "The diversity of alkane-degrading bacterial communities in seagrass <1 %

ecosystem of the South China Sea",

Ecotoxicology, 2021

Publication

- 13 Hawis Madduppa, Samsul Bahri, Ahmad Taufik Ghozali, Agus Soleh Atmadipoera et al. "Population genetic structure of Olive ridley (*Lepidochelys olivacea*) across Indonesian archipelago revealed by mitochondrial DNA: Implication for management", Regional Studies in Marine Science, 2020 <1 %
- Publication
-
- 14 Kazem Darvish Bastami, Majid Afkhami, Maryam Ehsanpour, Aida Khazaali, Farzane Soltani. "First report of two species of sea cucumbers from Qeshm Island (Persian Gulf)", Marine Biodiversity Records, 2012 <1 %
- Publication
-
- 15 Kouki Nakata, Hiroki Nagashima, Haruka Yamashita, Natsuki Inaba et al. "Overcoming Hybrid Lethality Induced by Chromosomal Instability in an Interspecific Hybrid of Genus *Nicotiana*", Research Square, 2020 <1 %
- Publication
-
- 16 N O Yonatika, N Widiasih, M Hamidah, M D Nurhakim et al. "Genetic structure and phylogenetic relationships of *Phyllidiellapustulosa* species from Seribu Islands, North Sulawesi, Halmahera, and West <1 %

Papua", IOP Conference Series: Earth and Environmental Science, 2021

Publication

- 17 Luciana Martins, Marcos Tavares. "Ypsilothuria bitentaculata bitentaculata (Echinodermata: Holothuroidea) from the southwestern Atlantic, with comments on its morphology", Zoologia, 2018 <1 %
Publication
- 18 Mahbobe Mehrdost, Ehsan Kamrani, Fereidoon Owfi, Maryam Ehsanpour. "New observation of two species of sea cucumbers from the intertidal habitats of Hengam Island (Persian Gulf, Iran)", Marine Biodiversity Records, 2015 <1 %
Publication
- 19 Sau Pinn Woo, Zulfigar Yasin, Shau Hwai Tan, Hiroshi Kajihara, Toshihiko Fujita. "Sea cucumbers of the genus *Stichopus* Brandt, 1835 (Holothuroidea, Stichopodidae) in Straits of Malacca with description of a new species", ZooKeys, 2015 <1 %
Publication
- 20 "Advances in Sea Cucumber Processing Technology and Product Development", Springer Science and Business Media LLC, 2023 <1 %
Publication

21

Ran-Ran Xing, Ran-Ran Hu, Nan Wang, Jiu-Kai
Zhang, Yi-Qiang Ge, Ying Chen.

<1 %

"Authentication of Sea Cucumber Products
using NGS-based DNA Mini-Barcoding", Food
Control, 2021

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Redescription of *Stichopus monotuberculatus* (Echinodermata, Holothuroidea, Stichopodidae) of Parang island, Karimunjawa Archipelago, Central Java, Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
