

Phytoplankton community structure and it's relationships with water quality in Bangka Island, Indonesia

by Widianingsih Widianingsih

Submission date: 17-Jan-2023 08:56AM (UTC+0700)

Submission ID: 1993859948

File name: 34436-112517-1-PB_1.pdf (341.3K)

Word count: 4735

Character count: 24007

Phytoplankton community structure and it's relationships with water quality in Bangka Island, Indonesia

Muh Yusuf^{1*}, Robin Robin², Wahyu Adi³, Mu'alimah Hudatwi⁴, Widianingsih⁵, Retno Hartati⁵, Robertus Triaji Mahendrajaya⁵, Cristiana Manullang⁶

¹Department of Oceanography, Faculty of Fisheries and Marine Sciences, Diponegoro University
Jl. Prof. Sudarto, SH, Tembalang, Semarang, Jawa Tengah, 50275 Indonesia

²Department of Aquaculture, Faculty of Agriculture, Fisheries and Biology, University of Bangka Belitung
Balunujuk Campus Centre, Merawang, Bangka, Bangka Belitung 33172, Indonesia

³Department of Water Resource Management, Faculty of Agriculture Fisheries and Biology, University of Bangka Belitung
Balunujuk Campus Centre, Merawang, Bangka, Bangka Belitung 33172, Indonesia

⁴Department of Marine Science, Faculty of Agriculture, Fisheries and Biology, University of Bangka Belitung
Balunujuk Campus Centre, Merawang, Bangka, Bangka Belitung 33172, Indonesia

⁵Department of Marine Science, Faculty of Fisheries and Marine Sciences, Diponegoro University
Jl. Prof. Sudarto, SH, Tembalang, Semarang, Jawa Tengah, 50275 Indonesia

⁶Graduate School of Engineering and Science, University of the Ryukyus
Nishihara, Okinawa, Japan

Email: yusuff.undip@gmail.com

Abstract

Phytoplankton plays an important role in primary productivity in marine environment. Various environmental changes in coastal area will impact the water quality and their phytoplankton compositions. The purpose of this study is to examine the abundance of phytoplankton from two different sites, i.e Tanah Merah (close to mining site) and Semujur Island (away from mining site) in Bangka Island. Phytoplankton and water sample were collected on June- August 2018. Water quality was measured using water quality checker, whereas the phytoplankton was identified under the microscope with a magnification of 100x. Non-parametric Kruskal test and T-test analysis was performed to determine the abundance, diversity, uniform, and dominance of phytoplankton between Sites, respectively. Statistical analyses showed the abundance of phytoplankton at Semujur Island was significantly higher than that at Tanah Merah ($p = 0.003$). In additions the diversity, uniform, and dominance were also significantly different between sites (all $p < 0.05$). In Semujur Island, Diatoms (*Thalassiothrix*, *Chaetoceros* and *Thalassionema*) were more dominants than the *Dinophyceae* group. However, in Tanah Merah, the genera *Ceratium* belong to class *Dinophyceae* was more dominant than the class *Bacillariophyceae*. These results performed that the phytoplankton in Tanah Merah and Semujur Island was affected by environment, in this case the mining area. The water quality in Semujur Island (non-mining Area) might have good quality than in Tanah Merah (mining area). The average value of turbidity and Total Suspended Solid in Tanah Merah Waters causes low abundance of phytoplankton. It can be concluded that tin mining can disrupt the abundance and composition of phytoplankton as a primary producer of waters.

Keywords: Tin Mining, Phytoplankton, Tanah Merah, Semujur Island, Bangka, Indonesia

Introduction

Phytoplankton compositions are determined by environmental parameters and their physiological characteristics. Compositions and abundances of phytoplankton will change as a result to environmental variance conditions both physical, chemical, and biological (Reynolds et al., 2002). Several factors are controlling the phytoplankton distributions, such as marine physical process and

nutrient inputs. Moreover, the change of water quality impacts of marine ecosystem included the composition of phytoplankton. (Nguyen and Nhien, 2020). The research on phytoplankton community response to these changes is useful to analyze the hydrological variations in coastal waters (Troccoli et al., 2004). The water quality assessment by using biological material approach, such phytoplankton, will reflect changes caused by a decrease in quality a water.

*) Corresponding author
© Ilmu Kelautan, UNDIP

ijms.undip.ac.id
DOI: 10.14710/ik.ijms.26.1.37-44

Received: 24-11-2020
Accepted: 01-02-2021

Bangka Belitung Islands Province has great tin potentials both in land and coastal areas. Mining activities are carried out from the plains to the coastal and marine areas. As the result of the heavy metal mining, it is inevitable that the occurrence of ecological damage on land to sea waters (Rachman, 2019). Exploitation of tin mining in coastal areas and oceans can cause toxin and waste. It causes low concentration of phosphate and nitrate, which are one of the limiting elements for water quality and ecosystem elements. (Lee and Jones-Lee, 2005; Haya, 2012).

Phosphate and nitrate have a vital role in the growth of phytoplankton and other aquatic plants, which are commonly used as indicators of water quality and productivity level in waters (Fachrul et al., 2005; Haya, 2012). The purpose of this study was to examine the abundance of phytoplankton in two different sites, which are Tanah Merah waters (as the mining site) and Semujur Island (away from the mining site).

Material and Methods

Phytoplankton sampling and water quality parameters were conducted in June, July, August 2018 in the Tanah Merah and Semujur Island, Bangka Belitung Islands. Tanah Merah Waters is close to the tin mining while Semujur Island Waters is far from tin mining (Figure 1.). Seawater's sampling was determined in 6 stations in each site, respectively.

Phytoplankton collection was taken vertically by using Kitahara nets (mesh size of 80 μm). Sea water measurement such as temperature, salinity, dissolved oxygen content was performed by using the water quality checker. Water samples for measurement of nitrate, phosphate, and TSS using Niskin bottle. Transparency and depth measurements were carried out in situ. While water turbidity were measured by using turbidity meter.

Nitrate concentration analysis was determined by Brusin method using spectrophotometer in 410 nm wavelength range. Meanwhile phosphate analysis was reduced using 0-1 mg.L^{-1} ascorbic acid and created phosphor-molybdate complex that had a blue colour. The colour intensity as the result of the measurement has a similar concentration with phosphor. The identification was done by spectrophotometer in 700- 880 nm range.

Phytoplankton identification and calculation were carried out using the fraction method by transferring 1 ml of the phytoplankton sample into the Sedgewick Rafter Counting Cell (SRCC). The identification was executed under a light microscope with a magnification of 100x by using Sedgewick Rafter Counting Cell (SRCC). The identification was performed up to genus level based on Tomas (1997) and Yamaji (1979).




Figure 1. Location Sampling on Tanah Merah Waters and Semujur Island Waters

Data analysis

The statistical analysis by non-parametric and parametric test was performed in this study. Kruskal-Wallis test is appropriate in cases in which the error distribution of a response variable is not normal (Zuur et al., 2009). T-test was carried out in case the data was normal and homogeny. The diversity index represents the condition of the ecosystem (Zaghloul et al., 2020). Diversity, Evenness and Dominance Indices were calculated based on Odum (1988). The analysis was conducted by using SPSS version 17, to determine the difference in abundance, diversity, uniform, and dominance of phytoplankton between Tanah Merah Waters and Semujur Island Waters.

Result and Discussion

Phytoplankton abundance

This study found that the abundance of phytoplankton in Semujur Island Waters was significantly higher than Tanah Merah ($\alpha = 0.01$; Figure 2). In Semujur Island Waters, the highest abundance of phytoplankton was found at station 4 that had abundance 318,000 cells.L⁻¹ (Figure 2; Table 1 and 2). The range of average abundance values of Phytoplankton in Tanah Merah Waters was 182,000 - 200,000 cells.L⁻¹ (Table 1; Figure 2). Besides, the range of the average value of phytoplankton abundance in Semujur Island Waters was 286,000-318,000 cells.L⁻¹ (Table 2; Figure 2).

By following the results, the classification system represented in the Tanah Merah Waters (mining area) and Semujur Island Waters (non-mining area) are included in the oligotrophic category because the abundance values in these

14
two waters have an abundance of phytoplankton <4.16x10⁶ cells.L⁻¹. (Spatharis and Tsirtsis, 2010)

The result of observations and data analysis shows the relative abundance (%) of genera Ceratium was greater in Tanah Merah Waters than at the Semujur Island waters, and then followed by genera Thalassiothrix and Chaetoceros (Figure 3 and Table 1 and 2). Widianingsih et al. (2007) reported that at East Belitung Island Waters, there were many Ceratium and Rhizosolenia but Thalassiothrix is only found at one station. In non-mining and mining areas, Chaetoceros shows a considerable abundance at each station at the Bangka Waters (Rachman, 2019). On the other hand, the abundance Thalassiothrix was greater in Semujur Island Waters than in the Tanah Merah Waters, followed by Chaetoceros and Thalassionema (Table 1; Figure 3).

The abundance of diatoms in the Semujur Island Waters (non-mining area) was significantly higher than in Tanah Merah Waters (mining area), this case indicated that the seawater quality in Semujur Island Waters better than Tanah Merah Waters. The average value of the diversity index range at Semujur Island Waters ranges from 1.971-2.229 (Figure 4.). This shows that Semujur Island Waters has more diverse phytoplankton than the Tanah Merah Waters. T-test shown that the diversity in Semujur Island was significantly higher than that in Tanah Merah (t-test, $p = 0.009$), followed by uniform and dominance also were significantly higher in Semujur Island than that in Tanah Merah (t.test, both $p = 0.05$, respectively). The average value of the diversity index range at Semujur Island Waters ranges from 1.971-2.229 (Figure 4.). This shows that Semujur Island had more diverse phytoplankton than that in Tanah Merah.




Figure 2. The Average of value abundant of Phytoplankton (x1000 cells.L⁻¹) on Tanah Merah Waters and Semujur Island Waters




Figure 3. The Average of Relative Abundance (%) Phytoplankton at Tanah Merah Waters and Semujur Island Waters




Figure 4. The Average Value of Diversity Index, Evenness Index and Dominance Index from June, July and August 2018 in the Tanah Merah Waters and the Semujur Island Waters.

Water quality parameter

The seawater temperature during sample collection was significantly different between Sites (t-test, $P=0.02$). The temperature was consistent at 31°C in Tanah Merah while it more fluctuated in Semujur Island (30-31°C). The temperature was colder in Semujur Island than that in Tanah Merah. Studies have revealed that phytoplankton growth depends on available sunlight, temperature, and nutrient levels. Cold temperature tends to have more nutrients than warm water, phytoplankton tends to be more plentiful where seawater was cold. Also, when surface seawater is cold, it is easier for deeper

water to rise to the surface, bringing nutrients to sunlit areas where phytoplankton can use them. Range of salinity still suitable for phytoplankton growth.

The diffusion rate of O_2 in sea water has been treated physical-chemically process. It is shown that although the activity of oxygen in the seawater remains constant when pressure O_2 and temperature are constant (equilibrium conditions), the diffusion rate of oxygen will nevertheless vary proportionally to the solubility of oxygen in sea water, which decreases with increasing salinity (Millero and Sohn, 1992).

Table 1. Average of Abundance of Phytoplankton (cells.L⁻¹) in the Tanah Merah Waters

No	Genera	Station in Tanah Merah Waters					
		1	2	3	4	5	6
1	Skeletonema	-	-	-	-	16,000	12,000
2	Melosira	-	-	-	-	-	-
3	Coscinodiscus	8,000	8,000	12,000	8,000	8,000	8,000
4	Rhizosolenia	16,000	16,000	12,000	16,000	12,000	8,000
5	Bacteriastrium	-	4,000	4,000	-	-	-
6	Chaetoceros	24,000	28,000	24,000	28,000	16,000	20,000
7	Biddulphia	-	-	8,000	4,000	-	-
8	Leptocylindrus	12,000	8,000	8,000	4,000	4,000	8,000
9	Thalassiothrix	28,000	24,000	24,000	28,000	16,000	16,000
10	Thalassionema	24,000	24,000	20,000	20,000	20,000	16,000
11	Pleurosigma	-	-	4,000	-	-	12,000
12	Nitzschia	16,000	20,000	16,000	16,000	24,000	20,000
13	Noctiluca	-	-	-	-	-	-
14	Ceratium	64,000	60,000	68,000	64,000	70,000	76,000
Total (cells.L ⁻¹)		192,000	192,000	200,000	188,000	186,000	196,000

This is evidence that there is a relationship between salinity levels and DO content in marine waters.

Based on the results of DO measurements in June, July and August in the Semujur Island Waters, it has an average value of 6.87-6.96 ppm (Table 3). Whereas, at the Tanah Merah Waters, the measured DO average value are 6.59 - 6.89 ppm. There is difference the mean of DO between Tanah Merah Waters (Mining Area) and Semujur Island Waters (t-test; one tail; t count = 4.778; df =5; P= 0.05) (Table 4.). However, the DO values in both waters are still within normal levels. Dissolved oxygen is important role to many forms of life including fish, invertebrates, bacteria and others marine organisms. These organisms use oxygen in respiration. Phytoplankton require dissolved oxygen for respiration when there is no light for photosynthesis. The amount of dissolved oxygen needed varies from creature to creature. Bottom feeders, crabs, oysters and worms need minimal amounts of oxygen (1-6 mg.L⁻¹), while shallow water fish need higher levels (4-15 mg.L⁻¹).

Microbes such as bacteria and fungi also require dissolved oxygen. These organisms use DO to decompose organic material at the bottom of a body of water. Microbial decomposition is an important contributor to nutrient recycling. However, if there is an excess of decaying organic material (from dying algae and other organisms), in a body of water with infrequent or no turnover (also known as stratification), the oxygen at lower water levels will get used up quicker.

Likewise, the average pH value in Tanah Merah Waters has a range of 8.21-8.30 and at the

Semujur Island Waters are 8.31-8.39 (Table 3). The range of the average pH value in Semujur Island Waters is higher than Tanah Merah waters. Statistical test results show that there is difference the mean of pH value between Tanah Merah Waters and Semujur Island Waters (t-test one tail; t count = 6.751; df = 5; P= 0.05) (Table 4.) The average pH value in the Semujur Island Waters is more alkaline than the average pH value in The Tanah Merah Waters. The average pH value in the Tanah Merah Waters and Semujur Island Waters is within normal limits. The average pH for sea water is 8.2 but can range between 7.5 and 8.5 depending on the local conditions. The average pH for sea water is 8.2 but can range between 7.5 and 8.5 depending on the local conditions (Dickson, 1993). Fluctuation in pH value can stress marine organisms and may ultimately lead many species to leave the area or die.

The results of transparency measurements in the Tanah Merah Waters had an average value range of 0.09-1.20 m, while those in the Semujur Island Waters had an average value range of 3.90-5.90 m (Table 3.). According to statistica test, there is difference the mean of transparency value between the Tanah Merah Waters and the Semujur Island Water (Table 4.).

As we know, turbidity is a measure of light scattering caused mainly by suspended sediment, microalgae, microorganisms and other particulate matter in the seawater column (Macdonald et al., 2013). Based on the results of turbidity measurements in June, July and August 2018, the results showed that Tanah Merah Waters had an average value range of 12.2-22.5 NTU and in

Semujur Island Waters were 3.60-7.80 NTU (Table 3.). There is statistically difference the mean of turbidity between the Tanah Merah Waters and the Semujur Island Waters. The average value of turbidity in the Tanah Merah waters was higher than in the Semujur Island Waters. This proves that the higher the turbidity value the higher the level of pollution in a waters (Jones *et al.*, 2015). The turbidity value in the Tanah Merah Waters is still the same compared to the research results (Jones *et al.*, 2015) which show that the mean turbidity for Barrow Island, Café Lambert and Burrup Peninsula, Australia had value range 3 - 5 NTU. The value of turbidity can be gave effect on Total Suspended Solid (TSS). Total suspended solid (TSS) and turbidity is one of the important indicators that could be determining the quality of the seawater. The greater the turbidity value, the greater the TSS value in a water.

In the Semujur Island Waters, the highest mean TSS (Total Suspended Solid) values were found at stations 2 and 3 (15.8 and 16.2 mg.L⁻¹) (Table 3.). Statistical results show there is no

significant difference the mean value of TSS between Tanah Merah Waters and Semujur Island Waters. (t-test one tail; t count= 0.177; df = 5; P= 0.05) (Table 4.). Based on measurements in the Tanah Merah Waters, the average nitrate concentration was 0.15-0.18 mg.L⁻¹ (Table 3). Meanwhile, in the Semujur Island Waters had range 0.14-0.33 mg.L⁻¹. The highest nitrate concentration was found in Station 1 in the Semujur Island Waters (0.33 mg.L⁻¹). This indicates that there is nutrient enrichment that causes phytoplankton growth. The range of nitrate in natural waters is between 0.01-0.7 mg.L⁻¹ and nitrate levels in natural waters are almost never more than 0.1 mg.L⁻¹, but if the nitrate level is greater than 0.2 mg.L⁻¹, it will result in eutrophication (enrichment) which further stimulates the rapid growth of algae and water plants. (Gypens *et al.*, 2009). However, based on analysis statistic, there is no significant difference the mean value of nitrate between the Tanah Merah Waters and the Semujur Island Waters. (t-test one tail; t count 0.753; df= 5; P= 0.05) (Table 4).

Table 2. Average of Abundance of Phytoplankton (cells.L⁻¹) in the Semujur Island Waters

No	Genus	Station in Semujur Island Waters					
		1	2	3	4	5	6
1	Skeletonema	4,000	-	8,000	8,000	-	12,000
2	Melosira	16,000	-	8,000	10,000	-	-
3	Coscinodiscus	8,000	8,000	12,000	8,000	8,000	8,000
4	Rhizosolenia	8,000	8,000	16,000	12,000	12,000	16,000
5	Bacteriastrum	12,000	12,000	8,000	8,000	8,000	10,000
6	Chaetoceros	56,000	68,000	60,000	56,000	80,000	56,000
7	Biddulphia	8,000	8,000	12,000	12,000	8,000	8,000
8	Leptocylindrus	8,000	8,000	12,000	12,000	4,000	4,000
9	Thalassiothrix	64,000	88,000	72,000	96,000	60,000	64,000
10	Thalassionema	40,000	44,000	44,000	36,000	56,000	44,000
11	Pleurosigma	-	-	-	-	-	-
12	Nitzschia	24,000	24,000	16,000	16,000	20,000	24,000
13	Noctiluca	8,000	8,000	4,000	4,000	4,000	4,000
14	Ceratium	32,000	36,000	44,000	40,000	36,000	36,000
Total (cells.L ⁻¹)		288,000	312,000	316,000	318,000	296,000	286,000

Table 3. The Value Average of Water Quality from June, July and August 2018 at Tanah Merah Waters and Semujur Island.

Location	Temperature (°C)	Salinity (ppt)	pH	Transparency (m)	DO (ppm)	Nitrate (ppm)	Phosphate (ppm)	Turbidity (NTU)	TSS (mg.L ⁻¹)
Tanah Merah									
1	31	31.00	8.27	0.76	6.60	0.15	0.03	19.5	11.8
2	31	34.00	8.23	0.09	6.63	0.16	0.05	22.5	8.2
3	31	35.00	8.30	1.20	6.89	0.16	0.03	15	6.9
4	31	34.00	8.21	0.09	6.70	0.16	0.04	14.9	9.8
5	31	32.00	8.30	1.10	6.54	0.17	0.03	12.2	7.6
6	31	32.00	8.23	1.20	6.79	0.18	0.02	12.5	6.8
Semujur Island									
1	30	33.00	8.32	5.00	6.87	0.33	0.01	4.82	3.8
2	31	35.00	8.36	4.00	6.89	0.14	0.02	7.8	15.8
3	30	37.00	8.39	4.60	6.92	0.15	0.02	4.14	16.2
4	30	35.00	8.31	5.90	6.96	0.13	0.02	7.22	5.2
5	31	33.00	8.36	4.90	6.89	0.14	0.02	3.6	3.8
6	30	35.00	8.36	3.90	6.94	0.14	0.01	4.88	3.2

Table 4. The result of statistic analysis water quality parameter between Tanah Merah Waters and Semujur Island Waters

Water Quality Parameter	Statistic analysis t-test one-tail ; df = 5; P= 0.05
Temperature	t count = 3.162; Different significant
Salinity	t count = 5.000; Different significant
DO	t count = 4.788; Different significant
pH	t count = 6.751; Different significant
Tranparancy	t count = 9.348; Different significant
Turbidity	t count = 7.904; Different significant
Nitrate	t count = 0.753; No different significant
Phosphate	t count = 5.000; Different significant
TSS	t count = 0.177; No different significant

Phosphate and Nitrate are essential nutrients for marine plant in the marine ecosystem. All living marine organisms require the nutrients nitrogen and phosphorus for their growth, metabolism and reproduction. Additionally, nitrogen is a component of amino acids, nucleic acids and other cell components, while phosphorus is found primarily in nucleic acids, phospholipids and adenosine triphosphate (ATP) (Patey et al., 2008).

There is statistically difference the mean of phosphate value between the Tanah Merah Water and the Semujur Island Water (one tail t-test; t-count = 5,000; df = 5; P= 0.05) (Table 4.). The high phosphate content at the Tanah Merah Waters (0.02-0.05 mg.L⁻¹) is thought to be because this station is close to the coastal tin mining location. This area gets a flow of nutrients from the land in the form of sediment flow or sediment from rivers or land when it rains, and also from the results of human activities. Apart from this, the stirring process at the bottom of the waters and the circulation process from the surface will greatly affect the amount of phosphate content. Meanwhile, on the Semujur Island Waters, which is far from the mainland, has a lower phosphate content (0.01-0.02 mg.L⁻¹). The low concentration of phosphate in the Semujur Island Waters is due to the utilization by phytoplankton, therefore the average phytoplankton abundance value in the Semujur Island Waters is higher than the Tanah Merah Waters. Patey et al. (2008) stated that the phosphorus content is generally declining further and further out to sea (off shore). In coastal waters and continental shelves, mining activities and land flows will carry away waste, tailings, and other inland phosphate sources and result in greater concentration from the surroundings.

Conclusion

The average abundance of phytoplankton in Semujur Island Waters (286,000-318,000 cells.L⁻¹) is higher than Tanah Merah Waters (188,000-200,000 cells.L⁻¹). The magnitude of the abundance value in the Semujur Island Waters is also followed by the large average value of the diversity index and

the low average value of the dominance index. However, the evenness index for both waters is included in the medium category. In the Semujur Island Waters, Diatoms (Thalassiothrix, Chaetoceros and Thalassionema) are more dominant than the Dinophyceae group. Whereas in the Tanah Merah Waters, the genera Ceratium belong to class Dinophyceae was more dominant than the class Bacillariophyceae. The influence of tin mining on the Tanah Merah Waters results in a lower average value of phytoplankton abundance compared to the Semujur Island Waters, where is far from the tin mining area. The magnitude of the concentration of Phosphate, nitrate, DO, pH, transparency and salinity in the Semujur Island Waters is compared to the Tanah Merah Waters, which results in the large average value of phytoplankton abundance in Semujur Island Waters. On the contrary, the average value of turbidity and Total Suspended Solid in Tanah Merah Waters can cause low abundance of phytoplankton. It can be concluded that tin mining can disrupt the abundance and composition of phytoplankton as a primary producer of waters.

References

- Dickson, A.G. 1993. The Measurement of seawater pH. *Mar. Chem.* 44:131-142. doi: 10.1016/0304-4203(93)90198-W
- Fachrul, M.F., Haeruman, H. & Sitepu, L.C. 2005. Komunitas Fitoplankton Sebagai Bio-Indikator Kualitas Perairan Teluk Jakarta, *Proceeding Seminar Nasional MIPA UI, Fakultas MIPA Universitas Indonesia, Depok*. 25-26 November 2005.
- Gypens, N., Borges, A.V. & Lancelot, C. 2009. Effect of eutrophication on air-sea CO₂ fluxes in the coastal Southern North Sea: a model study of the past 50 years. *Global Change Biology*, 15:1040-1056. doi: 10.1111/j.1365-2486.2008.01773.x
- Haya, L.O.M.Y. 2012. Distribution Patterns of Phosphate and Nitrate and Relation to Development of Seaweed farming In North Moramo Waters. *J. Fish. Mar. Sci.*, 1(2):24-32.

- Jones, R., Fisher, R., Stark, C. & Ridd, P. 2015. Temporal Patterns in Seawater Quality from Dredging in Tropical Environments. *Plos One*. 10(10):e0137112. doi: 10.1371/journal.pone.0137112.
- Lee, G.F. & Jones-Lee, A. 2005. Eutrophication (excessive fertilization). *Water Encyclopedia*, 3:107-114. doi: 10.1002/047147844X.sw1606
- Macdonald, R.K., Ridd, P.V., Whinney, J.C., Larcombe, P. & Neil, D.T. 2013. Towards Environmental Management of Water Turbidity Within Open Coastal Waters of The Great Barrier Reef. *Mar. Poll. Bull.*, 72:82-94. doi: 10.1016/j.marpolbul.2013.07.026PMID:23948091.
- Millero, F.S. & Sohn, M.L. 1992. Chemical Oceanography. CRC Press. London
- Nguyen, G.T. & Nhlen, H.T.H. 2020. Phytoplankton-Water Quality Relationship in Water Bodies in the Mekong Delta, Vietnam. *App. Environ. Res.*, 42(2):1-12. doi: 10.35762/AER.2020.42.2.1
- Odum, W.E. 1988. Comparative ecology of tidal freshwater and salt marshes. *Ann. Rev. Ecol. Sys.*, 19(1):147-176. doi: 10.35762/AER.2020.42.2.1.
- Patey, M.D., Rijkenberg, M.J.A., Statham, P.J., Stinchcombe, M.C., Achterberg, E.P. & Mowlem, M. 2008. Determination of nitrate and phosphate in seawater at nanomolar concentrations. *Trends in Anal. Chem.*, 27(2): 169-182. doi: 10.1016/j.trac.2007.12.006
- Reynolds, C., Huszar, V., Kruk, C., Naselli-Flores, L. & Melo, S. 2002. Towards a functional classification of the freshwater phytoplankton. *J. Plankton. Res.* 24:417-428. doi: 10.1093/plankt/24.5.417
- Rachman, A. 2019. Phytoplankton Community Structure in the Waters Around The Coastal Tin Mining of West Bangka. *J. Teknol. Ling.*, 20(2): 189-203. doi: 10.29122/jtl.v20i2.2938
- Spatharis, S. & Tsirtsis, G. 2010. Ecological Quality Scales Based on Phytoplankton for the Implementation of Water Framework Directive in the Eastern Mediterranean. *Ecol. Indic.*, 10(4):840-847. doi: 10.1016/j.ecolind.2010.01.005
- Tomas, C.R. 1997. Identifying Marine Phytoplankton. Academic Press. San Diego. California. USA. 858
- Troccoli, G.L., Herrera-Silveira, J.A. & Comín, F.A. 2004. Structural variations of phytoplankton in the coastal seas of Yucatan, Mexico. *Hydrobiologia*, 519: 85-102. doi: 10.1023/B:HYDR.0000026487.78497.b6
- Widianingsih, R. Hartati, A. Djamali & Sugestini. 2007. Abundant and Horizontal Distribution of Phytoplankton at East Coastal Belitung Island Waters. *Ilmu Kelautan: Indonesian Journal of Marine Science*, 12(1): 6-11.
- Yamaji, I. 1979. Illustrations of the Marine Plankton of Japan. Hoikusha. Publisher: Hoikusha, Japan.
- Zaghloul, F.A.R., Khairy, H.M., & Hussein, N.R. 2020. Assessment of phytoplankton community structure and water quality in the Eastern Harbor of Alexandria, Egypt. *Egypt. J. Aqua. Res.*, 46(2):145-151. doi: 10.1016/j.ejar.2019.11.008
- Zuur, A.F., Ieno, E.N., Walker, N.J., Saveliev, A.A., & Smith, G.M. 2009. Mixed effects models and extensions in ecology with R. *Springer*. 71-100. doi: 10.1007/978-0-387-87458-6_4.

Phytoplankton community structure and it's relationships with water quality in Bangka Island, Indonesia

ORIGINALITY REPORT

12%	%	12%	%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	Daoliang Li, Shuangyin Liu. "Sensors in Water Quality Monitoring", Elsevier BV, 2019 Publication	2%
2	E Juniardi, Sulistiono, S Hariyadi, A Ervinia, A Asriansyah, T Hestiranoto, B Subhan, A Ibrahim. "Macrozoobenthos community structure in sea waters of Panjang and Pamujan Besar Islands, Banten Bay, Indonesia", IOP Conference Series: Earth and Environmental Science, 2022 Publication	2%
3	Rolf Lange, Hans Staaland, Arvid Mostad. "The effect of salinity and temperature on solubility of oxygen and respiratory rate in oxygen-dependent marine invertebrates", Journal of Experimental Marine Biology and Ecology, 1972 Publication	1%
4	Udhi E. Hernawan, Susi Rahmawati, Rohani Ambo-Rappe, Nurul D.M. Sjafrie et al. "The	1%

first nation-wide assessment identifies valuable blue-carbon seagrass habitat in Indonesia is in moderate condition", Science of The Total Environment, 2021

Publication

5

Matthew D. Patey, Micha J.A. Rijkenberg, Peter J. Statham, Mark C. Stinchcombe, Eric P. Achterberg, Matthew Mowlem.

"Determination of nitrate and phosphate in seawater at nanomolar concentrations", TrAC Trends in Analytical Chemistry, 2008

Publication

1 %

6

Muhammad Arif Salsabil, Astri Rinanti, Melati Ferianita Fachrul. "Analysis of water quality in Maninjau Lake, West Sumatera, Indonesia using phytoplankton", MATEC Web of Conferences, 2018

Publication

1 %

7

V A Fabiani, H Aldila, Anggraeni, Nur'aini. "Green synthesis and characterization of Zinc Ferrite (ZnFe O) nanocomposite via Griff. natural extract ", IOP Conference Series: Earth and Environmental Science, 2020

Publication

1 %

8

Cristiana Manullang, Intan Herwindra Millyaningrum, Akira Iguchi, Aika Miyagi, Yasuaki Tanaka, Yukihiro Nojiri, Kazuhiko Sakai. " Responses of branching reef corals

<1 %

and to elevated temperature and pCO²,
PeerJ, 2020

Publication

9

Alvarez-Gongora, C.. "Variations of
phytoplankton community structure related
to water quality trends in a tropical karstic
coastal zone", Marine Pollution Bulletin,
200601

Publication

<1 %

10

Hamka Lodang, Nani Kurnia. "Distribution and
Abundance of Plankton in The Downstream of
Jeneberang River", Journal of Physics:
Conference Series, 2019

Publication

<1 %

11

Arief Rubiana Basarah, Patmawati, Muhamad
Amin, Ahmad Fahrul Syarif, Mu'alimah
Hudatwi, Sapto Andriyono. "Nutritional Fact of
Traditional Cracker (Amplang and Getas) from
Bangka Belitung", IOP Conference Series:
Earth and Environmental Science, 2022

Publication

<1 %

12

M F Fachrul, D I Hendrawan, A Rinanti. "The
study of N:P ratio for estimating nutrient
limitation at shallow lakes waters in Depok
City West Java, Indonesia", IOP Conference
Series: Earth and Environmental Science, 2021

Publication

<1 %

13

Hasim Hasim, Mohammad Akram, Yuniarti Koniyo. "Kinerja Kepadatan Spirulina Sp. yang diberi Salinitas Berbeda pada Meda Kultur Walne", Jurnal Sumberdaya Akuatik Indopasifik, 2022

Publication

<1 %

14

Rozirwan, Fauziah, Puspa Indah Wulandari, Redho Yoga Nugroho et al. "Assessment distribution of the phytoplankton community structure at the fishing ground, Banyuasin estuary, Indonesia", Acta Ecologica Sinica, 2022

Publication

<1 %

15

Tran Thi Kim Hong, Nguyen Thanh Giao. "Analysis of Surface Water Quality in Upstream Province of Vietnamese Mekong Delta Using Multivariate Statistics", Water, 2022

Publication

<1 %

16

Andrea García-Rojas, Hannia Vega Bolaños. "Cambios estacionales del fitoplancton en el área marina de pesca responsable de Paquera-Tambor, Golfo de Nicoya, Costa Rica", Revista Ciencias Marinas y Costeras, 2016

Publication

<1 %

17

Julián Plaza, Evelyn Castro, Camila Sayes, Yanett Leyton, Carlos Riquelme. "cultivated

<1 %

on integrated system with water recirculation and photovoltaic energy in the coastal area of Atacama desert ", Journal of Applied Aquaculture, 2021

Publication

18

E Susanti, M R Widoretno, E Prihatinningtyas, I Akhdiana, R Riffiani, C Henny, R L Toruan.

"Phytoplankton Diversity in Jakarta Bay Estuary, Indonesia", IOP Conference Series: Earth and Environmental Science, 2022

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Phytoplankton community structure and it's relationships with water quality in Bangka Island, Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8