

# Exploration of Bioactive Compounds Potency of Extract Namanereis sp. (polychaeta: Annalida) as an Antibacterial Agent Against Escherichia coli and Staphylococcus aureus.

*by* Widianingsih Widianingsih

---

**Submission date:** 17-Jan-2023 09:09AM (UTC+0700)

**Submission ID:** 1993866240

**File name:** 35695-127750-1-PB.pdf (184.68K)

**Word count:** 4446

**Character count:** 24538

## Exploration of Bioactive Compounds Potency of Extract *Namanereis* sp. (Polychaeta: Annelida) as an Antibacterial Agent Against *Escherichia coli* and *Staphylococcus aureus*

Delianis Pringgenies<sup>1\*</sup>, Endah Sari<sup>1</sup>, Widianingsih Widianingsih<sup>1</sup>, Anuwat Nateewathana<sup>2</sup>

<sup>1</sup>Department of Marine Sciences, Faculty of Fisheries and Marine Sciences, Diponegoro University  
Jl. Prof. H. Soedarto, S.H., Tembalang, Semarang, Jawa Tengah, 50275 Indonesia

<sup>2</sup>Department of Marine and Coastal Resources, Bangkok, Thailand  
Thung Song Hong, Lak Si, Bangkok 10210, Thailand  
Email: delianispringgenies@lecturer.undip.ac.id

### Abstract

<sup>13</sup>  
*Namanereis* sp. belongs to class Polychaeta, a group of main benthic community in marine ecosystems. The purpose of this study to determine the potential of bioactive compounds from *Namanereis* sp. extract as an antibacterial agent and to determine its extract content through Gas Chromatography–Mass Spectrometry (GC-MS) analysis. Samples of *Namanereis* sp. were collected from Demak Waters, Central of Java. This research used experimental laboratory method. Samples were extracted with non-polar (N-Hexane), semi-polar (Ethyl acetate) and polar (Methanol) solutions. Different concentrations of extraction, i.e. 5, 10, 20, 40, 80, and 100 µg.disk<sup>-1</sup>. then were tested against pathogenic bacteria *Escherichia coli* and *Staphylococcus aureus*. The extract content analysis was obtained using the GC-MS method. The results showed that *Namanereis* sp. extract had a potential as an antibacterial agent. The highest inhibition zone was seen in the fraction with ethyl acetate solvent at a concentration of 100%, namely 13±3.05 mm in *E. coli* and 14±3.04 mm in *S. aureus*, which then was analysed with GC-MS. Its results found 20 bioactive compounds in the extract. The five most dominant compounds determined by the highest peak were Dihydrojasmonone (12.31%), Hexanoic acid (11.49%), Cholesterol (7.66%), Cyclohexanol (7.53%) and Isopropyl Palmitate (7.52%). The research concluded that *Namanereis* sp. possessed antibacterial potency and contained several bioactive compounds.

**Keywords:** Polychaeta, *Namanereis* sp., antibacterial, bioactive compounds

### Introduction

<sup>2</sup>  
Secondary metabolites, also called natural products of organisms, are organic compounds that are not directly involved in the normal growth, development, or reproduction of an organism. These compounds function as individual defenses against predators, parasites, diseases, used in competition between species, and to facilitate the reproduction processes such as pigmentation, production and release of pheromones, and bioactive compounds (Pringgenies, 2019).

Bioactive compounds are found in immobile biota, such as invertebrate, and have a variety of potential applications in human life, such as a source of antioxidants, antibacterial agents, anti-inflammatory substances and anticancer agents. (Pringgenies et al., 2018) Antibacterial agents are substances that can inhibit bacterial growth and can be used in the treatment of infections. Based on how it works, antibacterial agents can work as bacteriostatic and bactericidal (Ariyanti et al 2012).

*Namanereis* sp. are abundant along the coastal area of Demak, which is generally used as shrimp feed and fish bait. *Namanereis* sp. is known to be nutritionally complete that contained protein, fat, carbohydrate, ash, fatty acids and amino acids, vitamins A, B<sub>1</sub>, B<sub>6</sub>, B<sub>12</sub>, E, and P, I<sub>2</sub>, Ca, Mg, C minerals comparable to those found in fish. (Hadiyanto, 2013). Therefore, this study aims to determine the potential of *Namanereis* sp. as an antibacterial agent and determine the content of its bioactive compounds by GC-MS analysis.

### Materials and Methods

#### Sample collections

*Namanereis* sp. samples were collected from waters of Demak, Central Java province, Indonesia. By using gloves, the samples were taken from its habitat, put into plastic bags, placed in a cool box filled with ice cubes, and were immediately transported to the laboratory. The samples were then rinsed with running water and soaked in aquabidest to ensure sterile samples. The sterile samples were

\* ) Corresponding author  
© Ilmu Kelautan, UNDIP

sun-dried for 14 days. Dried samples were put in a 150 °C oven for 120 mins for ideal dehydration.

#### Extraction of *Namanereis* sp. samples

*Namanereis* sp. extraction was initiated with a multilevel maceration process. 100 grams of samples were immersed in 350 ml of n-hexane solvent for 1 x 24 hours. The results of the immersion were filtered while the residue was separated. Then the residue of n-hexane immersed was re-immersed with ethyl acetate and methanol sequentially by the same process as with the n-hexane solvent. The extraction in this study uses the solid-liquid method with rotary evaporator for extraction (Doughari, 2012). After being filtered, the filtrate was evaporated with rotary vapor at 25 °C. The extract was transferred from the Erlenmeyer flask into the vial, then the sample with ethyl acetate filtrate and with methanol filtrate were alternately extracted using rotary evaporator (Burgess et al., 2003). The extract in vials was then stored in a refrigerator at 4 °C. (Vaikundavasagom et al., 2015).

#### Positive and negative control tests

The antibacterial activity test was conducted following the procedures of Balouiri et al. (2016), Pringgenies (2010) and Setyati and Subagyo (2012). The positive control test was carried out using *Amoxicillin*, in which using bacteria at 80 µg.disk<sup>-1</sup> concentration. This test aimed to determine and measure the diameter of inhibition zone formed by *Amoxicillin* antibiotics (Pringgenies, 2013). The form of inhibition zones was observed after 24 hours of incubation, then was measured. Negative control tests were carried out on test bacteria using n-hexane, ethyl acetate, and methanol as extract solvents. This test was carried out to determine the effect of the solvent used in the formation of inhibition zone (Hasanah et al., 2012).

#### Antibacterial activity test on *Namanereis* sp.

The antibacterial activity test of *Namanereis* sp. extract was carried out using the Kirby-Bauer disc diffusion method at different concentrations. Sterile NA media in a petri dish were allowed to stand until solidified. Pure extract at a concentration of 100% was then tested against pathogenic bacteria. The dissolved extract in ethyl acetate, in the form of a paste, was diluted with a DMSO solution to make an 80 µg.µl<sup>-1</sup> stock solution at each concentration. The concentrations needed were 5, 10, 20, 40, 80, and 100 µg.disk<sup>-1</sup> (Pringgenies, 2013). The paper disk was then placed on each test medium. A ten µg.µl<sup>-1</sup> diluted extract of each treatment was dropped using a micropipette on the paper disk. The test media were stored in an incubator at 37 °C for 24 h. After 24 h,

confirmation of the formation and measurement of the inhibition zone is done using a caliper. The inhibition zone formed was a clear zone formed around the paper disk. In this experiment, each treatment was triplicated.

#### GC-MS test

*Namanereis* sp. extract was put into the instrument, which was then separated into components and solutions. All compounds in the component were identified. Afterwards, quantitative analysis of each component was conducted. The amount of compound contained in the extract was indicated by the peak on the chromatogram, while the type of compound was interpreted from the spectrum data from each peak using the referential approach using the GC-MS database (Pringgenies, 2010).

Gas chromatography-mass spectrometry analysis was performed using the Shimadzu GC-MS-TQ8030 engine with the following analysis conditions: linear velocity: 36.7cm.sec<sup>-1</sup>, purge flow: 3.0 mL.min<sup>-1</sup>, Helium carrier gas with injector temperature level at 345 °C, pressure 100.4 kPa, total flow 50 mL.min<sup>-1</sup>, column flow 0.90 mL.min<sup>-1</sup>, purge flow 3 mL.min<sup>-1</sup>, split ratio -1 and programmed column temperature of 200 °C (maintained for 1 min) to 340 °C (maintained for 10 mins) at a rate of temperature rise of 14 °C.min<sup>-1</sup>, ion source temperature of 200 °C and interface temperature of 200 °C.

## Results and Discussion

#### Extraction of *Namanereis* sp. samples

The resulting extract texture was paste-like. The extraction results also indicated that the methanol solvent produced more extracts (6.8 mg), compared to the lowest yield obtained from the ethyl acetate solvent (1.25 mg) (Table 1.).

#### Positive and negative control tests

A positive control test was carried out against the amoxicillin antibiotic, to determine the activity of the test bacteria. Amoxicillin was confirmed to form a zone of inhibition against test bacteria. The highest activity of Amoxicillin was found in *S. aureus*, while the lowest was found in *E. coli* (Table 2.). Amoxicillin is a broad-spectrum, β-lactamase antibiotic and is often used to treat various infectious diseases caused by Gram-positive and Gram-negative bacteria such as ear infections, pneumonia, streptococcal pharyngitis, skin, urinary tract, Salmonella, Chlamydia infections and Lyme disease. The indication of the formation of inhibitory zones can be observed from

**Table 1.** Extraction of *Namanereis* sp.

Solvent	Extract weight (mg)	Shape	Color
N-Hexane	3.09	paste	Blackish brown
Ethyl acetate	1.25	paste	Blackish brown
Methanol	6.8	paste	Blackish brown

the clear zone and was used as a reference for determining the level of bacterial resistance to antibiotics used, where the diameter of the clear zone formed was inversely proportional to the growth rate of pathogenic bacterial colonies. There are three levels of bacterial resistance, namely sensitive, intermediate and resistant. A bacterial species was sensitive to an antibacterial agent if a clear zone was formed and resistant if no clear zone is formed from the test results. While the level of intermediate resistance is characterized by clear zones with small diameters (Jawetz et al., 2001).

Inadequate concentration of antibiotics will eventually enhance the resistance of *S. aureus* and *E. coli*. The concentration used in this study was 80 µg.disk<sup>-1</sup>, because at that concentration *Amoxicillin* was able to penetrate the outer membrane of each test bacterium. Inhibition zones of 32±1 mm in *S. aureus* and 22.33±15.37 mm in *E. coli* (Table 2.) reinforce the findings of previous studies, which stated that the greater the concentration of the test, the greater the inhibition zone formed (Azis and Cahyadi, 2020).

The results of negative control tests with solvents showed no significant antibacterial activity. The three solvents produced no inhibition zone. Therefore, it can be concluded that the solvent does not affect the activity of the test bacteria (Table 3.). Negative control tests were also carried out to determine the level of contamination in bacterial culture media. This test was carried out in accordance

**Table 2.** Diameter of Inhibition Zone on test bacteria by *Amoxicillin*

Test Bacteria	Inhibition Zone (mm)
<i>Escherichia coli</i>	22.33±15.37
<i>Staphylococcus aureus</i>	32±1

**Table 3.** Inhibition Zone of Extracts with Various Solvents on Test Bacteria

Test Bacteria	Diameter of Inhibition Zone (mm)		
	N-Hexane	Ethyl acetate	Methanol
<i>Escherichia coli</i>	0	0	0
<i>Staphylococcus aureus</i>	0	0	0

with the solvent used in the extraction process, namely n-hexane, ethyl acetate and methanol. In the negative control test, no inhibition zone formation was found. These results can explain that the antibacterial activity of each extract was not influenced by the solvent used, but rather by the active compound contained in the extract of *Namanereis* sp. (Salwan et al., 2020).

**Antibacterial activity test on *Namanereis* sp. against test bacteria**

The qualitative test of antibacterial activity of *Namanereis* sp. extract was carried out at concentrations of 5, 10, 20, 40, 80, and 100 µg.disk<sup>-1</sup>. The extracts with six different concentrations were tested against *E. coli* and *S. aureus*. A qualitative test with a 100% extract concentration was conducted with an observation period of 1x24 h. The results of this test indicated that the fraction with ethyl acetate solvent gave the largest inhibition zone compared to the fraction with n-hexane and methanol solvents (Table 4.).

*S. aureus* and *E. coli* were used as test bacteria based on the representation of the two bacteria as gram-positive bacteria (*Methicillin-resistant S. aureus*) and gram-negative bacteria (*E. coli*). The potential of antibacterial compounds in *Namanereis* sp. extract in this research was determined by antibacterial activity test against the selected test bacteria. The inhibition zone formed around the paper disk indicated its antibacterial activity. Observations revealed that different fractions showed different antibacterial activities at the same extract concentration, at 100%. The methanol extract produced no inhibitory zone, which meant that the test bacteria were resistant to the fraction of *Namanereis* sp. extract with methanol solvent. This finding indicated a different result than previous studies on *Polychaeta*, where the best results obtained came from the methanol fraction (Elayaraja et al., 2010) in which the inhibition zones formed was 7 mm in *E. coli* and 8 mm in *S. aureus*. Bacteria were not resistant to methanol fraction extracts. One of the factors that might affect the difference in results was the difference in sample species, where previous studies used *Perinereis cultrifera* while this study using *Namanereis* sp. Differences in treatment of the sample can also affect the results; in previous studies

used fresh samples while in this study dry samples was tested.

The n-hexane fraction indicated the antibacterial activity with inhibition zones of  $2\pm 3.46$  mm in *S. aureus* and  $6\pm 6$  mm in *E. coli*. The ethyl acetate fraction produced the most significant inhibition zone with  $13\pm 3.05$  mm in *E. coli* and  $14\pm 3.04$  mm in *S. aureus*. These findings indicated that the inhibition of the ethyl acetate fraction was better when compared to the n-hexane and methanol fractions.

The extract with ethyl acetate solvent was selected to be used in further testing at different concentrations. The smallest concentration of the extract ( $5\mu\text{g.disk}^{-1}$ ) resulted in the absence of inhibition zone on all pathogenic bacteria (Table 5). Antibacterial tests were carried out to determine the efficacy of administered antibacterial substances in different concentrations. This test was only carried out on the extract of *Namanereis* sp. from the ethyl acetate fraction. The extract was diluted with DMSO solution, since it is able to dissolve polar and non-polar compounds, as well as to act as a negative control. The use of DMSO has relatively no effect on cell proliferation, so it will not interfere with the results of observation and testing of antibacterial activity with the agar diffusion method (Tjernberg *et al.*, 2006).

Table 5 showed that the results of  $5\mu\text{g.disk}^{-1}$  dilution test showed that both test bacteria were not resistant to the antibacterial compounds contained in the extract of *Namanereis* sp. At a concentration of  $80\mu\text{g.disk}^{-1}$ , gave the highest inhibitory zones formed, i.e.  $7\pm 0.86$ mm and  $7.33\pm 6.42$  for *E. coli* and *S. aureus* respectively. The resulting reaction of the test bacteria to the extract of *Namanereis* sp. was caused by differences in the structure of cell wall of test bacteria. *S. aureus* is a gram-positive bacterium, which has a relatively simpler cell wall structure compared to that of gram-negative bacteria (Ariyanti *et al.*, 2012). *E. coli* is a gram-negative bacterium that has more complex, three-layered cell wall structure with an outer layer of lipoprotein, a thick middle layer of peptidoglycan and an inner layer of lipopolysaccharide. The simpler wall structure of the gram-positive bacterial cell wall makes it easy for antibacterial compounds to penetrate the cell wall and find targets to work. Conversely, complex cell walls inhibit bioactive compounds to penetrate bacterial cell membranes, so that *E. coli* is less sensitive to these bioactive compounds (Tjernberg *et al.*, 2006). This is shown by the difference in diameter of inhibition zones of *S. aureus* and *E. coli*

The concentration of the extract used gave significantly impact to the levels of antibacterial

activity. The less concentration administered, the smaller diameter of inhibition zone formed. The antibiotics compound in *Namanereis* sp. extract worked in a similar fashion to that in *Amoxicillin* antibiotics. Inadequate concentration used in the treatment may result in the absence of inhibition zone (Rahmawati *et al.*, 2016) due to the fact that antibacterial compound will only be able to penetrate the outer membrane of pathogenic species cell walls at high concentration.

#### GC-MS test

GC-MS analysis was performed on active isolates with ethyl acetate as a fraction of the best antibacterial activity. The compounds of the extract were determined using Gas Chromatography. GC-MS analysis obtained 20 peak compounds with different percentages and retention times (Table 6.). GC-MS analysis on the sample ethyl acetate fraction showed that, of the 20 main peaks that were detected, there were only 5 with high and dominant main compounds, namely Dihydrojasmane 12.31%, Hexanoic acid 11.49%, Cholesterol 7.66%, Cyclohexanol 7.53% and Isopropyl Palmitate 7.52%. GC-MS results in the sample fraction with ethyl acetate solvent showed that 20 major peaks were detected, but there were only the more dominant five compounds detected, i.e. Dihydrojasmane 12.31%, Hexanoic acid 11.49%, Cholesterol 7.66%, Cyclohexanol 7.53% and Isopropyl Palmitate 7.52%.

*Namanereis* sp. has neuropodia with two bundles of setae. The spinigianas superior part and the falcigerous inferior portion of a skeleton collaborate in turning the CIS-jasmone into a dihydrojasmane precursor (Zealand, 1999). This process was thought to be a contributor to the high peak of Dihydrojasmane in the GC-MS test. In the last 3 decades, the demand for Dihydrojasmane in the perfume industry has been very high, making the extract of *Namanereis* sp. a viable alternative for raw material in the perfume industry. Dihydrojasmane can also be used as a test target in the development of cyclopentenone synthesis (Ho, 1972).

Hexanoic acid and cholesterol are important compounds for the human body in the process of forming brain cells and hormones. Adequate intake of hexanoic acid can prevent dementia while cholesterol can form immune system hormones (Cheon *et al.*, 2014). These benefits are believed to be a driving factor of *Namanereis* sp. consumption in the coastal community of Jepara in traditional medicine. The efficacy of hexanoic acid in accelerating gonad maturation made *Namanereis* sp. widely used natural shrimp feed by local aqua farmers (Hadiyanto, 2013).

**Table 4.** Test Results of 100% Extract with Various Solvents

Test Bacteria	Diameter of Inhibition Zone (mm)		
	N-Hexane	Ethyl acetate	Methanol
<i>Escherichia coli</i>	6±6	13±3.05	0
<i>Staphylococcus aureus</i>	2±3.46	14±3.04	0

**Table 5.** Qualitative Test Results of *Namanereis* sp. Extract with Ethyl Acetate against Test Bacteria on Various Concentrations

Test Bacteria	Diameter of Inhibition Zone (mm)					
	100%	80 µg.disk <sup>-1</sup>	40 µg.disk <sup>-1</sup>	20 µg.disk <sup>-1</sup>	10 µg.disk <sup>-1</sup>	5 µg.disk <sup>-1</sup>
<i>E. coli</i>	13.33±3.04	7±0.86	6.66±1.04	5.16±0.28	3.73±3.23	0
<i>S. aureus</i>	14±3.04	7.33±6.42	1.33±2.03	1.66±2.88	1.5±2.59	0

**Table 6.** GC-MS Analysis Results

Peak	Retention Time	Area	Concentration (%)	Compound
1	3,140	925451565	7.52	Isopropyl Palmitate
2	3,201	1413084787	11.49	Hexanoic acid, 6-(acetylamino)-3-[[3,6-bis(acetylamino)-1-oxohexyl] amino]-, methyl ester
3	3,265	766169732	6.3	Pyridin, 2-methoxy-5-nitro
4	3,644	134778365	1.10	Benzene, 1-(chloromethyl)-2-nitro-
5	4,055	1514210322	12.31	Dihydroasmone
6	4,191	673168973	5.47	Hexadecanoic acid, 14-methyl-, methyl ester
7	4,314	899788762	7.32	Acetamide, 2-[2 (hexyloxy)ethoxy]-N,N-dipropyl
8	4,395	299034246	2.43	Cyclohexene, 4-ethenyl-4-methyl-3-(1-methylethenyl)-1-(1-methylethyl)-, (3R-trans)-
9	4,610	465710426	3.79	2-Naphthalenol, 1-nitro-
10	4,798	803944875	6.54	Oxirane, hexadecyl
11	5,394	424634820	3.45	1,2-Ethanediamine, N-(2-aminoethyl)-N'-[2-[(2-aminoethyl)amino]ethyl]-
12	5,820	925649776	7.53	Cyclohexanol, 5-methyl-2-(1-methylethenyl)-, [1R-(1.alpha.,2.beta.,5.alpha.)]-
13	5,899	510604149	4.15	Oxirane, hexadecyl-
14	6,220	502474677	4.08	Di-n-octyl phthalate
15	6,290	154520178	1.26	Octadecanoic acid, 3-hydroxy-, methyl ester
16	6,345	118390854	0.96	9-Octadecenamamide, (Z)-
17	6,504	168981701	1.37	Dodecan-1-yl acetate
18	6,785	530293525	4.31	1,3-Benzenedimethanol, 2-hydroxy-5-methyl-
19	9,209	941923357	7.66	Cholesterol
20	9,765	127687644	1.04	Cyclohexanol, 2-methyl-5-(1-methylethenyl)-, (1.alpha.,2.alpha.,5.beta.)-

Cyclohexanol is the main compound in nylon production. This compound can be used as an industrial-grade solvent and plasticizer base material. Cyclohexanol also has wide range of applications, among them are in disinfectants, seasonings, pesticides, disinfectants, wood preservatives and skin softeners (Pei *et al.*, 2015). Isopropyl palmitate is most often found in skincare or beauty products. Isopropyl palmitate (IPP) ester is a dry, soft, non-oily emollient. This compound is used as an excellent solvent for mineral oil, silicon and lanolin IPP, and is widely used in cosmetics and topical drug preparations which are well-absorbed through skin contact (Liu *et al.*, 2015). The content of isopropyl palmitate in the extract of *Namanereis* sp. is thought

to be a factor that gave a benefit results of the tests on Methicillin-resistant *S. aureus* (MRSA). *S. aureus* is a pathogenic species whose infestations in humans and animals cause diseases ranging from mild skin infections, such as folliculitis and furunculosis, to life-threatening infections such as sepsis, pyogenic, deep abscesses, to fatal septicemia (Nursidika *et al.*, 2014). Finally, the *Namanereis* sp. extract can be developed into a medicine for treating skin infections caused by *S. aureus*.

**Conclusion**

*Namanereis* sp. extract with ethyl acetate fraction has potential antibacterial compounds. This

finding is evidenced by the inhibition zone of  $13 \pm 3.05$  mm against pathogenic bacteria *E. coli* and  $14 \pm 3.04$  mm to *S. aureus*. GC-MS analysis indicated that the extract of *Namanereis* sp. with ethyl fraction had 20 bioactive compounds and five were dominant compounds, namely Dihydrojasmane 12.31%, Hexanoic acid 11.49%, Cholesterol 7.66%, Cyclohexanol 7.53% and Isopropyl Palmitate 7.52%.

### Acknowledgement

We would like to thank the technicians in Laboratory Biotechnology, Dept. of Marine Sciences, Faculty of Fisheries and Marine Sciences, University of Diponegoro.

### References

- Azis, A. & Cahyadi, J. 2020. Benefits of Tiwai Onion (*Eleutherine americana*) Extract as Phytopharmaceutical Plant to Inhibit the Growth of *Vibrio harveyi* Through in-Vitro and in-Vivo. *J. Ilmiah Perikanan dan Kelautan*. 12(1): 105-112. doi: 10.20473/jipk.v12i1.128 26
- Ariyanti, N.K., Darmayasa, I.B.G. & Sudirga, S.K. 2012. Daya hambat ekstrak kulit daun lidah buaya (*Aloe barbadensis* Miller) terhadap pertumbuhan bakteri *Staphylococcus aureus* ATCC 25923 dan *Escherichia coli* ATCC 25922. *J. Biologi*. 16(1):1-4.
- Burgess, J.G., Boyd, K.G., Armstrong, E., Jiang, Z., Yan, L., Berffren, M., May, U., Psacane, T., Granmo, A. & Adams, D.R. 2003. Development of a Marine natural Product-based Antifouling Paint. *Biofouling*, 19: 197-205
- Balouri, M., Sadiki, M. & Ibsouda, S.K. 2016. Methods for in vitro evaluating antimicrobial activity: A review. *J. Pharm. Anal.* 6(2): 71-79. doi: 10.1016/j.jpha.2015.11.005.
- Cheon, J.Y., Kim, J.H., Kim, J.H., Goddeti, K.C., Park, J.Y. and Joo, S.H., 2014. Intrinsic relationship between enhanced oxygen reduction reaction activity and nanoscale work function of doped carbons. *J. Am. Chem. Soc.*, 136(25): 8875-8878.
- Doughari, J.H. 2012. Phytochemicals: Extraction methods, basic structures and mode of action as potential chemotherapeutic agents. In: Rao, V. (Ed.). *Phytochemical – A Global Perspective of Their Role in Nutrition and Health*, INTECH Open Access Publisher. doi: 10.5772/26052
- Elayaraja, S., Murugesan, P., Vijayalakshmi, S. & Balasubramanian, T. 2010. Antibacterial and antifungal activities of polychaete *Perinereis cultrifera*. *Indian J. Mar. Sci.*, 39(2):257-261
- Hadiyanto. 2013. Nilai Ekonomis Cacing Laut (Annelida: Polychaeta). *Oseana*. 28(3): 23-31
- Hasanah, N.F., Delianis, P. & Wulandari, S.Y. 2012. Karakterisasi Metabolit Sekunder Bakteri Symbion Gastropoda *Conus miles* dengan Metode GC-MS Sebagai Antibakteri MDR (Multi Drug Resisten). *J. Mar. Res.*, 1(2): 197-202
- Ho, C. 1972. Synthesis of cis-Jasmone and Dihydrojasmane. *Can. J. Chem.*, 50(16): 2718-2721.
- Jawetz, E., Melnick, J.L. & Adelberg, E.A. 2001. Mikrobiologi Kedokteran. Nani W, penerjemah. Jakarta (ID): Salemba Medika. Translate from Medical Microbiology. 196-198p.
- Liu, M., Wen, J. & Sharma, M., 2020. Solid lipid nanoparticles for topical drug delivery: Mechanisms, dosage form perspectives, and translational status. *Curr. Pharm. Des.*, 26(27): 3203-3217.
- Nursidika, P., Saptarini, O. & Rafiqua, N. 2014. Aktivitas Antimikroba Fraksi Ekstrak Etanol Buah Pinang (*Areca catechu* L.) pada Bakteri Methicillin Resistant *Staphylococcus aureus*. *Majalah Kedokteran Bandung*, 46(2): 94-99.
- Pringgencies, D. 2010. Karakteristik Senyawa Bioaktif Bakteri Symbion Mollusca Dengan GC-MS. *J. Ilmu dan Teknologi Kelautan Tropis*, 2(2):34-40. doi: 10.28930/jitkt.v2i2.7850
- Pringgencies, D. 2013. Antibacterial Activity of Sea Cucumbers Harvested from Karimunjawa. *Squalen Bulletin of Marine & Fisheries Postharvest & Biotechnology*, 8(2): 87-94
- Pringgencies, D. 2019. Bioprospeksi bahan hayati laut untuk pengembangan industri Farmasi di Indonesia. Undip Press Semarang. ISBN: 978-979-097-637-5.
- Pringgencies, D., Rudiyaniti, S. & Yudiati, E. 2018. Exploration of Sea Cucumbers *Stichopus hermanii* from Karimunjawa Islands as Production of Marine Biological Resources. *3rd International Conference on Tropical and Coastal Region Eco Development 2017. IOP Conf. Series: Earth Environ. Sci.*, 116: 012039 doi: 10.1088/1755-1315/116/1/012039
- Pei, Y., Wang, Q., Gong, X., Lei, F. & Shen, B. 2015. Distribution of cyclohexanol and cyclohexanone between water and cyclohexane. *Fluid Phase*

- Equilibria*. 394: 129-139. doi: 10.1016/j.fluid.2015.02.029
- Rahmawati, U., Agustini T.W. & Romadhon. 2016. Kajian Aktivitas Antibakteri Ekstrak Lamun (*Thalassia hemprichii*) Terhadap Bakteri *Staphylococcus aureus* dan *Escherichia coli*. In: *Prosiding Seminar Nasional Tahunan Ke V 2015 Hasil Hasil Penelitian Perikanan dan Kelautan. Fakultas Perikanan dan Ilmu Kelautan UNDIP*, pp. 466-473.
- Salwan, R. & Sharma, V. 2020. Bioactive compounds of *Streptomyces*: Biosynthesis to applications. In: Atta-ur-Rahman (Ed.) *Studies in Natural Products Chemistry* (Vol. 64; 1<sup>st</sup> ed.). Elsevier. doi: 10.1016/B978-0-12-817903-1.000 15-2.
- Setyati, W.A. & Subagyo. 2012. Isolasi dan Seleksi Bakteri Penghasil Enzim Ekstraseluler (Proteolitik, amilolitik, lipolitik dan selulolitik) yang Berasal dari Sedimen Kawasan Mangrove. *Ilmu Kelautan: Indonesian Journal of Marine Sciences*, 17(3): 164-168
- Tjernbergh, A., Natalia, M., William, J.G., Griffiths., & Dan. H. 2006. DMSO-Related Effects in Protein Characterization. *J. Biomol. Screen.*, 11(2): 131-137.
- Vaikundavasago, C., Shankar, S., Satheesh, S., Viju, N., Mary, S., & Punitha, J. 2015. Antibacterial and biofilm inhibitory activities of bacteria associated with Polychaetas. *J. Coast. Life Med.*, 3(6): 495-502. doi: 10.12980/jclm.3.2015jclm-2015-0012



# Exploration of Bioactive Compounds Potency of Extract Namanereis sp. (polychaeta: Annalida) as an Antibacterial Agent Against Escherichia coli and Staphylococcus aureus.

---

## ORIGINALITY REPORT

---

10%

SIMILARITY INDEX

%

INTERNET SOURCES

10%

PUBLICATIONS

%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- 1 Nursyirwani Nursyirwani, Widya Asmara, Agnesia Endang Tri Hastuti Wahyuni, Triyanto Triyanto et al. "Phenotype and genotype of lactic acid bacteria (LAB) isolated from the tiger grouper *Epinephelus fuscoguttatus* alimentary tract", F1000Research, 2017  
Publication 1%

---
- 2 Venugopal, V. "Bioprospecting of Marine Microalgae, Corals and Microorganisms", Aquaculture Microbiology and Biotechnology Volume Two, 2011.  
Publication 1%

---
- 3 Trang, Vu Thu, Hiroaki Takeuchi, Hayato Kudo, Shinya Katsuno, Tomoko Shimamura, Takehiro Kashiwagi, Vu Hong Son, Tetsuro Sugiura, and Hiroyuki Ukeda. "In Vitro Antimicrobial Activity of Aminoreductone against the Pathogenic Bacteria Methicillin-Resistant *Staphylococcus aureus* (MRSA)", 1%

Journal of Agricultural and Food Chemistry,  
2011.

Publication

---

4

"ION TRAP MASS SPECTRA OF COMPOUNDS",  
Elsevier BV, 1989

Publication

---

1 %

5

S. Bhatia, A.L. Ahmad, A.R. Mohamed, S.Y.  
Chin. "Production of isopropyl palmitate in a  
catalytic distillation column: Experimental  
studies", Chemical Engineering Science, 2006

Publication

---

1 %

6

Rahmi Rahayu, Atok Miftachul Hudha,  
Sukarsono, Fendy Hardian Permana. "  
Analysis of Nutritional Content of Fresh Sea  
Worm Honingka as a Potential Food Source  
for Communities ", IOP Conference Series:  
Earth and Environmental Science, 2019

Publication

---

1 %

7

Jiří Lapčík, Olinda Gimello, Vincent Ladmiral,  
Chadron Mark Friesen, Bruno Ameduri. "A  
new oligo(hexafluoropropylene oxide)-b-  
oligo(ethylene oxide) diblock surfactant  
obtained by radical reactions", Polymer  
Chemistry, 2015

Publication

---

<1 %

8

Akbar Basha Roshan, Hosur Narayanappa  
Venkatesh, Nawal Kishore Dubey, Devihalli  
Chikkaiah Mohana. "Chitosan-based

<1 %

nanoencapsulation of *Toddalia asiatica* (L.) Lam. essential oil to enhance antifungal and aflatoxin B1 inhibitory activities for safe storage of maize", *International Journal of Biological Macromolecules*, 2022

Publication

---

9

Edible Medicinal and Non Medicinal Plants, 2015.

Publication

---

10

Trianto, Agus, Idam Hermawan, Nicole Joy de Voogd, and Junichi Tanaka. "Halioxepine, a New Meroditerpene from an Indonesian Sponge *Haliclona* sp.", *CHEMICAL & PHARMACEUTICAL BULLETIN*, 2011.

Publication

---

11

I Wijayanti, E S Susilo. " Proximate content of wild and cultured eel ( ) in different part of body ", *IOP Conference Series: Earth and Environmental Science*, 2018

Publication

---

12

T M Iqbalsyah, C Nisak, F Sartika, N Saidi, F Febriani. "Identification of Antibiotic in Ethyl Acetate Fraction Produced by A Local Isolate PLS 80 Isolated from Shallow Sea Fumaroles", *IOP Conference Series: Earth and Environmental Science*, 2019

Publication

---

<1 %

<1 %

<1 %

<1 %

13

A P Wijaya, K G Bondar, E H Frederick, Y Igarashi, M T Sibero. "Identification of marine bacteria HPP.4A and HPP.T13 and its anticancer activity against P388 murine leukaemia cell", IOP Conference Series: Earth and Environmental Science, 2020

Publication

---

<1 %

14

ChemADVISOR®, Inc., . "Cross-Reference Indexes of Chemical Names and Synonyms", Regulated Chemicals Directory 1995, 1995.

Publication

---

<1 %

15

Karthika Prasad, G. S. Lekshmi, Kola Ostrikov, Vanessa Lussini et al. "Synergic bactericidal effects of reduced graphene oxide and silver nanoparticles against Gram-positive and Gram-negative bacteria", Scientific Reports, 2017

Publication

---

<1 %

16

Roomana Aafaqi. "Kinetics of esterification of palmitic acid with isopropanol using p-toluene sulfonic acid and zinc ethanoate supported over silica gel as catalysts", Journal of Chemical Technology & Biotechnology, 10/2004

Publication

---

<1 %

17

Martin Sjögren, Mia Dahlström, Erik Hedner, Per R. Jonsson, Anders Vik, Lise-Lotte Gundersen, Lars Bohlin. " Antifouling activity

<1 %

of the sponge metabolite agelasine D and synthesised analogs on ", Biofouling, 2008

Publication

---

18

Pathikrit Saha, Beom Soo Kim. "Plant extract and agricultural waste-mediated synthesis of silver nanoparticles and their biochemical activities", Elsevier BV, 2022

Publication

---

19

Susilowati, Ragil, Agus Sabdono, and Ita Widowati. "Isolation and Characterization of Bacteria Associated with Brown Algae Sargassum spp. from Panjang Island and their Antibacterial Activities", Procedia Environmental Sciences, 2015.

Publication

---

20

"Grand Challenges in Marine Biotechnology", Springer Science and Business Media LLC, 2018

Publication

---

21

"Mechanism of Pancreatic Islet Number Improvement on Hyperglycemic Wistar Rats (Rattus norvegicus) After Golden Sea Cucumber (Stichopus hermanii) Extract Administration", International Journal of Innovative Technology and Exploring Engineering, 2020

Publication

---

<1 %

<1 %

<1 %

<1 %

22

Alwar Ramanujam Padmavathi, Bose Abinaya, Shunmugiah Karutha Pandian. " Phenol, 2,4-bis(1,1-dimethylethyl) of marine bacterial origin inhibits quorum sensing mediated biofilm formation in the uropathogen ", Biofouling, 2014

Publication

---

<1 %

23

Borras, E.. "Characterisation of polycyclic aromatic hydrocarbons in atmospheric aerosols by gas chromatography-mass spectrometry", Analytica Chimica Acta, 20070205

Publication

---

<1 %

24

Enas F. Abdel Hamed, Nahed E. Mostafa, Eman M. Fawzy, Mohamed N. Ibrahim, Rasha Attia, Marwa A. Salama. "The delayed death-causing nature of Rosmarinus officinalis leaf extracts and their mixture within experimental chronic toxoplasmosis: Therapeutic and prophylactic implications", Acta Tropica, 2021

Publication

---

<1 %

25

Fuji Astuti Febria, Yossi Rahmadeni, Amri Bachtiar. "ANTIBACTERIAL POTENTIAL ETHANOL EXTRACT OF KAYU RACUN LEAF (Rhinacanthus nasutus) AGAINST Staphylococcus aureus AND METHICILLIN

<1 %

# RESISTANT Staphylococcus aureus", Jurnal Pendidikan Matematika dan IPA, 2021

Publication

26

Juan Rodríguez-Hernández. "Polymers against Microorganisms", Springer Science and Business Media LLC, 2017

Publication

<1 %

27

Martin M. Gakuubi, Kiraithe N. Micheni, Wycliffe Wanzala. " Antibacterial Activity of Essential Oil from the Fruits of (L) Lam. (Rutaceae) ", Journal of Biologically Active Products from Nature, 2017

Publication

<1 %

28

Rizwan Ahmad, Niyaz Ahmad, Atta Abbas Naqvi. "" Ziziphus oxyphylla" : Ethnobotanical, ethnopharmacological and phytochemical review", Biomedicine & Pharmacotherapy, 2017

Publication

<1 %

29

"Blue Biotechnology", Springer Science and Business Media LLC, 2017

Publication

<1 %

Exclude quotes  On

Exclude matches  Off

Exclude bibliography  On

# Exploration of Bioactive Compounds Potency of Extract Namanereis sp. (polychaeta: Annalida) as an Antibacterial Agent Against Escherichia coli and Staphylococcus aureus.

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---