

Critical review on the immunomodulatory activities of carrot's β -carotene and other bioactive compounds

by Gemala Anjani

Submission date: 06-Dec-2022 08:52AM (UTC+0700)

Submission ID: 1972671252

File name: 1-s2.0-S1756464622003735-main.pdf (1.01M)

Word count: 12042

Character count: 69287



Critical review on the immunomodulatory activities of carrot's β -carotene and other bioactive compounds

Gemala Anjani^{a,b,1,*}, Fitriyono Ayustaningwarno^{a,b}, Rafika Eviana^a

^a Nutrition Department, Faculty of Medicine, Diponegoro University, Semarang, Indonesia

^b Center of Nutrition Research (CENURE), Faculty of Medicine, Diponegoro University, Semarang, Indonesia

ARTICLE INFO

Keywords:

Daucus carota L.
Immunomodulator
Oxidative stress
Inflammation
Bioactive compound

ABSTRACT

Carrots (*Daucus carota* L.) are vegetables proven to have nutraceutical effects and beneficial for health due to its natural bioactive substances. One of the potential of carrots to maintain optimum health status is by regulating immune response. This literature review summarized the proposed immunomodulatory mechanisms of the antioxidant properties of carrot's β -carotene and other bioactive compounds such as phenolic acid, flavonoid, polyacetylene and ascorbic acid via the anti-inflammatory, antioxidant and overall (innate and adaptive) immune response modulation. Overall, carrot's bioactive compounds regulated pro-inflammatory and anti-inflammatory cytokines, reduced oxidative stress by decreasing the reactive oxygen species accumulation and improving antioxidant capacity and the expression of genes in order to prevent more damaging oxidative destruction. Carrots also modulated the immune components by regulating leukocytes, antigens, immunoglobulins, and histamine levels. Thus, the immunomodulatory activity makes carrots as a functional food source that has the potential to prevent and treat various diseases.

1. Introduction

The immune system is critical to the body's physiological and immunological function, therefore any disturbance in the immune system can lead to various diseases (Agita & Thaha, 2017; Coussens & Werb, 2002). The immune is a complex framework designed primarily of leukocytes and various immune components such as antibodies, proteins, and cytokines that act as the first defense mechanism against numerous harmful substances in the environment. This interplay of diverse immune components facilitates the establishment of an optimum

immune response (Delves & Roitt, 2000). The immune system identifies and eliminates pathogens while inducing inflammation, cell or tissue damage, cell death, and wound healing in order to preserve redox balance in immune cells in order to prevent various disorders. If homeostasis is not maintained, immune-mediated diseases, metabolic disorders, and inflammatory diseases will develop, as well as susceptibility to infectious diseases such as corona virus disease of 2019 (COVID-19) (Adapa et al., 2011; Q. Li et al., 2022; Nardy et al., 2015; Paragh et al., 2014).

Nutrition is one of the factors that influences the immune system. For

Abbreviations: ABTS, 2,2'-azino-bis(3-ethylbenzthiazolin-6-sulfonic acid); ALT, alanine transaminase; APAP, acetaminophen; ARDS, acute respiratory distress syndrome; ASCVD, atherosclerotic factors of cardiovascular disease; AST, aspartate transaminase; Bcm1^{-/-}, β -carotene 15,15'-monooxygenase 1 knockout; CD4, clusters of differentiation; con-A, concavalin-A; COVID-19, corona virus disease of 2019; COX2, cyclooxygenase-2; CUPRAC, cupric reducing antioxidant capacity; FFA, free fatty acid; FRAP, ferric reducing antioxidant power; GPx, glutathione peroxidase; HO-1, heme oxygenase-1; hscRP, high-sensitivity C-reactive protein; IBD, inflammatory bowel disease; IFN- γ , interferon γ ; IgA, immunoglobulin A; IgE, immunoglobulin E; IL, interleukin; JAK-STAT, janus kinase-signal transducer and activator of transcription; JNK, jun N-terminal kinase; LPS, lipopolysaccharide; MAPK, mitogen-activated protein kinases; MCP-1, monocyte chemoattractant protein-1; MDA, malondialdehyde; NFkB, nuclear factor kappa-B; NK-cell, natural killer-cell; NLRP3, the nucleotide-binding domain (NOD)-like receptor protein 3; NO, nitric oxide; Nrf2, nuclear factor erythroid 2-related factor 2; PGE2, prostaglandin E2; RA, rheumatoid arthritis; ROS, reactive oxygen species; SARS-CoV-2, severe acute respiratory syndrome coronavirus 2; SLE, systemic lupus erythematosus; SOD, superoxide dismutase; STAT3, signal transducer and activator of transcription 3; Th, T helper; TLR4, toll-like receptor 4; TNF- α , tumor necrosis factor- α ; Treg, regulatory T cells; USDA, United States Department of Agriculture.

* Corresponding author.

E-mail address: gemaanjani@gmail.com (G. Anjani).

¹ Present Address: Jl. Prof. Mr. Sunario, Diponegoro University, Tembalang, Semarang City, Central Java, Postal Code 50275, Indonesia, Phone/Fax: (024) 76402881.

<https://doi.org/10.1016/j.jff.2022.105303>

Received 12 August 2022; Received in revised form 2 October 2022; Accepted 19 October 2022

Available online 26 October 2022

1756-4646/© 2022 The Authors. Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).




Fig. 1. Immunomodulator Mechanism via Antioxidant and Anti-Inflammation Properties. Th1, T helper type 1; Th2, T helper type 2; TNF- α , tumor necrosis factor- α ; IFN- γ , interferon- γ ; IL, interleukin.

instance, nutritional deficit may result in decreased immune responses, and supplementation of those specific components would often restore the compromised immune responses (Venter et al., 2020). Optimal nutrition intake affects the immune response through regulating various genes expression and signaling molecules (Farhan Aslam et al., 2017). Immunomodulators are components that regulate the body's immune system (Venkatalakshmi et al., 2016). Immunomodulators are not only available in endogenous form, but also from exogenous sources whose mechanism of action is often based on antioxidant properties. Studies have been able to prove that some natural antioxidants from plants can act as natural immunomodulators. Immunomodulators from antioxidant source maintain health status through regulation of the oxidant-antioxidant balance and immune response. Antioxidants operate either directly or indirectly, by eliminating reactive oxygen-nitrogen species or enhancing the body's antioxidant balance, so that immune and inflammatory responses may be regulated (Ortuño-Sahagún et al., 2017).

Currently, many bioactive compounds from plants have been investigated for their potential benefits as immunomodulators (Venkatalakshmi et al., 2016). Turmeric, Echinacea, green tea, and chili pepper are examples of plant sources with immunomodulatory effects (Catanzaro et al., 2018; Grüter et al., 2020; Hasan et al., 2021; Rahayu et al., 2018). However, intake from some of these sources are still limited. Carrots (*Daucus carota* L.) are beneficial vegetables in the world due to its massive production and its usage as food for public daily intake, both in fresh and processed form. This vegetable has nutraceutical effects and is beneficial for health due to its high natural bioactive substances (Ahmad et al., 2019).

Carrots' main carotenoid is β -carotene, a well-known and extensively researched carotenoid. Carotenoids are phytonutrients that give fruits and vegetables their unique yellow, orange, and red hues. β -carotene provides a variety of health benefits in addition to providing color in food. It has a high antioxidant capacity and provides a number of health advantages, including decreasing the risk of certain illnesses and cancers, strengthening the immune system, and protecting against age-

related eye degeneration (Gul et al., 2015).

A recent review has shown that carrots act as antioxidants, antidiabetic, antimicrobial, nephroprotective, hepatoprotective, cardioprotective, anti-inflammatory and wound healing properties (Al-Snafi, 2017). Moreover, carrots can also improve the immune system and eye health (da Silva Dias, 2014; Mahammad Shakheel et al., 2017). The previous review examined the protective effect of β -carotene against gastric cancer based on epidemiological surveys (Chen et al., 2021). In addition, β -carotene also plays a role in stimulating lipid catabolism, which metabolizes lipids, which is one of the atherosclerotic factors of cardiovascular disease (ASCVD) (Miller et al., 2020).

Although prior review have addressed the health benefits of carrot bioactive compounds (da Silva Dias, 2014), especially β -carotene, the specific immunomodulatory activities have not been well investigated as the mechanism is known to be exceedingly complicated and remain unclear. An experimental research discovered that administration of carrot juice improved antioxidant capacity and decreased malondialdehyde after 3 months consumption of carrot juice (Potter et al., 2011). A randomized cross-over study also noted an improvement in interleukin-2 (IL-2) and tumor necrosis factor- α (TNF- α) in healthy men after consumption of carrot juice or tomato juice in healthy men after conducting 2-weeks low carotenoid periods (Watzl et al., 2003). Therefore, the purpose of this literature review is to explore into the potential immunomodulatory mechanism of carrots' β -carotene and its other bioactive compound.

2. Immunomodulator

Immunomodulator is defined as substance that modulate the immunological response, either decreasing the immune response as a type of therapy for autoimmune diseases or enhancing the immune response to immuno-compromised conditions (Ilyas et al., 2016). Immune system is defined as the body's natural defense system against disease and disorders (Sharma et al., 2017). The immune system is

divided into innate and adaptive immune systems, both of which determine a person's ability to fight harmful agents. Innate immunity is known as the first defense against pathogens and consists of epithelial protection, antimicrobial proteins and peptides, humoral and cellular components. Then, the adaptive immune system was activated by the innate immune systems, which contributed in the foreign antigen recognitions (Tyagi, 2016). The adaptive immune responses consist of B lymphocytes with antigens, T lymphocytes and regulatory T cells (Treg) (Sharpe & Mount, 2015).

The main source of cytokines in the body is the T lymphocytes, in which these cells have specific antigen receptors that enable pathogen recognitions. T lymphocytes expressing the clusters of differentiation 4 (CD4) are known as T helper (Th) cells. This subset is further subdivided into Th1 and Th2 cells, which produce Th1 and Th2 cytokines, respectively. Th1 cytokines tend to act as pro-inflammatory agents for intracellular pathogens elimination. Th1 cytokines include TNF- α , interferon γ (IFN- γ), interleukin 1 β (IL-1 β), IL-2 and interleukin 18 (IL-18). Th2 cytokines, on the other hand, are known to be more complex. Th2 cytokines include the immunoglobulin response-related cytokines such as interleukin 4 (IL-4), interleukin 5 (IL-5), interleukin 9 (IL-9), and interleukin (IL-13), as well as pro-inflammatory interleukin 6 (IL-6) and anti-inflammatory interleukin (IL-10). As a result, a balance of Th1 and Th2 responses is required in order to combat pathogens effectively while avoiding uncontrollable inflammation (Berger, 2000).

Inflammation is known as a normal physiological response of the body's immune system to counter harmful stimuli such as infections, damaged cells and toxic compounds. Inflammatory responses, whether acute or chronic, can cause tissue damage or disorders in the heart, pancreas, liver, kidneys, lungs, brain, gastrointestinal tract, and reproductive system. Infectious and non-infectious agents, as well as cell damage, usually activate inflammatory signaling pathways such as nuclear factor kappa-B (NF κ B), mitogen-activated protein kinases (MAPK), and janus kinase-signal transducer and activator of transcription (JAK-STAT) (Chen et al., 2018). Excessive inflammation exacerbates functional disorders and tissue injuries (Bartold & Dyke, 2017). For example, the hyper-inflammation that occurs in COVID-19 causes a "cytokine storm" leading to acute respiratory distress syndrome (ARDS), pneumonia, organ failures, fibrosis, or even death (Channappanavar & Perlman, 2017; Shi et al., 2020).

During the homeostatic state, several endogenous and external factors contribute to immunosuppression and immuno-stimulation (Jantan et al., 2015), thus affecting the innate and adaptive immune response (Fig. 1). Immune response modulation occurred to regulate the body's immune response in therapeutic and preventive attempts against causative agents (Rasheed et al., 2016; Razali et al., 2016). The use of immunomodulators is presently one of the most pressing issues in the treatment of numerous diseases (Sharma et al., 2015).

Application of plant-derived bioactive compounds has become one of the primary areas of exploration for the natural immunomodulators (Shukla et al., 2014). The use of natural immunomodulators is currently increasingly being accepted in the world because of their wide range in preventing and treating disease via changes in immune response or oxidant-antioxidant status (Ajith et al., 2017).

The antioxidant defense system involves free radical reduction, antioxidant enzyme activation and maintenance, enhancement of other antioxidant mechanisms, and specific immunomodulatory effects (Ajith et al., 2017). Plants that serve as natural immunomodulators have the capacity to boost immunity by activating innate immune responses such as activating immune cellular components, modifying cytokine profiles, decreasing infection and inflammation. The immunomodulatory properties of these plants cannot be separated from their bioactive compounds (Jantan et al., 2015).

3. Nutrition and bioactive contents of carrots

Daucus carota L., or commonly known as carrot, is a vegetable

Table 1
Nutrition and Bioactive Contents in Carrots (per 100 g FW).

Component	Amounts	Unit	Reference	RDA (age 10–12 y.o)
Energy	41	kcal	USDA, 2015	2.05 %
Proximate		g		
Moisture	88,3	g	USDA, 2015	4,77 %
Carbohydrate	9,58	g	USDA, 2015	3,19 %
Protein	0,93	g	USDA, 2015	1,86 %
Fat	0,24	g	USDA, 2015	0,37 %
Fibre	2,80	g	USDA, 2015	10 %
Ash	0,97	g	USDA, 2015	4,77 %
Iron	0,3	mg	USDA, 2015	3,75 %
Magnesium	12	mg	USDA, 2015	7,5 %
Phosphorus	35	mg	USDA, 2015	2,8 %
Potassium	320	mg	USDA, 2015	8,21 %
Sodium	69	mg	USDA, 2015	3,63 %
Calcium	33	mg	USDA, 2015	2,75 %
Zinc	0,24	mg	USDA, 2015	3 %
Manganese	0,14	mg	USDA, 2015	26,6%
Vitamin B2	0,06	mg	USDA, 2015	4,61 %
Vitamin B3	0,98	mg	USDA, 2015	8,17 %
Folate	19	μ g	USDA, 2015	4,75 %
Vitamin C	5,90	mg	USDA, 2015	11,8 %
Total Flavonoid	4,50	mg	Bahorun et al., 2004	–
Quercetin	1,50	mg	Bahorun et al., 2004	–
Luteolin	0,80	mg	Bahorun et al., 2004	–
Kaempferol	0,60	mg	Bahorun et al., 2004	–
Total Phenolic	89,30	mg	Helström et al., 2020	–
Caffeic acid	41,10	mg	Helström et al., 2020	–
Ferulic acid	7,10	mg	Helström et al., 2020	–
p-coumaric acid	73,10	μ g	Rozan, 2017	–
Carotenoids	12,71	mg β CE	Rozan, 2017	–
α -carotene	3,38	mg	Rozan, 2017	–
β -carotene	8,21	mg	Rozan, 2017	–
Falcarinol	359–1900	μ g	Rawson et al., 2010	–
Falcarindiol	154–18000	μ g	Acworth et al., 2011	–
Falcarinyl-acetate	77–140	μ g	Acworth et al., 2011	–

Abbreviation: USDA, United States Department of Agriculture.

belonging to the *Apiaceae* family (Bussmann et al., 2020). Based on FAOSTAT 2021 data from the Food and Agriculture Organization (FAO), in 2019, nearly 44.8 million tons of carrots and radishes were widely produced for daily consumption (FAOSTAT, 2021). Carrots consist of stems and roots in which the roots consist of the periderm, outer cortex (phloem) and nucleus (xylem). The part of the carrot that is commonly consumed is the root. High quality carrots have a greater proportion of phloem than xylem (Char, 2018). The taxonomy of carrots from the Integrated Taxonomic Information System (2022) database are as follows:

Kingdom: *Plantae*.
Sub Kingdom: *Viridiplantae*.
Division: *Tracheophyta*.
Sub Division: *Spermatophytina*.
Class: *Magnoliopsida*.
Order: *Apiales*.
Family: *Apiaceae*.
Genus: *Daucus* L.
Species: *Daucus carota* L.

Orange carrots are the dominant type of carrots and are mostly used in various cuisines both fresh or cooked. The rate of carrot consumption

Table 2
The Roles of Carrot and its bioactive compound as immunomodulatory.

Subjects	Study Design	Preparation	Dose	Route	Duration	Immunomodulatory Activity	Modulation	Parameters/Mediator Affected	Ref
BALB/c mice	<i>In vivo</i>	Organic and conventional carrots	70 g/kg of feed	Oral	30 days	immunostimulant in intestinal and peripheral immunity	↑	Lymphocyte population, including Treg	Roselli et al., (2012)
APAP-induced oxidative stress BALB/c mice	<i>In vivo</i>	Carrot deuterium-depleted aqueous extract	500 mg/kg	Oral	7 days	Oxidative stress indicators Antioxidant enzyme	↓ ↑	ALT, AST and 8-OHdG SOD	Saba et al., (2019)
Cadmium-induced oxidative cytotoxic damage in rats	<i>In vivo</i>	Carrot juice	<i>ad libitum</i>	Oral	5 days	Improve endogenous antioxidant	↑	Catalase, SOD, GPx	Embugushiki et al., (2013)
High fructose diet-induced inflammation in weaning wistar rats	<i>In vivo</i>	Carrot juice	0.6 mL	Oral	8 weeks	Pro-inflammatory mediator	↓	MCP1 and hsCRP	Maresh et al., (2017)
LPS-induced CB57BL/6 mice	<i>In vivo</i>	Carrot's falcarinol	5 mg/kgBW	Oral	7 days	Immune homeostasis's regulator	↑	Heme oxygenase-1 8.42 times in mRNA levels and 10.7 times in protein levels	Stefansson & Bakovic, (2020)
						Cytokine release	↑	Plasma IL-4, IL-9, IL-10, IL-13, illustrating the characteristics in the improvement of type 1 inflammation	
LPS-induced CB57BL/6 mice	<i>In vivo</i>	Carrot's falcarinol	5 mg/kgBW	Oral	7 days	Pro-inflammatory cytokines	↓	IL-6, TNF-α/TNF-αr, and IFNγ gene expression	Stefansson & Bakovic, (2018)
Albino wistar rats	<i>In vivo</i>	Carotenoid extract from carrots	559 mg/kg	Oral	21 days	White blood cells	↑	lymphocytes, eosinophils, monocyte counts	Ekam et al., (2006)

↓, decrease; ↑, increase;

Abbreviations: APAP, Acetaminophen; LPS, lipopolisaccharide; Treg, T regulator; SOD, superoxide dismutase; GPx, glutathione peroxidase; MCP-1, monocyte chemoattractant protein-1; hsCRP, high-sensitivity C-reactive protein; ALT, alanine transaminase; AST, aspartate transaminase; IL, interleukin; TNF-α, tumor necrosis factor-alpha

increased due to an increase in public awareness of carrots as functional food that is useful in supporting healthy lifestyle (Ergun, 2018). The antioxidant activity of orange carrots based on the Ferric Reducing Antioxidant Power (FRAP), 2,2'-Azinobis-(3-Ethylbenzthiazolin-6-Sulfonic Acid) (ABTS) and Cupric Reducing Antioxidant Capacity (CUPRAC) were 1.22 mol TE/g, 1.22 mol TE/g and 3.07 mol TE/g fresh weight, respectively (Singh et al., 2018). The following Table 1 presents data on the nutritional and bioactive content of carrots.

4. Potential of carrots as immunomodulator

Carrots and their phytochemical compounds have been shown to have potential as a natural immunomodulator through both antioxidant and anti-inflammation properties, or by their modulation to other immune components (Table 2). For example, Roselli et al. (2012) indicated that organic carrot consumption at a rate of 70 mg/kg of feed for 30 days stimulated intestinal and peripheral immunity, with an increase in lymphocytes, including Treg. Saba et al. (2019) showed an improvement in superoxide dismutase (SOD) after administration of carrot extract at the dose of 500 mg/kg for 7 days in oxidative stress rats model. SOD controls extracellular reactive oxygen species (ROS), which are known to cause collateral tissue damage even though they are essential for the immune system's defense against pathogens (Break et al., 2012). In a cytotoxic oxidative tissue damage model, Embugushiki et al. (2013) also observed an improvement in the activity of antioxidant enzymes after 5

days of carrot juice intervention (as drinking water). As previously stated, increasing the body's antioxidant status by administering natural antioxidant counteracts the impacts of inflammation and further tissue injury through improvement of oxidative stress (Arulselvan et al., 2016).

The body is constantly in a condition of equilibrium between the formation and elimination of free radicals. The catalase, glutathione reductase, and SOD antioxidant enzymes, as well as glutathione, urate, coenzyme Q, or exogenous stimuli such bioactive substances like β-carotene, are all parts of the body's natural defensive system against free radicals. Because of the neutrophil-produced free radicals, the immune system's defense against infections serves as one of the sources of ROS generation. Micronutrient deficiency was known to cause a deterioration in immune function, both innate and adaptive, thereby rendering the body susceptible to pathogen attack as a consequence of immune imbalance. Thus, it was critical to consume bioactive compounds that serve as antioxidants in order to maintain a balanced immune response (Brambilla et al., 2008).

Carrots contain a variety of bioactive compounds, including flavonoids, phenolic acids, carotenoids, polyacetylene, and ascorbic acid. Those bioactive compound was shown to have immunomodulatory properties due to their antioxidant, anti-inflammatory, and regulatory properties of innate and adaptive immune response components. These modulatory activities included the suppression and stimulation of inflammatory cytokines, antioxidative activity through ROS scavenging




Fig. 2. Summary of Potential Immunomodulatory Mechanisms of Bioactive Compounds in Carrots. NK-cell, natural killer-cell; CD4, cluster of differentiation 4; IgA, Immunoglobulin A; IgE, Immunoglobulin E; NF- κ B, nuclear factor kappa-light-chain-enhancer of activated B cells; MAPK, mitogen-activated protein kinase; NLRP3, the nucleotide-binding domain (NOD)-like receptor protein 3; PGE2, prostaglandin E2; TLR4, toll-like receptor 4; STAT3, signal transducer and activator of transcription 3; Nrf2, nuclear factor erythroid 2-related factor 2; ROS, reactive oxygen species; SOD, superoxide dismutase; MDA, malondialdehyde; TNF- α , tumor necrosis factor- α ; IFN- γ , interferon- γ ; IL-, interleukin.

activity and improvement in endogenous antioxidant defense, as well as modulating pro-inflammatory and anti-inflammatory signaling pathways (Fig. 2).

Carotenoids are isoprenoids present in a variety of plants, including carrots. Carotenoids are acyclic molecules with five to six C rings at one or both ends (Sharma et al., 2012). Carotenoids' color variation is caused by conjugated double bonds in polyene chains that function as chromophore (Rodríguez-Concepción & Stange, 2013; Ruiz-Sola & Rodríguez-Concepción, 2012). The main carotenoids present in orange carrots are α -carotene and β -carotene (Soltoft et al., 2011).

β -carotene is a secondary metabolite produced by plants that belongs to the non-oxidized carotenoids group. β -carotene possesses a long chain of conjugated double bonds due to its role as a polyene compound of the derived acyclic structure (Fratianni et al., 2010; Meléndez-Martínez et al., 2010). Carotenoids, which have the same structure as retinol molecules, exhibit vitamin A activity, with β -carotene having the highest bioactivity as a vitamin A precursor. It is the most potent vitamin A precursor and occurs naturally as a combination of several isomers (cis and trans) of the β -carotene molecule (Toti et al., 2018). *In vivo* model indicated that long-term consumption of β -carotene reduced the production of malondialdehyde (MDA), the end product of lipid peroxidation (Hosseini et al., 2010).

MDA is an oxidative stress biomarker that contributes in maintaining balance of physiological functions, including innate and adaptive immune responses, thus affecting pathological conditions such as neurodegenerative diseases, metabolic diseases, cancers and also infectious diseases (Papac-Milicevic et al., 2016). For instance, in some autoimmune disorders, such as systemic lupus erythematosus (SLE), MDA levels are elevated due to metabolic dysregulation and increased ROS. Suppressing excess MDA in interventions is therefore advantageous (Hardt et al., 2018).

4.1. Immunomodulatory effect of β -carotene

Katsuura et al. (2009) showed that β -carotene administration in a time and dose-dependent manner was positively correlated with lipid peroxidation and glutathione synthesis, as well as negatively correlated with the transcription of cytokines such as IL-1 β , IL-6, and IL-12, in lipopolysaccharide (LPS) and INF- γ -induced RAW264 cells. This indicated that β -carotene triggers changes in intracellular redox status thereby regulating the immune function of macrophages.

Monocyte chemoattractant protein-1 (MCP-1) is a CC-chemokine that plays an important role in the inflammatory process by attracting or increasing the expression of inflammatory factors/cells. MCP-1 is involved in the pathogenesis of a variety of diseases, both directly and indirectly, including COVID-19, cancer, neuroinflammatory diseases, rheumatoid arthritis, and cardiovascular disease (Deshmane et al., 2009). In high-fructose diet-induced weaning rats, Mahesh et al. (2017) demonstrated that 0.3 mL of β -carotene from carrot juice reduced the level of free fatty acid (FFA), MCP-1 and high-sensitivity C-reactive protein (hsCRP) by 50 %, 32 % and 40 % respectively. Ekam et al., (2006) analyzed the immunomodulatory effect of carotenoid extracts (α -carotene and β -carotene) of carrots at a dose of 559 mg/kg for 21 days, increased levels of lymphocytes, eosinophils, monocytes and platelets in male Wistar rats was observed.

In their investigation on weanling mice, Nishida et al. (2014) discovered that β -carotene supplementation at a dose of 50 mg/kg feed for 14 weeks elevated the synthesis of immunoglobulin A (IgA) in the jejunum or ileum. Retinoic acid-mediated immune response was responsible for this outcome. Considering neonate and children are susceptible to diarrhea and gastrointestinal infections, that study also found an improvement in the immune response in the gastrointestinal tract (Nishida et al., 2014). Nishiyama et al. (2011) investigated the effect of β -carotene supplementation in pregnant mice during pregnancy

and lactation on IgA parameters from mammary glands, guts, and transfer of IgA from milk to neonate mice. Administration of 50 mg/kg of β -carotene supplements from 6.5 days postcoitus to 14 days postpartum showed an increased presence of IgA in the mammary glands and ileum during lactation, thereby triggering increased transfer of maternal milk IgA to neonate mice. IgA is the most abundant type of antibody in the body, and it protects mucous glands from microbial invasion while also maintaining immune homeostasis in the gastrointestinal, respiratory, and vaginal tracts. The protective immune response against intestinal viral enteropathogens was dependent on the presence of this immunoglobulin, which demonstrated the importance of IgA (Blutt et al., 2012).

The JAK-STAT pathway is critical in combating the immune system's challenges, extending from infection prevention to cancer protection. This function, however, must be maintained. Excessive and prolonged JAK-STAT signaling occurs in practically all types of autoimmune disorders (Villarino et al., 2015), as well as arthritis, Alzheimer disease, and osteosarcoma (Chiba et al., 2009; Saravanan et al., 2014; Yan et al., 2015). Signal transducer and activator of transcription 3 (STAT3), specifically, also contributes in the development of inflammatory disease such as periodontitis and cancers (De Souza et al., 2011; Geiger et al., 2016). One of the primary functions of the *jun* N-terminal kinase (JNK)/p38 MAPK is to regulate the immune response, particularly in activating stress. Therefore, pharmacological interventions capable of inhibiting these pathways have been shown to be effective in alleviating inflammation in the those diseases (Jeffrey et al., 2007; Kumar et al., 2003; Manning & Davis, 2003). Yang et al. (2021) showed that administration of β -carotene for 7 days remarkably reduced production of nitric oxide (NO), prostaglandin E2 (PGE2), TNF- α , and IL-1 β in LPS-induced intestinal inflammation. Moreover, there was suppression of mRNA expression of IL-1 β and TNF- α was observed after intervention. The postulated mechanisms for these findings included autophagy regulation as well as JAK2/STAT3 and JNK/p38 MAPK inflammatory pathways.

According to a prior review, β -carotene inhibits ROS-mediated inflammatory signaling, including the MAPK pathway and redox-sensitive transcription factors, as well as the production of inflammatory mediators including IL-8, nitric oxide synthase (NOS), and cyclooxygenase-2 (COX2) (Kang & Kim, 2017). The anti-inflammation mechanism of β -carotene was related to the reduction of the NF κ B inflammatory pathway and suppression of ROS production (Di Tomo et al., 2012). A variety of inflammatory disorders, including rheumatoid arthritis (RA), inflammatory bowel disease (IBD), multiple sclerosis, atherosclerosis, SLE, type I diabetes, chronic obstructive pulmonary disease, and "cytokine storm" in COVID-19, have been linked to NF κ B activation (Hariharan et al., 2021; Pai & Thomas, 2008). Since uncontrolled NF κ B activation is linked to these multiple inflammatory conditions, targeting NF κ B signaling is a viable approach for anti-inflammatory therapy (Liu, Zhang et al., 2017; Liu, Shen et al., 2017). Zhou et al. (2018) discovered that β -carotene blocked the NF κ B activation pathway and diminished the production of pro-inflammatory cytokines. Furthermore, this study revealed that β -carotene treatment markedly reduced oxidative stress by lowering ROS, MDA, and NO while increasing SOD. van Helden et al. (2013) showed a depletion in mRNA expression of pro-inflammatory genes associated with the NF κ B pathway involved in the production and regulation of IFN in lungs, liver and adipose tissue in β -carotene 15,15'-monooxygenase 1 knockout (Bcmo1^{-/-}) mice after 14 days of β -carotene supplement intervention.

4.2. Other bioactive compound in carrots as immunomodulator

4.2.12 Phenolic acid and flavonoid

The most abundant phenolic acid in carrots is chlorogenic acid, such as caffeic, ferulic and p-coumaric acids (Arscott & Tanumihardjo, 2010). Caffeic acid influenced antibody production and inflammatory signaling, improving oxidative state in severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) infected individuals (Adem et al.,

2021). Ferulic acid pre-treatment (50 mg/kg) for 30 days reduced LPS-induced inflammation by upregulating IL-1, IL-6, TNF- α , and IL-10 production. Acute ARDS was improved by ferulic acid's antioxidative and anti-inflammatory effects via inhibition of the MAPK inflammatory pathway, which subsequently resulted in decreased MDA and myeloperoxidase, as well as an increase in antioxidant status (Zhang et al., 2018). Other study had shown that ferulic acid intervention for 4 weeks regulated NF κ B pathway at the dose of 40 mg/kg and the nucleotide-binding domain (NOD)-like receptor protein 3 (NLRP3) inflammasome at the doses of 40 and 80 mg/kg, characterized by decreased levels of TNF- α , IL-1 β , and IL-6 mRNA expression in stressed-induced mice (Liu, Zhang et al., 2017; Liu, Shen et al., 2017).

Like NF κ B, the NLRP3 inflammasome promotes inflammation. In several inflammatory conditions like Alzheimer's, atherosclerosis, atrial fibrillation, osteoarthritis, cancer, and autoimmune illnesses like SLE and IBD, NLRP3 modulate the inflammation response in innate immune system (Wang et al., 2020). On the contrary, NLRP3's role in inflammation makes it essential for the elimination of viruses (Zhao & Zhao, 2020). Thus, modulating these pathways is vital for therapeutic treatment. P-coumaric acid had been shown to ameliorate LPS-induced oxidative damage through ROS scavenging activity and oxidative stress improvement in inflammatory state of the lungs (Kheiry et al., 2020).

The main flavonoids detected in orange carrots were quercetin, luteolin and kaempferol (Baharun et al., 2004). Quercetin inhibited NF κ B and IL-6 signaling, which exhibited an anti-inflammatory impact. The administration of quercetin remarkably reduced LPS-induced lung inflammation, as seen by lower levels of neutrophils, lymphocytes, and pro-inflammatory cytokines in the lungs (Huang et al., 2015). Quercetin also inhibited the activation of the NLRP3 inflammasome, IL-1 α , IL-6, and IL-18 secretion (Jiang et al., 2016). Luteolin at a dose of 50 mg/kg reduced TNF- α , IL-1 β and IL-6 production (Lodhi et al., 2020). In LPS-induced bronchopneumonia model, luteolin reduces inflammatory injury by regulating the expression of microRNA-132, thereby inhibiting NF κ B pathway (Liu & Meng, 2018). microRNA-132 is RNA molecule the involved in the inflammation suppression during wound healing (Li et al., 2015). Kaempferol acted as an anti-inflammatory agent by suppressing the binding of NF κ B from DNA decreasing the secretion of pro-inflammatory cytokines, inhibiting toll-like receptor 4 (TLR4) which acted as a trigger in pro-inflammatory response, and increasing expression of nuclear factor target gene mRNA and nuclear factor E2-related factor 2 (Nrf2) which was involved in oxidative damage prevention (Nam et al., 2017; Saw et al., 2014; Tang et al., 2015; R. Zhang et al., 2017). It may be inferred that the majority of these flavonoids and phenolic acids' immunological actions are connected to their anti-inflammatory properties and facilitate the protection of oxidative damage. This is important since many inflammatory and infectious diseases depend on the inflammatory process, as demonstrated by its influence on lung disorders like as ARDS, bronchopneumonia, and lung inflammation.

4.2.2. Polyacetylene

Polyacetylene is a non-volatile phytochemical group consisting of at least two C-C triple bonds. Carrots contain polyacetylene such as falcariol, falcariindiol, and falcariindiol 3-acetate (Christensen, 2012; Dawid et al., 2015). Recent review showed that polyacetylene has the potential in improving health status due to its antitumor, antifungal, anti-inflammatory, antibacterial and serotogenic properties (Dawid et al., 2015).

The transcription factor nuclear factor (erythroid-derived 2)-like 2 (Nrf2) defend against intestinal inflammation and oxidative damage. Falcariol was shown to be a dietary immunosuppressant in individuals with gastrointestinal inflammation, such as IBD, celiac disease, colitis, and enteritis (Stefanson & Bakovic 2020). Stefanson & Bakovic (2020) proved that falcariol extract supplementation of 5 mg/kg given twice daily for a week enhanced the heme oxygenase-1 (HO-1) expression,

which is the target enzyme of Nrf2. This research also observed an increase in IL-4, IL-9, IL-10, and IL-13, indicating a decrease in inflammation after falcarnol ethanolic extracts administration in LPS-induced oxidative stress mice. Stefansson & Bakovic (2018) also demonstrated that pre-treatment of falcarnol extract at a dose of 5 mg/kg twice a day for 7 days prevented pro-inflammatory genes elevation proving its protective effect against systemic and intestinal inflammation in LPS-induced C57BL/6 mice. That study discovered that LPS induction increased the expression of pro-inflammatory genes such as IL-6 up to 103.6-fold, *Tnfa* up to 48.8-fold, *Tnfa* receptor (*Tnfr*) up to 5.1-fold, *Ifn γ* up to 22.3-fold, and STAT3 3.6 times. In the presence of pre-treatment of the falcarnol ethanol extract, the expression level of pro-inflammatory genes (IL-6, *Tnfa*/*Tnfr* and *Ifn γ*) was equivalent to that of the negative control (not LPS induced), and the expression of STAT3 increased 2-fold. Gastrointestinal inflammation is an important issue because disruption of the gastrointestinal barrier is an indication of early deviations from body's homeostasis state, which lead to the development of more serious and various diseases such as autoimmune, food allergy, obesity, and chronic disorders (Stefansson & Bakovic, 2018). According to the findings of these research, falcarnol's ability to reduce inflammation may be useful in the treatment and prevention of disorders of the gastrointestinal tract or other systems.

4.2.3. Ascorbic acid

Apart from being rich in provitamin A, carrots also rich in vitamin C (Char, 2018). Vitamin C is a water-soluble vitamin founded in various fruits and vegetables (Devaki & Raveendran, 2017). Vitamin C in carrots is available in the form of ascorbic acid. This vitamin also functions as an antioxidant by scavenging free radicals, thus improving oxidative stress conditions. Vitamin C exerts antioxidative activity by transferring electrons in both enzymatic and non-enzymatic processes to generate less reactive molecules (Akbari et al., 2016).

Ascorbic acid has antiviral properties, and a recent meta-analysis suggested that administering large doses of ascorbic acid relieved common cold symptoms and duration of sickness (Ran et al., 2018). An enhanced immune response and resistance against viral infections by increasing IFN- α/β was observed in the lungs infected with the Influenza A virus in vitamin C deficient mice after the intervention of sodium L-ascorbate (3.3 g/L) for 3 weeks (Kim et al., 2013).

As an antioxidant, ascorbic acid protected lymphocytes from oxidative impairment (Lenton et al., 1999). Supplementation of 1 g of ascorbic acid for 38 days also decreased the level of IgE and histamine in a swine model induced with soybean glycinin hypersensitivity, indicating a reduce in anaphylactic reactions (Sun et al., 2009). A review summarized another role of ascorbic acid in modulating the immune system is by influencing the functioning and growth of T lymphocytes and NK cells (Van Gorkom et al., 2018). Administration of 100 μ M of vitamin C also regulated the cytokines by decreasing IFN- γ and TNF- α , as well as increasing the production of IL-10 in human lymphocytes *in vitro* model (Molina et al., 2014). At 48 and 72 h after ascorbic acid intervention, splenocytes of C57BL/6 mice induced by LPS or concavalin-A (con-A) showed an upregulation of anti-inflammatory cytokines such as IL-4 and IL-10, as well as downregulation in the expression of pro-inflammatory cytokines such as IL-6, IL-12, and TNF- α , indicating a decrease in pain caused by inflammation (Kong et al., 2015). Mechanism of ascorbic acid in reducing TNF- α and IL-6 production is also involving regulatory effects on the NF κ B pathway by suppressing the production and signaling of ROS in NF κ B transcription (Bowie & O'Neill, 2000; Cárcamo et al., 2002; Peng et al., 2005).

5. Challenges and future development

This review shows the potential of carrot has a huge variety of bioactive compounds that can be used as nutritional/therapeutic agents to modulate the immune system, either on inflammatory processes, antioxidant or redox balance, or on immune components in general.

Nowadays, because of the escalating cost of immune therapy, natural medicines produced from plants are becoming more popular. Carrots, in particular, have a significant potential in the prevention or treatment of immune-related disorders due to their widespread availability and low cost. This suggests that developing carrot-based products, such as functional foods or supplements, might be a viable therapy for various immune-mediated diseases, metabolic disorders, inflammatory diseases, and infectious diseases.

However, there are several issues that must be addressed in future study. First, it is important to emphasize the study to characterize the specific phytochemical composition of carrots using established analytical methods and determine the best extraction methods to optimize the yield. The origin, plant identification and cultivation condition such as climate of the raw material (carrot) must be verified. Such parameters are essential in such research to produce replicable outcomes in pharmacological or clinical trials, enabling appropriate comparability. Though, many studies have yet to specifically examine the biochemical components of carrots for bioactive qualities such as antioxidant, anti-inflammatory, and immunomodulatory activity. It is recommended that analysis of bioactive components be used to evaluate the immunomodulatory activity of carrots, because, in this literature review, one of the bioactive chemical components that have been specifically identified for their effect on modulating the immune system is β -carotene and falcarnol.

Second, many studies used carrot interventions in the form of juice or extract and the use of isolated bioactive compound was still limited. This is a concern since it is difficult to determine which bioactive compounds have been shown to be effective in immunomodulatory pathways when utilizing an intervention in the form of a mixture. Specific bioactive compound identification can be used to prevent these misleading results. Then, in relation to experimental studies, it is crucial to extensively define the experimental procedures and compare the intervention with standard therapy in order to fully understand the effects of carrot's bioactive compounds on immune system modulation. Additionally, toxicological investigation is also necessary to ensure the continuity of clinical studies. Thus, further research was suggested to conduct Phase I to III clinical trials to determine the mechanisms, pharmacokinetics, safety and effectiveness of the immunomodulatory effects of carrots.

The ability of immune system to prevent diseases strongly influenced by the nutritional status of host. The discussion in this review suggested that the carrots can be used as a potent immunomodulator and when ingested could enhance the both cell-mediated and humoral immunity of host, either through their antioxidant or anti-inflammation properties. Carrots have the capacity to enhance the pharmaceutical, food, medical, and agricultural industries. Carrots are becoming increasingly popular as a result of increased public health awareness due to its high nutritional content and health advantages.

It has been explained that carrots contain bioactive compounds such as carotenoids, flavonoids, phenolic acids, polyacetylenes, and ascorbic acid, which have been proposed to have immunomodulatory effects; therefore, chemists may develop bioactive compounds isolated from carrots in the future. There has been a lot of development of genetically modified plant to enhance the amount of phytonutrients and biomarkers for dietary health, for example, red-fleshed apples rich in anthocyanins (Espley et al., 2013; Sekido et al., 2010), therefore researchers in this area of expertise such as biologists may develop genetically modified carrots high in β -carotene in order to enriched the nutritional benefit of carrots.

Nutraceuticals can be used as alternative supplementation in disease management as well as to maintain a healthy lifestyle. Carrot-based nutraceutical products enriched with β -carotene and antioxidant properties have been developed in the form of carrot-mix juice, carrot jam, and carrot candies (Riaz et al., 2022). Other carrot-based product development is also encouraged in order to create a more diverse functional foods, therefore it can reach a broader consumer target. Furthermore, carrot-based supplements can also be developed in




Fig. 3. Carrot's immunomodulatory effect future research and development.

pharmaceutical industry as single compounds or as phytocomplexes. Following pre-clinical and clinical research, carrot-based supplements or nutraceuticals can be used as human interventions as a complementary treatment in immune-related diseases.

Therefore, future collaborations between researchers in various fields such as chemists, biologists, clinicians, and pharmacists as well as food industry are needed for further exploration regarding the influence of carrots on the human immunity (Fig. 3). Collaboration between researchers and industry can be a benefit in order to fasten the discoveries of this vegetable's benefits.

6. Conclusion

Carrot bioactive compounds such as β -carotene, phenolic acids, flavonoids, polyacetylenes, and ascorbic acid were found to have immunomodulatory activities via anti-inflammatory, antioxidant, and immune system pathways. On the anti-inflammatory pathway, the bioactive substances in carrots inhibition of pro-inflammatory responses and improvement of the anti-inflammatory cytokine levels, and promoted signaling pathways related to inflammation prevention. On the antioxidant pathway, carrots' bioactive substances carrots ameliorated oxidative stress by eliminating ROS and thus reduced oxidative stress parameters, increased antioxidant enzymes and oxidative stress prevention mechanism to achieve an improvement in antioxidant status. Immunomodulating effects of carrots also included the regulation of antibodies, leukocytes, immunoglobulins and histamine.

β -carotene is one of the most commonly encountered provitamin A and was abundant in carrots. β -carotene acts as an immunomodulator by regulating leukocyte formation and activity, enhancing immunoglobulin production, modulating inflammatory cytokine release, and decreasing pro-inflammatory mediators. One of the ways β -carotene controls the immune system is through redox balance alterations related to lipid

peroxidation and antioxidant enzymes.

These immunomodulatory characteristics is one of the efforts to maintain the balance of immune response and oxidant-antioxidant status in the body. Thus, it could be suggested that carrots have the potential to maintain the optimal immunity. Many studies have yet to precisely study carrot's biochemical components for bioactive properties such as antioxidant, anti-inflammatory, and immunomodulatory activity. Collaborative works are suggested to further investigate bioactive components as well as underlying processes, specifically carrot immunomodulatory action and future accessibility of carrot-based products.

122

CRediT authorship contribution statement

28 Gemala Anjani: Conceptualization, Methodology, Validation, Writing – review & editing, Supervision, Funding acquisition. Fitriyono Ayustaningwarno: Conceptualization, Methodology, Validation, Writing – review & editing, Supervision. Rafika Eviana: Methodology, Formal analysis, Investigation, Data curation, Writing – original draft.

4

Declaration of Competing Interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Gemala Anjani reports financial support was provided by The Ministry of Education, Culture, Research, and Technology of Indonesia.

39

Data availability

No data was used for the research described in the article.

Acknowledgment

The authors acknowledge the financial support from the Indonesia's The Ministry of Education, Culture, Research, and Technology "Matching Fund" Funding Assistance Program Year 2021.

Gemala Anjani reports financial support was provided by The Ministry of Education, Culture, Research, and Technology of Indonesia.

Funding statement

This work was funded from the "Matching Fund" Funding Assistance Program Year 2021, based on the Cooperation Agreement between the Directorate General of Higher Education Institutions Ministry of Education, Culture, Research, and Technology of Indonesia's Ministry of Education, Culture, Research, and Technology with Diponegoro University Number 3077/E3/PKS.08/KL/2021.

References

- Acworth, I., Plante, M., Bailey, B., Crafts, C., & Waraska, J. (2011). Simple and Direct Analysis of Falcarninol and other polyacetylenic Oxylipins in Carrots by Reverse Phase HPLC and Charged Aerosol Detection. *Planta Medica*, 77(12), 1–5. <https://doi.org/10.1055/s-0031-1282199>
- Adapa, D., Sai, Y., Anand, S., Mehaboobi, S., & Aramall, E. (2011). A Brief Review on Immune Mediated Diseases. *Journal of Clinical & Cellular Immunology*, 02(03). <https://doi.org/10.4172/2155-9899.s11-001>
- Adem, S., Eyupoglu, V., Sarfraz, I., Rasul, A., Zahoor, A. F., Ali, M., Abdalla, M., Ibrahim, I. M., & Elfiky, A. A. (2021). Caffeic acid derivatives (CAFDs) as inhibitors of SARS-CoV-2: CAFDs-based functional foods as a potential alternative approach to combat COVID-19. *Phytomedicine*, 85(August), Article 153310. <https://doi.org/10.1016/j.phymed.2020.153310>
- Agita, A., & Thaha, M. (2017). Inflammation, Immunity, and Hypertension. *Acta Med. Indones.* 2017; 49:158–165. *Acta Medica Indonesiana*, 49(2), 158–165.
- Ahmad, T., Cawood, M., Iqbal, Q., Ariño, A., Batool, A., Sabir Tariq, R. M., Azam, M., & Akhtar, S. (2019). Phytochemicals in daucus carota and their health benefits—review article. *Foods*, 8(9), 1–22. <https://doi.org/10.3390/FOODS8090424>
- Ajith, Y., Dimri, U., Dixit, S. K., Singh, S. K., Gopalakrishnan, A., Madhesh, E., Rajesh, J. B., & Sangeetha, S. G. (2017). Immunomodulatory basis of antioxidant therapy and its future prospects: An appraisal. *Inflammopharmacology*, 25(5), 487–498. <https://doi.org/10.1007/s10787-017-0393-5>
- Akbari, A., Jelodar, G., Nazifi, S., & Sajedianfard, J. (2016). An Overview of the Characteristics and Function of Vitamin C in Various Tissues: Relying on its Antioxidant Function. *Zahedan. In Press Journal of Research in Medical Sciences*. <https://doi.org/10.17795/zjrm-4037>
- Al-Snafi, A. E. (2017). Nutritional and therapeutic importance of *Daucus carota*- A review. *IOSR Journal of Pharmacy (IOSRPHR)*, 07(02), 72–88. <https://doi.org/10.9790/3013-0702017288>
- Arscott, S. A., & Tanumihardjo, S. A. (2010). Carrots of many colors provide basic nutrition and bioavailable phytochemicals acting as a functional food. *Comprehensive Reviews in Food Science and Food Safety*, 9(2), 223–239. <https://doi.org/10.1111/j.1541-4337.2009.00103.x>
- Arulselvan, P., Fard, M. T., Tan, W. S., Gothai, S., Fakurazi, S., Norhaizan, M. E., & Kumar, S. S. (2016). Role of Antioxidants and Natural Products in Inflammation. *Oxidative Medicine and Cellular Longevity*, 2016. <https://doi.org/10.1155/2016/5276130>
- Bahorun, T., Luximon-Ramma, A., Crozier, A., & Aruoma, O. I. (2004). Total phenol, flavonoid, proanthocyanidin and vitamin C levels and antioxidant activities of Mauritian vegetables. *Journal of the Science of Food and Agriculture*, 84(12), 1553–1561. <https://doi.org/10.1002/jsfa.1820>
- Bartold, P. M., & Dyke, T. E. Van. (2017). Host modulation: controlling the inflammation to control the infection. *Periodontology* 2000, 75(1), 317–329. <https://doi.org/10.1111/prd.12169>
- Berger, A. (2000). Science commentary: Th1 and Th2 responses: What are they? *British Medical Journal*, 321(7258), 424. <https://doi.org/10.1136/bmj.321.7258.424>
- Blutt, S. E., Miller, A. D., Salmon, S. L., Metzger, D. W., & Conner, M. E. (2012). IgA is Important for Clearance and Critical for Protection from Rotavirus Infection. *Mucosal Immunology*, 5(6), 712–719. <https://doi.org/10.1038/mi.2012.51>
- Bowie, A. G., & O'Neill, L. A. J. (2000). Vitamin C Inhibits NF- κ B Activation by TNF Via the Activation of p38 Mitogen-Activated Protein Kinase. *The Journal of Immunology*, 165(12), 7180–7188. <https://doi.org/10.4049/jimmunol.165.12.7180>
- Brambilla, D., Mancuso, C., Scuderi, M. R., Bosco, P., Cantarella, G., Lempereur, L., Di Benedetto, G., Pezzino, S., & Bernardini, R. (2008). The role of antioxidant supplement in immune system, neoplastic, and neurodegenerative disorders: A point of view for an assessment of the risk/benefit profile. *Nutrition Journal*, 7(29), 1–9. <https://doi.org/10.1186/1475-2891-7-29>
- Break, T. J., Jun, S., Indramohan, M., Carr, K. D., Sieve, A. N., Dory, L., & Berg, R. E. (2012). Extracellular Superoxide Dismutase Inhibits Innate Immune Responses and Clearance of an Intracellular Bacterial Infection. *The Journal of Immunology*, 188(7), 3342–3350. <https://doi.org/10.4049/jimmunol.1102341>
- Busmann, R. W., Batsatsashvili, K., & Kikvidze, Z. (2020). *Daucus carota* L. Apiaceae. In *Ethnobotany of the Mountain Regions of Central Asia and Altai* (pp. 259–266). https://doi.org/10.1007/978-3-030-28947-8_47
- Cárcamo, J. M., Pedraza, A., Bórquez-Ojeda, O., & Golde, D. W. (2002). Vitamin C suppresses TNF α -induced NF κ B activation by inhibiting I κ B phosphorylation. *Biochemistry*, 41(43), 12995–13002. <https://doi.org/10.1021/bi0263210>
- Catanzaro, M., Corsini, E., Rosini, M., Racchi, M., & Lanni, C. (2018). Immunomodulators inspired by nature: A review on curcumin and Echinacea. *Molecules*, 23(11), 1–17. <https://doi.org/10.3390/molecules23112778>
- Channappanavar, R., & Perlman, S. (2017). Pathogenic human coronavirus infections: Causes and consequences of cytokine storm and immunopathology. *Seminars in Immunopathology*, 39(5), 529–539. <https://doi.org/10.1007/s00281-017-0629-x>
- Char, C. D. (2018). Carrots. In *Fruit and Vegetable Phytochemicals* (p. 969).
- Chen, L., Deng, H., Cui, H., Fang, J., Zuo, Z., Deng, J., Li, Y., Wang, X., & Zhao, L. (2018). Oncotarget 7204 www.impactjournals.com/oncotarget Inflammatory responses and inflammation-associated diseases in organs. *Oncotarget*, 9(6), 7204–7218. www.impactjournals.com/oncotarget/
- Chen, Q. H., Wu, B. K., Pan, D., Sang, L. X., & Chang, B. (2021). Beta-carotene and its protective effect on gastric cancer. *World Journal of Clinical Cases*, 9(23), 6591–6607. <https://doi.org/10.12998/wjcc.v9.i23.6591>
- Chiba, T., Yamada, M., & Aiso, S. (2009). Targeting the JAK2 / STAT3 axis in Alzheimer's disease. *Expert Opinion on Therapeutic Targets*, 13(10), 1155–1168.
- Christensen, L. P. (2012). Aliphatic C17-Polyacetylenes of the Falcarninol Type as Potential Health Promoting Compounds in Food Plants of the Apiaceae Family. *Recent Patents on Food, Nutrition & Agriculture*, 3(1), 64–77. <https://doi.org/10.2174/2212798411103010064>
- Coussens, L. M., & Werb, Z. (2002). Inflammation and cancer. *Nature*, 420(6917), 860–867. <https://doi.org/10.1038/nature01322>
- da Silva Dias, J. C. (2014). Nutritional and Health Benefits of Carrots and Their Seed Extracts. *Food and Nutrition Sciences*, 05(22), 2147–2156. <https://doi.org/10.4236/fns.2014.52227>
- Dawid, C., Dunemann, F., Schwab, W., Nothnagel, T., & Hofmann, T. (2015). Bioactive C17-Polyacetylenes in Carrots (*Daucus carota* L.): Current Knowledge and Future Perspectives. *Journal of Agricultural and Food Chemistry*, 63(42), 9211–9222. <https://doi.org/10.1021/acs.jafc.5b04357>
- De Souza, J. A. C., Nogueira, A. V. B., De Souza, P. P. C., Cirelli, J. A., Garlet, G. P., & Rossa, C. (2011). Expression of suppressor of cytokine signaling 1 and 3 in ligature-induced periodontitis in rats. *Archives of Oral Biology*, 56(10), 1120–1128. <https://doi.org/10.1016/j.archoralbio.2011.03.022>
- Delves, P. J., & Roitt, I. M. (2000). Advances in Immunology. *The New England Journal of Medicine*, 343(1), 37–49. <https://doi.org/10.12968/bjha.2021.15.10.492>
- Deshmane, S. L., Kremlev, S., Amini, S., & Sawaya, B. E. (2009). Monocyte chemoattractant protein-1 (MCP-1): An overview. *Journal of Interferon and Cytokine Research*, 29(6), 313–325. <https://doi.org/10.1089/jir.2008.0027>
- Devaki, S. J., & Raveendran, R. L. (2017). Vitamin C: Sources, Functions, Sensing and Analysis. In *Vitamin C* (Issue August, p. 3). <https://doi.org/10.5772/intechopen.70162>
- Di Tomo, P., Canali, R., Ciavardelli, D., Di Silvestre, S., De Marco, A., Giardinelli, A., Pipino, C., Di Pietro, N., Virgili, F., & Pandolfi, A. (2012). β -Carotene and lycopene affect endothelial response to TNF- α reducing nitro-oxidative stress and interaction with monocytes. *Molecular Nutrition and Food Research*, 56(2), 217–227. <https://doi.org/10.1002/mnfr.201100500>
- Ekam, V. S., Udosen, E. O., & Chigbu, A. E. (2006). Comparative effect of carotenoid complex from Golden Neo-Life Dynamite (GNLD) and carrot extracted carotenoids on immune parameters in albino Wistar rats. *Nigerian Journal of Physiological Sciences : Official Publication of the Physiological Society of Nigeria*, 21(1–2), 1–4.
- Embugushiki, Elisha, R., Gabriel, M. S., & Zebulun, O. (2013). Protective effect of carrot juice pretreatment on cadmium-induced oxidative cytotoxic damage to some rat tissues. *IOSR Journal of Pharmacy and Biological Sciences*, 7(6), 55–62. <https://doi.org/10.9790/3008-0765562>
- Ergun, M. (2018). Evaluating Carrot As a Functional Food. *Middle East Journal of Science*, 4(2), 113–119. <https://doi.org/10.23884/mejs.2018.4.2.07>
- Espley, R. V., Bovy, A., Bava, C., Jaeger, S. R., Tomes, S., Norling, C., Crawford, J., Rowan, D., McGhie, T. K., Brendolise, C., Putterill, J., Schouten, H. J., Hellens, R. P., & Allan, A. C. (2013). Analysis of genetically modified red-fleshed apples reveals effects on growth and consumer attributes. *Plant Biotechnology Journal*, 11(4), 408–419. <https://doi.org/10.1111/pbi.12017>
- FAOSTAT. (2021). *Crops and Livestock Products*. Food and Agriculture Organization. <http://www.fao.org/faostat/en/#data/QCL>
- Farhan Aslam, M., Majeed, S., Aslam, S., & Irfan, J. A. (2017). Vitamins: Key Role Players in Boosting Up Immune Response-A Mini Review. *Vitamins & Minerals*, 06(01), 1–8. <https://doi.org/10.4172/2376-1318.1000153>
- Fratianne, A., Cinquanta, L., & Panfili, G. (2010). Degradation of carotenoids in orange juice during microwave heating. *LWT - Food Science and Technology*, 43(6), 867–871. <https://doi.org/10.1016/j.lwt.2010.01.011>
- Geiger, J. L., Grandis, J. R., & Bauman, J. E. (2016). The STAT3 pathway as a therapeutic target in head and neck cancer: Barriers and innovations. *Oral Oncology*, 56, 84–92. <https://doi.org/10.1016/j.oraloncology.2015.11.022>
- Grüter, T., Blusch, A., Motte, J., Sgodzai, M., Bachir, H., Klimas, R., Ambrosius, B., Gold, R., Ellrichmann, G., & Pitarokoli, K. (2020). Immunomodulatory and anti-oxidative effect of the direct TRPV1 receptor agonist capsaicin on Schwann cells. *Journal of Neuroinflammation*, 17(1), 1–16. <https://doi.org/10.1186/s12974-020-01821-5>
- Gul, K., Tak, A., Singh, A. K., Singh, P., Yousuf, B., & Wani, A. A. (2015). Chemistry, encapsulation, and health benefits of β -carotene - A review. *Cogent Food & Agriculture*, 1, 1–12. <https://doi.org/10.1080/23311932.2015.1018596>

- Hardt, U., Larsson, A., Gunnarsson, I., Clancy, R. M., Petri, M., Buyon, J. P., Silverman, G. J., Svenungsson, E., & Grönwall, C. (2018). Autoimmune reactivity to malondialdehyde adducts in systemic lupus erythematosus is associated with disease activity and nephritis. *Arthritis Research and Therapy*, 20(1), 1–12. <https://doi.org/10.1186/s13075-018-1530-2>
- Hariharan, A., Hakeem, A. R., Radhakrishnan, S., Reddy, M. S., & Rela, M. (2021). The Role and Therapeutic Potential of NF-kappa-B Pathway in Severe COVID-19 Patients. *Inflammopharmacology*, 29(1), 91–100. <https://doi.org/10.1007/s10787-020-00773-9>
- Hasan, Z. Y. M., AlHalbosi, M. M. F., & Al-Lihaibi, R. K. (2021). The Immunomodulation Effect Of Purified Resveratrol Extracted From Black Grape Skin Cultivated In Iraq. *Journal of Natural Volatiles and Essential Oils*, 8(4), 4490–4504.
- Hellström, J., Granato, D., & Mattila, P. H. (2020). Accumulation of phenolic acids during storage over differently handled fresh carrots. *Foods*, 9(10), 1–13. <https://doi.org/10.3390/foods9101515>
- Hosseini, F., Naseri, M. K. G., Badavi, M., Ghaffari, M. A., Shahbazian, H., & Rashidi, I. (2010). Effect of beta carotene on lipid peroxidation and antioxidant status following renal ischemia/reperfusion injury in rat. *Scandinavian Journal of Clinical and Laboratory Investigation*, 70(4), 259–263. <https://doi.org/10.3109/00365511003777810>
- Huang, R., Zhong, T., & Wu, H. (2015). Quercetin protects against lipopolysaccharide-induced acute lung injury in rats through suppression of inflammation and oxidative stress. *Archives of Medical Science*, 11(2), 427–432. <https://doi.org/10.5114/aoms.2015.50975>
- Ilyas, U., Katar, D. P., Aeri, V., & Naseef, P. P. (2016). A review on hepatoprotective and immunomodulatory herbal plants. *Pharmacognosy Reviews*, 10(19), 66–70. <https://doi.org/10.4103/0973-7847.176544>
- Integrated Taxonomic Information System - Report. *Daucus carota L. (2022). Integrated Taxonomic Information System - Report*. https://www.itis.gov/servlet/SingleRpt/SingleRpt?search_topic=TSN&search_value=29477#null
- Jantan, I., Ahmad, W., & Bukhari, S. N. A. (2015). Plant-derived immunomodulators: An insight on their preclinical evaluation and clinical trials. *Frontiers in Plant Science*, 6, 1–18. <https://doi.org/10.3389/fpls.2015.00655>
- Jeffrey, K. L., Camps, M., Rommel, C., & Mackay, C. R. (2007). Targeting dual-specificity phosphatases: Manipulating MAP kinase signalling and immune responses. *Nature Reviews Drug Discovery*, 6(5), 391–403. <https://doi.org/10.1038/nrd2289>
- Jiang, W., Huang, Y., Han, N., He, F., Li, M., Bian, Z., Liu, J., Sun, T., & Zhu, L. (2016). Quercetin suppresses NLRP3 inflammasome activation and attenuates histopathology in a rat model of spinal cord injury. *Spinal Cord*, 54(8), 592–596. <https://doi.org/10.1038/sc.2015.227>
- Kang, H., & Kim, H. (2017). Astaxanthin and β -carotene in *Helicobacter pylori*-induced Gastric Inflammation: A Mini-review on Action Mechanisms. *Journal of Cancer Prevention*, 22(2), 57–61. <https://doi.org/10.15430/jcp.2017.22.2.57>
- Katsuura, S., Imamura, T., Bando, N., & Yamanishi, R. (2009). β -Carotene and β -cryptoxanthin but not lutein evoke redox and immune changes in RAW264 murine macrophages. *Molecular Nutrition and Food Research*, 53(11), 1396–1405. <https://doi.org/10.1002/mnfr.200800566>
- Kheiry, M., Diantar, M., Badavi, M., Mard, S. A., & Bayati, V. (2020). Does p-coumaric acid improve cardiac injury following LPS-induced lung inflammation through miRNA-146a activity? *Avicenna Journal of Phytomedicine*, 10(1), 50–57. <https://doi.org/10.22038/ajp.2019.13526>
- Kim, Y., Kim, H., Bae, S., Choi, J., Lim, S. Y., Lee, N., Kong, J. M., Hwang, Y., Kang, J. S., & Lee, W. J. (2013). Vitamin C Is an Essential Factor on the Anti-viral Immune Responses through the Production of Interferon- α at the Initial Stage of Influenza A Virus (H3N2) Infection. *Immune Network*, 13(2), 70. <https://doi.org/10.4110/in.2013.13.2.70>
- Kong, E. H., Ma, S. Y., Jeong, J. Y., & Kim, K. H. (2015). Effects of L-ascorbic acid on the production of pro-inflammatory and anti-inflammatory cytokines in C57BL/6 mouse splenocytes. *Kosin Medical Journal*, 30(1), 41–49. <https://doi.org/10.7180/kmj.2015.30.1.41>
- Kumar, S., Boehm, J., & Lee, J. C. (2003). p38 MAP kinases: Key signalling molecules as therapeutic targets for inflammatory diseases. *Nature Reviews Drug Discovery*, 2(9), 717–726. <https://doi.org/10.1038/nrd1177>
- Lenton, K. J., Theriault, H., Fülöp, T., Payette, H., & Wagner, J. R. (1999). Glutathione and ascorbate are negatively correlated with oxidative DNA damage in human lymphocytes. *Carcinogenesis*, 20(4), 607–613. <https://doi.org/10.1093/carcin/20.4.607>
- Li, D., Wang, A., Liu, X., Meisen, F., Grünler, J., Botusan, I. R., Narayanan, S., Erieki, E., Li, X., Blomqvist, L., Du, L., Pivaresi, A., Sonkoly, E., Chowdhury, K., Catrina, S. B., Ståhle, M., & Landén, N. X. (2015). MicroRNA-132 enhances transition from inflammation to proliferation during wound healing. *Journal of Clinical Investigation*, 125(8), 3008–3026. <https://doi.org/10.1172/JCI79052>
- Li, Q., Wang, Y., Sun, Q., Knopf, J., Hermann, M., Lin, L., ... Shi, Y. (2022). Immune response in COVID-19: What is next? *Cell Death and Differentiation*, 29(6), 1107–1122. <https://doi.org/10.1038/s41418-022-01015-x>
- Liu, T., Zhang, L., Joo, D., & Sun, S. C. (2017). NF- κ B signaling in inflammation. *Signal Transduction and Targeted Therapy*, 2(April). <https://doi.org/10.1038/sigtrans.2017.23>
- Liu, X., & Meng, J. (2018). Luteolin alleviates LPS-induced bronchopneumonia injury in vitro and in vivo by down-regulating microRNA-132 expression. *Biomedicine and Pharmacotherapy*, 106(6), 1641–1649. <https://doi.org/10.1016/j.biopha.2018.07.094>
- Liu, Y. M., Shen, J. D., Xu, L. P., Li, H. B., Li, Y. C., & Yi, L. T. (2017). Ferulic acid inhibits neuro-inflammation in mice exposed to chronic unpredictable mild stress. *International Immunopharmacology*, 45, 128–134. <https://doi.org/10.1016/j.intimp.2017.02.007>
- Lodhi, S., Vadnere, G. P., Patil, K. D., & Patil, T. P. (2020). Protective effects of luteolin on injury induced inflammation through reduction of tissue uric acid and pro-inflammatory cytokines in rats. *Journal of Traditional and Complementary Medicine*, 10(1), 60–69. <https://doi.org/10.1016/j.jtcm.2019.02.004>
- Mahammad Shakheel, B., Saliyan, T., Satish, S., & Hedge, K. (2017). Therapeutic Uses of *Daucus carota*: A Review. *International Journal of Pharma And Chemical Research*, 3(2), 138–143. https://www.researchgate.net/publication/316430041_Therapeutic_Uses_of_Daucus_carota_A_Review
- Maresh, M., Bharathi, M., Reddy, M. R. G., Pappu, P., Vajreswari, U. K. P. A., & Jayakumar, S. M. (2017). Carrot juice ingestion attenuates high fructose-induced circulatory pro-inflammatory mediators in weanling Wistar rats. *Journal of the Science of Food and Agriculture*, 97(25), 1582–1591. <https://doi.org/10.1002/jsfa.7906>
- Manning, A. M., & Davis, R. J. (2003). Targeting JNK for therapeutic benefit: From junk to gold? *Nature Reviews Drug Discovery*, 2(7), 554–565. <https://doi.org/10.1038/nrd1132>
- Meléndez-Martínez, A. J., Escudero-Gilete, M. L., Vicario, I. M., & Heredia, F. J. (2010). Study of the influence of carotenoid structure and individual carotenoids in the qualitative and quantitative attributes of orange juice colour. *Food Research International*, 43(5), 1289–1296. <https://doi.org/10.1016/j.foodres.2010.03.012>
- Miller, A. P., Coronel, J., & Amengual, J. (2020). The role of β -carotene and vitamin A in atherogenesis: Evidence from preclinical and clinical studies. *Biochimica et Biophysica Acta - Molecular and Cell Biology of Lipids*, 1865(11), 1–32. <https://doi.org/10.1016/j.bbalip.2020.158635>
- Molina, N., Morandi, A. C., Bolin, A. P., & Otton, R. (2014). Comparative effect of fucoxanthin and vitamin C on oxidative and functional parameters of human lymphocytes. *International Immunopharmacology*, 22(1), 41–50. <https://doi.org/10.1016/j.intimp.2014.06.026>
- Nam, S. Y., Jeong, H. J., & Kim, H. M. (2017). Kaempferol impedes IL-32-induced monocyte-macrophage differentiation. *Chemico-Biological Interactions*, 274, 107–115. <https://doi.org/10.1016/j.cbi.2017.07.010>
- Nardy, A. F., Freire-de-Lima, G. G., & Morrot, A. (2015). Immune responses to infectious pathogens. *Innovative Immunology*, 1–11.
- Nishida, K., Sugimoto, M., Ikeda, S., & Kume, S. (2014). Effects of supplemental β -carotene on mucosal IgA induction in the jejunum and ileum of mice after weaning. *British Journal of Nutrition*, 111(2), 247–253. <https://doi.org/10.1017/S000714513002195>
- Nishiyama, Y., Sugimoto, M., Ikeda, S., & Kume, S. (2011). Supplemental β -carotene increases IgA-secreting cells in mammary gland and IgA transfer from milk to neonatal mice. *British Journal of Nutrition*, 105(1), 24–30. <https://doi.org/10.1017/S000714510003089>
- Ortuno-Sahagún, D., Zänker, K., Rawat, A. K. S., Kaveri, S. V., & Hegde, P. (2017). Natural Immunomodulators. *Journal of Immunology Research*, 2017, 1–2. <https://doi.org/10.1155/2017/7529408>
- Pai, S., & Thomas, R. (2008). Immune deficiency or hyperactivity-Nf-kb illuminates autoimmunity. *Journal of Autoimmunity*, 31(3), 245–251. <https://doi.org/10.1016/j.jaut.2008.04.012>
- Papac-Milicevic, N., Busch, C. J.-L., & Binder, C. J. (2016). Malondialdehyde Epitopes as Targets of Immunity and the Implications for Atherosclerosis. *Advances in Immunology*, 131, 1–59. <https://doi.org/10.1016/bs.ai.2016.02.001>
- Paragh, G., Seres, I., Harangi, M., & Fülöp, P. (2014). Dynamic interplay between metabolic syndrome and immunity. *Advances in Experimental Medicine and Biology*, 824, 171–190. https://doi.org/10.1007/978-3-319-7320-0_13
- Peng, Y., Kwok, K. H. H., Yang, P. H., Ng, S. M., Liu, J., Wong, O. G., He, M. L., Kung, H. F., & Lin, M. C. M. (2005). Ascorbic acid inhibits ROS production, NF- κ B activation and prevents ethanol-induced growth retardation and microcephaly. *Neuropharmacology*, 48(3), 426–434. <https://doi.org/10.1016/j.neuropharm.2004.10.018>
- Potter, A. S., Foroudi, S., Stamatikos, A., Patil, B. S., & Deyhim, F. (2011). Drinking carrot juice increases total antioxidant status and decreases lipid peroxidation in adults. *Nutrition Journal*, 10(1), 1–6. <https://doi.org/10.1186/1475-2891-10-96>
- Rahayu, R. P., Prasetyo, R. A., Purwanto, D. A., Kresnadi, U., Iskandar, R. P. D., & Rubianto, M. (2018). The immunomodulatory effect of green tea (*Camellia sinensis*) leaves extract on immunocompromised Wistar rats infected by *Candida albicans*. *Veterinary World*, 11(6), 765–770. <https://doi.org/10.14202/vetworld.2018.765-770>
- Ran, L., Zhao, W., Wang, J., Wang, H., Zhao, Y., Tseng, Y., & Bu, H. (2018). Extra Dose of Vitamin C Based on a Daily Supplementation Shortens the Common Cold: A Meta-Analysis of 9 Randomized Controlled Trials. *BioMed Research International*, 2018, 1–12. <https://doi.org/10.1155/2018/1837634>
- Rasheed, H. M. F., Rasheed, F., Qureshi, A. W., & Jabeen, Q. (2016). Immunostimulant activities of the aqueous methanolic extract of *Leptadenia pyrotechnica*, a plant from Cholistan desert. *Journal of Ethnopharmacology*, 186, 244–250. <https://doi.org/10.1016/j.jep.2016.03.039>
- Rawson, A., Koidis, A., Patras, A., Tuohy, M. G., & Brunton, N. P. (2010). Modelling the effect of water immersion thermal processing on polyacetylene levels and instrumental colour of carrot disks. *Food Chemistry*, 121(1), 62–68. <https://doi.org/10.1016/j.foodchem.2009.12.003>
- Razali, F. N., Sinniah, S. K., Hussin, H., Zainal Abidin, N., & Shuib, A. S. (2016). Tumor suppression effect of *Solanum nigrum* polysaccharide fraction on Breast cancer via immunomodulation. *International Journal of Biological Macromolecules*, 92, 185–193. <https://doi.org/10.1016/j.ijbiomac.2016.06.079>
- Riaz, N., Yousaf, Z., Yasmin, Z., Munawar, M., Younas, A., Rashid, M., Aftab, A., Shamsheer, B., Yasin, H., Najeebullah, M., & Simon, P. W. (2022). Development of Carrot Nutraceutical Products as an Alternative Supplement for the Prevention of Nutritional Diseases. *Frontiers in Nutrition*, 8(January), 1–19. <https://doi.org/10.3389/fnut.2021.787351>

- Rodríguez-Concepción, M., & Stange, C. (2013). Biosynthesis of carotenoids in carrot: An underground story comes to light. *Archives of Biochemistry and Biophysics*, 539(2), 110–116. <https://doi.org/10.1016/j.abb.2013.07.009>
- Roselli, M., Finamore, A., Brasili, E., Capuani, G., Kristensen, H. L., Micheloni, C., & Mengheri, E. (2012). Impact of organic and conventional carrots on intestinal and peripheral immunity. *Journal of the Science of Food and Agriculture*, 92(14), 2913–2922. <https://doi.org/10.1002/jsfa.5667>
- Rozan, M. (2017). Carotenoids, Phenolics, Antioxidant Activity and Sensory Attributes of Carrot Jam Effect of Turmeric Addition. *Egypt. J. Food Sci.*, 45, 113–123. http://www.researchgate.net/publication/335452601_Carotenoids_Phenolics_Antioxidant_Activity_and_Sensory_Attributes_of_Carrot_Jam_Effect_of_Turmeric_Addition
- Ruiz-Sola, M.Á., & Rodríguez-Concepción, M. (2012). Carotenoid Biosynthesis in Arabidopsis: A Colorful Pathway. *The Arabidopsis Book*, 10, 1–28. <https://doi.org/10.1199/tab.0158>
- Saba, E., Lee, Y. Y., Kim, M., Kim, H.-K., & Rhee, M. H. (2019). Comparison of the Antioxidant Activities of Various Processed Fruits and Vegetables in APAP-induced Oxidative Stress in BALB/c Mice. *Biomedical Science Letters*, 25(3), 211–217. <https://doi.org/10.15616/bsl.2019.25.3.211>
- Saravanan, S., Hairul Islam, V. I., Prakash Babu, N., Pandikumar, P., Thiruganasambantham, K., Chellappandian, M., Simon Durai Raj, C., Gabriel Paulraj, M., & Ignacimuthu, S. (2014). Swertiamarin attenuates inflammation mediators via modulating NF- κ B/I κ B and JAK2/STAT3 transcription factors in adjuvant induced arthritis. *European Journal of Pharmaceutical Sciences*, 56(1), 70–86. <https://doi.org/10.1016/j.ejps.2014.02.005>
- Saw, C. L. L., Guo, Y., Yang, A. Y., Paredes-Gonzalez, X., Ramirez, C., Pung, D., & Kong, A. N. T. (2014). The berry constituents quercetin, kaempferol, and pterostilbene synergistically attenuate reactive oxygen species: Involvement of the Nrf2-ARE signaling pathway. *Food and Chemical Toxicology*, 72(August), 303–311. <https://doi.org/10.1016/j.fct.2014.07.038>
- Sekido, K., Hayashi, Y., Yamada, K., Shiratake, K., Matsumoto, S., Maejima, T., & Komatsu, H. (2010). Efficient breeding system for red-fleshed apple based on linkage with s3-rnaase allele in "Pink pearl". *HortScience*, 45(4), 534–537. <https://doi.org/10.21273/hortsci.45.4.534>
- Sharma, K. D., Karki, S., Thakur, N. S., & Attri, S. (2012). Chemical composition, functional properties and processing of carrot-A review. *Journal of Food Science and Technology*, 49(1), 22–32. <https://doi.org/10.1007/s13197-011-0310-7>
- Sharma, P., Kumar, P., Sharma, R., Gupta, G., & Chaudhary, A. (2017). Immunomodulators: Role of medicinal plants in immune system. *National Journal of Physiology, Pharmacy and Pharmacology*, 7(6), 552–556. <https://doi.org/10.5455/njppp.2017.7.0203808032017>
- Sharma, V. K., Zboril, R., & Vama, R. S. (2015). Ferrates: Greener oxidants with multimodal action in water treatment technologies. *Accounts of Chemical Research*, 48(2), 182–191. <https://doi.org/10.1021/ar5004219>
- Sharpe, M., & Mount, N. (2015). Genetically modified T cells in cancer therapy: Opportunities and challenges. *DMM Disease Models and Mechanisms*, 8(4), 337–350. <https://doi.org/10.1242/dmm.018036>
- Shi, Y., Wang, Y., Shao, C., Huang, J., Gan, J., Huang, X., Bucci, E., Piacentini, M., Ippolito, G., & Melino, G. (2020). COVID-19 infection: The perspectives on immune responses. *Cell Death and Differentiation*, 27(5), 1451–1454. <https://doi.org/10.1038/s41418-020-0530-3>
- Shukla, S., Bajpai, V. K., & Kim, M. (2014). Plants as potential sources of natural immunomodulators. *Reviews in Environmental Science and Biotechnology*, 13(1), 17–33. <https://doi.org/10.1007/s11157-012-9303-x>
- Singh, B. K., Koley, T. K., Maurya, A., Singh, P. M., & Singh, B. (2018). Phytochemical and antioxidative potential of orange, red, yellow, rainbow and black coloured tropical carrots (*Daucus carota* subsp. *sativus* Schubl & Martens). *Physiology and Molecular Biology of Plants*, 24(5), 899–907. <https://doi.org/10.1007/s12298-018-0574-8>
- Søltoft, M., Bysted, A., Madsen, K. H., Mark, A. B., Bügel, S. G., Nielsen, J., & Knuthsen, P. (2011). Effects of organic and conventional growth systems on the content of carotenoids in carrot roots, and on intake and plasma status of carotenoids in humans. *Journal of the Science of Food and Agriculture*, 91(4), 767–775. <https://doi.org/10.1002/jsfa.4248>
- Stefansson, A., & Bakovic, M. (2020). Dietary polyacetylene falcarninol upregulated intestinal heme oxygenase-1 and modified plasma cytokine profile in late phase lipopolysaccharide-induced acute inflammation in CB57BL/6 mice. *Nutrition Research*, 80, 89–105. <https://doi.org/10.1016/j.nutres.2020.06.014>
- Stefansson, A. L., & Bakovic, M. (2018). Falcarninol is a potent inducer of heme oxygenase-1 and was more effective than sulforaphane in attenuating intestinal inflammation at diet-achievable doses. *Oxidative Medicine and Cellular Longevity*, 2018, 1–15. <https://doi.org/10.1155/2018/3153527>
- Sun, P., Li, D., Dong, B., Qiao, S., Ma, X., & Chen, X. (2009). Vitamin C: An immunomodulator that attenuates anaphylactic reactions to soybean glycinin hypersensitivity in a swine model. *Food Chemistry*, 113(4), 914–918. <https://doi.org/10.1016/j.foodchem.2008.08.018>
- Tang, X. L., Liu, J. X., Dong, W., Li, P., Li, L., Hou, J. C., Zheng, Y. Q., Lin, C. R., & Ren, J. G. (2015). Protective Effect of Kaempferol on LPS plus ATP-Induced Inflammatory Response in Cardiac Fibroblasts. *Inflammation*, 38(1), 94–101. <https://doi.org/10.1007/s10753-014-0011-2>
- Toti, E., Oliver Chen, C. Y., Palmery, M., Valencia, D. V., & Peluso, I. (2018). Non-provitamin A and provitamin A carotenoids as immunomodulators: Recommended dietary allowance, therapeutic index, or personalized nutrition? *Oxidative Medicine and Cellular Longevity*, 2018. <https://doi.org/10.1155/2018/4637861>
- Tyagi, R. K. (2016). Introductory Chapter: Immunity and Immunomodulation. In *Immune Response Activation and Immunomodulation* (pp. 1–3). <https://doi.org/https://doi.org/10.5772/intechopen.85299>
- USDA FoodData Central. (2015). FoodData Central. <https://fdc.nal.usda.gov/fdc-app.html#/food-details/170393/nutrients>
- Van Gorkom, G. N. Y., Klein Wolterink, R. G. J., Van Elsen, C. H. M. J., Wieten, L., Gemeraad, W. T. V., & Bos, G. M. J. (2018). Influence of Vitamin C on lymphocytes: An overview. *Antioxidants*, 7(3), 1–14. <https://doi.org/10.3390/ANTIOX7030041>
- van Helden, Y. G. J., Godschalk, R. W. L., van Schooten, F. J., & Keijer, J. (2013). Organ specificity of beta-carotene induced lung gene-expression changes in B6m1-/- mice. *Molecular Nutrition and Food Research*, 57(2), 307–319. <https://doi.org/10.1002/mnfr.201200277>
- Venkatalakshmi, P., Vadivel, V., & Brindha, P. (2016). Role of phytochemicals as immunomodulatory agents: A review. *International Journal of Green Pharmacy*, 10(1), 1–18.
- Venter, C., Eyerich, S., Sarin, T., & Klatt, K. C. (2020). Nutrition and the immune system: A complicated tango. *Nutrients*, 12(3). <https://doi.org/10.3390/nu12030818>
- Villario, A. V., Kanno, Y., & O'Shea, J. J. (2015). Mechanisms of Jak/STAT signaling in immunity and disease. *The Journal of Immunology*, 194(1), 21–27. <https://doi.org/10.4049/jimmunol.1401867>
- Wang, Z., Zhang, S., Xiao, Y., Zhang, W., Wu, S., Qin, T., Yue, Y., Qian, W., & Li, L. (2020). NLRP3 Inflammasome and Inflammatory Diseases. *Oxidative Medicine and Cellular Longevity*, 2020. <https://doi.org/10.1155/2020/4063562>
- Watzl, B., Bub, A., Briviba, K., & Rechkemmer, G. (2003). Supplementation of a Low-Carotenoid Diet with Tomato or Carrot Juice Modulates Immune Functions in Healthy Men. *Annals of Nutrition and Metabolism*, 47(6), 255–261. <https://doi.org/10.1159/000072397>
- Yan, J., Wang, Q., Zou, K., Wang, L., Schwartz, E. B., Fuchs, J. R., Zheng, Z., & Wu, J. (2015). Inhibition of the JAK2/STAT3 signaling pathway exerts a therapeutic effect on osteosarcoma. *Molecular Medicine Reports*, 12(1), 498–502. <https://doi.org/10.3892/mmr.2015.3439>
- Yang, Y., Li, R., Hui, J., Li, L., & Zheng, X. (2021). β -Carotene attenuates LPS-induced rat intestinal inflammation via modulating autophagy and regulating the JAK2/STAT3 and JNK/p38 MAPK signaling pathways. *Journal of Food Biochemistry*, 45(1), 1–12. <https://doi.org/10.1111/jfbc.13544>
- Zhang, R., Ai, X., Duan, Y., Xue, M., He, W., Wang, C., Xu, T., Xu, M., Liu, B., Li, C., Wang, Z., Zhang, R., Wang, G., Tian, S., & Liu, H. (2017). Kaempferol ameliorates H9N2 swine influenza virus-induced acute lung injury by inactivation of TLR4/MyD88-mediated NF- κ B and MAPK signaling pathways. *Biomedicine and Pharmacotherapy*, 89, 660–672. <https://doi.org/10.1016/j.biopha.2017.02.081>
- Zhang, S., Wang, P., Zhao, P., Wang, D., Zhang, Y., Wang, J., Chen, L., Guo, W., Gao, H., & Jiao, Y. (2018). Pretreatment of ferulic acid attenuates inflammation and oxidative stress in a rat model of lipopolysaccharide-induced acute respiratory distress syndrome. *International Journal of Immunopathology and Pharmacology*, 32(16). <https://doi.org/10.1177/0394632017750518>
- Zhao, C., & Zhao, W. (2020). NLRP3 Inflammasome—A Key Player in Antiviral Responses. *Frontiers in Immunology*, 11(February), 1–8. <https://doi.org/10.3389/fimmu.2020.00211>
- Zhou, L., Ouyang, L., Lin, S., Chen, S., Liu, Y. J., Zhou, W., & Wang, X. (2018). Protective role of β -carotene against oxidative stress and neuroinflammation in a rat model of spinal cord injury. *International Immunopharmacology*, 61(May), 92–99. <https://doi.org/10.1016/j.intimp.2018.05.022>

Critical review on the immunomodulatory activities of carrot's β -carotene and other bioactive compounds

ORIGINALITY REPORT

23%

SIMILARITY INDEX

16%

INTERNET SOURCES

18%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|------|
| 1 | Hafiza Mahreen Mehwish, Muhammad Shahid Riaz Rajoka, Yongai Xiong, Kai Zheng et al. "Moringa oleifera – A Functional Food and Its Potential Immunomodulatory Effects", Food Reviews International, 2020
Publication | 1 % |
| 2 | repo-dosen.ulm.ac.id
Internet Source | 1 % |
| 3 | Khalid Gul, Afshan Tak, A. K. Singh, Preeti Singh, Basharat Yousuf, Ali Abas Wani. "Chemistry, encapsulation, and health benefits of β -carotene - A review", Cogent Food & Agriculture, 2015
Publication | <1 % |
| 4 | www.pubfacts.com
Internet Source | <1 % |
| 5 | Cielo D. Char. " Carrots (L.) ", Wiley, 2017
Publication | <1 % |
| 6 | docshare.tips
Internet Source | <1 % |

7	animalsciencejournal.usamv.ro Internet Source	<1 %
8	jneuroinflammation.biomedcentral.com Internet Source	<1 %
9	www.cambridge.org Internet Source	<1 %
10	"Anticancer plants: Properties and Application", Springer Science and Business Media LLC, 2018 Publication	<1 %
11	Tanveer Ahmad, Maria Cawood, Asmat Batool, Rana Muhammad Sabir Tariq, Muhammad Awais Ghani, Muhammad Azam. " Phytochemicals in and their importance in nutrition Review article ", PeerJ, 2017 Publication	<1 %
12	Submitted to University of Sydney Student Paper	<1 %
13	Yu Yang, Ruonan Li, Junnan Hui, Lingqian Li, Xin Zheng. "β - Carotene attenuates LPS - induced rat intestinal inflammation via modulating autophagy and regulating the JAK2/STAT3 and JNK/p38 MAPK signaling pathways", Journal of Food Biochemistry, 2020 Publication	<1 %

14	cdr.lib.unc.edu Internet Source	<1 %
15	sutir.sut.ac.th:8080 Internet Source	<1 %
16	www.idexlab.com Internet Source	<1 %
17	Submitted to Çankırı Karatekin University Student Paper	<1 %
18	oro.open.ac.uk Internet Source	<1 %
19	Ivan Milovanovic, Gokhan Zengin, Svetolik Maksimovic, Vanja Tadic. " Supercritical and ultrasound - assisted extracts from mushroom: Chemical profiles, antioxidative and enzyme inhibitory properties ", Journal of the Science of Food and Agriculture, 2020 Publication	<1 %
20	Nadia Riaz, Zubaida Yousaf, Zarina Yasmin, Muneeb Munawar et al. "Development of Carrot Nutraceutical Products as an Alternative Supplement for the Prevention of Nutritional Diseases", Frontiers in Nutrition, 2022 Publication	<1 %
21	bmcmmedgenet.biomedcentral.com Internet Source	<1 %

22	pericles.pericles-prod.literatumonline.com Internet Source	<1 %
23	biblio.ugent.be Internet Source	<1 %
24	escholarship.org Internet Source	<1 %
25	journals.lww.com Internet Source	<1 %
26	novapublishers.com Internet Source	<1 %
27	spandidos-publications.com Internet Source	<1 %
28	pure.rug.nl Internet Source	<1 %
29	www.tandfonline.com Internet Source	<1 %
30	Budi Setiawan, Cecilia Oktaria Permatadewi, Baringin de Samakto, Ashar Bugis et al. "Von Willebrand Factor: Antigen and Adamts-13 Level, but not Soluble P-Selectin, are Risk Factors for the First Asymptomatic Deep Vein Thrombosis in Cancer Patients Undergoing Chemotherapy", Research Square Platform LLC, 2020 Publication	<1 %

- | | | |
|----|--|------|
| 31 | Junrui Cheng, Emilio Balbuena, Baxter Miller, Abdulkerim Eroglu. "The Role of β -Carotene in Colonic Inflammation and Intestinal Barrier Integrity", Frontiers in Nutrition, 2021
Publication | <1 % |
| 32 | www.dovepress.com
Internet Source | <1 % |
| 33 | www.isnff-jfb.com
Internet Source | <1 % |
| 34 | www.semenska.org
Internet Source | <1 % |
| 35 | "Oxidative Stress in Lung Diseases", Springer Science and Business Media LLC, 2020
Publication | <1 % |
| 36 | Farzana Sabir. "Chapter 17 Biotechnological Approaches for the Production of Immunomodulating Phytomolecules", Springer Science and Business Media LLC, 2022
Publication | <1 % |
| 37 | Ma, Xi, Pingli He, Peng Sun, and Pengfei Han. "Lipoic Acid: An Immunomodulator That Attenuates Glycinin-Induced Anaphylactic Reactions in a Rat Model", Journal of Agricultural and Food Chemistry, 2010.
Publication | <1 % |

38	Sakurako Katsuura, Tomomi Imamura, Noriko Bando, Rintaro Yamanishi. " β -Carotene and β -cryptoxanthin but not lutein evoke redox and immune changes in RAW264 murine macrophages", Molecular Nutrition & Food Research, 2009 Publication	<1 %
39	Xuan Wang, Zhenxia Liu, Xiaohui Jiang, Liangmin Yu. "Self-polishing antifouling coatings based on benzamide derivatives containing capsaicin", Marine Pollution Bulletin, 2022 Publication	<1 %
40	jultika.oulu.fi Internet Source	<1 %
41	nutritionj.biomedcentral.com Internet Source	<1 %
42	thno.org Internet Source	<1 %
43	www.sciencegate.app Internet Source	<1 %
44	Submitted to Chester College of Higher Education Student Paper	<1 %
45	Submitted to City College Brighton and Hove Student Paper	<1 %

46	Submitted to University of Edinburgh Student Paper	<1 %
47	Submitted to University of Leeds Student Paper	<1 %
48	Yang Li, Hongjie Hou, Haiyan Yang. "Lack of Marked Association Between Gastrointestinal Symptoms and COVID-19 Mortality: An Updated Meta-analysis Based on Adjusted Effect Estimates", Mayo Clinic Proceedings, 2021 Publication	<1 %
49	aran.library.nuigalway.ie Internet Source	<1 %
50	cora.ucc.ie Internet Source	<1 %
51	discovery.ucl.ac.uk Internet Source	<1 %
52	hal-univ-rochelle.archives-ouvertes.fr Internet Source	<1 %
53	oatao.univ-toulouse.fr Internet Source	<1 %
54	repositorio.tec.mx Internet Source	<1 %
55	www.medwave.cl Internet Source	<1 %

56 "Diabetes and Aging-related Complications", Springer Science and Business Media LLC, 2018
Publication <1 %

57 Amel Saidi Merzouk, Bouchra Loukidi, Réda Bettioui, Hafida Merzouk. "Effects of Vitamin C and E Against Oxidative Stress: Is Antioxidant Supplementation Efficient?", Current Nutraceuticals, 2020
Publication <1 %

58 Pastor, Patricia, Pedro Cisternas, Katterine Salazar, Carmen Silva-Alvarez, Karina Oyarce, Nery Jara, Francisca Espinoza, Agustín D. Martínez, and Francisco Nualart. "SVCT2 vitamin C transporter expression in progenitor cells of the postnatal neurogenic niche", Frontiers in Cellular Neuroscience, 2013.
Publication <1 %

59 www.intechopen.com
Internet Source <1 %

60 www.jove.com
Internet Source <1 %

61 www.siftdesk.org
Internet Source <1 %

62 Annegowda H.V, Pulak Majumder. "Valuable bioactives from vegetable wastes", Elsevier <1 %

63

Maresh, Malleswarapu, Bharathi Munugala, Mooli Raja Gopal Reddy, Pranati Pappu, Uday Kumar Putcha, Ayyalasomayajula Vajreswari, and Shanmugam M Jeyakumar. "Carrot juice ingestion attenuates high fructose-induced circulatory pro-inflammatory mediators in weanling Wistar rats : Carrot juice attenuates inflammation", Journal of the Science of Food and Agriculture, 2016.

Publication

<1 %

64

Neetu Kumra Taneja, Amritpal Singh, D.P. Shivaprasad, Pankaj Taneja, Divya Sachdev. "Nutraceuticals and Natural - Product Derivatives for Disease Prevention", Wiley, 2022

Publication

<1 %

65

Obesity, 2016.

Publication

<1 %

66

Perrine Bortolotti, Emmanuel Faure, Eric Kipnis. "Inflammasomes in Tissue Damages and Immune Disorders After Trauma", Frontiers in Immunology, 2018

Publication

<1 %

67

cmjournal.biomedcentral.com

Internet Source

<1 %

68	drum.lib.umd.edu Internet Source	<1 %
69	jbino.com Internet Source	<1 %
70	livrepository.liverpool.ac.uk Internet Source	<1 %
71	old.aok.pt Internet Source	<1 %
72	patents.justia.com Internet Source	<1 %
73	www.agricultforest.ac.me Internet Source	<1 %
74	www.ars.usda.gov Internet Source	<1 %
75	www.swift.in Internet Source	<1 %
76	"Cannabis Therapy in Veterinary Medicine", Springer Science and Business Media LLC, 2021 Publication	<1 %
77	"Foodborne Parasites", Springer Science and Business Media LLC, 2018 Publication	<1 %
78	"Healing Properties of Food", Traditional Herbal Medicines for Modern Times, 2015.	<1 %

- 79 H. Miyakawa, K. Sato, T. Shinbori, T. Okamoto, Y. Gushima, M. Fujiki, M. Suga. "Effects of inducible nitric oxide synthase and xanthine oxidase inhibitors on SEB-induced interstitial pneumonia in mice", European Respiratory Journal, 2002

<1 %

Publication

- 80 Keywan Mortezaee, Neda Khanlarkhani, Cordian Beyer, Adib Zendedel. "Inflammasome: Its role in traumatic brain and spinal cord injury", Journal of Cellular Physiology, 2018

<1 %

Publication

- 81 P. Menu. "The NLRP3 inflammasome in health and disease: the good, the bad and the ugly : Inflammasomes: the good, the bad and the ugly", Clinical & Experimental Immunology, 10/2011

<1 %

Publication

- 82 Wang, Mengdong, Miki Sugimoto, Shuntaro Ikeda, and Shinichi Kume. "Effects of coumestrol administration to maternal mice during pregnancy and lactation on immunoblogulin A-secreting cells in mammary glands : Coumestrol on IgA Cells in Mammary Gland", Animal Science Journal, 2013.

<1 %

Publication

83	aidic.it Internet Source	<1 %
84	all-imm.com Internet Source	<1 %
85	corpus.ulaval.ca Internet Source	<1 %
86	crossmark.crossref.org Internet Source	<1 %
87	edepot.wur.nl Internet Source	<1 %
88	jompd.org Internet Source	<1 %
89	pubag.nal.usda.gov Internet Source	<1 %
90	qspace.library.queensu.ca Internet Source	<1 %
91	researchers.adm.niigata-u.ac.jp Internet Source	<1 %
92	www.adb.org Internet Source	<1 %
93	"Functional Food and Human Health", Springer Science and Business Media LLC, 2018 Publication	<1 %

94

Cui-Cui Yin, Biao Ma, Derek Phillip Collinge, Barry James Pogson et al. "Ethylene Responses in Rice Roots and Coleoptiles Are Differentially Regulated by a Carotenoid Isomerase-Mediated Absciscic Acid Pathway", The Plant Cell, 2015

Publication

<1 %

95

Fan Yang, Yue Zhang, Akash Tariq, Xiaolan Jiang et al. " Food as medicine: A possible preventive measure against coronavirus disease (- 19) ", Phytotherapy Research, 2020

Publication

<1 %

96

N. Papac-Milicevic, C.J.-L. Busch, C.J. Binder. "Malondialdehyde Epitopes as Targets of Immunity and the Implications for Atherosclerosis", Elsevier BV, 2016

Publication

<1 %

97

Qian Chen, Zheng Xiao, Qing-Ying He, Rui-Rong Zhang, Shu-Xian Chen, Jia-Wei Dong, Hua Zhang, Xiao-Fan Chen. "Effect of Shenling Baizhu powder on immunity to diarrheal disease: A systematic review and meta-analysis", Frontiers in Pharmacology, 2022

Publication

<1 %

98

Somnath Basak, Jyoti Gokhale. "Immunity boosting nutraceuticals: Current trends and challenges", Journal of Food Biochemistry, 2021

<1 %

99

Wei Song, Christopher M. Derito, M. Keshu Liu, Xiangjiu He, Mei Dong, Rui Hai Liu.
"Cellular Antioxidant Activity of Common Vegetables", Journal of Agricultural and Food Chemistry, 2010

Publication

<1 %

100

Yongchul Jang, Insu Kwon, Wankeun Song, Ludmila M. Cosio-Lima, Youngil Lee.
"Endurance Exercise Mediates Neuroprotection Against MPTP-mediated Parkinson's Disease via Enhanced Neurogenesis, Antioxidant Capacity, and Autophagy", Neuroscience, 2018

Publication

<1 %

101

bsdwebstorage.blob.core.windows.net

Internet Source

<1 %

102

core.ac.uk

Internet Source

<1 %

103

curis.ku.dk

Internet Source

<1 %

104

elifesciences.org

Internet Source

<1 %

105

etheses.whiterose.ac.uk

Internet Source

<1 %

106

gup.ugal.ro

Internet Source

<1 %

107

journals.biologists.com

Internet Source

<1 %

108

ora.ox.ac.uk

Internet Source

<1 %

109

pdffox.com

Internet Source

<1 %

110

repository.ju.edu.et

Internet Source

<1 %

111

research.sabanciuniv.edu

Internet Source

<1 %

112

research.wur.nl

Internet Source

<1 %

113

rsdjournal.org

Internet Source

<1 %

114

rucore.libraries.rutgers.edu

Internet Source

<1 %

115

www.all-imm.com

Internet Source

<1 %

116

www.foodstandards.gov.au

Internet Source

<1 %

117

www.minervamedica.it

Internet Source

<1 %

118	www.miragenews.com Internet Source	<1 %
119	www.tdx.cat Internet Source	<1 %
120	"Infectious Microecology", Springer Science and Business Media LLC, 2014 Publication	<1 %
121	Amanda Stefanson, Marica Bakovic. "Dietary polyacetylene falcarinol upregulated intestinal heme oxygenase-1 and modified plasma cytokine profile in late phase lipopolysaccharide (LPS)-induced acute inflammation in CB57BL/6 mice", Nutrition Research, 2020 Publication	<1 %
122	Hatice Neval Özbek. "Radio frequency-assisted hot air drying of carrots for the production of carrot powder: Kinetics and product quality", LWT, 2021 Publication	<1 %
123	Thanuja D Herath, Gunaranjan Paturi, Christine A Butts, Catherine E Sansom, Marco P Morgenstern. "In vitro and in vivo effects of carrot on human blood platelet aggregation", International Journal of Food Science & Technology, 2020 Publication	<1 %

124	Victor Vicent Matabura. "Impact of temperature fluctuations on quality changes of frozen green beans and carrots during storage", Food Science and Technology International, 2021 Publication	<1 %
125	"Bioactive Dietary Factors and Plant Extracts in Dermatology", Springer Science and Business Media LLC, 2013 Publication	<1 %
126	"Plant Metabolites: Methods, Applications and Prospects", Springer Science and Business Media LLC, 2020 Publication	<1 %
127	Nadia Calabriso, Marika Massaro, Egeria Scoditti, Antonella Pasqualone, Barbara Laddomada, Maria Annunziata Carluccio. "Phenolic extracts from whole wheat biofortified bread dampen overwhelming inflammatory response in human endothelial cells and monocytes: major role of VCAM-1 and CXCL-10", European Journal of Nutrition, 2019 Publication	<1 %
128	Qi Xie, Changhong Wang. "Polyacetylenes in herbal medicine: A comprehensive review of its occurrence, pharmacology, toxicology, and	<1 %

-
- 129 Rein Verbeke, Michael J. Hogan, Karin Loré, Norbert Pardi. "Innate immune mechanisms of mRNA vaccines", Immunity, 2022 $<1\%$
- Publication
-

- 130 Torsten Bohn, Charles Desmarchelier, Lars O. Dragsted, Charlotte S. Nielsen et al. "Host-related factors explaining interindividual variability of carotenoid bioavailability and tissue concentrations in humans", Molecular Nutrition & Food Research, 2017 $<1\%$
- Publication
-

- 131 Ya-Shuo Feng, Zi-Xuan Tan, Man-Man Wang, Ying Xing, Fang Dong, Feng Zhang. "Inhibition of NLRP3 Inflammasome: A Prospective Target for the Treatment of Ischemic Stroke", Frontiers in Cellular Neuroscience, 2020 $<1\%$
- Publication
-

- 132 theses.gla.ac.uk $<1\%$
- Internet Source
-

Critical review on the immunomodulatory activities of carrot's β -carotene and other bioactive compounds

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11