

The abundance of *Holothuria* (*Halodeima*) *atra* (Jaeger, 1833) in Karimunjawa and Sintok Island, Karimunjawa National Park, Jepara, Indonesia related to density of seagrass

by Widianingsih Widianingsih

Submission date: 18-Jan-2023 01:53PM (UTC+0700)

Submission ID: 1994638279

File name: ional_Park,_Jepara,_Indonesia_related_to_density_of_seagrass.pdf (710.14K)

Word count: 4392

Character count: 23317

The abundance of *Holothuria (Halodeima) atra* (Jaeger, 1833) in Karimunjawa and Sintok Island, Karimunjawa National Park, Jepara, Indonesia related to density of seagrass

Meylisa S. Ayu¹, Retno Hartati^{1*}, Sunaryo¹, Raden Ario¹, Widianingsih Widianingsih¹ and Agoes Soegianto^{2*}

¹Department of Marine Science, Faculty of Fisheries and Marine Science, Diponegoro University, Jl. Prof. Dr. Soedharto, SH Kampus Tembalang, Semarang, Indonesia

²Department of Biology, Faculty of Sciences and Technology, Universitas Airlangga Surabaya, Indonesia

(Received 30 July, 2020; Accepted 15 September, 2020)

ABSTRACT

Seagrasses have been known as one of the most important marine resources. They provide a great ecological important marine ecosystem service. They also have function to provide habitat for a variety of life forms in coastal waters as well as serving as nursery, shelter and feeding area for many species, including sea cucumber. The objectives of the present work was to determine the relationship between *Holothuria (Halodeima) atra* (Jaeger, 1833) abundance and seagrass density in Alang-alang waters and Pancuran waters of Karimunjawa and Sintok Island of Karimunjawa National Park. Quadrant transects of 50x50 cm² were applied to observe the seagrass species, density and coverage as well as to determine the abundance of *H. atra* in different locations with dense, medium, low density of seagrass. The results showed that there were medium coverage conditions of seagrass in research locations, where in Pancuran waters found more percentage coverage than other locations. It also had more (8) species of seagrass, i.e. *Enhalusacoroides*, *Thalassia hemprichii*, *Halophila ovalis*, *Halodule uninervis*, *Halodule pinifolia*, *Cymodocea rotundata*, *Cymodocea serrulata*, and *Syringodium isoetifolium* while in Alang-Alang and Sintok Waters found only five species. Based on density of seagrass the denser seagrass the more abundance the *H. atra*. So that in Karimunjawa National Park, the abundance of *H. atra* found to have strong to very strong relationship with seagrass density that showed its importance as habitat as well as food source for *H. atra* populations.

Key words : Alang-alang waters, Habitat, *Holothuria atra*, Pancuran Waters, Sintok waters

Introduction

Situated some 90 km northwest off the northern coast of Jepara Regency in Central Java Province, Indonesia, the Karimunjawa Archipelago consists of numerous islands and outspread reefs and seagrass ecosystem. The Karimunjawa National Park (KNP),

a nature reserve which has an original ecosystem and a richly diverse coral reef, seagrass, and mangrove, is managed by Karimunjawa National Park Authority (KNPA). In 2001, all marine waters of KNP were designated as a marine conservation area by the Ministry of Forestry Decree No.74/Kpts-II/2001 (KNPA, 2014; Campbell *et al.*, 2013). The KNP

area is divided into nine zones which have different functions and allocations, i.e. core (444.63 ha), forest (1,451.77 ha), marine protection (2,599.77 ha), land utilization (55.99 ha), marine tourism utilization (2,733.74 ha), mariculture (1,370.73 ha), religious, cultural, and historical (0.86 ha), rehabilitation (68.33 ha), and traditional fisheries zone (102,899.25 ha) (Yuliana *et al.*, 2016).

The Karimunjawa Islands have high biodiversity natural resources from land to waters that need to be maintained and utilized sustainably and wisely (BTNKKJ, 2018). There are three important ecosystems to maintain balance in the coastal areas, namely seagrass, mangrove and coral reef ecosystems. As one of the important ecosystems, seagrass beds have primary productivity as autotrophic plants that bind carbon dioxide into energy in the food chain which will be eaten by herbivorous animals and through decomposition as litter, it also plays a role in nutrient recycling, as a food source that contains nutrients, as a shelter from predators (Kawaroe *et al.*, 2016). Many marine biotas are associated with this ecosystem, one of them is sea cucumber which uses the seagrass meadow a place of refuge, spawning and foraging. Sea cucumber prefers habitat with fine sandy substrates that are overgrown with protective plants, such as seagrass (Tanita and Yamada, 2019). The existence of sea cucumbers in nature is influenced by the availability of food. *H. atra* can be found in areas containing organic material from microphytobenthic organisms (Hartati *et al.*, 2020b). Ecologically, sea cucumbers have an important role as a deposit feeder that can process the substrate in their environment and provide food in the form of eggs, larvae and juveniles for another marine biota.

H. atra commonly known as black sea cucumber, is one of the most abundant and widely distributed sea cucumber species in most parts of the Indo-Pacific region (Conand, 2008) as well as in Indonesia, especially in Karimunjawa Islands (Purwanti *et al.*, 2010; Mustagpirin and Hartati, 2016). They inhabit a wide range of depths and a broad variety of habitat ranging from rocky reefs to mudflats (Conand, 2008; Purcell *et al.*, 2009). Before exploitation went to high-value sea cucumber, but recently shifted towards low value species (Anderson *et al.*, 2011), such as *H. atra* (Purwanti *et al.*, 2010; Hartati *et al.*, 2019). *H. atra* preferably lives in coastal seagrass beds, soft and hard substrates of coral reefs as having been previously reported by Dissanayake and Stefansson

(2012); Setyastuti (2014), Asha *et al.* (2015) and Hartati *et al.* (2017). As deposit feeder, *H. atra* consume detritus, uneaten food, and algae in the substrate (Hartati *et al.*, 2017; 2020a, b). It also ingests and grains, digests the nutrient, and then expels and pellets both in day and night time. Hartati *et al.*, (2020b) found that in Panjang Island, Jepara, *H. atra* had efficiency in feeding by taking advantage of the high TOM content and high abundance of microphytobenthic organisms which is related to their natural microhabitats with different seagrass coverage. Buckius *et al.* (2010) also emphasized that MPAs have the ability to maintain higher sea cucumber abundance; therefore the present research was aimed to establish the relationship between the abundance of *H. atra* with the density of seagrass in different waters of Karimunjawa and Sintok Island, in Karimunjawa National Park.

2 Materials and Methods

The research was conducted in Karimunjawa National Park, i.e., in Alang-alang and Pancuran Waters of Karimunjawa Island and Sintok Waters of Sintok island, Jepara Regency, Indonesia during October 2019. The sampling map is presented in Figure 1 and the position of each location is presented in Table 1.

Sea cucumber of *Holothuria (Halodeima) atra* (Jae-gier, 1833) samples were observed and determined along with seagrass observations in different density and no seagrass (none) location as control. Seagrass density was categorized based on Rahmawati *et al.* (2014), i.e. Seagrass percent cover of 0-25, 26-50, 51-75 and 76-100% as low, medium, dense, and very dense, respectively. The quadrat consisted of 50 × 50 cm² of PVC-pipe separated into 50 squares. If any seagrass was found in a square, then it was identified and counted for their shoot. Percent cover for each species was calculated from the shoot density according to Rahmawati *et al.* (2014); McCloskey and Unsworth (2015). Sea cucumber found in the same seagrass transect were counted as abundance of sea cucumber. A regression analysis was applied in order to establish the relationship between abundance of sea cucumber *H. atra* and density of seagrass in three different locations in Karimunjawa National Parks.

Water quality were measured in situ, temperature with hand thermometer, salinity with Atago N1 Hand Held Refractometer, pH with Ohaus ST10 Starter




Fig. 1. Map of sampling stations of Karimunjawa and Sintok Island, Karimunjawa National Park, Jepara, Central Java, Indonesia

pH meter, dissolved oxygen with HANNA HI 91466 DO meter. Sediment sample was taken using sediment core with the depth of 20 mm for organic matter and chlorophyll-a analysis. Total Organic Matter (TOM) analysis was done using the 'ash method' (Wang *et al.* (2011)). Biomass of microphytobenthic was measured as concentration of chlorophyll-a following procedures of DeJonge *et al.* (2012) and Johan *et al.*, 2015) using a Perkin-Elmer Lambda 3BUV/VIS spectrophotometer. The chlorophyll-a concentrations were calculated with the formula of Montani *et al.* (2012).

Results and Discussion

In the Karimunjawa National Park, the research was

conducted in Alang-Alang and Pancuran Waters of Karimunjawa Island and in Sintok Waters of Sintok Island. Alang-Alang waters were located close to residential areas, so the water quality was influenced by the activities of coastal community which can have an impact on the surrounding ecosystem and can cause a decrease in the potential for biodiversity. Pancuran waters were one of tourist destination with quite rich seagrass beds and coral reef ecosystems. Meanwhile, the Sintok Island is being designed as tourist destination; many biotas found and associated with seagrass ecosystems and coral reef ecosystems. As seagrass bed in Sintok waters located in piers, this location has potential damage to due to influence of ship traffic activities.

Table 1. Geographical position of sampling locations in Karimunjawa and Sintok Island, Karimunjawa National Park, Jepara, Central Java, Indonesia

Locations	Seagrass Density	South	East
Alang-Alang Waters Karimunjawa Island	Low (L)	5°50'17,57"	110°25'08,29"
	Medium (M)	5°50'17,72"	110°25'09,06"
	Dense (D)	5°50'18,11"	110°25'10,89"
	None (N)	5°50'14,58"	110°25'10,50"
Pancuran Waters Karimunjawa Island	Low (L)	5°52'52,04"	110°26'56,84"
	Medium (M)	5°52'54,57"	110°26'54,94"
	Dense (D)	5°52'53,33"	110°26'58,07"
	None (N)	5°52'55,76"	110°26'57,94"
Sintok Waters Sintok Island	Low (L)	5°47'07,86"	110°30'41,47"
	Medium (M)	5°47'09,67"	110°30'45,91"
	Dense (D)	5°47'08,54"	110°30'43,28"
	None (N)	5°47'06,30"	110°30'42,73"

Overall, the seagrass coverage in research locations varied, i.e. their average value in Alang-Alang, Pancuran and Sintok waters was 34.92, 51.5% and 41.59% respectively. According to Rahmawati *et al.* (2014), this condition belonged to moderate conditions. There were eight species of seagrass were found in the research location, i.e. *Enhalus acoroides*, *Thalassia hemprichii*, *Halophila ovalis*, *Halodule uninervis*, *Halodule pinifolia*, *Cymodocea tundata*, *Cymodocea serrulata*, and *Syringodium isoetifolium*, in which Pancuran waters have more species of seagrass than two other locations. Five species of seagrass were found in the dense seagrass bed at all locations, i.e. *E. acoroides*, *T. hemprichii*, *H. ovalis*, *H. uninervis*, and *C. rotundata* (Table 3). Density of *T. hemprichii* showed higher (172-781 ind.m⁻²) than other species in all locations. This seagrass species is able to grow in any condition (Riniatsih, 2016). The composition of the types of seagrass density (ind.m⁻²) and the abundance of *H. atra* (ind.m⁻²) in Alang-Alang, Pancuran and Sintok waters were showed in Table 2.

Seagrasses have been known as one of the most important marine resources. They provide a great ecological and economic important marine ecosystem services, including their capacity in sediment stabilization, water quality improvement, and their role in carbon and nutrient cycling (Nadiarti *et al.*, 2012). They also have function to provide habitat for a variety of life forms in coastal waters as well as serving as nursery, shelter and feeding area for many species, including sea cucumber (Dissanayake and Stefansson, 2012).

One of the marine organisms associated with seagrass ecosystem is black sea cucumbers, *H. atra*. It had characteristic of elongated body, medium flesh, and reddish black color (Purcell *et al.*, 2014). The whole body was covered with long, small, tight papillae on the dorsal surface, and long densely small tube feet on the ventral surface of the body. The sea cucumbers found in Alang-Alang, Pancuran and Sintok waters in this study were relatively low. The research was conducted in the morning until noon, *H. atra* was probably buried itself in the sand,

Table 2. Seagrass composition and density of seagrass (ind.m⁻²) and abundance of *H. atra*(ind.m⁻²) in Alang-Alang, Pancuran waters (Karimunjawa Island) and Sintok Waters (Sintok Island)

Seagrass species	Dense Ind.m ⁻²	Medium Ind.m ⁻²	Low Ind.m ⁻²	None Ind.m ⁻²
Alang-Alang Waters, Karimunjawa Island				
<i>Enhalus acoroides</i>	122	39	2	0
<i>Thalassia hemprichii</i>	770	210	64	0
<i>Halophila ovalis</i>	148	18	0	0
<i>Halodule uninervis</i>	193	0	0	0
<i>Cymodocea rotundata</i>	30	8	0	0
Pancuran Waters, Karimunjawa Island				
<i>Enhalus acoroides</i>	66	11	0	0
<i>Thalassia hemprichii</i>	781	147	54	0
<i>Halophila ovalis</i>	119	25	0	0
<i>Halodule uninervis</i>	37	0	0	0
<i>Cymodocea rotundata</i>	740	197	27	0
<i>Cymodocea serrulata</i>	89	0	0	0
<i>Syringodium isoetifolium</i>	27	0	0	0
<i>Halodule pinifolia</i>	95	0	0	0
Sintok Waters, Sintok Island				
<i>Enhalus acoroides</i>	5	0	0	0
<i>Thalassia hemprichii</i>	172	158	47	0
<i>Halophila ovalis</i>	48	23	17	0
<i>Halodule uninervis</i>	11	16	19	0
<i>Cymodocea rotundata</i>	1508	145	26	0
Sea cucumber <i>H. atra</i>				
Alang-Alang Waters	2.67	1.23	0.36	0.18
Pancuran Waters	4.0	2.25	0.85	0
Sintok Waters	1.43	1.23	0.31	0

so that the number of *H. atra* was relatively low. According to Dissanayakea and Stefanssona (2012), most sea cucumbers were active foraging at night and hiding during the day and according to Setyastuti (2014) sea cucumbers usually live immersing themselves in sand.

Habitats with different density seagrasses revealed to have varied *H. atra* abundance. The abundance of *H. atra* was varied among the locations and densities of seagrass bed in the range of 0.31-4.0 ind.m⁻². The higher abundance found in Pancuran water over the different density of seagrass. As same location of marine protected area, it is almost the same abundance of *H. atra* of Cabansag *et al.* (2014) studies in MPAs of Guiuan Eastern Samar and Palompon Leyte, Philippines. It is said that *H. atra* are found to show preference for specific habitat characteristics that are associated with their feeding and protection (Dissanayakea and Stefanssona, 2012). The relationship between abundance of *H. atra* and density of seagrass were presented in Figure 2. Abundance of *H. atra* and density of seagrass in different locations were connected in a linear equation. In Karimunjawa Island (Alang-alang and Pancuran Waters) abundance of *H. atra* had very strong positive relation with density of seagrass (range of 0.91-0.93) and there was strong relationship between the min Sintok Waters with correlation coefficient of 0.78 (Figure 2). So the abundance of *H. atra* in three locations showed higher in denser seagrass than lower density seagrass (Table 3). Even in open none seagrass area, *H. atra* was only found

in one location, i.e. in Alang-alang waters. These results indicate the important role of seagrass for the life of the associated biota, especially sea cucumbers as a shelter. Denser seagrass is preferred place by sea cucumbers, so they use them as shelter from predators, waves and as a place foraging. In accordance with Tanita and Yamada (2019) sea cucumbers like a soft sandy base that is covered with protective plants, such as seagrass and the like and is free from the waves. The factors that influence the high abundance of sea cucumbers in Pancuran are thought to be due to the location that lacks local activity, tourist sites that are visited by few tourists, these waters are overgrown with various types of seagrass with sand substrate types and chemical physics factors that support sea cucumber life. In accordance with Asha *et al.* (2015), sea cucumbers were adaptive to change of chemical physicals environmental factors that support sea cucumber life. In Alang-Alang waters and Sintok Island waters, the abundance of sea cucumbers was lower than in Pancuran waters, presumably due to ecological pressure or disturbances from the surrounding environment, such as community and boat traffic activities which disturb the seagrass beds and associated biota, so that the density and type of seagrass in these waters is relatively low. In accordance with Kawaroe *et al.* (2016) the seagrass ecosystem is dynamic, which is influenced by changes in environmental conditions, if there is a decline, indicating a pressure or threat to the ecosystem.

Water quality in the form of physical and chemi-

Table 3. Water quality parameter of Karimunjawa and Sintok Island, Karimunjawa National Park, Jepara, Indonesia

Parameter	Seagrass density	Alang-Alang waters	Pancuran Waters	Sintok Waters
Temperature (°C)		31.9 ± 2	30.5 ± 2	31.2 ± 2
pH		8.13 ± 2	7.76 ± 2	7.36 ± 2
Salinity (ppt)		28.67 ± 2	30.33 ± 2	33.67 ± 2
Dissolved oxygen (mg.L ⁻¹)		5.13 ± 2	5.57 ± 2	6.1 ± 2
Transparency (%)		100	100	100
Depth (cm)		85-105	88-100	80-110
Sediment		Sandy	Sandy	Sandy
Organic matter (%) in seagrass density	Low	7.57	6.16	6.86
	Medium	6.01	5.19	7.18
	Dense	6.85	5.28	7.54
	None	5.92	6.39	5.82
Chlorophyll- a (mg.g ⁻¹) in seagrass density	Low	0.23	1.15	0.51
	Medium	0.63	1.01	0.72
	Dense	0.70	0.55	0.69
	None	1.59	0	0

cal parameters was presented in Table 3. The bottom substrate of different seagrass densities in all locations were dominated by sand which is preferred by sea cucumbers. According to Tanita and Yamada (2019), *H. atra* preferred seagrass beds with a sand-based substrate. Apart from being influenced by the seagrass ecosystem, the availability of food in the ecosystem also affects the abundance of sea cucumbers. According to Hartati *et al.* (2020a), ecologically, *H. atra* were deposit feeders also acted as bioturbator or sediment processor in shallow ecosystem waters. The organic matter and chlorophyll-a in Alang-Alang, Pancuran and Sintok waters at different seagrass densities tended to be low and medium, ranging from 5.19- 7.57%. Low and moderate organic matter content is thought to have been used by sea cucumbers as a food source. According to Hartati *et al.* (2017), black sea cucumbers were also omnivorous, ate algae, detritus (non-living matter). *H. atra* were usually found in habitats with




Fig. 2. The relationship between abundance of *H. Atra* with density of seagrass in Alang-Alang Waters (A) and Pancuran Waters (B) in Karimunjawa Island and Sintok Waters in Sintok Island (C).

organic matter content of 2.1-2.9% (Asha *et al.*, 2015). The chlorophyll-a content in the three study sites with different seagrass densities showed low value which were thought to have been eaten by sea cucumbers. According to Hartati *et al.* (2017), in addition to organic matter, chlorophyll-a can represent algal biomass and microphytobenthic which are a source of food for sea cucumbers. Water quality parameters also affect the condition of sea cucumber habitat in seagrass ecosystem. The water temperature in research locations were 30.3-32°C. According to Asha *et al.* (2015), sea cucumbers can adjust to sea water temperatures between 28-32°C. Salinity in the research locations were in the range of 26.67-35.67‰ which was still suitable for the life of sea cucumbers (Sun *et al.*, 2018). High temperatures cause an increase in oxygen consumption by biota, but dissolved oxygen content was still good for *H. atra*. Transparency of water was 100% meant that the sunlight was down reached the bottom and it allowed microphytobenthic to do photosynthesis and grow that will support the sea cucumbers (Pitt and Duy, 2004). As the water quality of Alang-Alang, Pancuran waters in Karimunjawa Island and Sintok Waters in Sintok Island were not varied much and in acceptable level of *H. atra*, so it did not influence the abundance of *H. atra* therefore it confirmed the relation with density of seagrass.

Conclusion

The seagrass coverage in Alang-Alang, Pancuran (Karimunjawa Island) and Sintok waters (Sintok island) of Karimunjawa National Park showed in moderate condition (34.92-51.5%) consisted of eight species, i.e. *Enhalus sacoroides*, *Thalassia hemprichii*, *Halophila ovalis*, *Halodule uninervis*, *Halodule pinifolia*, *Cymodocea rotundata*, *Cymodocea serrulata*, and *Syringodium isoetifolium*. The abundance of *Holothuria atra* found to have strong to very strong relationship with seagrass density that showed its importance as habitat as well as food source for *Holothuria atra* populations.

Acknowledgements

Authors wish to extend our gratitude to Karimunjawa National Park Authority for the research permits, as well as to Suket Laut Team of Marine Science Department of Diponegoro University (Dewi, Valmay, Kiky, Bowo, Hanaf, Ayu, and

Firda) for their cooperation and helps during field work.

References

- Anderson, S.C., Flemming, J.M., Watson, R. and Lotze, H.K. 2011. Serial exploitation of global sea cucumber fisheries. *Fish and Fisheries*. 12 (3) : 317–339. DOI: 10.1111/j.1467-2979.2010.00397.x
- Asha, P.S., Diwakar, K., Santhanavalli, G. and Manissery, M.K. 2015. Comparative distribution and habitat preference of the sea cucumber *Holothuria atra* Jaeger at protected and unprotected sites in Thoothukudi Region of Gulf of Mannar, South-East Coast of India. *Indian J. Fish.* 1(62) : 52-57.
- Balai Taman Nasional Karimunjawa (BTNKJ). 2018, *Statistik Balai Taman Nasional Karimunjawa 2018*. Balai Taman Nasional Karimunjawa, Semarang, 50 hlm. (In Indonesian)
- Buckius, C., Albert, S., Tibbetts, I. and Udy, J. 2010. Effect of diel activity patterns and harvesting pressure on the diversity and biomass of sea cucumbers in Marovo Lagoon, Solomon Islands. *Environmental Management*. 45 : 963-973.
- Cabansag, J.B.P., de la Cruz, M.T., Gajelan-Samson, B.P. and Diaz, F.A. 2014. Abundance and density of *Holothuria atra* and *Bohadschia marmorata* in Marine Protected Areas (MPAs) of Guiuan Eastern Samar and Palompon Leyte, Philippines. *Journal of Agricultural Technology*. 10(1) : 119-131.
- Campbell, S.J., Kartawijaya, T., Yulianto, I., Prasetia, R. and Clifton, J. 2013. Co-management approaches and incentives improve management effectiveness in the Karimunjawa National Park, Indonesia. *Marine Policy*. 41 : 72-79.
- Conand, C. 2008. Population status, fisheries and trade of sea cucumbers in Africa and the Indian Ocean. In: Toral-Granda, V., Lovatelli, A. and Vasconcellos, M. (Eds.), *Sea cucumbers- A global review of fisheries and trade*. FAO fisheries and aquaculture technical paper, No. 516, FAO, Rome, p. 143–193.
- Dissanayake, D.C.T. and Stefansson, G. 2012. Habitat preference of sea cucumbers: *Holothuria atra* and *Holothuria edulis* in the coastal waters of Sri Lanka. *Journal of the Marine Biological Association of the United Kingdom*. 92(3) : 581-590.
- De Jonge, V.N., De Boer, W.F., de Jong, D. J. and Brauer, V. S. 2012. Long-term mean annual microphytobenthos chlorophyll a variation correlated with air temperature. *Mar. Ecol. Prog. Ser.* 468 : 43-56.
- Johan, F., Jafri, M.Z., Lim, H.S. and Maznah, W.W.O. 2015. Laboratory Measurement: Chlorophyll-a concentration measurement with acetone method in spectrophotometer. *Proc. 2014 IEEE IEEM*. pp. 744-748.
- Hartati, R., Widianingsih, A., Trianto, M., Zainuri and Ambariyanto, 2017. The abundance of prospective natural food for sea cucumber *Holothuria atra* at Karimunjawa Island waters, Jepara, Indonesia. *Biodiversitas*. 3 (18) : 947-953.
- Hartati, R., Zainuri, M., Ambariyanto, A., Ayodya, F.P., Widianingsih, W., Mustagpirin, M. and Soegianto, A. 2019. Initial assessment of *Holothuria atra* in Panjang Island, Jepara, Indonesia. *Eco. Env. & Cons.* 25 (Suppl. Issue) : S1-S6.
- Hartati, R., Zainuri, M., Ambariyanto, A., Widianingsih, W., Supriyo, E. and Soegianto, A. 2020a. Organic Matter, Chlorophyll and Grain Size Features of the Sediment in the Culture Sea Pens of *Holothuria atra* (Holothuroidea, Echinodermata). *Annals of Biology*. 36(2) : 311-316.
- Hartati, R., Zainuri, M., Ambariyanto, A. and Widianingsih, W. 2020b. Feeding selectivity of *Holothuria atra* in different microhabitat in Panjang Island, Jepara (Java, Indonesia). *Biodiversitas* 5(21): 2233-2239.
- McCloskey, R.M. and Unsworth, R.K.F. 2015. Decreasing seagrass density negatively influences associated fauna. *PeerJ*. 3 : e1053. <https://doi.org/10.7717/peerj.1053>
- Kawaroe, M., Nugraha, A.H., Juraij, Tasabaramo, I.A. 2016. 2016 Pages: DOI: 10.13057/biodiv/d170228 Seagrass biodiversity at three marine ecoregions of Indonesia: Sunda Shelf, Sulawesi Sea, and Banda Sea. *Biodiversitas*. 17(2) : 585-591.
- KNPA, 2014. Karimunjawa National Park Statistics 2014. Semarang: BTNKJ (In Indonesian)
- Montani, S., Tsutsumi, H. and Komorita, T. 2012. Measurement of benthic microalgal chlorophyll a and pheopigments by fluorometric analysis. *Workshop of The First Asian Marine Biology Symposium at Phuket, Thailand on Dec. 16-17, 2012*.
- Mustagpirin and Hartati, R. 2016. Biodiversity of Seacucumber in Nyamuk Waters, Karimunjawa Islands. *Proceeding National Seminar of Fisheries and Marine Science Research VI – 2016*. Universitas Diponegoro, Semarang, 12 November 2016 (In Indonesian).
- Nadiarti, R.E., Djuwita, I., Budiharsono, S., Purbayanto, A., and Asmus, H. 2012. Challenging for seagrass management in Indonesia. *Journal of Coastal Development* 15(3) : 234-242
- Pitt, R. and Duy, N.D.Q. 2004. Breeding and rearing of the sea cucumber *Holothuria scabra* in Viet Nam. In: Lovatelli A, Conand C, Purcell S, Uthicke S, Hamel JF, Mercier A. (eds) *Advances in sea cucumber aquaculture and management*. Fisheries Technical Paper No 463, FAO, Rome. pp 333-346
- Purwati, P., Hartati, R. and Widianingsih, 2010. Eighteen sea cucumber species fished in Karimunjawa Islands, Java Sea. *Mar. Res. Ind.* 35 (2) : 10-15.
- Purcell, S.W., Polidoro, B.A., Hamel, J.F., Gamboa, and

- Mercier, A. 2014. The cost of being valuable: predictors of extinction risk in marine invertebrates exploited as luxury seafood. *Biological Sciences*. 281(1781): 20133296. <https://doi.org/10.1098/rspb.2013.3296>
- Rahmawati, S., Irwan, A., Supriyadi, L.H. and Azkab, M.H. 2014. *Panduan Monitoring Padang Lamun*, COREMAP-CTI, LIPI, 21 hlm. (In Indonesian)
- Riniatsih, I. 2016. Struktur komunitas larva ikan pada ekosistem padang lamun di perairan Jepara. *Jurnal Kelautan Tropis*. 1(19) : 21-28.
- Setyastuti A. 2014. Echinodermata, *Holothurians*, in an intertidal seagrass bed off the Bama Beach, Baluran National Park, East Java, Indonesia. *J. Ilmu and Tek Kelautan Tropis*. 6 (1) : 31-39.
- Sun, J., Zhang, L., Pan, Y., Lin, C., Wang, F. and Yang, H. 2018. Effect of water temperature on diel feeding, locomotion and digestive physiology in the sea cucumber *Apostichopus japonicus*. *Journal of Experimental Biology*. 221: jeb177451. DOI: 10.1242/jeb.177451
- Tanita, I. and Yamada, H. 2019. Distribution of sea cucumbers in relation to sediment characteristics in coral reef lagoons and adjacent waters around Ishigaki Island, Southern Japan. *Japan. Marine Ecology Wiley*. 1-13. DOI: 10.1111/maec.12564.
- Wang, Q., Li, Y. and Wang, Y. 2011. Optimizing the weight loss on ignition methodology to quantify organic and carbonate carbon of sediments from diverse sources. *Environ. Monitoring and Assessment*. 174: 241-257.
- Yuliana, E., Fahrudin, A., Boer, M., Kamal, M.M. and Pardede, S.T. 2016. The effectiveness of the zoning system in the management of reef fisheries in the marine protected area of Karimunjawa National Park, Indonesia. *AACL Bioflux*. 9 (3) : 483-497.

The abundance of *Holothuria (Halodeima) atra* (Jaeger, 1833) in Karimunjawa and Sintok Island, Karimunjawa National Park, Jepara, Indonesia related to density of seagrass

ORIGINALITY REPORT

14%

SIMILARITY INDEX

%

INTERNET SOURCES

14%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1

Retno Hartati, Agus Trianto, Widianingsih. "Habitat characteristic of two selected locations for sea cucumber ranching purposes", IOP Conference Series: Earth and Environmental Science, 2017

Publication

3%
 - 2

R Hartati, T S Alya, M Zainuri, A Ambariyanto, W Widianingsih, R Ario, I Pratikto, R Sulisyati. "Modeling of habitat suitability for sea cucumber ranching in Sintok Island, Karimunjawa National Park, Indonesia", IOP Conference Series: Earth and Environmental Science, 2021

Publication

2%
 - 3

R Hartati, M Zainuri, A Ambariyanto, W Widianingsih, Mustagpirin, F Panji. "Ranching of sea cucumber in Indonesia : A study case of *Holothuria atra*", IOP Conference Series: Earth and Environmental Science, 2021

Publication

1%
-

4

Pramaditya Wicaksono, Muhammad Afif Fauzan, Ignatius Salivian Wisnu Kumara, Rifka Noviaris Yogyantoro et al. "Analysis of reflectance spectra of tropical seagrass species and their value for mapping using multispectral satellite images", International Journal of Remote Sensing, 2019

Publication

1 %

5

Salahuddin Bachmid, Laura Siahainenina, Charlotha I Tupan. "HUBUNGAN KEPADATAN TERIPANG (HOLOTHUROIDEA) DENGAN KERAPATAN LAMUN DI PERAIRAN PULAU BUNTAL-TELUK KOTANIA, KABUPATEN SERAM BAGIAN BARAT", TRITON: Jurnal Manajemen Sumberdaya Perairan, 2020

Publication

1 %

6

D P Wijayanti, E Indrayanti, H Nuryadi, S A Rintiantono, A Sabdono. " DNA barcode of of Karimunjawa Archipelago ", IOP Conference Series: Earth and Environmental Science, 2018

Publication

1 %

7

D'Souza, Elrika, Vardhan Patankar, Rohan Arthur, Núria Marbà, and Teresa Alcoverro. "Seagrass Herbivory Levels Sustain Site-Fidelity in a Remnant Dugong Population", PLoS ONE, 2015.

Publication

1 %

8

Luong, Robert J., Teruhisa Komatsu, Nguyen Dac Ve, Dam Duc Tien, Naoto Ebuchi, Delu Pan, Toshiro Saino, and Nguyen Van Thao. "Status and threats on seagrass beds using GIS in Vietnam", Remote Sensing of the Marine Environment II, 2012.

Publication

1 %

9

Fatma Bouhlel, Hazar Mliki, Mohamed Hammami. "Abnormal crowd density estimation in aerial images based on the deep and handcrafted features fusion", Expert Systems with Applications, 2021

Publication

1 %

10

Sani Mohammed Adams, Franklin Onyekachi Anianwu, Victor Sunday Aigbodion. "Ecofriendly new nanocomposites coating formulation of zinc reinforced with calcium oxide nanoparticles synthesis from oyster shell", Journal of the Indian Chemical Society, 2022

Publication

1 %

11

D.C.T. Dissanayake, G. Stefansson. "Habitat preference of sea cucumbers: and in the coastal waters of Sri Lanka ", Journal of the Marine Biological Association of the United Kingdom, 2011

Publication

<1 %

12

A H Nugraha, M Kawaree, E S Srimariana, I Jaya, D Apdillah, S R Deswati. "Carbon storage in seagrass meadow of Teluk Bakau – Bintan Island", IOP Conference Series: Earth and Environmental Science, 2019

Publication

<1 %

13

Rohani Ambo-Rappe, Abigail M. Moore. "Sulawesi Seas, Indonesia", Elsevier BV, 2019

Publication

<1 %

14

"Seagrasses of Australia", Springer Science and Business Media LLC, 2018

Publication

<1 %

15

Christine Buckius, Simon Albert, Ian Tibbetts, James Udy. "Effect of Diel Activity Patterns and Harvesting Pressure on the Diversity and Biomass of Sea Cucumbers in Marovo Lagoon, Solomon Islands", Environmental Management, 2010

Publication

<1 %

16

Jiuwen Cao, Long Chen, Dinghan Hu, Fang Dong, Tiejia Jiang, Weidong Gao, Feng Gao. "Unsupervised Eye Blink Artifact Detection From EEG With Gaussian Mixture Model", IEEE Journal of Biomedical and Health Informatics, 2021

Publication

<1 %

17

Lisa Fajar Indriana, Muhammad Firdaus, -
Supono, Fawzan Bhakti Soffa. "FISSIPAROUS
SEA CUCUMBER (*Holothuria atra*)
POPULATION IN NORTH LOMBOK,
INDONESIA", Marine Research in Indonesia,
2018

Publication

<1 %

18

Luca Scapin, Francesco Cavraro, Stefano
Malavasi, Federico Riccato, Matteo Zucchetta,
Piero Franzoi. "Linking pipefishes and
seahorses to seagrass meadows in the Venice
lagoon: Implications for conservation",
Aquatic Conservation: Marine and Freshwater
Ecosystems, 2018

Publication

<1 %

19

Matthew J. Slater, Andrew G. Jeffs. "Do
benthic sediment characteristics explain the
distribution of juveniles of the deposit-feeding
sea cucumber *Australostichopus mollis*?",
Journal of Sea Research, 2010

Publication

<1 %

20

Mercier, A., and J.-F. Hamel. "Sea cucumber
aquaculture: hatchery production, juvenile
growth and industry challenges", Advances in
aquaculture hatchery technology, 2013.

Publication

<1 %

21

R Machrizal, K Khairul, I Chastanti, N F Sari, N
Ritonga, Y Sepriani. "A study on the density

<1 %

and the cover of seagrass species along the West Coast of Natal", IOP Conference Series: Earth and Environmental Science, 2019

Publication

22

S. Uthicke, J. A. H. Benzie, E. Ballment.
"Genetic structure of fissiparous populations of *Holothuria* (*Halodeima*) atra on the Great Barrier Reef", Marine Biology, 1998

<1 %

Publication

23

M P Rizqi. "The Condition of Seagrass Beds in North Sulawesi following the implementation of Community-Based Coastal Management Program", IOP Conference Series: Earth and Environmental Science, 2021

<1 %

Publication

24

Nur Anisa Eka Ariyani, Kismartini.
"Implementation Of Conservation Policy Through The Protection Of Life Support System In The Karimunjawa National Park", E3S Web of Conferences, 2018

<1 %

Publication

25

R Hartati, A Ambariyanto, W Widianingsih, R T Mahendrajaya, M Mustagfirin, P Prihatinningsih. "Stichopodidae (Holothuroidea: Echinodermata) from Nyamuk Island, Karimunjawa National Park, Central of Java, Indonesia", IOP Conference Series: Earth and Environmental Science, 2021

<1 %

Publication

26

Rohani Ambo-Rappe. "Differences in Richness and Abundance of Species Assemblages in Tropical Seagrass Beds of Different Structural Complexity", Journal of Environmental Science and Technology, 2016

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

The abundance of *Holothuria* (*Halodeima*) *atra* (Jaeger, 1833) in Karimunjawa and Sintok Island, Karimunjawa National Park, Jepara, Indonesia related to density of seagrass

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8