

Comparative Estimation of Pleural Effusion Volume Based on Lateral Decubitus Position of Chest X-ray and CT Scan Imaging

by Bambang Satoto

Submission date: 09-Feb-2023 09:57AM (UTC+0700)

Submission ID: 2009786631

File name: CLEAN_MANUSCRIPT_03.07.2022.pdf (285.66K)

Word count: 3667

Character count: 18983

Comparative Estimation of Pleural Effusion Volume Based on Lateral Decubitus Position of Chest X-ray and CT Scan Imaging

¹Bambang Satoto, ²Wahyu S. Budi, ²Ali Khumaeni ³Yuyun Yueniwati, ³Noorhamdani Amuinsiradj

¹ Radiology Staff of Dr. Kariadi General Hospital/ Medical Faculty of Diponegoro University Semarang, ²Dep of Physics Faculty of Sciences and Mathematics, Diponegoro University Semarang. ³Medical Faculty of Brawijaya University Malang

¹bambang99satoto@gmail.com ²wahyu.sb@fisika.fsm.undip.ac.id ²khumaeni@fisika.fsm.undip.ac.id
³yuyun@ub.ac.id ³dr.noorhamdani@gmail.com

ABSTRACT

Previous study using thoracic phantom for estimating fluid volume has been obtained which represents the case of pleural effusion based on the size of the x-ray radiograph. The models are obtained in the form of three equations, the pleural effusion volume as a function of height, length times the height, and area of the radiograph image. The three models of estimation have high linearity with ratio value more than 0.988, higher than the modelling measurement using ultrasonography modality. The modelling is expected to give a contribution on developing method for helping clinicians estimate the pleural effusion volume as a basic for performing fluid aspiration and to monitor the therapy. However, because modelling is developed using phantoms, then to be applied clinically, further research is needed for its application to patients. The height function model yields correlation value of 0.966 and paired T-test value of 0.892. The height times length function model yields correlation value of 0.982 and paired T-test value of 0.611. The area function model yields correlation value of 0.997 and paired T-test value of 0.647. From the three equations, measurement of estimated pleural effusion volume using area function on chest x-ray lateral decubitus position is the most appropriate equation. Corresponding to the results of the measurement of gold standard using a CT scan. Height measurement is the measurement that is the fastest and easiest in the application. Limitation of the study is it only can be done in ⁵right lateral decubitus position of the patient, and also cannot be applied to patients with condition such as post lung surgery, massive subpulmonic/ supradiaphragmatic pleural effusion, empyema, an atypical pleural effusion such as septated, encapsulated, loculated pleural effusion and anatomical deformity, scoliosis, or abnormalities of thoracic cavity.

Keywords: estimation modelling, pleural effusion volume (PEV), chest X-ray, CT scan

INTRODUCTION

Modelling the estimated of pleural effusion volume using imaging modalities such as ultrasonography (USG) or computerized tomography (CT) scan is generally a mathematical model of a relationship between volume and dimensions of the image [1,2]. In a previous study using a unique thoracic phantom [3], a model for estimating fluid volume in phantom has been obtained which represents the case of pleural effusion based on the size of the x-ray radiograph. A phantom was made with acrylic by measuring mid clavicular line, mid axillary line, medial line of the body, and depth of thoracic cavity for the model. The phantom was filled with water with volumes ranging from 20 ml to 800 ml, at each increase of 20 ml. The model obtained in the form of three equations that connect in a row, the pleural effusion volume as a height function of radiograph image, volume as a function of length times the height of the image and volume as an area function of the image. Those three models of estimation have high linearity with ratio value more than 0.988 [4], higher than the modelling measurement using ultrasonography modality [5]. Another advantage of the model developed is the easy and fast of estimating the pleural effusion volume, as well as the image of the radiograph needed that can be obtained at health facilities with x-ray equipment for chest radiographs.

The modelling is expected to give a contribution on developing method for helping clinicians estimate the pleural effusion volume as a basic for performing fluid aspiration and to monitor the success of therapy. However, because modelling is developed using phantoms, it cannot perfectly represent the real thoracic cavity that may have many variation compared to the phantom. Then to be applied clinically, further research is needed for its application to patients.

In this study, an estimation of the pleural effusion volume (PEV) was based on the size of thoracic imaging of patients with lateral decubitus position [5,6], using three estimation models, namely ; $PEV = 126.9 \times h$, $PEV = 6.584 \times (h \times l)$ and $PEV = 6.574 \times A$, h: height of pleural effusion on image, l: length of pleural effusion on image and A: area of pleural effusion on image [4]. The estimation of pleural effusion volume for each patients was measured by three models, then compared or correlated with measurements

using CT scans. As a comparison, measurements using CT scan were chosen because it was ¹⁶ accurate and detailed in evaluating the thoracic wall, pulmonary parenchyma, and mediastinum, and was able to distinguish between fluid and tissue clearly and produced image in the form of two-dimensional images, three-dimensional images, and various organ composition images as well as gold standard measuring the pleural effusion volume [1,2].

METHODS

Chest X-ray with right lateral decubitus position and thoracic CT scan were done for a number of patients. Right lateral decubitus position is chosen to avoid interference from the heart that occurs when tilted to the left. The height, length and area of pleural effusion were measured from photostimulabel plate computed radiography (CR) of chest X-ray imaging. Then, ¹² estimation of pleural effusion volume was developed based on three models. [4] On CT scans imaging, the ¹² estimation of pleural effusion volume was done using a standard program on CT modalities.




Figure 1: Height measurement of image of pleural effusion (h) of ⁷ lateral decubitus position of chest x-ray.




Figure 2: Length measurement of image of pleural effusion (l) of lateral decubitus position of chest X-ray.




Figure 3: Area measurement of image of pleural effusion (A) of lateral decubitus position of chest X-ray.

The Siemens somatom emotion CT scans 64 slices was used in this study, using 0.6-5mm thickness slices. The protocol of CT scans was used in this study, using kV 120 and mAs 66, pitch 1.4. The picture criteria should be included chest image from the apex to the posterior costiphrenic region, thoracic wall image, as well as the adjacent muscles, the major mediastinal vessels, pleuromediastinal margin, endotracheal and endobronchial margins, the intrathoracic oesophagus, endotracheal and endobronchial margins. The procedure that has been performed are the patients lay on lateral decubitus position on the examination table (head first or feet first), create a topogram with slices from lung's apex to the costae phrenicus sinus, create slices each with 5 mm thickness, form 5 mm thickness slices then should be reconstructed into 1 mm thickness slices, from 1 mm thickness slices could be reconstructed into 3D picture so we can make axial,

coronal, and sagittal cuts, at the time of scanning with the command “inhale, hold your breath” until the examination is complete, and when printing was made with the condition of lung and mediastinal window. Contrast media was needed to differentiate pleural effusion and lung parenchyma or consolidation. Osiris software was used to analyze the height, length, and area measurement.

¹¹ This study used an Observational design with a Cross Sectional approach. The results of the measurement of pleural effusion volume estimation with the model in patients performed based on thoracic x ray images of decubitus position were correlated (compared) with the results of measurements of pleural effusion volume from CT scan. Furthermore, each estimation of PEV value from three models based on lateral decubitus position of chest X-ray were statistically tested using T- test against the results from CT scan.

RESULTS

Of the 25 patients with pleural effusion, 5 patients did not fit the inclusion and exclusion criteria, only data from 20 patients with pleural effusion were used for analysis. Most ⁶ of the study subjects were women (66.7%). The average age of the subjects was (49.9 ± 11.3) years, with ¹⁰ the youngest was 20 years old and the oldest was 69 years old. The average age of male subjects was older than female subjects.

⁸ This research has been approved by ethical committee, as shown in ethical clearance from Health Research Ethics Committee, Faculty of Medicine, Diponegoro University No. 884/EC/KEPK/FK-UNDIP/IX/2016.

Results of PEV Estimation Based on Height on Lateral Decubitus Chest X-Ray Images in Figure 4, a graph shows the correlation between the estimated pleural effusion volumes in patients as a height function image of pleural effusion with 95% confident interval (CI), it appears that there is only one point outside the region with a value close to the border of the highest deviation value. Assessment of PEV calculation on CT scans compared with formula of $PEV = 126.9 \times \text{height}$ (height based on chest X-ray with lateral decubitus position) has a very significant correlation value, namely: 0.966, sig 0,000. ⁴ There was a significant correlation between the volume of pleural effusion calculated in CT scans and the volume calculated with the $PEV = 126.9 \times h$ (height, based on chest X-ray with lateral decubitus position).




Figure 4. Graph of the correlation between the estimated results of PEV in patients as a function of pleural effusion image height with 95% CI.

Results of PEV Estimation Based on Height Times the Length on the Lateral Decubitus Chest X-Ray Images. Figure 5 shows the graph of the correlation between the estimated results of pleural effusion volume in patients as a function of height times the length of pleural effusion images with a 95% CI, it appears that all measured values are within the determined limits. Assessment of PEV calculation on CT scans compared with formula of $PEV = 6.584 \times (\text{height} \times \text{length})$ which based on chest X-ray with lateral decubitus position has a very significant correlation value, namely: 0.982, sig 0,000. There is a significant correlation between the volume calculated on CT scan with the volume calculated with the equation of 6.584 multiply (h x l) pleural effusion in patients with lateral decubitus chest X-ray examination




Figure 5. Graph of the correlation between the estimated results of PEV in patients as a function of height times the length of pleural effusion images with 95% CI.

Results of PEV Estimation Based on Area on the Lateral Decubitus Chest X-Ray Images.

Figure 6 shows the graph of the correlation between the estimated results of pleural effusion volume in patients as a function of the area of pleural effusion images with a 95% CI, it appears that all measured values are within the determined limits. Assessment of PEV calculation on CT scan compared with formula of $PEV = 6.574 \times \text{area}(A)$ (area, based on chest X-ray with lateral decubitus position) has a very significant correlation value, namely: 0.997, sig 0,000. There is a significant correlation between the volume calculated on CT scan with the volume calculated with the equation of 6.574 multiply the area of pleural effusion in patients with chest X-ray examination in the lateral decubitus position




Figure 6. Graph of the correlation between the estimated results of PEV in patients as a function of area of pleural effusion images with 95% CI.

Results of effusion volume using the 3 models and CT are compared and shown in Figure 7.




Figure 7. Graph of pleural effusion volume on CT Scans and chest X-ray of lateral decubitus position by measuring height, height times length, and area of image of the patient.

PEV CT is the pleural effusion volume that measured in the phantom; PEV h x l is the pleural effusion volume on lateral decubitus thoracic x-examination using the equation of $PEV = 6.584 \times (h \times l)$; PEV h is the pleural effusion volume on lateral decubitus thoracic x-examination using the equation of $PEV = 126.9$

x h; PEV area is the pleural effusion volume on lateral decubitus thoracic x-examination using the equation of $PEV = 6.574 \times A$.

Paired T-test results for the volume value of the equation by measuring the height, height times length and area of pleural effusion on chest X-ray lateral decubitus with volume of pleural effusion on CT scan were respectively sig 0.892, sig 0.611, and sig 0.647, shown in Table 1.

Table 1. Paired T-test results

Models	Value
$PEV = 126.9 \times h$	0.892
$PEV = 6.584 \times (h \times l)$	0.611
$PEV = 6.574 \times A$	0.647

DISCUSSION

Initially there were 25 patients included in this study, with 5 patients later excluded, because of their health condition. Subjects with lobulated effusion, massive effusion and unmeasured of effusion image were excluded from the study. Use of the three equations has limitation that all patients in this study were examined in the right lateral decubitus position, to avoid superposition with the heart contour that occur if the patient was positioned in the left lateral decubitus position. This calculation cannot be used if patient is in an examination position other than the right lateral decubitus. In addition, the three equation also can not be applied to patients with condition such as post lung surgery, massive subpulmonic/ supradiaphragmatic pleural effusion, empyema, an atypical pleural effusion such as septated, encapsulated, loculated pleural effusion and anatomical deformity, scoliosis, or abnormalities of thoracic cavity, and anatomical abnormalities of thoracic cavity. In these condition it will be difficult to determine the measurement limit so that the result of the volume calculation will not be accurate.

The average age of subjects was 49.9 ± 11.3 years, with the youngest age being 20 years and the oldest being 69 years. The average age of male subjects was older compared to female subjects. There was no significant difference of number between men and women. Those finding is in accordance with research that

states that there is no difference in incidence of pleural effusion based on sex although some causes of pleural effusion have sex predilection [6,7].

In this study, pleural effusion is relatively more common in young adults and older people. There is no difference with the previous studies because the etiology of pleural effusion comes from several diseases in adults. The study found that men experience more malignant effusion than women. Lung cancer itself was more commonly found in men than women [8]. Another study assessed chest X-ray images of patients with radiological diagnosis of pleural effusion, found that pleural effusion was obtained most of the age of 45 - 59 years, male more often than female, and pleural effusion location mostly on the right side compared to the left and bilateral lungs. [9-12]

In a previous study using a unique thoracic phantom [3], a model for estimating fluid volume in phantom has been obtained which represents the case of pleural effusion based on the size of the x-ray radiograph. The model obtained in the form of three equations that connect in a row, the pleural effusion volume as a height function of radiograph image, volume as a function of length times the height of the image and volume as an area function of the image. Those three models of estimation have high linearity with ratio value more than 0.988 [4], higher than the modelling measurement using ultrasonography modality that use balik formula [5]. The pleural effusion volume can be estimated with ultrasonography, by measuring the maximal distance between parietal and visceral pleura (Sep) in end-expiration, where the patient was positioned supine with mild trunk elevation at 15°, probe was moved upwards in posterior axillary line and transverse section perpendicular to the body axis with pleural separation visible at the lung base, evaluation in ultrasonography use balik formula : $V \text{ (ml)} = 20 \times \text{Sep (mm)}$ [14]. Also the value of linearity when use USG is 95%




Figure 8 Graph of the relationship between volume of pleural effusion and maximum separation of pleural layers (Sep).

The new three equations model or formulas developed using the unique phantom in the previous study, but now in this work already applied to real human patients so that proves that the formulas could be applied for measurement the pleural effusion volume of the human. This equation could be applied even in the hospital with limited equipment such as only used a conventional x ray, due the models is simple but has high linearity. Comparison measurement results of pleural effusion volume of the human by these models that only use conventional x-ray with measurement using CT-scan as the gold standard to calculate using an area of pleural effusion image are in accordance.

From the calculation of the estimated volume of pleural effusion in patients, the calculation using an area of pleural effusion image has the value closest to the standard value of gold that is CT scan with the highest correlation value as well as the spread is 95% CI. This is in accordance with the definition of volume or can also be called capacity is the calculation of how much space can be occupied in an object. Measurement of area images using the freehand method has already exist in the CR software. For measurement of height times length is still below area due to irregularity of thoracic cavity shape on phantoms or patients. Anatomically, a space between the parietal pleura and the visceral pleura are called pleural cavities [13]. In

the chest cavity there are several organs that will affect the shape of the pleural space or pleural cavity. The parietal pleura limits ² the thoracic wall, covering the thoracic surface of the diaphragm and the lateral surface of the mediastinum, extending to the base of the neck to limit the surface of the suprapleural membrane to the superior thoracic aperture. The visceral pleura encloses ² the entire outer surface of the lungs and extends into the interlobar fissure [13].

On validation results, the data distribution on differences of PEV from CT scans with PEV using area on chest x-ray imaging is closer to mean value compared to using height or height times length. Based on the distribution in the graph at height there is one subject outside the lower boundary line of the CI 95% acceptable range. This supports the results of the conformity analysis, the PEV from the area equation is the closest approximation to PEV from the CT scan. In the paired T test for PEV by CT scans and PEV by chest X-ray lateral decubitus position with height, height times length, area, the value was significantly above 0.05, with sig value 0.892, 0.611 and 0.647 respectively.

Another advantage of developing the models measurement are it can be easy to used and fast for estimating the pleural effusion volume at health facilities that didn't have CT-scan modalities it can be evaluate with x-ray equipment for chest radiographs or if the health facilities doesn't have the software for calculating the pleural effusion volume.

CONCLUSIONS

Estimated pleural effusion volume in patient's thoracic cavity can be measured by measuring the height, length and area of the pleural effusion image on lateral decubitus position of chest X-ray with the PEV equation = $126.9 \times \text{height}$, $\text{PEV} = 6.584 \times (\text{height} \times \text{length})$, $\text{PEV} = 6.574 \times A$. From the three equations, measurement of estimated pleural effusion volume using function of area on chest x-ray lateral decubitus position is the most appropriate equation. Measurement of the height on chest x-ray lateral decubitus position is the fastest measurement and also to measure pleural effusion volume in its application. The three equations can not be applied to patients with condition such as post lung surgery, massive subpulmonic/

supradiaphragmatic pleural effusion, empyema, an atypical pleural effusion such as septated, encapsulated, loculated pleural effusion and anatomical deformity, scoliosis, or abnormalities of thoracic cavity, and anatomical abnormalities of thoracic cavity.

REFERENCE

1. Grimberg A, David CA, Alvaro NA, Sergio A, Wagner L. 2010. *Diagnostic Accuracy of Sonography for Pleural Effusion: Systematic Review*. Sao Paulo Med J ,128(2).
2. Ramerand F, Mao Z, Ferrari F. 2010. Multiplane Ultrasound Approach to Quantify Pleural Effusions at the Bedside. *Intensive Care Med*. 36: 656 – 664.(2):90-95
3. Satoto B, Mohandas FS, Wahyu SB, Yuyun Y, Noor H. 2018. Imaginary Anatomy Line Measurements on the Thorax. *IJIRAE/RS/Vol.05/Issue01/ DCAE10090DOI:10.26562/IJIRAE.2017.DCAE10090*
4. Satoto B, Wahyu SB, Yuyun Y, Noor H. 2017. Estimation of the Volume of Pleural Effusion in the Thoracic Phantom by Examination on the Lateral Position of the Decubitus Photo Thoraks. *Transylvaniaan Review :Vol XXV,no.23* 6170-6174
5. Balik M, Plasis P, Waldauf P. 2006. Ultrasound Estimation of Volume of Plural Fluid in Ventilated Patients. *Intensive care Med*, 32; 318-21
6. Selly JM, Ayyappan AP. 2008. Pleural Effusion. In *Imaging of The Chest Volume II*. Editor: Nestor L, Isabella S. Philadelphia, Saunders Elsevier. p1336-1366.
7. Republic Indonesia Ministry of Health. 2015. Cancer diseases situation. *Buletin Jendela. Data and information of Health*. Jakarta.
8. Alberg AJ, Samet JM. Epidemiology of lung cancer. *J Semin Interv Radiol*. 2013;30(1): 173–177.
9. Melinda G. L, Elvie L, Ramli H A. 2016. Image of the chest radiograph on pleural effusion in Radiology Department, Medical Faculty of Sam Ratulangi University – Prof. Dr. R.D. Kandouw Manado Hospital, Periode November 2014 – Oktober 2015. *Jurnal e-Clinic (eCI)*: 272-274
10. Desalew M, Amanuel A, Addis A,Zewdu H, Jemal A. 2012. Pleural effusion: Presentation, Causes and Treatment Outcome in a Resource Limited Area, Ethiopia. *Health*. 4:15-19.
11. Mcgrath EE.,Anderson BP. 2011. Diagnosis of Pleural Effusion: A Systematic Approach. *Am J Crit Care*, 20: 119-128.

12. Khairani R, Syahrudin E, Partakusuma LG. 2012. Pleural effusion characteristic in Persahabatan Hospital. *J Respir Indo*. 32:155-160.
13. Ferreira A, Filho FM, Braga T, 2006. The Role of Ultrasound in the Assesment of Pleural Effusion. *Radiologia Brasileira*. 39: 2.
14. M, Balik; P, Plasil; P, Waldauf; JpAZOUT; M, Fric; M, Otahal; J, Pahl;. (2006). Ultrasound estimation of volume of pleural fluid in mechanically ventilated patients. *Intensive Care Med*, 318-321.

Comparative Estimation of Pleural Effusion Volume Based on Lateral Decubitus Position of Chest X-ray and CT Scan Imaging

ORIGINALITY REPORT

10%
SIMILARITY INDEX

5%
INTERNET SOURCES

8%
PUBLICATIONS

1%
STUDENT PAPERS

PRIMARY SOURCES

1 link.springer.com **2%**
Internet Source

2 jddtonline.info **1%**
Internet Source

3 Ulf Teichgräber, Judith Hackbarth.
"Sonographic Bedside Quantification of
Pleural Effusion Compared to Computed
Tomography Volumetry in ICU Patients",
Ultrasound International Open, 2018
Publication **1%**

4 Yoshihiro Yamada, Jun Tanno, Shintaro
Nakano, Takatoshi Kasai, Takaaki
Senbonmatsu, Shigeyuki Nishimura. "Clinical
implications of pleural effusion in patients
with acute type B aortic dissection", European
Heart Journal: Acute Cardiovascular Care,
2016
Publication **1%**

5 "Operative Dictations in General and Vascular
Surgery", Springer Science and Business **1%**

6	www.ncbi.nlm.nih.gov Internet Source	1 %
7	Seijiro Sonoda, Yuko Kumagawa, Eiichi Inada. "A case of cardiac herniation after extrapleural pneumonectomy for malignant thymoma", Journal of Anesthesia, 2010 Publication	1 %
8	Submitted to Universitas Diponegoro Student Paper	<1 %
9	jurnal.unimus.ac.id Internet Source	<1 %
10	tip.fusabil.org Internet Source	<1 %
11	Komang Juwita Endrawati, Ni Kadek Mulyantari, Sianny Herawati, Anak Agung Wiradewi Lestari et al. "The Correlation of CD4+ T-Lymphocyte Count and Chemokine Ligand 13 Levels in Human Immunodeficiency Virus Patients Receiving Anti-retrovirus Therapy in Sanglah Central General Hospital", Open Access Macedonian Journal of Medical Sciences, 2022 Publication	<1 %
12	Bolanle Olubunmi Ibitoye, Bukunmi Michael Idowu, Akinwumi Babatunde Ogunrombi,	<1 %

Babalola Ishmael Afolabi. "Ultrasonographic quantification of pleural effusion: comparison of four formulae", Ultrasonography, 2018

Publication

13

www.frontiersin.org

Internet Source

<1 %

14

Jayita Das. "Trauma in Otolaryngology", Springer Science and Business Media LLC, 2018

Publication

<1 %

15

John J. Eicken, Michael Billington, Vicki E. Noble. "Pulmonary Ultrasound Examination for Edema, Effusion, and Thromboembolism", Global Heart, 2013

Publication

<1 %

16

www.scielo.br

Internet Source

<1 %

17

Le Neindre, Aymeric, Silvia Mongodi, François Philippart, and Bélaïd Bouhemad. "Thoracic ultrasound: Potential new tool for physiotherapists in respiratory management. A narrative review", Journal of Critical Care, 2016.

Publication

<1 %

18

M Balik, P Plasil, P Waldauf, J Pazout, M Fric, M Otahal, J Pachi. "Ultrasound estimation of volume of pleural fluid in mechanically

<1 %

ventilated patients", Intensive Care Medicine,
2006

Publication

19

Murk Rehman, Pertab Rai. "QUANTIFICATION
OF PLEURAL EFFUSION ON CT IMAGES BY
AUTOMATIC AND MANUAL SEGMENTATION",
International Journal of Engineering
Technologies and Management Research,
2020

<1 %

Publication

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Comparative Estimation of Pleural Effusion Volume Based on Lateral Decubitus Position of Chest X-ray and CT Scan Imaging

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13

PAGE 14