

Reproducibility and Validity of a Food Frequency Questionnaire (FFQ) Developed for Middle-Aged and Older Adults in Semarang, Indonesia

by Ahmad Syauqy

Submission date: 08-May-2023 10:04PM (UTC+0700)

Submission ID: 2087627908

File name: C-1_Journal_-_Nutrients,_A._Syauq.pdf (621.15K)

Word count: 7052

Character count: 35527

Article

Reproducibility and Validity of a Food Frequency Questionnaire (FFQ) Developed for Middle-Aged and Older Adults in Semarang, Indonesia

Ahmad Syaury^{1,2}, Diana Nur Afifah^{1,2}, Rachma Purwanti^{1,2}, Choirun Nissa^{1,2}, Deny Yudi Fitranti^{1,2} and Jane C.-J. Chao^{3,4,5,*}

¹ Department of Nutrition Science, Faculty of Medicine, Diponegoro University, Jl. Prof. H. Soedarto, S.H., Tembalang, Semarang 50275, Indonesia; syaury@fk.undip.ac.id (A.S.); diananurafifah@live.undip.ac.id (D.N.A.); rachmapurwanti@fk.undip.ac.id (R.P.); choirun.nissa@live.undip.ac.id (C.N.); denyyudi@gmail.com (D.Y.F.)

² Center of Nutrition Research (CENURE), Diponegoro University, Jl. Prof. H. Soedarto, S.H., Tembalang, Semarang 50275, Indonesia

³ School of Nutrition and Health Sciences, College of Nutrition, Taipei Medical University, 250 Wu-Hsing Street, Taipei 11031, Taiwan

⁴ Master Program in Global Health and Development, College of Public Health, Taipei Medical University, 250 Wu-Hsing Street, Taipei 11031, Taiwan

⁵ Nutrition Research Center, Taipei Medical University Hospital, 252 Wu-Hsing Street, Taipei 11031, Taiwan

* Correspondence: chenju@tmu.edu.tw; Tel.: +886-2-2736-1661 (ext. 6548); Fax: +886-2-2737-3112



Citation: Syaury, A.; Afifah, D.N.; Purwanti, R.; Nissa, C.; Fitranti, D.Y.; Chao, J.C.-J. Reproducibility and Validity of a Food Frequency Questionnaire (FFQ) Developed for Middle-Aged and Older Adults in Semarang, Indonesia. *Nutrients* **2021**, *13*, 4163. <https://doi.org/10.3390/nu13114163>

Academic Editor: Patrick R.M. Lauwers

Received: 27 October 2021

Accepted: 19 November 2021

Published: 20 November 2021

11

Publisher's Note: MDPI stays neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Copyright: © 2021 by the authors. Licensee MDPI, Basel, Switzerland. This article is an open access article distributed under the terms and conditions of the Creative Commons Attribution (CC BY) license (<https://creativecommons.org/licenses/by/4.0/>).

Abstract: We assessed the reproducibility and validity of a food frequency questionnaire (FFQ) among middle-aged and older adults in Semarang, Indonesia. A total of 259 subjects aged 40–80 years completed two FFQs (nine-month apart) and nine 24 h dietary recalls (24HDRs, reference method). The reproducibility of the FFQ was analyzed using correlation coefficient, intra-class correlation coefficient (ICC), weighted kappa statistics and misclassification analysis. The validity was estimated by comparing the data acquired from FFQ1 and 24HDRs. The crude Pearson's correlation coefficients and ICC for total energy and nutrients between FFQ1 and FFQ2 ranged from 0.50 to 0.81 and 0.44 to 0.78, respectively. Energy adjustment decreased the correlation coefficients for most nutrients. The crude, energy-adjusted and de-attenuated correlation coefficients for FFQ1 and 24HDRs ranged from 0.41 to 0.70, 0.31 to 0.89 and 0.54 to 0.82, respectively. The agreement rates for the same or adjacent quartile classifications were 81.1–94.6% for two FFQs and 80.7–89.6% for FFQ1 and 24HDRs. The weighted kappa values were 0.21 to 0.42 for two FFQs and 0.20 to 0.34 for FFQ1 and 24HDRs. A positive mean difference was found in the Bland–Altman analyses for energy and macronutrients. The FFQ could be acceptable for nutritional epidemiology study among Indonesians.

Keywords: reproducibility; validity; dietary assessment; dietary recalls; food frequency questionnaire; methodological study; middle-age and older adults; Indonesia

1. Introduction

The prevalence of chronic diseases is growing rapidly and have become public health burden worldwide [1]. Certain chronic diseases such as diabetes and cardiovascular disease may be preventable with diet and lifestyle modification [2]. Previous evidence has suggested that diet and nutrients were correlated with the development of chronic diseases [3,4]. Hence, it is necessary to accurately evaluate dietary and nutrient intakes. Evaluation of dietary intake requires a valid assessment instrument.

Food frequency questionnaires (FFQs) have been extensively used to evaluate dietary and nutrient intakes in most epidemiological studies [5–7]. This questionnaire is easy to administer and inexpensive to conduct in a major population and gives valuable data on dietary intake over a long period of time [8]. Nevertheless, the implementation of FFQ


is susceptible²⁹ to the socio-cultural background and ethnicity of the study population [9]. Therefore, it is important to assess the reproducibility and validity of a FFQ among a specific population for studies. The most frequently used reference method to validate FFQ is 24 h dietary recall (24HDR) [10–12].

Although a number of FFQs have been developed in some countries including the Asia region [13–15], to the best of our knowledge, the reproducibility and validity of nutrition surveys conducted in Indonesia have not been reported. It is important to precisely measure the dietary assessment tools among Indonesians since Indonesia is the most populated country in Southeast Asia, typically characterized by many mixed dishes and foods with several different³² cooking methods that affect the composition of nutrients [16,17]. Therefore, the objective of the study was to evaluate the reproducibility and validity of a FFQ to be used for epidemiological studies in Indonesia.

2. Materials and Methods

2.1. Study Population

The subjects were recruited using a multi-stage cluster random sampling in Semarang (Figure 1). First, three out of sixteen subdistricts were chosen randomly. Then, two suburbs/villages within the three subdistricts were randomly elected as the final areas. Finally, we randomly recruited 300 eligible individuals to join this study. The inclusion criteria were healthy local residents aged between 40 and 80 years and lived in Semarang for at least two years, not following a specific diet such as a weight loss diet and not pregnant. Among the 300 chosen subjects, 265 individuals were approved to join our study and conducted the study (response rate = 88.3%). Some subjects did not participate in our study because of refusal, poor health, or not attending during the study period. Semarang, the capital city of Central Java in Indonesia, is divided into lowland and highland areas and slum areas in the urban region. Semarang represents Indonesian characteristics, including demographics and lifestyle and provides a good overview of Indonesian people [18].



³⁷ **Figure 1.** Flow diagram of subject recruitment.

2.2. Assessment of Dietary Intake

All subjects completed their usual dietary intake twice using the same FFQ. Two FFQs (FFQ1 and FFQ2) surveys were assessed nine months apart. The FFQ contained 137 food items and 24 food categories based on the Indonesian Food Composition Data and the eating habits of Indonesian people (Table 1) [7,19]. The subjects gave information about the frequency of consumption (never, daily, weekly, monthly, 44 yearly) and the portion size of all food items they had eaten. The reported consumption of each food item was converted to grams per day for further evaluation.

33
Table 1. Food groups and food items used in the food frequency questionnaire.

Food Groups	Food Items
Rice/flour products	Rice, noodles, vermicelli, plain bread
Root crops	Boiled/steamed potato, taro, cassava
Whole grains	Whole grains, whole wheat, mixed grains, brown rice, oatmeal
Staples cook with oil	Fried rice, fried noodle, <i>ketiau</i> , <i>uduk</i> rice, <i>kebuli</i> rice, yellow rice
Legumes and nuts	Peas, nuts, beans, peanuts coated with flour
Soybeans	Steamed <i>tempe</i> , <i>tahu</i> , tofu
Milk and dairy products	Milk, cheese, yoghurt
Light-colored vegetables	Cabbage, Chinese cabbage, cucumber, pechay, chayote, squash, radish, bean sprouts, pumpkin, mushroom
Dark-colored vegetables	Spinach, carrots, kale, <i>buncis</i> , caisin, lotus root leaves, papaya leaves, long beans, mustard greens, glossy nightshade
Eggs	Chicken eggs, duck eggs, quail eggs
Fish and seafood	Fish, squid, shrimp, octopus, crab
Meat	Beef, veal, lamb, goat, pork
Poultry	Chicken, duck, goose, pigeon
Fast food	Instant noodles, chicken nugget, pizza, hamburger, doughnut, <i>martabak</i> , <i>bakso</i>
Processed food	Sausage, canned food, instant food
Fried food	Fried chicken, dried fish-tofu (<i>batagor</i>), <i>bakwan</i> , <i>risol</i> , <i>cakwe</i> , <i>pastel</i> , <i>cireng</i> , <i>gorengan</i> , fried fish (<i>pempek</i>), fried banana, chips, cassava chips
Organs of animals	Liver, kidney, heart, intestines
Fruits	Banana, orange, pear, mango, papaya, avocado, watermelon, apple, grape, starfruit, dragon fruit, <i>duku</i> , <i>rambutan</i> , rose apple, lemon, coconut, mangosteen, giant granadilla, jack fruit, snack fruit, soursop, breadfruit
Processed fruit	Canned fruit, <i>rujak</i> , <i>asinan</i>
Traditional snacks	<i>Gethuk</i> , <i>serabi</i> cake, <i>putu</i> cake, <i>gemblong</i> , <i>pukis</i>
Jam/honey	Jam, honey
Sugary drinks	Soft drinks, soda, energy drinks, flavored fruit drinks
Sweet dessert	Butter bread, sweet bread, cake, cookies, biscuit, crackers
Tea and coffee	Green tea, black tea, black coffee, traditional coffee

Nine multiple pass 24HDRs were collected every month for successive nine months. Nine 24HDRs contained three days of the weekend and six days of the weekdays. The first 24HDR was accomplished one month after the administration of the first FFQ (in August 2020) and the last 24HDR was recorded one month before the administration of the second FFQ (in April 2021). We asked the subjects to recall their consumption of all foods and beverages, including the names and quantities, during the previous 24 h. The previous 24 h

was defined as subsequent 24 h from the bedtime to the following bedtime in a day before 24HDR assessment. We then calculated the mean intake from 24HDR data for each subject.

The FFQ and 24HDR data were collected by the trained nutritionists at the subjects' homes. The trained nutritionists assisted the subjects to evaluate the portion size of food consumption using a book of photographs containing each food item with different portion sizes and kitchen utensils (i.e., spoons, tablespoons, scoops, glasses and cups). We used the Indonesian Food Composition Data to estimate the daily intakes of energy, macro- and micro-nutrients [19]. Additionally, we also used the food composition data of the United States Department of Agriculture database for few specific micronutrients due to lacking information from the Indonesian Food Composition Data [20].

2.3. Other Variables

We collected demographic and lifestyle characteristics including age, gender (male and female), marital status (married and not married/divorce) and smoking status (current smoker, ex-smoker and never smoke). We also measured body weight and height. Body mass index (BMI) was determined as weight (kg) divided by height squared (m^2).

2.4. Statistical Analysis

We used the SPSS statistical software package version 25 (SPSS Inc., Chicago, IL, USA) for statistical analyses. The normality of distributions of dietary data was analyzed by the Kolmogorov–Smirnov test. Variables not normally distributed (carbohydrate, cholesterol, vitamin A, thiamin, vitamin E, sodium and potassium) were natural log-transformed to reach a normal distribution and to allow the use of parametric tests. Means and standard deviations were counted for energy, nutrients and food group intakes for both FFQ and 24HDR. Reproducibility was evaluated by comparing the intakes between FFQ1 and FFQ2. We compared the data of FFQ1 with the mean of 24HDRs to assess the validity of the FFQ.

The reproducibility was assessed to compare the intakes between two FFQs using paired *t*-test, Pearson's correlation coefficient, intra-class correlation coefficient (ICC), weighted kappa statistic and misclassification analysis. The validity of the FFQ1 comparable with the mean of 24HDRs was analyzed by paired *t*-test, Pearson's correlation coefficient, ICC, weighted kappa statistic and misclassification analysis. We calculated the percentages of agreement (classification in the same or adjacent quartile) and disagreement (classification in one quartile apart or opposite quartile). De-attenuated correlation coefficients were counted using Rosner and Willett's formula to improve within-person variation in the mean of 24HDRs [21,22]. We analyzed Bland–Altman plots to compare the differences between FFQ1 and the mean of 24HDRs across energy, carbohydrate, fat and protein intakes. The differences between FFQ1 and the mean of 24HDR were plotted (FFQ1—the mean of 24HDRs; *y*-axis) against the mean of the two methods for energy, carbohydrate, fat and protein intakes $[(\text{FFQ1} + \text{the mean of 24HDRs})/2]$; *x*-axis) [23].

3. Results

Among 265 subjects who initially participated in our study, 16 subjects were excluded because they did not complete two FFQs or nine 24HDRs. Therefore, a total of 259 subjects were included in the final analysis. Total energy intake of all subjects in our study ranged between 500 and 5000 kcal. Table 2 shows the characteristics of the subjects. There were 57.9% male subjects and 55.2% current smokers. The mean age was 54.8 ± 9.6 years and the mean body mass index was $24.0 \pm 3.2 \text{ kg/m}^2$.

Table 2. Characteristics of the subjects ($n = 259$)¹.

Characteristics	All Subjects
Age, years	54.8 ± 9.6
Gender, %	
Male	57.9
Female	42.1
Marital status, %	
Married	81.5
Not married/divorce	18.5
Smoking status, %	
Current smoker	55.2
Ex-smoker	17.0
Never smoke	27.8
Body mass index, kg/m ²	24.0 ± 3.2

¹ Data are presented as means ± SD for continuous variables or % for categorical variables.

Table 3 describes the mean intakes of total energy and nutrients derived from two FFQs and the mean of 24HDRs, the comparisons from the paired *t*-test and the percentage of mean differences between two FFQs and between FFQ1 and the mean of 24HDRs. A paired *t*-test indicated that the intakes of energy and most nutrients, except for monounsaturated fatty acids (MUFA), β -carotene, niacin, sodium and copper, assessed by two FFQs were significantly different. The mean intakes for energy and nutrients evaluated by FFQ1 were higher than the data acquired by FFQ2 and the differences in mean intakes ranged from 1.7% for niacin to 27.8% for thiamin. The paired *t*-test also showed that the intakes of energy and all nutrients evaluated by FFQ1 were statistically different from the intakes evaluated by the mean of 24HDRs. Compared with the mean of 24HDRs as a reference method, the data of FFQ1 tended to overestimate intakes of all nutrients and food groups.

Table 4 illustrates the crude and energy-adjusted correlation coefficients for FFQ1 and FFQ2. These results gave the evaluation of the reproducibility of two FFQs. The crude Pearson's correlation coefficients for total energy and nutrients ranged from 0.50 for fiber to 0.81 for potassium and the crude ICC ranged from 0.44 for fiber to 0.78 for sodium and phosphorus. However, the correlation coefficients were changed after adjusting for energy. The energy-adjusted Pearson's correlation coefficients ranged from 0.30 for fiber to 0.78 for calcium and energy-adjusted ICC ranged from 0.31 for fiber to 0.66 for retinol and calcium. Table 4 also describes the crude and energy-adjusted and de-attenuated Pearson's correlation coefficients between FFQ1 and the mean of 24HDRs to evaluate the validity of the FFQ. The crude Pearson's correlation coefficients for FFQ1 and the mean of 24HDRs ranged from 0.41 for thiamin to 0.70 for β -carotene. The energy-adjusted coefficients ranged from 0.31 for phosphorus to 0.89 for copper, while the de-attenuated coefficients ranged from 0.54 for thiamin to 0.82 for zinc.

Table 5 shows the misclassification and weighted kappa values between FFQ1 and FFQ2 and between FFQ1 and the mean of 24HDRs. After we categorized the intakes into quartiles, the ranges of the agreement rates for the same or adjacent quartile classifications were from 81.1% for thiamin to 94.6% for carbohydrate as compared between FFQ1 and FFQ2 and 80.7% for vitamin D to 89.6% for β -carotene as compared between FFQ1 and the mean of 24HDRs. Extreme misclassification into opposite quartile was <6% for energy and all nutrients. The weighted kappa values described moderate conformity, ranging from 0.21 (fiber, cholesterol and riboflavin) to 0.42 (retinol and iron) between two FFQs and 0.20 (carbohydrate and phosphorus) to 0.34 (vitamin C) between FFQ1 and the mean of 24HDRs.

Table 3. Comparisons of nutrient intakes between two FFQs and between FFQ1 and the mean of 24HDRs.

Variables	FFQ1		FFQ2		24HDRs		<i>p</i> -Value ¹		Percentage of Mean Difference	
	Mean	SD	Mean	SD	Mean	SD	FFQ1 vs. FFQ2	FFQ1 vs. 24HDRs	FFQ1 vs. FFQ2	FFQ1 vs. 24HDRs
Energy (kcal) ²	1921	423	1877	386	1751	413	0.008	<0.001	2.4	9.7
Carbohydrate (g) ²	288	67	282	64	264	66	<0.001	<0.001	2.0	8.8
Fiber ³	18	4	16	3	15	3	<0.001	<0.001	8.9	19.5
Fat (g) ²	61	14	59	12	55	13	0.015	<0.001	2.5	11.0
MUFA (g) ³	22	6	21	5	21	6	0.055	<0.001	2.9	6.5
PUFA (g) ³	11	2	10	2	8	2	<0.001	<0.001	8.1	31.3
Cholesterol (mg) ³	129	33	123	32	109	29	0.002	<0.001	5.4	19.0
Protein (g) ²	55	13	52	11	50	12	0.005	<0.001	2.7	8.6
Retinol (μg) ²	438	121	405	142	331	131	<0.001	<0.001	7.9	32.1
β-Carotene (μg) ²	1884	531	1834	530	1716	502	0.08	<0.001	2.7	9.8
Vitamin C (mg) ²	116	41	112	33	90	34	0.038	<0.001	3.7	29.0
Vitamin D (μg) ³	2.81	0.56	2.59	0.66	2.49	0.73	<0.001	<0.001	8.5	12.9
Vitamin E (mg) ³	2.29	0.63	2.11	0.64	2.03	0.58	<0.001	<0.001	8.5	12.8
Thiamin (mg) ²	1.15	0.58	0.90	0.54	0.75	0.50	<0.001	<0.001	27.8	53.3
Riboflavin (mg) ²	1.87	0.60	1.76	0.40	1.64	0.41	<0.001	<0.001	6.3	14.0
Niacin (mg) ²	13.5	3.3	13.3	3.7	12.7	3.6	0.203	<0.001	1.7	6.6
Sodium (mg) ²	1705	466	1674	443	1580	453	0.09	<0.001	1.9	7.9
Potassium (mg) ²	3795	948	3538	945	3328	968	<0.001	<0.001	7.3	14.0
Calcium (mg) ²	579	122	540	124	477	127	<0.001	<0.001	7.2	21.4
Phosphorus (mg) ²	623	141	590	124	536	137	<0.001	<0.001	5.6	16.3
Magnesium (mg) ³	377	91	370	100	364	101	<0.001	<0.001	1.9	3.5
Iron (mg) ²	11.8	3.3	10.7	3.4	10.2	3.3	<0.001	<0.001	10.1	16.2
Copper (mg) ²	634	183	619	135	578	163	0.06	<0.001	2.5	9.6
Zinc (mg) ²	10.1	2.3	8.4	1.8	6.9	1.8	<0.001	<0.001	20.1	46.2

FFQ: food frequency questionnaire, 24HDR: 24 h dietary recall, MUFA: monounsaturated fatty acids, PUFA: polyunsaturated fatty acids.

¹ Differences were tested using paired *t*-test. ² Nutrient intakes were analyzed using the Indonesian Food Composition Data. ³ Nutrient intakes were analyzed using the United States Department of Agriculture (USDA).

Table 4. Correlation coefficients for nutrient and food group intakes between FFQ1 and FFQ2 and between FFQ1 and the mean of 24HDRs.

Variables	FFQ1 vs. FFQ2				FFQ1 vs. 24HDRs		
	PCC		ICC		PCC		
	Crude	Energy Adjust	Crude	Energy Adjust	Crude	Energy Adjust	De-Attenuated
Energy (kcal)	0.78	-	0.77	-	0.63	-	0.70
Carbohydrate (g)	0.76	0.47	0.76	0.48	0.56	0.49	0.67
Fiber (g)	0.50	0.30	0.44	0.31	0.51	0.33	0.69
Fat (g)	0.71	0.43	0.70	0.45	0.53	0.76	0.65
MUFA (g)	0.60	0.41	0.59	0.44	0.58	0.68	0.67
PUFA (g)	0.67	0.50	0.62	0.34	0.46	0.35	0.58
Cholesterol (mg)	0.58	0.56	0.55	0.52	0.52	0.46	0.71
Protein (g)	0.74	0.42	0.71	0.43	0.54	0.39	0.66

Table 4. Cont.

Variables	FFQ1 vs. FFQ2				FFQ1 vs. 24HDRs		
	PCC		ICC		PCC		
	Crude	Energy Adjust	Crude	Energy Adjust	Crude	Energy Adjust	De-Attenuated
Retinol (µg)	0.72	0.70	0.70	0.66	0.57	0.58	0.74
6 Carotene (µg)	0.64	0.44	0.63	0.38	0.70	0.66	0.78
Vitamin C (mg)	0.70	0.72	0.68	0.65	0.60	0.59	0.73
Vitamin D (µg)	0.65	0.44	0.58	0.44	0.46	0.46	0.60
Vitamin E (mg)	0.69	0.67	0.66	0.64	0.56	0.56	0.68
Thiamin (mg)	0.60	0.72	0.75	0.56	0.41	0.50	0.54
Riboflavin (mg)	0.66	0.62	0.60	0.54	0.55	0.70	0.68
2 Niacin (mg)	0.62	0.62	0.61	0.63	0.56	0.73	0.67
Sodium (mg)	0.61	0.44	0.78	0.53	0.65	0.59	0.74
Potassium (mg)	0.81	0.53	0.69	0.42	0.67	0.52	0.77
Calcium (mg)	0.78	0.78	0.74	0.66	0.60	0.83	0.76
Phosphorus (mg)	0.79	0.48	0.78	0.45	0.61	0.31	0.74
Magnesium (mg)	0.60	0.58	0.58	0.38	0.43	0.40	0.55
Iron (mg)	0.62	0.63	0.59	0.60	0.48	0.48	0.59
Copper (mg)	0.71	0.70	0.67	0.62	0.69	0.89	0.78
Zinc (mg)	0.55	0.53	0.57	0.55	0.48	0.48	0.82

FFQ: food frequency questionnaire, 24HDR: 24 h dietary recall, ICC: intra-class correlation coefficient, PCC: Pearson's correlation coefficient, MUFA: monounsaturated fatty acids, PUFA: polyunsaturated fatty acids.

Table 5. Agreement rates (%) for the same quartile or adjacent quartile classifications, disagreement rates (%) for one quartile apart or opposite quartile classifications and weighted kappa values between FFQ1 and FFQ2 and between FFQ1 and the mean of 24HDRs.

Variables	FFQ1 vs. FFQ2					FFQ1 vs. 24HDRs				
	Same Quartile	8 Adjacent Quartile	One Quartile Apart	Opposite Quartile	Weighted Kappa	Same Quartile	Adjacent Quartile	One Quartile Apart	Opposite Quartile	Weighted Kappa
Energy (kcal)	50.6	42.1	7.0	0.4	0.34	42.1	44.0	10.4	3.5	0.23
Carbohydrate (g)	51.4	43.2	4.6	0.8	0.35	39.3	45.2	11.2	4.3	0.20
18 Fat (g)	41.6	48.8	7.7	1.9	0.21	43.3	42.9	10.4	3.5	0.24
Fat (g)	52.1	39.4	7.0	1.5	0.37	44.4	40.5	10.8	4.3	0.26
MUFA (g)	50.6	36.3	9.3	3.9	0.34	49.4	37.1	10.8	2.7	0.32
PUFA (g)	43.6	50.2	4.6	1.5	0.25	46.7	37.8	11.2	4.3	0.29
Cholesterol (mg)	40.5	49.8	7.7	1.9	0.21	43.3	42.9	10.4	3.5	0.24
Protein (g)	48.7	42.1	8.9	0.4	0.30	44.8	38.2	15.8	1.2	0.25
Retinol (µg)	56.8	35.1	7.7	0.4	0.42	43.3	37.8	18.5	0.4	0.24
6 Carotene (µg)	50.6	36.3	9.3	3.9	0.34	49.8	39.8	6.9	3.5	0.33
Vitamin C (mg)	55.2	35.5	8.5	0.8	0.40	50.6	36.3	10.4	2.7	0.34
Vitamin D (µg)	54.1	34.4	8.8	2.7	0.39	45.6	35.1	13.5	5.8	0.27
Vitamin E (mg)	55.2	34.4	9.7	0.8	0.41	49.1	37.1	9.7	4.2	0.32
Thiamin (mg)	44.8	36.3	17.8	1.2	0.26	44.0	41.3	11.2	3.5	0.25
Riboflavin (mg)	40.5	49.8	7.7	1.9	0.21	43.3	42.9	10.4	3.5	0.24
2 Niacin (mg)	42.1	40.5	16.6	0.8	0.22	44.0	41.3	11.2	3.5	0.25
Sodium (mg)	55.2	37.1	7.0	0.8	0.40	47.9	41.7	7.7	2.7	0.32
Potassium (mg)	49.0	44.0	6.6	0.4	0.33	44.8	44.4	8.5	2.3	0.27
Calcium (mg)	46.8	42.9	7.0	3.4	0.39	44.8	44.0	10.4	0.8	0.24
Phosphorus (mg)	50.6	42.5	6.6	0.4	0.32	43.3	43.2	11.2	2.3	0.20
Magnesium (mg)	55.2	32.4	8.1	4.3	0.39	43.6	44.0	6.6	5.8	0.24
Iron (mg)	56.8	31.3	9.3	2.7	0.42	47.9	33.2	14.3	4.6	0.30
Copper (mg)	52.9	31.3	14.7	1.2	0.37	44.0	43.6	8.9	3.5	0.25
Zinc (mg)	42.9	42.1	13.9	1.2	0.22	42.5	39.4	13.1	5.0	0.23

FFQ: food frequency questionnaire, 24HDR: 24 h dietary recall, MUFA: monounsaturated fatty acids, PUFA: polyunsaturated fatty acids.

Figure 2 describes the level of discrepancy for energy and macronutrient intakes using the Bland-Altman plot method. A positive mean difference was shown in the analyses for energy and macronutrients. We also found that less than 10% of the subjects were outside the confidence intervals for all nutrients.




Figure 2. Bland-Altman plots for the intakes of energy (kcal) (a), carbohydrate (g) (b), fat (g) (c) and protein (g) (d). The difference between the mean estimate of energy and macronutrient intakes by two dietary assessment methods (y-axis) was plotted against the mean of nutrients measured by two dietary assessment methods (x-axis). FFQ: food frequency questionnaire, 24HDR: 24 h dietary recall.

4. Discussion

The reproducibility and validity of a 137-item FFQ with Indonesian dietary patterns were investigated in our study. Based on a previous study, the number of food items in FFQ might vary between 5 and 350 [8]. Our results indicated that the reproducibility and validity of the FFQ could be acceptable in relation to the reference method for nutritional epidemiology study among Indonesians.

The mean intakes for all nutrients from FFQ1 were higher compared to the data from FFQ2. This could be elucidated by the learning effect of the subjects. The subjects might estimate dietary intake more accurately after the survey of FFQ1 [24]. Crude Pearson's correlations and crude ICC for reproducibility between FFQ1 and FFQ2 in this study ranged between 0.50 and 0.81 and between 0.44 and 0.78, respectively. The coefficient correlation in our study was higher compared to that with a range of 0.20 to 0.80 in the previous studies [24–26]. Our results may reflect that this FFQ was relatively stable to assess dietary habits among the subjects. After energy adjustment, the correlation coefficients were higher only for few nutrients, but lower for most nutrients. The reason for increased correlation

coefficients after energy adjustment could be explained by the existed association between nutrient intake and energy intake. While decreased correlation coefficients after energy adjustment could because of systematic overestimation or underestimation [14]. The systematic error was also found in other results that energy adjustment did not increase the correlation coefficients between two FFQs [13,24,25].

Numerous time intervals from FFQ1 to FFQ2 have been recorded in other studies from several days to several years [11,27,28]. The short-term interval can cause high correlation coefficients as the subjects might easily memorize and restate the similar answers. The long-term interval can lead to weak correlation coefficients because of the variations in answers that reflect an alteration in dietary habits for a certain period of time [24]. In this study, to narrow the error and reduce the variation, we used nine-month interval between FFQ1 and FFQ2.

The percentage of the subjects categorized into the same, adjacent, or opposite quartiles and the weighted kappa values between FFQ1 and 24HDRs were similar to the previous results [15,24,29]. A study found that the agreement for grouping nutrient intakes into the same or adjacent category ranged approximately from 50 to 75% for macronutrient [33] and 48 to 70% for micronutrients [15]. Another study also showed that the weighted kappa values for energy and nutrients ranged from 0.20 to 0.45 between FFQ1 and FFQ2 and 0.07 to 0.42 between FFQ1 and 24HDRs [24], which were comparable with the data in our study. A large positive kappa value reflects great agreement among the tools. The kappa values between 0.21 and 0.40 were classified as fair agreement and between 0.41 and 0.60 were classified as moderate agreement [29], while the value ≤ 0 was indicated as no agreement [29].

Our study observed relative validity analyzed by comparing energy and nutrient intakes derived from FFQ1 with those derived from the mean of 24HDRs. We used nine dietary recalls during the study period to reduce the effect of seasonal variation of food consumption on dietary evaluation. Our results revealed that the intakes of all nutrients evaluated by FFQ1 showed a tendency to be overestimated compared with those assessed by the mean of 24HDRs. Positive mean differences were also observed using the Bland-Altman method. It could be explained that certain food items could be reported more than once when the subjects consumed the foods in a mixed dish [24].

Our study found moderate correlation coefficients between FFQ1 and the mean of 24HDRs according to the category of “tolerable” with Pearson’s correlations between 0.30 and 0.49 and “preferable” with Pearson’s correlations ≥ 0.50 for validation studies [30]. Our results were consistent with the previous reports [13,15]. A study in China revealed that the energy-adjusted correlations ranged between 0.19 and 0.58 [13]. Another study in Malaysia showed that the energy-adjusted correlations varied between 0.22 and 0.68 [15]. After adjusting for energy, we observed slightly decreased or no changed validity correlation between FFQ1 and the mean of 24HDRs for most nutrients. This could be because of the between-person variation in nutrient intakes. However, we found that $\geq 80.7\%$ of the subjects were categorized in the same or adjacent quartile, which was also similar to the previous results [13,24,25,31–33]. The weighted kappa values in this study achieved an acceptable agreement for most nutrients [29]. Our results were comparable with other studies with the weighted kappa values for nutrient intakes from 0.20 to 0.45 between two FFQs and from 0.07 to 0.42 between FFQ1 and the mean of 24HDRs [24], or from 0.35 to 0.53 between two FFQs and from 0.37 to 0.52 between FFQ1 and the mean of 24HDRs [25].

Strengths and Limitations

The present study had some strengths. To the best of our knowledge, this is the first study to discuss the validity and reproducibility of nutrition surveys conducted among Indonesian adults. Moreover, the characteristics of the subjects including demographics and lifestyle represented the Indonesian population. However, this study also had several limitations. We used 24HDR as the reference method. Both 24HDR and FFQ had the same error due to subjects’ incomplete memory and social-desirability bias [34]. Previous studies

stated that biomarkers could be considered as an alternative reference method [14,34]. However, no biomarkers were measured in this study. Some studies showed that the correlation of food intake with nutrient status and its biomarker was not exactly direct because the absorption of the nutrients in the body should also be considered [35,36]. Moreover, 24HDR was often used in the validity study of FFQ [10–12,24,34] because 24HDR estimated dietary intake more precisely than FFQ [10,37]. In addition, this analysis was restricted only to middle-aged and older adults aged 40–80 years. It is uncertain whether our FFQ can also be appropriate for dietary assessment among children or younger adults.

5. Conclusions

In conclusion, the 137-item FFQ designed for this study shows acceptable reproducibility and validity. Hence, the FFQ can be utilized as a reliable tool in epidemiological studies among middle-aged and older adults in different settings in Indonesia. Further evaluation and modifications of food items in the proposed FFQ are needed to improve its validity and reproducibility for some nutrients.

Author Contributions: Conceptualization and methodology, A.S., D.N.A., R.P., C.N., D.Y.F. and J.C.-J.C.; formal analysis, A.S. and R.P.; investigation, A.S., C.N. and D.Y.F.; writing—original draft preparation, A.S.; writing—review and editing, A.S. and J.C.-J.C. All authors have read and agreed to the published version of the manuscript.

Funding: This research was funded by Institute for Research and Community Services (LPPM-RPIBT), Diponegoro University (233-36/UN7.6.1/PP/2020).

Institutional Review Board Statement: The study was conducted according to the guidelines of the Declaration of Helsinki and approved by the Health Research Ethics Committee of Medical Faculty of Diponegoro University (No.172/EC/KEPK/FK-UNDIP/VII/2020).

Informed Consent Statement: Informed consent was obtained from all subjects involved in the study.

Data Availability Statement: The data presented in this study are available from the first author on reasonable request.

Acknowledgments: The authors are grateful to the subjects who took part in this study and the research team for their cooperation and assistance.

Conflicts of Interest: The authors declare no conflict of interest. The funders had no roles in the design of the study, the collection, analyses, or interpretation of data, the writing of the manuscript, or the decision to publish the results.

References

1. Yach, D.; Hawkes, C.; Gould, C.L.; Hofman, K.J. The global burden of chronic diseases: Overcoming impediments to prevention and control. *JAMA* **2004**, *291*, 2616–2622. [\[CrossRef\]](#) [\[PubMed\]](#)
2. Ng, R.; Sutradhar, R.; Yao, Z.; Wodchis, W.P.; Rosella, L.C. Smoking, drinking, diet and physical activity-modifiable lifestyle risk factors and their associations with age to first chronic disease. *Int. J. Epidemiol.* **2020**, *49*, 113–130. [\[CrossRef\]](#) [\[PubMed\]](#)
3. Syauqy, A.; Hsu, C.Y.; Lee, H.A.; Rau, H.H.; Chao, J.C.J. Association between dietary patterns and kidney function parameters in adults with metabolic syndrome: A cross-sectional study. *Nutrients* **2020**, *13*, 40. [\[CrossRef\]](#) [\[PubMed\]](#)
4. Ramos-Lopez, O.; Milagro, F.I.; Allayee, H.; Chmurzynska, A.; Choi, M.S.; Curi, R.; De Caterina, R.; Ferguson, L.R.; Goni, L.; Kang, J.X.; et al. Guide for current nutrigenetic, nutrigenomic, and nutriepigenetic approaches for precision nutrition involving the prevention and management of chronic diseases associated with obesity. *J. Nutr. Nutr.* **2017**, *10*, 43–62. [\[CrossRef\]](#)
5. Shim, J.S.; Oh, K.; Kim, H.C. Dietary assessment methods in epidemiologic studies. *Epidemiol. Health* **2014**, *36*, e2014009. [\[CrossRef\]](#) [\[PubMed\]](#)
6. Syauqy, A.; Hsu, C.Y.; Rau, H.H.; Chao, J.C.J. Association of dietary patterns with components of metabolic syndrome and inflammation among middle-aged and older adults with metabolic syndrome in Taiwan. *Nutrients* **2018**, *10*, 143. [\[CrossRef\]](#)
7. Syauqy, A.; Noer, E.R.; Fajrani, A.M.; Kurniawati, D.M.a.; Purwanti, R.; Rahadiyanti, A.; Rahma, D.E. Dietary patterns were associated with obesity parameters among healthy women. *J. Nutr. Coll.* **2020**, *9*, 6. [\[CrossRef\]](#)
8. Cade, J.; Thompson, R.; Burley, V.; Warm, D. Development, validation and utilisation of food-frequency questionnaires—A review. *Public Health Nutr.* **2002**, *5*, 567–587. [\[CrossRef\]](#)
9. Cui, Q.; Xia, Y.; Wu, Q.; Chang, Q.; Niu, K.; Zhao, Y. A meta-analysis of the reproducibility of food frequency questionnaires in nutritional epidemiological studies. *Int. J. Behav. Nutr. Phys. Act.* **2021**, *18*, 12. [\[CrossRef\]](#)

10. Dehghan, M.; Ilow, R.; Zatonska, K.; Szuba, A.; Zhang, X.; Mente, A.; Regulska-Ilow, B. Development, reproducibility and validity of the food frequency questionnaire in the Poland arm of the Prospective Urban and Rural Epidemiological (PURE) study. *J. Hum. Nutr. Diet.* **2012**, *25*, 225–232. [CrossRef]
11. Xue, Y.; Yang, K.; Wang, B.; Liu, C.; Mao, Z.; Yu, S.; Li, X.; Wang, Y.; Sun, H.; Wang, C.; et al. Reproducibility and validity of an FFQ in the Henan Rural Cohort Study. *Public Health Nutr.* **2020**, *23*, 34–40. [CrossRef]
12. Visser, M.; Elstgeest, L.E.M.; Winkens, L.H.H.; Brouwer, I.A.; Nicolaou, M. Relative validity of the HELIUS food frequency questionnaire for measuring dietary intake in older adult participants of the longitudinal aging study Amsterdam. *Nutrients* **2020**, *12*, 1998. [CrossRef] [PubMed]
13. Ye, Q.; Hong, X.; Wang, Z.; Yang, H.; Chen, X.; Zhou, H.; Wang, C.; Lai, Y.; Sun, L.; Xu, F. Reproducibility and validity of an FFQ developed for adults in Nanjing, China. *Br. J. Nutr.* **2016**, *115*, 887–894. [CrossRef] [PubMed]
14. Doustmohammadian, A.; Amini, M.; Esmailzadeh, A.; Omidvar, N.; Abtahi, M.; Dadkhah-Piraghaj, M.; Nikooyeh, B.; Neyestani, T.R. Validity and reliability of a dish-based semi-quantitative food frequency questionnaire for assessment of energy and nutrient intake among Iranian adults. *BMC Res. Notes* **2020**, *13*, 95. [CrossRef] [PubMed]
15. Nurul-Fadhilah, A.; Teo, P.S.; Foo, L.H. Validity and reproducibility of a food frequency questionnaire (FFQ) for dietary assessment in Malay adolescents in Malaysia. *Asia Pac. J. Clin. Nutr.* **2012**, *21*, 97–103.
16. Ramadhani, P.P.; Dieny, F.F.; Kurniawati, D.; Sandi, H.; Fitranti, D.Y.; Rahadiyanti, A.; Tsani, A.F.A. Household food security and diet quality with chronic energy deficiency among preconception women. *Indones. J. Nutr.* **2021**, *9*, 111–122. [CrossRef]
17. Hermingtyas, E.H.; Ng, T.S. Prevalence and distribution of metabolic syndrome and its components among provinces and ethnic groups in Indonesia. *BMC Public Health* **2019**, *19*, 377. [CrossRef]
18. Statistics of Semarang Municipality. Available online: <https://semarangkota.bps.go.id/> (accessed on 1 August 2021).
19. Indonesian Food Composition Data. Available online: <https://www.panganku.org/en-EN/beranda> (accessed on 1 August 2021).
20. United States Department of Agriculture. Available online: <https://ndb.nal.usda.gov/ndb/search/list> (accessed on 1 August 2021).
21. Rosner, B.; Willett, W.C. Interval estimates for correlation coefficients corrected for within-person variation: Implications for study design and hypothesis testing. *Am. J. Epidemiol.* **1988**, *127*, 377–386. [CrossRef]
22. Willett, W. *Nutritional Epidemiology*; Oxford University Press: Oxford, UK, 2012; Volume 40.
23. Bland, J.M.; Altman, D. Statistical methods for assessing agreement between two methods of clinical measurement. *Lancet* **1986**, *327*, 307–310. [CrossRef]
24. Zhuang, M.; Yuan, Z.; Lin, L.; Hu, B.; Wang, X.; Yang, Y.; Chen, X.; Jin, L.; Lu, M.; Ye, W. Reproducibility and relative validity of a food frequency questionnaire developed for adults in Taizhou, China. *PLoS ONE* **2012**, *7*, e48341. [CrossRef]
25. Xia, W.; Sun, C.; Zhang, L.; Zhang, X.; Wang, J.; Wang, H.; Wu, L. Reproducibility and relative validity of a food frequency questionnaire developed for female adolescents in Suihua, North China. *PLoS ONE* **2011**, *6*, e19656. [CrossRef]
26. Marques-Vidal, P.; Ross, A.; Wynn, E.; Rezzi, S.; Paccaud, F.; Decarli, B. Reproducibility and relative validity of a food-frequency questionnaire for French-speaking Swiss adults. *Food Nutr. Res.* **2011**, *55*, 5905. [CrossRef] [PubMed]
27. De la Fuente-Arrillaga, C.; Ruiz, Z.V.; Bes-Rastrollo, M.; Sampson, L.; Martinez-González, M.A. Reproducibility of an FFQ validated in Spain. *Public Health Nutr.* **2010**, *13*, 1364–1372. [CrossRef] [PubMed]
28. Vereecken, C.A.; Maes, L. A Belgian study on the reliability and relative validity of the health behaviour in school-aged children food-frequency questionnaire. *Public Health Nutr.* **2003**, *6*, 581–588. [CrossRef] [PubMed]
29. McHugh, M.L. Interrater reliability: The kappa statistic. *Biochem. Med.* **2012**, *22*, 276–282. (In Zagreb) [CrossRef]
30. Masson, L.F.; McNeill, G.; Tomany, J.O.; Simpson, J.A.; Peace, H.S.; Wei, L.; Grubb, D.A.; Bolton-Smith, C. Statistical approaches for assessing the relative validity of a food-frequency questionnaire: Use of correlation coefficients and the kappa statistic. *Public Health Nutr.* **2003**, *6*, 313–321. [CrossRef] [PubMed]
31. Deschamps, V.; de Lauzon-Guillain, B.; Lafay, L.; Borys, J.M.; Charles, M.A.; Romon, M. Reproducibility and relative validity of a food-frequency questionnaire among French adults and adolescents. *Eur. J. Clin. Nutr.* **2009**, *63*, 282–291. [CrossRef]
32. Zhang, C.X.; Ho, S.C. Validity and reproducibility of a food frequency questionnaire among Chinese women in Guangdong province. *Asia Pac. J. Clin. Nutr.* **2009**, *18*, 240–250.
33. Villegas, R.; Yang, G.; Liu, D.; Xiang, Y.B.; Cai, H.; Zheng, W.; Shu, X.O. Validity and reproducibility of the food-frequency questionnaire used in the Shanghai men's health study. *Br. J. Nutr.* **2007**, *97*, 993–1000. [CrossRef]
34. Praagman, J.; Adolphi, A.P.; van Rossum, C.T.; Sluijs, I.; van der Schouw, Y.T.; Beulens, J.W. Reproducibility and relative validity of a FFQ to estimate the intake of fatty acids. *Br. J. Nutr.* **2016**, *115*, 2154–2161. [CrossRef]
35. Cade, J.E.; Burley, V.J.; Warm, D.L.; Thompson, R.L.; Margetts, B.M. Food-frequency questionnaires: A review of their design, validation and utilisation. *Nutr. Res. Rev.* **2004**, *17*, 5–22. [CrossRef] [PubMed]
36. Noor Hafizah, Y.; Ang, L.C.; Yap, F.; Nurul Najwa, W.; Cheah, W.L.; Ruzita, A.T.; Jumuddin, F.A.; Koh, D.; Lee, J.A.C.; Essau, C.A.; et al. Validity and reliability of a food frequency questionnaire (FFQ) to assess dietary intake of preschool children. *Int. J. Environ. Res. Public Health* **2019**, *16*, 4722. [CrossRef] [PubMed]
37. Haftenberger, M.; Heuer, T.; Heidemann, C.; Kube, F.; Krems, C.; Mensink, G.B. Relative validation of a food frequency questionnaire for national health and nutrition monitoring. *Nutr. J.* **2010**, *9*, 36. [CrossRef] [PubMed]

Reproducibility and Validity of a Food Frequency Questionnaire (FFQ) Developed for Middle-Aged and Older Adults in Semarang, Indonesia

ORIGINALITY REPORT

14%

SIMILARITY INDEX

10%

INTERNET SOURCES

12%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	Jessica Winarto, Dae-Geun Song, Cheol-Ho Pan. "The Role of Fucoxanthin in Non-Alcoholic Fatty Liver Disease", International Journal of Molecular Sciences, 2023 Publication	1 %
2	doczz.net Internet Source	1 %
3	Submitted to jku Student Paper	1 %
4	Se-Young Oh. "Reproducibility and validity of a semiquantitative food frequency questionnaire to assess dietary intakes for the Korean elderly", Ecology of Food and Nutrition, 2002 Publication	1 %
5	eprints.bmsu.ac.ir Internet Source	1 %
6	Submitted to University of Melbourne Student Paper	<1 %

7	Submitted to University of Reading Student Paper	<1 %
8	Sameera A. Talegawkar, T. Tanaka, J. E. Maras, L. Ferrucci, K. L. Tucker. "Validation of nutrient intake estimates derived using a semi-quantitative FFQ against 3 day diet records in the Baltimore Longitudinal Study of Aging", The journal of nutrition, health & aging, 2016 Publication	<1 %
9	avmajournals.avma.org Internet Source	<1 %
10	Chang Liu, Yuan Xue, Yan Wang, Yujing Zhang et al. "Association between Daily Egg Intake and Lipid Profiles in Adults from the Henan Rural Cohort Study", Nutrition, Metabolism and Cardiovascular Diseases, 2020 Publication	<1 %
11	Submitted to Michigan Technological University Student Paper	<1 %
12	www.unirepository.svkri.uniri.hr Internet Source	<1 %
13	Satalic, Z.. "Short food frequency questionnaire can discriminate inadequate and adequate calcium intake in Croatian	<1 %

postmenopausal women", Nutrition Research,
200709

Publication

14

Gabriele Nagel, Dorothee Zoller, Tilla Ruf, Sabine Rohrmann, Jakob Linseisen. "Long-term reproducibility of a food-frequency questionnaire and dietary changes in the European Prospective Investigation into Cancer and Nutrition (EPIC)-Heidelberg cohort", British Journal of Nutrition, 2007

Publication

<1 %

15

Jeanine L Marnewick, Irma Venter, Fanie Rautenbach, Henry Neethling, Maritha Kotze. "Rooibos: Effect on Iron Status in South African Adults at Risk for Coronary Heart Disease", American Chemical Society (ACS), 2013

Publication

<1 %

16

journals.sagepub.com

Internet Source

<1 %

17

scholarworks.iupui.edu

Internet Source

<1 %

18

www.nourishorganics.in

Internet Source

<1 %

19

Submitted to University of North Texas Health Science Center

Student Paper

<1 %

20	Submitted to Western Governors University Student Paper	<1 %
21	S De Henauw. "Evaluation of a 2-day food record to determine iron, calcium and vitamin C intake in young Belgian women", European Journal of Clinical Nutrition, 01/2008 Publication	<1 %
22	Chiba, Nagako, Nagako Okuda, Akira Okayama, Takashi Kadowaki, and Hirotugu Ueshima. "Development of a Food Frequency and Quantity Method for Assessing Dietary Habits of Japanese Individuals -Comparison with Results from 24hr Recall Dietary Survey", Journal of Atherosclerosis and Thrombosis, 2008. Publication	<1 %
23	Submitted to Leeds Beckett University Student Paper	<1 %
24	moam.info Internet Source	<1 %
25	uwe-repository.worktribe.com Internet Source	<1 %
26	Yi Cao, Cheng Wang, Jun Liu, Zhao-min Liu, Wen-hua Ling, Yu-ming Chen. "Greater serum carotenoid levels associated with lower	<1 %

prevalence of nonalcoholic fatty liver disease in Chinese adults", Scientific Reports, 2015

Publication

27

Constance A Gewa, Suzanne P Murphy, Charlotte G Neumann. "A comparison of weighed and recalled intakes for schoolchildren and mothers in rural Kenya", Public Health Nutrition, 2009

Publication

<1 %

28

Fatemeh TOORANG, Bahareh SASANFAR, Soodeh RAZEGHI JAHROMI, Soraiya EBRAHIMPOUR KOUJAN et al. "Validation of Diet History Questionnaire in Assessing Energy and Nutrient Intakes of Iranian Population", Iranian Journal of Public Health, 2020

Publication

<1 %

29

Submitted to University of Malaya

Student Paper

<1 %

30

WEI-CHUNG VIVIAN YANG, HENG-KIEN AU, CHING-WEN CHANG, HUEI-WEN CHEN et al. "Matrix remodeling and endometriosis", Reproductive Medicine and Biology, 2005

Publication

<1 %

31

acikbilim.yok.gov.tr

Internet Source

<1 %

32

bmcpediatr.biomedcentral.com

Internet Source

<1 %

33

citeseerx.ist.psu.edu

Internet Source

<1 %

34

kuscholarworks.ku.edu

Internet Source

<1 %

35

shapeamerica.tandfonline.com

Internet Source

<1 %

36

www.gery.fi

Internet Source

<1 %

37

Gross, D.P.. "Acute Opioid Administration Improves Work-Related Exercise Performance in Patients With Chronic Back Pain", *Journal of Pain*, 200809

Publication

<1 %

38

Liisa Vahatalo. "Relative validity of a dietary interview for assessing infant diet and compliance in a dietary intervention trial", *Maternal and Child Nutrition*, 7/2006

Publication

<1 %

39

iron-body.com

Internet Source

<1 %

40

pure.rug.nl

Internet Source

<1 %

41

studentsrepo.um.edu.my

Internet Source

<1 %

42

www.degruyter.com

Internet Source

<1 %

43

www.reviewofoptometry.com

Internet Source

<1 %

44

Alireza Sadeghi, Karim Parastouei, Sharareh Seifi, Adnan Khosravi et al. "Inflammatory Potential of Diet and Odds of Lung Cancer: A Case-Control Study", Nutrition and Cancer, 2022

Publication

<1 %

45

Chermaine T. Kwant, Frans A. L. van der Horst, Harm J. Bogaard, Frances S. de Man, Anton Vonk Noordegraaf. "Nutritional status in pulmonary arterial hypertension", Pulmonary Circulation, 2022

Publication

<1 %

46

Fiona Riordan, Kathleen Ryan, Ivan J Perry, Matthias B Schulze et al. "A systematic review of methods to assess intake of fruits and vegetables among healthy European adults and children: a DEDIPAC (DEterminants of Diet and Physical Activity) study", Public Health Nutrition, 2016

Publication

<1 %

47 Nor A Mohd-Shukri, Jennifer L Bolton, Jane E Norman, Brian R Walker, Rebecca M Reynolds. "Evaluation of an FFQ to assess total energy and nutrient intakes in severely obese pregnant women", Public Health Nutrition, 2012
Publication

48 Subrata D. Nath, Fatma G. Huffman. "Validation of a semiquantitative food frequency questionnaire to assess energy and macronutrient intakes of Cuban Americans", International Journal of Food Sciences and Nutrition, 2009
Publication

49 Wm. Cameron Chumlea, David B. Cockram, Johanna T. Dwyer, Haewook Han, Mary Pat Kelly. "Chapter 4 Nutrition Assessment in Chronic Kidney Disease", Springer Science and Business Media LLC, 2008
Publication

50 docplayer.com.br
Internet Source

51 epdf.pub
Internet Source

52 eprints.ucl.ac.uk
Internet Source

53

Internet Source

<1 %

54

www.imrpress.com

Internet Source

<1 %

55

Emily B Levitan, Camilla W Westgren, Simin Liu, Alicja Wolk. "Reproducibility and validity of dietary glycemic index, dietary glycemic load, and total carbohydrate intake in 141 Swedish men", The American Journal of Clinical Nutrition, 2007

Publication

<1 %

56

Ho, Y.-S.. "Removal of copper ions from aqueous solution by tree fern", Water Research, 200305

Publication

<1 %

57

M Vanderheyden. "Non-compaction of the left ventricle", Heart, 3/1/2004

Publication

<1 %

Exclude quotes

Off

Exclude matches

Off

Exclude bibliography

On

Reproducibility and Validity of a Food Frequency Questionnaire (FFQ) Developed for Middle-Aged and Older Adults in Semarang, Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11