

KORESPONDENSI PAPER

JUDUL : Bioactive components and antibacterial activity in Robusta coffee leaves (Coffea canephora)

JURNAL : International Journal of Pharmaceutical Research

STATUS : Scopus Q2 (0,28)

No	Aktivitas	Tanggal	Halaman
1	Submission Artikel	18 November 2019	2-12
2	Pengiriman Hasil Review	22 November 2019	13
3	Pengiriman Hasil Review OJS	15 Januari 2020	14
4	Hasil Perbaikan Artikel	4 Maret 2020	15-25
5	Letter of Accepted	23 Januari 2020	26
6	Artikel Published	18 Mei 2020	27
7	Artikel Final	Vol 12 No.3 : 1374-1382	28-37

Submission Artikel

The screenshot shows an email client interface. On the left is a sidebar with folders: Compose, Inbox (187), Starred, Snoozed, Important, Sent, Drafts (54), Categories, and More. Below these are Labels: [imap]/Archive, [imap]/Drafts, and Notes. The main area displays an email from 'ICHWB Submission and Review System' with subject '[ICHWB2019] Submission Acknowledgement - 14758'. The email content includes a thank you message and submission details:

Submitted on Monday, 18 November 2019 - 4:24pm

Thank you for your submission
Submitted values are:

==Corresponding Author Information==
Designation: Dr.
Full Name: Gemala Anjani
Affiliation: Universitas Diponegoro
Email: gemaanjani@gmail.com

==Author Information==
Authors Names: Gemala Anjani, Nurmasari Widyastuti, Zulfatul Masruroh, Rizka Ayu Dwi Yuliana, Vita Gustin Almira
Authors emails: gemaanjani@gmail.com, widyastutinurmasari@gmail.com, masrurohzulfatul@gmail.com, rickaradya10@gmail.com, vitagalmira@gmail.com

==Paper Information==

Bioactive components and antibacterial activity in Robusta coffee leaves (*cofea canephora*)

Gemala Anjani^{1,2,φ,*}, Nurmasari Widyastuti^{1,2,φ}, Zulfatul Masruroh¹, Rizka Ayu Dwi Yuliana¹, Vita Gustin Almira¹, A.Fahmy Arif Tsani^{1,2}, Choirun Nissa^{1,2}, M Iqbal Prawira-Atmaja³

¹ Department of Nutrition Science, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

² Center of Nutrition Research (CENURE), Universitas Diponegoro, Semarang, Indonesia

³ Research Institute for Tea and Cinchona, Bandung, Indonesia

Abstract

Purpose: To analyze bioactive components and antibacterial activity in brewed Robusta coffee leaves that processed by *Japanese Style Green Tea Process (JGTP)*.

Methodology: This was an experimental research with the sample of Robusta coffee leaves and brewed Robusta coffee leaves. The coffee leaves were brewed in water at 70°C during 10 minutes. The analysis of nutrition contents by proximate method, total flavonoids by calorimeter method, chlorogenic acid and caffeine by HPLC method, antioxidant activity by DPPH method, total chlorophyll using the arnon formula and antibacterial activity by disk diffusion method in 5, 10, 15, and 20 g.

Results: There were significant differences in color, taste, and acceptability of brewed Robusta coffee leaves. The best acceptance of brewed Robusta coffee leaves was 5 g of dried coffee leaves. Robusta coffee leaves contain 3,1% water, 6,03% ash, 17,76% protein, 2,37% fat, and 70,83% carbohydrate. The content of flavonoids, chlorogenic acid and caffeine in Robusta coffee leaves were 0.81 mgQE/g, 16.92 mg/g, 0.46 mg while in the brewed were 0.10 mgQE/g, 3.49 mg/g, 0.19 mg/g, respectively. The inhibition of free radical in Robusta coffee leaves were 90,2-92,7%, while in the brewed were 83,3-86,1%. Total chlorophyll of dried leaves were 63,62-69,39 mg/L and the brewed were 2,08-6,01 mg/L. Antibacterial activity *Escherichia coli* 7,65-9,89 mm and *Salmonella tiphy* 7,47-8,70 mm inhibition zone.

Applications/Originality/Value: The role of various bioactive components in Robusta coffee leaves on antioxidant and antibacterial activity.

^φ Both authors contributed equally to this manuscript

* Corresponding author: gemaanjani@gmail.com

Introduction

Coffee plant (*Coffea sp.*) is included in the family *Rubiaceae*. In general, there are three types of coffee, including Arabica coffee (*Coffea arabica*), Robusta coffee (*Coffea canephora*), and Liberica coffee (*Coffea liberica*). The most widely consumed coffee in Indonesia is Robusta coffee. Robusta coffee can be easily grown at lower altitude of less than 1000 meters above sea level and it resilient against disease and pests.(Kanisius, 1988)

The results of research on Arabica coffee leaves have nutritional content based on proximate components. Proximate components of Arabica coffee leaves from the Ethiopian region include 5.5-7.8% moisture, 8.8-12.4% ash content, 14.4-19.0% protein, 4.5-12.5% fat, and 51.0-63.9% carbohydrate.(Woldesenbet, 2005) Robusta coffee leaves contain phytochemicals that vary depending on leaf development.(X.-M. Chen, Ma, & Kitts, 2018) The younger the coffee leaves, the higher the chlorogenic acid content even 10 times higher than the old coffee leaves.(Campa et al., 2017) The content of chlorogenic acid in Robusta coffee is higher compared to Arabica coffee, respectively 11.3% and 4.1%.(Ky et al., 2001) The caffeine content in young Robusta coffee leaves (0.29-0.5%) is lower than the caffeine content in its fruit (1.6-2.4%).(Khotimah, 2014)

Robusta coffee leaves also contain phenolic acid and total flavonoids, respectively 27.04 µg/g and 10.90 µg/g higher than Arabica coffee leaves respectively 21.80 µg/g and 8.08 µg/g.(Nayeem, Denny, & Mehta, 2011) Flavonoids as natural antioxidants can prevent the occurrence of diseases, such as cancer, metabolic syndrome, diabetes mellitus, and atherosclerosis which associated to oxidative stress.(X. Chen, 2019; Manach, Scalbert, Morand, Rémésy, & Jiménez, 2004)

The content of caffeine, flavonoids, and chlorogenic acid of coffee leaves can also function as the prevention of diarrhea. Caffeine can inhibit protein and DNA synthesis by inhibiting the merging of adenine and thymidine. Caffeine also increases genotoxicity after DNA damage.(Pruthviraj, Shital, Suchita, & Shilpa, 2011) Flavonoids as an antibacterial by inhibiting nucleic acid synthesis, inhibiting energy metabolism, and changes in cytoplasmic membrane function.(Farhadi, Khameneh, Iranshahi, & Iranshahy, 2018) In addition to flavonoids, there are chlorogenic acid contents that function as antibacterial. Chlorogenic acid is a phenolic bioactive compound causes physiological changes in bacterial cell membranes that encourage damage to bacterial nucleotides.(Kabir, Katayama, Tanji, & Nakamura, 2014; Lou, Wang, Zhu, Ma, & Wang, 2011)

The processing of coffee leaves to obtain optimal antibacterial is done by minimizing chemical and enzymatic reactions to prevent damage in phytochemical content. The process can be obtained using green tea processing.(X.-M. Chen et al., 2018) The processing of green tea is able to produce phenolic components 6 times higher than black tea. Green tea processing aims to maintain the content of natural polyphenols which have beneficial effects on health. (Chacko, Thambi, Kuttan, & Nishigaki, 2010). Therefore, the aim of this research was to examine bioactive components, including caffeine, chlorogenic acid, total flavonoids, antioxidant activity, total chlorophyll, and nutritional contents and antibacterial activity in Robusta coffee leaves by green tea processing.

Materials and Methods

The samples used *Robusta canephora* Pierre ex Froehner from Gambung village. Coffee leaves were processed using the Japanese Style Green Tea Process (JGTP) method at the Tea Quality Processing and Testing Laboratory, Tea and Quinine Research Center (PPTK) Gambung, Bandung Regency, West Java, Indonesia. Preliminary research was conducted to obtain the JGTP procedure at the withering stage, including 2 minutes steaming method and 45 s, 60 s, and 75 s of blanching. The best withering method was chosen based on the results of the best organoleptic tests conducted by trained panelists. The best withering result that will be used in the processing of coffee leaves was the blanching method for 75 s.

This study was an experimental study with one factor completely randomized design (CRD) with 2 samples, dried Robusta coffee leaves and brewed Robusta coffee leaves. Dried Robusta coffee leaves were brewed with 200 ml of water in 70°C for 10 minutes. Samples were analyzed in duplicate with three replications.

Processing Robusta Coffee Leaves with JGTP Method

JGTP method was carried out through 5 steps, picking; blanching; cooling, crushing, and drying. Coffee leaves were picked from 2nd, 3rd, and 4th leaf of each branch of coffee plants, sorted, then carried out the blanching process for 75 s. Leaves were soaked in water and put in seducer for ± 15 minutes while separating the leaves from its midrib. The next process was crushing the leaves three times using a crushing-tearing-curling machine (CTC) (CNC, Sri Lanka). The last process was the drying process. The drying process was 4-5 hours long by a rack drier machine (CNC, Sri Lanka) at 80°C.

Proximate Analysis

Proximate analyses using dried Robusta coffee leaves samples processed by JGTP to analyze the moisture, ash, fat, protein, and carbohydrate contents with AOAC 2005 method.(Analysis., 2005)

Total Flavonoid Analysis

5 grams of dried Robusta coffee leaves in 200 ml of methanol were homogenized with a magnetic stirrer for \pm 30 minutes. The solution was taken 60 ml for centrifugation, then coffee leaves produced supernatant in methanol. Meanwhile, 5 grams of dried Robusta coffee leaves were brewed in 200 ml of hot water (70°C) for 10 minutes and stirred. After being dissolved in water, 60 ml was taken to be centrifuged, coffee leaves supernatant will be produced in brewed water. Afterwards, two samples were analyzed.

Total flavonoid analysis using aluminum chloride (AlCl₃) calorimetry method with absorbance measurement using UV-VIS spectrophotometer. 1 ml of sample solution was mixed with 9 ml of methanol. Pipette 2.8 ml sample solution; 0.4 ml AlCl₃ 5%; 6.8 ml of 5% acetic acid. Incubation for 30 minutes, measure the absorbance (λ 428 nm). Calibration curves used the standard quercetin in methanol (0.005 mg / ml) with varying concentrations of 0 μ g/ml, 5 μ g/ml, 10 μ g/ml, 20 μ g/ml, 40 μ g/ml, and 60 μ g/ml. Standard was carried out using the same procedures as samples. (Rhayanne T. M. Ramos, Ferreira, & Soares, 2017; Silva, Pezzini, & Soares, 2015)

Analysis of Chlorogenic Acid and Caffeine

Analysis of chlorogenic acid using 20 g of dried coffee leaves extracted with 150 mL methanol:water (50:50 v/v) in soxhlet. After 4-5 cycles, soxhletation was stopped and cooled down at room temperature. The extract was transferred into a 200 mL flask and the volume was adjusted to 200 mL with a mixture of methanol-water. Meanwhile, caffeine analysis using 10-20 g of coffee leaves were added to 5 g of MgO and put into 200 mL of water. Furthermore, the samples were heated in a water bath at 90°C and sterilized for 20 minutes. Then the extracts were filtered with 0.45 μ L, 25 mm cellulose filter cartridge. The filtrates were injected into the HPLC.(Naegele, 2016a, 2016b)

Total Chlorophyll Test

The total chlorophyll test was carried out by 5, 10, 15 and 20 grams of brewed Robusta coffee leaves compared to the samples dissolved with 80% acetone. Total chlorophyll test was conducted using 663 nm and 645 nm wave spectrophotometry. Chlorophyll levels were calculated using the Arnon formula:(Prawira-Atmaja et al., 2018)

Chlorophyll a = (12.7 x A663) - (2.69 x A645)

Chlorophyll b = (22.9 x A645) - (4.68 x A663)

Total chlorophyll = (20.2 x A645) + (8.02 x A663)

Analysis of Antioxidant Activity

Sample preparation was performed as a total flavonoid test. Antioxidant activity was analyzed using the DPPH method (1,1-diphenyl-2-picrylhydrazyl). A DPPH 0.4 mM solution was prepared by 0.00394 g of DPPH powder and dissolved in absolute methanol to 25 ml. The test was done by piping 0.1 ml of samples and sufficient volume with absolute methanol into a 5 ml measuring flask, then vortex. Measurement of antioxidant activity was done by adding 1 ml of DPPH 0.4 mM solution, then vortex for \pm 1 minute. Incubate the solutions for 30 minutes in a dark place, measured the absorbance by UV-VIS spectrophotometry (λ 517 nm). A blank solution was obtained by piping 5 ml absolute methanol and 1 ml DPPH solution, left for 30 minutes in a dark place and measuring the absorbance as with the samples.(Hasanah, Maharani, & Munarsih, 2017)

Antibacterial Activity Test

Antibacterial activity test was carried out by the disk diffusion Kirby-Bauer method.(Rohdiana, Arief, & Budiman, 2013) This method was used to see the bacterial growth inhibition response based on the diameter of the clear zone of *E. coli* and *S. typhi*. Antibacterial activity test began by dissolving nutrients with aquades. The media stock and the equipment used were put into a 121°C autoclave for 20 minutes. The next step was making a suspension by taking an ose of pure bacteria and implanting it in a solution of NaCl. Next, applied the suspension to the hard agar using a sterile cotton swab. Blank discs that have been soaked in 5, 10, 15, and 20 grams of brewed Robusta coffee leaves for 20 minutes were placed on hard agar and incubated at 37°C for 24-hour incubation periods. Classification of bacterial growth inhibition was observed by clear zone diameter consists of

4 groups: weak response (diameter ≤ 5 mm), moderate (diameter 5-10 mm), strong (diameter 10-20 mm), and very strong (diameter ≥ 20 mm). (Sadino, Sahidin, & Wahyuni, 2018)

Organoleptic Test

Organoleptic test was carried out by 30 semi-trained panelists of Nutrition Sciences students with variations in weight of 5, 10, 15, and 20 grams of dried coffee leaves brewed with water at 70°C. The assessment of quality attributes of brewed coffee leaves used 4 favorite scales, 1 (dislike very much), 2 (dislike slightly), 3 (like slightly), and 4 (like very much).

Statistic Analysis

Statistical analysis using univariate and bivariate analysis. If the data is normally distributed, use the paired t test, whereas if the data is normally distributed, use the Wilcoxon test. Analysis of antioxidant activity using the One Way Anova statistical test if normally distributed, and using the Kruskal Wallis test if the distribution is not normal. The analysis was performed with a 95% confidence level with p value 0.05 and $\alpha = 0.05$.

Results

Processing Robusta Coffee Leaves with JGTP Method

Based on the appearance and color aspects, brewed coffee leaves with the blanching method had an average value from 7.5 to 8 (1-10) while the steaming method got a value of 4 (1-10). The taste of brewed using the blanching method had an average value from 34 to 35 (20-49) and obtaining a value of 27 for brewed by steaming method. The aroma of brewed and pulp of coffee leaves had a value of 5-5.50 (1-10) for the method of blanching water and steaming. From the aspect of the aroma of coffee grounds pulp obtained a value of 9 for each blanching method and 4 for the steaming method (1-10). Based on organoleptic test, the 75-second blanching method was a processing method that had the highest average score.




Figure 1. Selection of Withering Methods

Nutrient Content

The material used in this study was Robusta coffee leaves which have been dried through the Japanese Style Green Tea Process (JGTP) method as seen in Table 1 .

Table 1. Nutrient Content of Dried Robusta Coffee Leaves

Proximate Component	Mean±SD (%)
Moisture	3.1±0.00
Ash	6.03±0.90
Protein	17.76±2.06
Fat	2.37±0.25
Carbohydrate	70.83±2.83

Note : SD = standard deviation

Bioactive Components (total flavonoids, chlorogenic acid, caffeine)

The results of the analysis showed that the content of bioactive components, as total flavonoids, chlorogenic acid and caffeine in Robusta coffee leaves as seen in Table 2.

Table 2. The Content of Bioactive Components

Treatment	Total flavonoids	Chlorogenic acid	Caffeine
	(mg QE/g)	(mg/g)	(mg/g)
	Mean±SD	Mean±SD	Mean±SD
Dried coffee leaves	0.81 ± 0.02	16.58 ± 0.48	0.456 ± 0.08
5 g of brewed coffee leaves	0.10± 0.01	3.49 ± 1.62	0.194 ± 0.00

Note: 5 g of dried coffee leaves were brewed with 200 mL of water at 70°C for 10 minutes

Total Chlorophyll

The results of chlorophyll analysis showed a significant difference in the groups of brewed coffee leaves ($p = 0.000$). While the results of chlorophyll analysis showed significant differences in the coffee leaves group ($p = 0.014$) and brewed coffee leaves ($p = 0.000$). The results of total chlorophyll analysis showed significant differences in each coffee leaves group ($p = 0.006$) and brewed coffee leaves group ($p = 0.000$). There were significant differences in chlorophyll a, chlorophyll b and total chlorophyll between coffee leaves and brewed coffee leaves ($p = 0,000$) as seen in Table 3.

Table 3. Total Chlorophyll

Samples	Chlorophyll a		Chlorophyll b		Total Chlorophyll	
	Dried coffee leaves	Brewed coffee leaves	Dried coffee leaves	Brewed coffee leaves	Dried coffee leaves	Brewed coffee leaves
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD
5 g	23.66±0.35 ^a	0.74±0.04 ^a	39.98±2.37 ^a	1.35±0.11 ^a	63.62±2.01 ^a	2.08±0.14 ^a
10 g	23.75±0.17 ^a	1.16±0.09 ^b	42.65±0.63 ^{ab}	2.19±0.13 ^b	66.37±0.67 ^{ab}	3.35±0.22 ^b
15 g	23.97±0.39 ^a	1.73±0.31 ^c	43.17±0.61 ^{ab}	3.19±0.44 ^c	67.13±0.99 ^{ab}	4.91±0.75 ^c
20 g	24.66±0.73 ^a	2.12±0.02 ^c	44.76±0.80 ^b	3.91±0.52 ^d	69.40±1.45 ^b	6.01±0.09 ^d
P^x	0.098	0.000*	0.014*	0.000*	0.006*	0.000*

Note: One-way-anova.test. Dried coffee leaves were brewed with 200 mL of water at 70°C for 10 minutes.. *Significant ($p < 0.05$). Numbers followed by superscript letters (a,b,c,d) differ to show significant differences.

Antioxidant Activity

The results of the analysis showed that there were no significant differences in the Robusta coffee leaves group ($p = 0.206$) and brewed ($p = 0.131$) as seen in **Table 4**.

Table 4. Antioxidant Activity with DPPH Method

Treatment	Inhibition (%)		p
	Dried coffee leaves	Brewed coffee leaves	
	Mean \pm SD(%)	Mean \pm SD (%)	
5 gram	92.7 \pm 0.61	84.9 \pm 0.23	0.109 ^p
10 gram	91.8 \pm 0.76	86.1 \pm 0.85	0.019 ^{*q}
15 gram	92.0 \pm 2.00	85.6 \pm 1.20	0.021 ^{*q}
20 gram	90.2 \pm 1.36	83.3 \pm 1.47	0.000 ^{*q}
p	0,206 ^z	0,131 ^y	

Note: Dried coffee leaves were brewed with 200 mL of water at 70°C for 10 minutes. ^{*}Significant ($p < 0.05$). ^pWilcoxon test, ^qPaired t test, ^zOne way Anova, ^yKruskal Wallis test

Antibacterial Activity Test

The results of the antibacterial activity analysis in *Escherichia coli* and *Salmonella typhi* showed significant differences in the groups of brewed coffee leaves ($p = 0.000$) as seen in Table 5.

Table 5. Antibacterial Activity

Samples	<i>Escherichia coli</i>		<i>Salmonella typhi</i>	
	Mean \pm SD	Inhibition	Mean \pm SD	Inhibition
5 g	7.65 \pm 0.10 ^a	Medium	7.47 \pm 0.28 ^a	Medium
10 g	8.47 \pm 0.15 ^b	Medium	8.20 \pm 0.05 ^b	Medium
15 g	9.17 \pm 0.15 ^c	Medium	8.55 \pm 0.26 ^b	Medium
20 g	9.89 \pm 0.26 ^d	Medium	8.70 \pm 0.19 ^c	Medium
p	0,000 [*]		0,000 [*]	

^{*}One Way Anova test. ^{*}Significant ($p < 0.05$). Numbers followed by superscript letters (a,b,c,d) differ to show significant differences.

Acceptability of Brewed Coffee Leaves

There was a significant difference in the color and Taste of brewed coffee leaves ($p = 0.000$) and did not show a significant difference in the aroma of brewed coffee leaves ($p = 0.392$) as seen in **Table 6**.

Table 6. Acceptability of Brewed Coffee Leaves

Treatment	Color		Aroma		Taste	
	Median	Note	Median	Note	Median	Note
5 g	3(1-4) ^a	Like slightly	3(1-4) ^a	Like slightly	3(2-4) ^a	Like slightly
10 g	3(2-4) ^{ab}	Like slightly	3(2-4) ^a	Like slightly	2(1-3) ^a	Dislike slightly
15 g	2,5(1-4) ^{bc}	Dislike slightly	3(1-4) ^a	Like slightly	1(1-4) ^b	Dislike very much
20 g	2(1-4) ^c	Dislike slightly	3(1-4) ^a	Like slightly	1(1-2) ^c	Dislike very much
p	0,000 [*]		0,392		0,000 [*]	

Note: Kruskal Wallis test. ^{*}Significant ($p < 0.05$). Numbers followed by superscript letters (a,b,c,d) differ to show significant differences.

Discussion

In this study, Robusta coffee leaves were processed by the Japanese Style Green Tea Process (JGTP). JGTP method is one of the tea processing without fermentation process. Previous study has explained that the JGTP method in Arabica coffee leaves has higher phytochemicals concentration compared to other tea-making methods.(X.-M. Chen et al., 2018). The blanching process was used at the withering stage. Blanching can deactivate the enzyme polyphenol oxidase (PPO) which can cause oxidation.(Pourcel, Routaboul, Cheynier, Lepiniec, & Debeaujon, 2007)

Oxidation is a degenerative process triggered by the presence of free radicals which causes rancidity and decreasing components in food. The oxidation process causes the loss of phenolic components of food.(X.-M. Chen et al., 2018) The blanching process increases the polyphenol content during extraction.(Nayak, Liu, & Tang, 2015) In addition, the processing of green tea maintain the composition of chlorophyll including chlorophyll a (Chl a) and chlorophyll b (Chl b) that acts as an antibacterial.(Li et al., 2018)

Based on acceptability of brewed coffee leaves, the best acceptance was 5 g of brewed coffee leaves. The more the amount of leaves that are brewed, it will produce a thick yellowish-green color. Coffee leaves products using the JGTP method do not undergo oxidation of polyphenol compounds so that the resulting color was greenish or yellowish-green. Green is the main color determined by the chlorophyll content. Chlorophyll a on the leaves will produce a dark green color and chlorophyll b will produce a yellowish-green color. In addition to the chlorophyll content, flavonol content also plays a role in producing a yellowish color on the coffee leaves by the JGTP method.(Chaturvedula & Prakash, 2011)

Based on the aroma described by panelists, the aroma of brewed coffee leaves such as the aroma of fresh leaves. The more the amount of leaves in brewed, the smell will be thicker and increasingly disliked by panelists. The aroma of the leaves is caused by the high compound of catechins.(Ho, Zheng, & Li, 2015) The taste of brewed coffee leaves was bitter. The bitter taste is influenced by the presence of chlorogenic acid and caffeine.(Belay, 2011; Gokul G & Lakshmanan, 2016) Caffeine is a natural compound in food and drinks that contributes to affect the bitter taste. Some studies show that the main taste of caffeine is bitter. Caffeine generally amounts to 2-5% of the dry weight of tea. Caffeine will cause a bitter taste on the tongue depending on the level of concentration.(Keast & Roper, 2007; Zou, Xiao, Wang, & Zhang, 2018) The study explained that the bitter taste began to emerge when the caffeine content was 75-155 mg/L. Chlorogenic acid also plays a role in acidity and bitterness.(Belay, 2011) The moisture and ash content in dried Robusta coffee leaves processed by the JGTP method met the Indonesian National Standard (SNI) green tea requirements, where the moisture of green tea is set to a maximum of 8% and ash content of 4-8%.(SNI, 2016) The moisture in accordance with SNI can prevents the growth of bacteria and fungi, and prevent oxidation process. Protein content in dried Robusta coffee leaves with JGTP method included in the range of green tea, where protein content in green tea about 15-20%. Protein has the potential as an additional antioxidant in food because it can inhibit fat oxidation through several pathways including inactivation of Reactive Oxygen Species (ROS) through antioxidant enzymes (superoxide dismutase (SOD) and catalase), scavenge free radicals, chelating prooxidative transition metals, reducing hydroperoxide, and reducing hydroperoxides, and enzymatically removes specific oxidants.(Elias, Kellerby, & Decker, 2008) Protein also acts as an antimicrobial. Previous study explained that protein extracted from lemon, orange and grapefruit seeds could inhibit the growth of some pathogenic bacteria.(Karabiber, Yilmaz, & Zorba, 2018)

The fat content in Robusta coffee leaves processed by the JGTP method was lower than in Arabica coffee leaves with the same method. Based on previous research, Arabica coffee leaves have higher fat content about 4.5-12.5%. Meanwhile, the carbohydrate content in Robusta coffee leaves was higher than in Arabica coffee leaves. In previous study, carbohydrate content of Arabica coffee leaves was about 51.99-63.93%.(Woldesenbet, 2005)

In this study, Robusta coffee leaves processed by JGTP method have bioactive components of flavonoids, chlorogenic acid, caffeine, and chlorophyll. Previous study has shown that coffee leaves contain flavonoids such as quercetin, rutin, and kaempferol.(Martins, Araújo, Tohge, Fernie, & DaMatta, 2014) Flavonoids can act as antioxidants to scavenge free radicals, prevent the formation of reactive oxygen species (ROS), and increase endogenous antioxidants (superoxide dismutase, catalase, and glutathione peroxidase). The mechanism of flavonoids in preventing damage by free radicals is by giving hydrogen atoms to free radicals to form stable flavonoxyl radicals (flavonoids (O•)). In addition, flavonoids can protect the phospholipid membrane by giving one of the hydrogen ions to the radical lipid peroxy, so that radical reactions can be stopped.(Hamid, Usman, Aiyelaagbe, & Ameen, 2010; Irina & Mohamed, 2012) Flavonoids can also function as antibacterial by inhibiting nucleic acid synthesis, inhibiting energy metabolism, changes in cytoplasmic membrane function, reducing cell adhesion and biofilm formation and causing damage to the cytoplasmic membrane.(Farhadi et al., 2018)

Chlorogenic acid is an ester formed from hydroxycinnamic acid (caffeic acid, ferulic acid, p-kumaric acid) and quinic acid and into the phenolic acid group.(Liang & Kitts, 2016) Chlorogenic acid can chelate transition metals like Fe²⁺ to scavenge free radicals and break radical chains. Chlorogenic acid scavenge DPPH radicals, superoxide anions, hydroxyl radicals, and protect DNA from damage caused by oxidative stress.(Azam, Hadi, Khan, & Hadi, 2003) Chlorogenic acid also function as antibacterial. Chlorogenic acid is a phenolic bioactive compound that can

cause physiological changes in bacterial cell membranes.(Kabir et al., 2014) Chlorogenic acid causes cell membrane rupture and release of the content inside the cellular. Chlorogenic acid changes in bacterial cell membrane permeability that will cause damage to bacterial nucleotides.(Lou et al., 2011)

Robusta coffee leaves with processed by JGTP method also contained caffeine. Caffeine prevents lipid oxidation by inhibiting the production of free radicals including hydroxyl radicals, peroxy radicals, and singlet oxygen.(Azam et al., 2003) Caffeine also acts as antibacterial agent. It can inhibit protein and DNA synthesis by inhibiting the merging of adenine and thymidine. Besides, caffeine also increases genotoxicity after DNA damage.(Pruthviraj et al., 2011)

The amount of chlorophyll in brewed Robusta coffee leaves was very low compared to the amount of chlorophyll in tea leaves. Chlorophyll acts as antioxidants, where it can prevent the Fenton reaction, by chelating Fe(II). Chlorophyll directly scavenges the hydroxyl free radicals generated from H₂O₂.(Hsu, Chao, Hu, & Yang, 2013) It can also act as antibacterial by increasing membrane permeability in bacteria.(Kustov et al., 2018) This will cause changes in membrane structure and accelerate intracellular leakage thereby destroying membrane integrity which will facilitate the entry of antibacterial.(Wu et al., 2016)

The results showed that Robusta coffee leaves with the JGTP method had a DPPH radical inhibition. Based on the results of this analysis, Robusta coffee leaves and brewed coffee leaves have high antioxidant activity, being able to capture free radicals by 50%. The results of this study are in line with research on green tea, where green tea has a DPPH radical inhibition more than 50%.

The results of antibacterial of brewed coffee leaves showed the inhibitory power of 5 g, 10 g, 15 g, and 20 g brewed Robusta coffee leaves against *Escherichia coli* and *Salmonella typhi* bacteria. *Escherichia coli* and *Salmonella typhi* belong to gram-negative bacteria. The greater the amount of leaves in the brewed process, the greater the diameter of the inhibition zone formed. The effectiveness of antibacterial substances will be influenced by the concentration of these substances. An increase in the concentration of the substance causes an increase in the content of active compounds such as flavonoids, caffeine and chlorogenic acid which functions as an antibacterial so that its ability to inhibit bacterial growth increases.(Mufti, Bahar, & Arisanti, 2017; Muslim & Dephinto, 2017) Previous study conducted on CTC brewed black tea with a comparison of tea weight and water volume (w/v) 1:50, 2:50, and 3:30 resulted in an area of 29.89 successive barriers; 33.64 and 45.02 mm².(Rohdiana et al., 2013)

Conclusion

Robusta coffee leaves in this research were processed by JGTP method with 75-second blanching. The best organoleptic based on color, aroma, and taste was 5 g of dried Robusta coffee leaves. Brewed Robusta coffee leaves processed by the Japanese Style Green tea Process (JGTP) have contain flavonoids 0.10 mg QE/g, chlorogenic acid 3.49 mg/g, caffeine 0.194 mg/g and high antibacterial activity in 20 g of brewed Robusta coffee leaves.

Conflict of Interest

No conflict of interest.

Acknowledgments

This research was supported by Research and Development (RPP) from the Faculty of Medicine, Universitas Diponegoro.

References

- Analysis., O. M. of. (2005). *Official Methods of Analysis*. (18th ed.). AOAC International: USA.
- Azam, S., Hadi, N., Khan, N. U., & Hadi, S. M. (2003). Antioxidant and Prooxidant Properties of Caffeine , Theobromine and Xanthine. *Med Sci Monit.*, 9(9), 325–331.
- Belay, A. (2011). Some biochemical compounds in coffee beans and methods developed for their analysis. *International Journal of the Physical Sciences*, 6(28), 6373–6378.
- Campa, C., Urban, L., Mondlot, L., Fabre, D., Roques, S., Lizzi, Y., ... Etienne, H. (2017). Juvenile Coffee Leaves Acclimated to Low Light Are Unable to Cope with a Moderate Light Increase. *Front Plant Sci.*, 8(1126), 1–16.

- Chacko, S. M., Thambi, P. T., Kuttan, R., & Nishigaki, I. (2010). Beneficial effects of green tea : A literature review. *Chin Med*, 5(15), 1–9.
- Chaturvedula, V. S. P., & Prakash, I. (2011). The aroma, taste, color and bioactive constituents of tea. *Journal of Medicinal Plants Research*, 5(11), 2110–2124.
- Chen, X. (2019). A Review on Coffee Leaves : Phytochemicals, Bioactivities and Applications. *Crit Rev Food Sci Nutr*, 59(6), 1008–1025.
- Chen, X.-M., Ma, Z., & Kitts, D. D. (2018). Effects of processing method and age of leaves on phytochemical profiles and bioactivity of coffee leaves. *Food Chemistry*, 249, 143–153.
- Elias, R. J., Kellerby, S. S., & Decker, E. A. (2008). Antioxidant Activity of Proteins and Peptides. *Critical Reviews in Food Science and Nutrition*, 48(5), 430–441.
- Farhadi, F., Khameneh, B., Iranshahi, M., & Iranshahy, M. (2018). Antibacterial activity of flavonoids and their structure – activity relationship : An update review. *Phyther Res*, 1–28.
- Gokul G, & Lakshmanan, R. (2016). Effect Of Chlorhexidie Mouthwash On Taste Alteration. *Asian J Pharm Clin Res*, 9(1), 102–104.
- Hamid, A. A., Usman, L. A., Aiyelaagbe, O., & Ameen, M. O. (2010). Antioxidants : Its Medicinal and Pharmacological Applications. *African J Pure Appl Chem*, 4(8), 142–151.
- Hasanah, M., Maharani, B., & Munarsih, E. (2017). Daya Antioksidan Ekstrak Dan Fraksi Daun Kopi Robusta (*Coffea robusta*) terhadap Pereaksi DPPH (2,2-difenil-1-pikrilhidrazil) (Antioxidant of Extract and Fraction *Coffea robusta* Leaves with Diphenylpicrylhydrazyl (DPPH) Method. *Indonesian Journal of Pharmaceutical Science and Technology*, 4(2), 42–49.
- Ho, C.-T., Zheng, X., & Li, S. (2015). Tea Aroma Formation. *Food Science and Human Wellness*, 4(1), 9–27.
- Hsu, C.-Y., Chao, P.-Y., Hu, S.-P., & Yang, C.-M. (2013). The Antioxidant and Free Radical Scavenging Activities of Chlorophylls and Pheophytins. *Food and Nutrition Sciences*, 4, 1–18.
- Irina, I., & Mohamed, G. (2012). Biological Activities and Effects of Food Processing on Flavonoids as Phenolic Antioxidants. In *Advances in Applied Biotechnology*. (pp. 112–114). Croatia.
- Kabir, F., Katayama, S., Tanji, N., & Nakamura, S. (2014). Antimicrobial Effects of Chlorogenic Acid and Related Compounds. *Journal of the Korean Society for Applied Biological Chemistry*, 57(3), 359–365.
- Kanisius, A. A. (1988). *Budidaya Tanaman Kopi* (16th ed.). Yogyakarta: Kanisius.
- Karabiber, E., Yilmaz, E., & Zorba, N. N. D. (2018). antimicrobial and functional properties of the proteins extracted from lemon, orange and grapefruit seeds press meals. *Quality Assurance and Safety of Crops & Foods*, 10(2), 1–10.
- Keast, R. S. J., & Roper, J. (2007). A Complex Relationship Among Chemical Concentration, Detection Threshold, and Suprathreshold Intensity of Bitter Compounds. *Chem Senses*, 32(3), 245–253.
- Khotimah, K. (2014). Karakteristik Kimia Kopi Kawa dari Berbagai Umur Helai daun Kopi yang Diproses dengan Metode Berbeda. *J Teknol Pertan*, 9(1), 40–45.
- Kustov, A. V., Belykh, D. V., Smirnova, N. L., Venediktov, E. A., Kudayarova, T. V., Kruchin, S. O., ... Berezin, D. B. (2018). Synthesis and investigation of water-soluble chlorophyll pigments for antimicrobial photodynamic therapy. *Dyes and Pigments*, 149, 553–559.
- Ky, C.-L., Louarn, J., Dussert, S., Guyot, B., Hamon, S., & Noiro, M. (2001). Caffeine, trigonelline, chlorogenic acids and sucrose diversity in wild *Coffea arabica* L. and *C. canephora* P. accessions. *Food Chemistry*, 75(2), 223–230.
- Li, X., Zhou, R., Xu, K., Xu, J., Jin, J., Fang, H., & He, Y. (2018). Rapid Determination of Chlorophyll and Pheophytin in Green Tea Using Fourier Transform. *Molecules*, 21, 1–13.
- Liang, N., & Kitts, D. D. (2016). Role of Chlorogenic Acids in Controlling Oxidative and Inflammatory Stress Conditions. *Nutrients*, 6, 16.
- Lou, Z., Wang, H., Zhu, S., Ma, C., & Wang, Z. (2011). Antibacterial Activity and Mechanism of Action of Chlorogenic Acid. *J Food Sci.*, 76(6), 398–403.
- Manach, C., Scalbert, A., Morand, C., Rémésy, C., & Jiménez, L. (2004). Polyphenols: Food Sources and Bioavailability. *The American Journal of Clinical Nutrition*, 79(5), 727–747.
- Martins, S. C. V., Araújo, W. L., Tohge, T., Fernie, A. R., & DaMatta, F. M. (2014). n High-Light-Acclimated Coffee Plants the Metabolic Machinery Is Adjusted to Avoid Oxidative Stress Rather than to Benefit from Extra Light Enhancement in Photosynthetic Yield. *PLoS ONE*, 9(4), 1–11.
- Mufti, N., Bahar, E., & Arisanti, D. (2017). Uji Daya Hambat Ekstrak Daun Sawo terhadap Bakteri *Escherichia coli* secara In Vitro (Inhibition Test of Inhibition Test of Sawo Leaf Extract Against *Escherichia coli* Bacteria in Vitro). *Jurnal Kesehatan Andalas*, 6(2), 289–294.
- Muslim, Z., & Dephinto, Y. (2017). Perbandingan Efektivitas Antimikroba Ekstrak Daun Kopi Robusta (*Coffea Canephora*) Dengan Variasi Pengeringan Terhadap *Escherichia coli* (Comparison of Antimicrobial Effectiveness of Robusta (*Coffea Canephora*) Leaf Extract with Drying Variations on *Escherichia coli*). *J Sains Dan Teknol Farm*, 19, 86–88.
- Naegel, E. (2016a). Determination of Caffeine in Coffee Products According to DIN 20481. *Agil Technol*, 1–6.

- Naegele, E. (2016b). Determination of Chlorogenic Acid in Coffee Products According to DIN 10767. *Agil Technol*, 1–8.
- Nayak, B., Liu, R. H., & Tang, J. (2015). Effect of Processing on Phenolic Antioxidants of Effect of Processing on Phenolic Antioxidants of Fruits, Vegetables, and Grains — A Review. *Crit Rev Food Sci Nutr.*, 55(7), 887–918.
- Nayeem, N., Denny, G., & Mehta, S. K. (2011). Comparative phytochemical analysis, antimicrobial and anti oxidant activity of the methanolic extracts of the leaves of *Coffea Arabica* and *Coffea Robusta*. *Der Pharmacia Lettre*, 3(1), 292–297.
- Permadi, M. R., Oktafa, H., & Agustianto, K. (2018). Perancangan Sistem Uji Sensoris Makanan dengan Pengujian Peference Test (Hedonik dan Mutu Hedonik), Studi Kasus Roti Tawar Menggunakan Algoritma Radial Basis Function Network (Design of Food Sensory Test System with Peference Test (Hedonic and Hedonic Qua. *Jurnal Mikrotik*, 8(1), 29–42.
- Pourcel, L., Routaboul, J.-M., Cheynier, V., Lepiniec, L., & Debeaujon, I. (2007). Flavonoid Oxidation in Plants : from Biochemical Properties to Physiological Functions. *Trends Plant Sci.*, 12(1), 29–36.
- Prawira-Atmaja, M. I., Shabri, Khomaini, H. S., Maulana, H., Harianto, S., & Rohdiana, D. (2018). Changes in chlorophyll and polyphenols content in *Camellia sinensis* var. *sinensis* at different stage of leaf maturity. *Earth Environmental Sci*, 131, 1–7.
- Pruthviraj, P., Shital, K., Suchita, B., & Shilpa, K. (2011). Evaluation of Antibacterial Activity of Caffeine. *Int J Res Ayurveda Pharm*, 2(4), 1354–1357.
- Rhayanne T. M. Ramos, I. C. F. B., Ferreira, M. R. A., & Soares, L. A. L. (2017). Spectrophotometric Quantification of Flavonoids in Herbal Material, Crude Extract, and Fractions from Leaves of *Eugenia uniflora* Linn. *Pharmacognosy Res.*, 9(3), 253–260.
- Rohdiana, D., Arief, D. Z., & Budiman, A. (2013). Aktivitas penghambatan pertumbuhan bakteri *Escherichia coli* oleh berbagai jenis teh dan seduhannya. (Inhibitory activity of *Escherichia coli* by type of teas and its liquors). *Jurnal Penelitian Teh Dan Kina*, 16(1), 37–44.
- Sadino, A., Sahidin, I., & Wahyuni, W. (2018). Antibacterial Activity of *Polygonum pulchrum* Blume Ethanol Extract on *Staphylococcus aureus* and *Escherichia coli*. *Pharmacol Clin Pharm Res.*, 3, 26–32.
- Silva, L. A. L. da, Pezzini, B. R., & Soares, L. (2015). Spectrophotometric Determination of The Total Flavonoid Content in *Ocimum basilicum* L.(Lamiaceae) Leaves. *Pharmacogn Mag*, 11(41), 96–101.
- SNI. (2016). SNI 3945:2016 Teh Hijau (Green Tea). Jakarta.
- Woldesenebet, A. (2005). Nutritional Composition, Phytochemical Screening, Processing Methods and Sensory Attributes of A Brew Made From Infusions of Matured Leaves of *Arabica* Coffee Tree Consumed In Sidama, Kambata and Harar Communities, Ethiopia. Addus Ababa University.
- Wu, Y., Bai, J., Zhong, K., Huang, Y., Qi, H., Jiang, Y., & Gao, H. (2016). Antibacterial Activity and Membrane-Disruptive Mechanism of 3-p-trans Coumaroyl-2-hydroxyquinic Acid, a Novel Phenolic Compound from Pine Needles of *Cedrus deodara*, against *Staphylococcus aureus*. *Molecules*, 21(8), 1–12.
- Zou, G., Xiao, Y., Wang, M., & Zhang, H. (2018). Detection Of Bitterness And Astringency Of Green Tea With Different Taste By Electronic Nose And Tongue. *PLoS ONE*, 13(12), 1–10.

Hasil Review

Inbox 187

- Starred
- Snoozed
- Important
- Sent
- Drafts** 54
- Categories
- More

Labels +

- [imap]/Archive
- [imap]/Drafts
- Notes

ICHWB Submission and Review System <conferences-noreply@ums.ac.id>
to me

Fri, 22/11/2019 - 9:27pm

This is to inform you that your submission has been changed by the track-director.

The status of your submitted paper no 14787 is Revision Required

We thank you for submitting your abstract titled "Bioactive components and antibacterial activity study in Robusta coffee leaves (coffea canephora)" for presenting and publishing in International Conference on Health and Wellbeing, going to be held at Muhammadiyah Surakarta University on 3-4 December 2019.

Your abstract is accepted for presentation, and reviewer's comments for publication in Scopus indexed journal will be sent later.

Revision Required

=====
Detail information of your paper and LETTER OF ACCEPTANCE can be found in your author account (login required).
For registration your presentation attendance, please visit
<http://jseth.ums.ac.id/2019/reg19>
=====
Please login to your author account

Fri, Nov 22, 2019, 9:27 PM

☆ Reply

Hasil Review OJS

The screenshot displays an email client interface. On the left, there is a sidebar with folders: 'Inbox' (187), 'Starred', 'Snoozed', 'Important', 'Sent', 'Drafts' (54), 'Categories', and 'More'. Below these are 'Labels' including '[imap]/Archive', '[imap]/Drafts', and 'Notes'. The main content area shows an email from 'ICHWB Submission and Review System' (conferences-noreply@ums.ac.id) dated 'Wed, Jan 15, 2020, 1:09 PM'. The subject is 'ICHWB2019 status your paper submission 14787 - Revision Required'. The email body states: 'The status of your submitted paper no 14787 is Revision Required'. It then provides feedback: 'The internal reviewers have assessed your submission and feel that your manuscript "Bioactive components and antibacterial activity study in Robusta coffee leaves (coffea canephora)" has potential for publication in journal indexed by Scopus with a minor revision. Please revise your manuscript as the following comments and resubmit the final revision to ICHWB website by Wednesday, 22 th January 2020. A submission after the deadline will be presume that you have withdrawn your submission. We look forward to hearing from you.' The email concludes with 'Revision Required' and a note that 'Detail information of your paper and LETTER OF ACCEPTANCE can be found in'.

Hasil Perbaikan Artikel

The screenshot displays an email client interface. On the left, a sidebar lists folders: 'Inbox' (187), 'Starred', 'Snoozed', 'Important', 'Sent', 'Drafts' (54), and 'Categories' (More). Below these are 'Labels' including '[imap]/Archive', '[imap]/Drafts', and 'Notes'. The main content area shows an email from 'Vita Almira <vitagalmira@gmail.com>' to 'lrr151, me' dated 'Wed, Mar 4, 2020, 11:42 AM'. The subject is 'Paper Gemala Anjani (revised)'. The email body contains the following text:

Dear Bu Luluk,

Berikut terlampir revisi paper kami yang berjudul "**Bioactive components and antibacterial activity in Robusta coffee leaves (coffea canephora)**" dengan Gemala Anjani, Nurmasari Widyastuti, Zulfatul Masruroh, Rizka Ayu Dwi Yuliana, Vita Gustin Almira, A.Fahmy Arif Tsani, Choirun Nissa, M Iqbal Prawira-Atmaja sebagai authors.

Terima kasih.
Best Regards,

Vita Gustin Almira, S.Gz

The interface also includes a language selector (Indonesian to English), a 'Translate message' link, and a 'Turn off for: Indonesian' option. On the right side, there are icons for printing, sharing, and a plus sign for additional actions.

Bioactive components and antibacterial activity in Robusta coffee leaves (*cofea canephora*)

Gemala Anjani^{1,2,φ,*}, Nurmasari Widyastuti^{1,2,φ}, Zulfatul Masruroh¹, Rizka Ayu Dwi Yuliana¹, Vita Gustin Almira¹, A.Fahmy Arif Tsani^{1,2}, Choirun Nissa^{1,2}, M Iqbal Prawira-Atmaja³

¹ Department of Nutrition Science, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

² Center of Nutrition Research (CENURE), Universitas Diponegoro, Semarang, Indonesia

³ Research Institute for Tea and Cinchona, Bandung, Indonesia

Abstract

Purpose: To analyze bioactive components and antibacterial activity in brewed Robusta coffee leaves that processed by *Japanese Style Green Tea Process* (JGTP).

Methodology: This was an experimental research with the sample of Robusta coffee leaves and brewed Robusta coffee leaves. The coffee leaves were brewed in water at 70°C during 10 minutes. The analysis of nutrition contents by proximate method, total flavonoids by calorimeter method, chlorogenic acid and caffeine by HPLC method, antioxidant activity by DPPH method, total chlorophyll using the arnon formula and antibacterial activity by disk diffusion method in 5, 10, 15, and 20 g.

Results: There were significant differences in color, taste, and acceptability of brewed Robusta coffee leaves. The best acceptance of brewed Robusta coffee leaves was 5 g of dried coffee leaves. Robusta coffee leaves contain 3,1% water, 6,03% ash, 17,76% protein, 2,37% fat, and 70,83% carbohydrate. The content of flavonoids, chlorogenic acid and caffeine in Robusta coffee leaves were 0.81 mgQE/g, 16.92 mg/g, 0.46 mg while in the brewed were 0.10 mgQE/g, 3.49 mg/g, 0.19 mg/g, respectively. The inhibition of free radical in Robusta coffee leaves were 90,2-92,7%, while in the brewed were 83,3-86,1%. Total chlorophyll of dried leaves were 63,62-69,39 mg/L and the brewed were 2,08-6,01 mg/L. Antibacterial activity *Escherichia coli* 7,65-9,89 mm and *Salmonella tiphy* 7,47-8,70 mm inhibition zone.

Applications/Originality/Value: The role of various bioactive components in Robusta coffee leaves on antioxidant and antibacterial activity.

^φ Both authors contributed equally to this manuscript

* Corresponding author: gemaanjani@gmail.com

Introduction

Coffee plant (*Coffea sp.*) is included in the family *Rubiaceae*. In general, there are three types of coffee, including Arabica coffee (*Coffea arabica*), Robusta coffee (*Coffea canephora*), and Liberica coffee (*Coffea liberica*). The most widely consumed coffee in Indonesia is Robusta coffee. Robusta coffee can be easily grown at lower altitude of less than 1000 meters above sea level and it resilient against disease and pests.(Kanisius, 1988)

The results of research on Arabica coffee leaves have nutritional content based on proximate components. Proximate components of Arabica coffee leaves from the Ethiopian region include 5.5-7.8% moisture, 8.8-12.4% ash content, 14.4-19.0% protein, 4.5-12.5% fat, and 51.0-63.9% carbohydrate.(Woldesenebet, 2005) Robusta coffee leaves contain phytochemicals that vary depending on leaf development.(X.-M. Chen, Ma, & Kitts, 2018) The younger the coffee leaves, the higher the chlorogenic acid content even 10 times higher than the old coffee leaves.(Campa et al., 2017) The content of chlorogenic acid in Robusta coffee is higher compared to Arabica coffee, respectively 11.3% and 4.1%.(Ky et al., 2001) The caffeine content in young Robusta coffee leaves (0.29-0.5%) is lower than the caffeine content in its fruit (1.6-2.4%).(Khotimah, 2014)

Robusta coffee leaves also contain phenolic acid and total flavonoids, respectively 27.04 µg/g and 10.90 µg/g higher than Arabica coffee leaves respectively 21.80 µg/g and 8.08 µg/g.(Nayeem, Denny, & Mehta, 2011) Flavonoids as natural antioxidants can prevent the occurrence of diseases, such as cancer, metabolic syndrome, diabetes mellitus, and atherosclerosis which associated to oxidative stress.(X. Chen, 2019; Manach, Scalbert, Morand, Rémésy, & Jiménez, 2004)

The content of caffeine, flavonoids, and chlorogenic acid of coffee leaves can also function as the prevention of diarrhea. Caffeine can inhibit protein and DNA synthesis by inhibiting the merging of adenine and thymidine. Caffeine also increases genotoxicity after DNA damage.(Pruthviraj, Shital, Suchita, & Shilpa, 2011) Flavonoids as an antibacterial by inhibiting nucleic acid synthesis, inhibiting energy metabolism, and changes in cytoplasmic membrane function.(Farhadi, Khameneh, Iranshahi, & Iranshahy, 2018) In addition to flavonoids, there are chlorogenic acid contents that function as antibacterial. Chlorogenic acid is a phenolic bioactive compound causes physiological changes in bacterial cell membranes that encourage damage to bacterial nucleotides.(Kabir, Katayama, Tanji, & Nakamura, 2014; Lou, Wang, Zhu, Ma, & Wang, 2011)

The processing of coffee leaves to obtain optimal antibacterial is done by minimizing chemical and enzymatic reactions to prevent damage in phytochemical content. The process can be obtained using green tea processing.(X.-M. Chen et al., 2018) The processing of green tea is able to produce phenolic components 6 times higher than black tea. Green tea processing aims to maintain the content of natural polyphenols which have beneficial effects on health. (Chacko, Thambi, Kuttan, & Nishigaki, 2010). Therefore, the aim of this research was to examine bioactive components, including caffeine, chlorogenic acid, total flavonoids, antioxidant activity, total chlorophyll, and nutritional contents and antibacterial activity in Robusta coffee leaves by green tea processing.

Materials and Methods

The samples used *Robusta canephora* Pierre ex Froehner from Gambung village. Coffee leaves were processed using the Japanese Style Green Tea Process (JGTP) method at the Tea Quality Processing and Testing Laboratory, Tea and Quinine Research Center (PPTK) Gambung, Bandung Regency, West Java, Indonesia. Preliminary research was conducted to obtain the JGTP procedure at the withering stage, including 2 minutes steaming method and 45 s, 60 s, and 75 s of blanching. The best withering method was chosen based on the results of the best organoleptic tests conducted by trained panelists. The best withering result that will be used in the processing of coffee leaves was the blanching method for 75 s.

This study was an experimental study with one factor completely randomized design (CRD) with 2 samples, dried Robusta coffee leaves and brewed Robusta coffee leaves. Dried Robusta coffee leaves were brewed with 200 ml of water in 70°C for 10 minutes. Samples were analyzed in duplicate with three replications.

Processing Robusta Coffee Leaves with JGTP Method

JGTP method was carried out through 5 steps, picking; blanching; cooling, crushing, and drying. Coffee leaves were picked from 2nd, 3rd, and 4th leaf of each branch of coffee plants, sorted, then carried out the blanching process for 75 s. Leaves were soaked in water and put in seducer for ± 15 minutes while separating the leaves from its midrib. The next process was crushing the leaves three times using a crushing-tearing-curling machine (CTC) (CNC, Sri Lanka). The last process was the drying process. The drying process was 4-5 hours long by a rack drier machine (CNC, Sri Lanka) at 80°C.

Proximate Analysis

Proximate analyses using dried Robusta coffee leaves samples processed by JGTP to analyze the moisture, ash, fat, protein, and carbohydrate contents with AOAC 2005 method.(Analysis., 2005)

Total Flavonoid Analysis

5 grams of dried Robusta coffee leaves in 200 ml of methanol were homogenized with a magnetic stirrer for \pm 30 minutes. The solution was taken 60 ml for centrifugation, then coffee leaves produced supernatant in methanol. Meanwhile, 5 grams of dried Robusta coffee leaves were brewed in 200 ml of hot water (70°C) for 10 minutes and stirred. After being dissolved in water, 60 ml was taken to be centrifuged, coffee leaves supernatant will be produced in brewed water. Afterwards, two samples were analyzed.

Total flavonoid analysis using aluminum chloride (AlCl_3) calorimetry method with absorbance measurement using UV-VIS spectrophotometer. 1 ml of sample solution was mixed with 9 ml of methanol. Pipette 2.8 ml sample solution; 0.4 ml AlCl_3 5%; 6.8 ml of 5% acetic acid. Incubation for 30 minutes, measure the absorbance (λ 428 nm). Calibration curves used the standard quercetin in methanol (0.005 mg / ml) with varying concentrations of 0 $\mu\text{g/ml}$, 5 $\mu\text{g/ml}$, 10 $\mu\text{g/ml}$, 20 $\mu\text{g/ml}$, 40 $\mu\text{g/ml}$, and 60 $\mu\text{g/ml}$. Standard was carried out using the same procedures as samples. (Rhayanne T. M. Ramos, Ferreira, & Soares, 2017; Silva, Pezzini, & Soares, 2015)

Analysis of Chlorogenic Acid and Caffeine

Analysis of chlorogenic acid using 20 g of dried coffee leaves extracted with 150 mL methanol:water (50:50 v/v) in soxhlet. After 4-5 cycles, soxhletation was stopped and cooled down at room temperature. The extract was transferred into a 200 mL flask and the volume was adjusted to 200 mL with a mixture of methanol-water. Meanwhile, caffeine analysis using 10-20 g of coffee leaves were added to 5 g of MgO and put into 200 mL of water. Furthermore, the samples were heated in a water bath at 90°C and sterilized for 20 minutes. Then the extracts were filtered with 0.45uL, 25 mm cellulose filter cartridge. The filtrates were injected into the HPLC.(Naegele, 2016a, 2016b)

Total Chlorophyll Test

The total chlorophyll test was carried out by 5, 10, 15 and 20 grams of brewed Robusta coffee leaves compared to the samples dissolved with 80% acetone. Total chlorophyll test was conducted using 663 nm and 645 nm wave spectrophotometry. Chlorophyll levels were calculated using the Arnon formula:(Prawira-Atmaja et al., 2018)

Chlorophyll a = $(12.7 \times A_{663}) - (2.69 \times A_{645})$

Chlorophyll b = $(22.9 \times A_{645}) - (4.68 \times A_{663})$

Total chlorophyll = $(20.2 \times A_{645}) + (8.02 \times A_{663})$

Analysis of Antioxidant Activity

Sample preparation was performed as a total flavonoid test. Antioxidant activity was analyzed using the DPPH method (1,1-diphenyl-2-picrylhydrazyl). A DPPH 0.4 mM solution was prepared by 0.00394 g of DPPH powder and dissolved in absolute methanol to 25 ml. The test was done by piping 0.1 ml of samples and sufficient volume with absolute methanol into a 5 ml measuring flask, then vortex. Measurement of antioxidant activity was done by adding 1 ml of DPPH 0.4 mM solution, then vortex for \pm 1 minute. Incubate the solutions for 30 minutes in a dark place, measured the absorbance by UV-VIS spectrophotometry (λ 517 nm). A blank solution was obtained by piping 5 ml absolute methanol and 1 ml DPPH solution, left for 30 minutes in a dark place and measuring the absorbance as with the samples.(Hasanah, Maharani, & Munarsih, 2017)

Antibacterial Activity Test

Antibacterial activity test was carried out by the disk diffusion Kirby-Bauer method.(Rohdiana, Arief, & Budiman, 2013) This method was used to see the bacterial growth inhibition response based on the diameter of the clear zone of *E. coli* and *S. typhi*. Antibacterial activity test began by dissolving nutrients with aquades. The media stock and the equipment used were put into a 121°C autoclave for 20 minutes. The next step was making a suspension by taking an ose of pure bacteria and implanting it in a solution of NaCl. Next, applied the suspension to the hard agar using a sterile cotton swab. Blank discs that have been soaked in 5, 10, 15, and 20 grams of brewed Robusta coffee leaves for 20 minutes were placed on hard agar and incubated at 37°C for 24-hour

incubation periods. Classification of bacterial growth inhibition was observed by clear zone diameter consists of 4 groups: weak response (diameter ≤ 5 mm), moderate (diameter 5-10 mm), strong (diameter 10-20 mm), and very strong (diameter ≥ 20 mm). (Sadino, Sahidin, & Wahyuni, 2018)

Organoleptic Test

Organoleptic test was carried out by 30 semi-trained panelists of Nutrition Sciences students with variations in weight of 5, 10, 15, and 20 grams of dried coffee leaves brewed with water at 70°C. The assessment of quality attributes of brewed coffee leaves used 4 favorite scales, 1 (dislike very much), 2 (dislike slightly), 3 (like slightly), and 4 (like very much).

Statistic Analysis

Statistical analysis using univariate and bivariate analysis. If the data is normally distributed, use the paired t test, whereas if the data is normally distributed, use the Wilcoxon test. Analysis of antioxidant activity using the One Way Anova statistical test if normally distributed, and using the Kruskal Wallis test if the distribution is not normal. The analysis was performed with a 95% confidence level with p value 0.05 and $\alpha = 0.05$.

Results

Processing Robusta Coffee Leaves with JGTP Method

Based on the appearance and color aspects, brewed coffee leaves with the blanching method had an average value from 7.5 to 8 (1-10) while the steaming method got a value of 4 (1-10) as seen in Figure 1. The taste of brewed using the blanching method had an average value from 34 to 35 (20-49) and obtaining a value of 27 for brewed by steaming method. The aroma of brewed and pulp of coffee leaves had a value of 5-5.50 (1-10) for the method of blanching water and steaming. From the aspect of the aroma of coffee grounds pulp obtained a value of 9 for each blanching method and 4 for the steaming method (1-10). Based on organoleptic test, the 75-second blanching method was a processing method that had the highest average score.




Figure 2. Selection of Withering Methods

Nutrient Content

The material used in this study was Robusta coffee leaves which have been dried through the Japanese Style Green Tea Process (JGTP) method as seen in Table 1 .

Table 7. Nutrient Content of Dried Robusta Coffee Leaves

Proximate Component	Mean±SD (%)
Moisture	3.1±0.00
Ash	6.03±0.90
Protein	17.76±2.06
Fat	2.37±0.25
Carbohydrate	70.83±2.83

Note : SD = standard deviation

Bioactive Components (total flavonoids, chlorogenic acid, caffeine)

The results of the analysis showed that the content of bioactive components, as total flavonoids, chlorogenic acid and caffeine in Robusta coffee leaves as seen in Table 2.

Table 8. The Content of Bioactive Components

Treatment	Total flavonoids (mg QE/g)	Chlorogenic acid (mg/g)	Caffeine (mg/g)
	Mean±SD	Mean±SD	Mean±SD
Dried coffee leaves	0.81 ± 0.02	16.58 ± 0.48	0.456 ± 0.08
5 g of brewed coffee leaves	0.10± 0.01	3.49 ± 1.62	0.194 ± 0.00

Note: 5 g of dried coffee leaves were brewed with 200 mL of water at 70°C for 10 minutes

Total Chlorophyll

The results of chlorophyll analysis showed a significant difference in the groups of brewed coffee leaves ($p = 0.000$). While the results of chlorophyll analysis showed significant differences in the coffee leaves group ($p = 0.014$) and brewed coffee leaves ($p = 0.000$). The results of total chlorophyll analysis showed significant differences in each coffee leaves group ($p = 0.006$) and brewed coffee leaves group ($p = 0.000$). There were significant differences in chlorophyll a, chlorophyll b and total chlorophyll between coffee leaves and brewed coffee leaves ($p = 0,000$) as seen in Table 3.

Table 9. Total Chlorophyll

Samples	Chlorophyll a		Chlorophyll b		Total Chlorophyll	
	Dried coffee leaves	Brewed coffee leaves	Dried coffee leaves	Brewed coffee leaves	Dried coffee leaves	Brewed coffee leaves
	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD	Mean ± SD
5 g	23.66±0.35 ^a	0.74±0.04 ^a	39.98±2.37 ^a	1.35±0.11 ^a	63.62±2.01 ^a	2.08±0.14 ^a
10 g	23.75±0.17 ^a	1.16±0.09 ^b	42.65±0.63 ^{ab}	2.19±0.13 ^b	66.37±0.67 ^{ab}	3.35±0.22 ^b
15 g	23.97±0.39 ^a	1.73±0.31 ^c	43.17±0.61 ^{ab}	3.19±0.44 ^c	67.13±0.99 ^{ab}	4.91±0.75 ^c
20 g	24.66±0.73 ^a	2.12±0.02 ^c	44.76±0.80 ^b	3.91±0.52 ^d	69.40±1.45 ^b	6.01±0.09 ^d
P^x	0.098	0.000*	0.014*	0.000*	0.006*	0.000*

Note: One-way-anova.test. Dried coffee leaves were brewed with 200 mL of water at 70°C for 10 minutes.. *Significant ($p < 0.05$). Numbers followed by superscript letters (a,b,c,d) differ to show significant differences.

Antioxidant Activity

The results of the analysis showed that there were no significant differences in the Robusta coffee leaves group ($p = 0.206$) and brewed ($p = 0.131$) as seen in **Table 4**.

Table 10. Antioxidant Activity with DPPH Method

Treatment	Inhibition (%)		p
	Dried coffee leaves Mean ± SD(%)	Brewed coffee leaves Mean ± SD (%)	
5 gram	92.7 ± 0.61	84.9 ± 0.23	0.109 ^p
10 gram	91.8 ± 0.76	86.1 ± 0.85	0.019 ^{*q}
15 gram	92.0 ± 2.00	85.6 ± 1.20	0.021 ^{*q}
20 gram	90.2 ± 1.36	83.3 ± 1.47	0.000 ^{*q}
p	0,206 ^z	0,131 ^y	

Note: Dried coffee leaves were brewed with 200 mL of water at 70°C for 10 minutes. *Significant ($p < 0,05$). ^pWilcoxon test, ^qPaired t test, ^zOne way Anova, ^yKruskal Wallis test

Antibacterial Activity Test

The results of the antibacterial activity analysis in *Escherichia coli* and *Salmonella typhi* showed significant differences in the groups of brewed coffee leaves ($p = 0.000$) as seen in Table 5.

Table 11. Antibacterial Activity

Samples	<i>Escherichia coli</i>		<i>Salmonella typhi</i>	
	Mean \pm SD	Inhibition	Mean \pm SD	Inhibition
5 g	7.65 \pm 0.10 ^a	Medium	7.47 \pm 0.28 ^a	Medium
10 g	8.47 \pm 0.15 ^b	Medium	8.20 \pm 0.05 ^b	Medium
15 g	9.17 \pm 0.15 ^c	Medium	8.55 \pm 0.26 ^b	Medium
20 g	9.89 \pm 0.26 ^d	Medium	8.70 \pm 0.19 ^c	Medium
<i>p</i>	0,000*		0,000*	

*One Way Anova test. *Significant ($p < 0.05$). Numbers followed by superscript letters (a,b,c,d) differ to show significant differences.

Acceptability of Brewed Coffee Leaves

There was a significant difference in the color and Taste of brewed coffee leaves ($p = 0,000$) and did not show a significant difference in the aroma of brewed coffee leaves ($p = 0.392$) as seen in Table 6.

Table 12. Acceptability of Brewed Coffee Leaves

Treatment	Color		Aroma		Taste	
	Median	Note	Median	Note	Median	Note
5 g	3(1-4) ^a	Like slightly	3(1-4) ^a	Like slightly	3(2-4) ^a	Like slightly
10 g	3(2-4) ^{ab}	Like slightly	3(2-4) ^a	Like slightly	2(1-3) ^a	Dislike slightly
15 g	2,5(1-4) ^{bc}	Dislike slightly	3(1-4) ^a	Like slightly	1(1-4) ^b	Dislike very much
20 g	2(1-4) ^c	Dislike slightly	3(1-4) ^a	Like slightly	1(1-2) ^c	Dislike very much
<i>p</i>	0,000*		0,392		0,000*	

Note: Kruskal Wallis test. *Significant ($p < 0.05$). Numbers followed by superscript letters (a,b,c,d) differ to show significant differences.

Discussion

In this study, Robusta coffee leaves were processed by the Japanese Style Green Tea Process (JGTP). JGTP method is one of the tea processing without fermentation process. Previous study has explained that the JGTP method in Arabica coffee leaves has higher phytochemicals concentration compared to other tea-making methods.(X.-M. Chen et al., 2018). The blanching process was used at the withering stage. Blanching can deactivate the enzyme polyphenol oxidase (PPO) which can cause oxidation.(Pourcel, Routaboul, Cheynier, Lepiniec, & Debeaujon, 2007)

Oxidation is a degenerative process triggered by the presence of free radicals which causes rancidity and decreasing components in food. The oxidation process causes the loss of phenolic components of food.(X.-M. Chen et al., 2018) The blanching process increases the polyphenol content during extraction.(Nayak, Liu, & Tang, 2015) In addition, the processing of green tea maintain the composition of chlorophyll including chlorophyll a (Chl a) and chlorophyll b (Chl b) that acts as an antibacterial.(Li et al., 2018)

Based on acceptability of brewed coffee leaves, the best acceptance was 5 g of brewed coffee leaves. The more the amount of leaves that are brewed, it will produce a thick yellowish-green color. Coffee leaves products using the JGTP method do not undergo oxidation of polyphenol compounds so that the resulting color was greenish or yellowish-green. Green is the main color determined by the chlorophyll content. Chlorophyll a on the leaves will produce a dark green color and chlorophyll b will produce a yellowish-green color. In addition to the chlorophyll content, flavonol content also plays a role in producing a yellowish color on the coffee leaves by the JGTP method.(Chaturvedula & Prakash, 2011)

Based on the aroma described by panelists, the aroma of brewed coffee leaves such as the aroma of fresh leaves. The more the amount of leaves in brewed, the smell will be thicker and increasingly disliked by panelists. The aroma of the leaves is caused by the high compound of catechins.(Ho, Zheng, & Li, 2015) The taste of brewed coffee leaves was bitter. The bitter taste is influenced by the presence of chlorogenic acid and caffeine.(Belay, 2011; Gokul G & Lakshmanan, 2016) Caffeine is a natural compound in food and drinks that contributes to affect the bitter taste. Some studies show that the main taste of caffeine is bitter. Caffeine generally amounts to 2-5% of the dry weight of tea. Caffeine will cause a bitter taste on the tongue depending on the level of concentration.(Keast & Roper, 2007; Zou, Xiao, Wang, & Zhang, 2018) The study explained that the bitter taste began to emerge when the caffeine content was 75-155 mg/L. Chlorogenic acid also plays a role in acidity and bitterness.(Belay, 2011)

The moisture and ash content in dried Robusta coffee leaves processed by the JGTP method met the Indonesian National Standard (SNI) green tea requirements, where the moisture of green tea is set to a maximum

of 8% and ash content of 4-8%.(SNI, 2016) The moisture in accordance with SNI can prevent the growth of bacteria and fungi, and prevent oxidation process. Protein content in dried Robusta coffee leaves with JGTP method included in the range of green tea, where protein content in green tea about 15-20%. Protein has the potential as an additional antioxidant in food because it can inhibit fat oxidation through several pathways including inactivation of Reactive Oxygen Species (ROS) through antioxidant enzymes (superoxide dismutase (SOD) and catalase), scavenge free radicals, chelating prooxidative transition metals, reducing hydroperoxide, and reducing hydroperoxides, and enzymatically removes specific oxidants.(Elias, Kellerby, & Decker, 2008) Protein also acts as an antimicrobial. Previous study explained that protein extracted from lemon, orange and grapefruit seeds could inhibit the growth of some pathogenic bacteria.(Karabiber, Yilmaz, & Zorba, 2018)

The fat content in Robusta coffee leaves processed by the JGTP method was lower than in Arabica coffee leaves with the same method. Based on previous research, Arabica coffee leaves have higher fat content about 4.5-12.5%. Meanwhile, the carbohydrate content in Robusta coffee leaves was higher than in Arabica coffee leaves. In previous study, carbohydrate content of Arabica coffee leaves was about 51.99-63.93%.(Woldesenbet, 2005)

In this study, Robusta coffee leaves processed by JGTP method have bioactive components of flavonoids, chlorogenic acid, caffeine, and chlorophyll. Previous study has shown that coffee leaves contain flavonoids such as quercetin, rutin, and kaempferol.(Martins, Araújo, Tohge, Fernie, & DaMatta, 2014) Flavonoids can act as antioxidants to scavenge free radicals, prevent the formation of reactive oxygen species (ROS), and increase endogenous antioxidants (superoxide dismutase, catalase, and glutathione peroxidase). The mechanism of flavonoids in preventing damage by free radicals is by giving hydrogen atoms to free radicals to form stable flavonoxyl radicals (flavonoids (O•)). In addition, flavonoids can protect the phospholipid membrane by giving one of the hydrogen ions to the radical lipid peroxy, so that radical reactions can be stopped.(Hamid, Usman, Aiyelaagbe, & Ameen, 2010; Irina & Mohamed, 2012) Flavonoids can also function as antibacterial by inhibiting nucleic acid synthesis, inhibiting energy metabolism, changes in cytoplasmic membrane function, reducing cell adhesion and biofilm formation and causing damage to the cytoplasmic membrane.(Farhadi et al., 2018)

Chlorogenic acid is an ester formed from hydroxycinnamic acid (caffeic acid, ferulic acid, p-kumaric acid) and quinic acid and into the phenolic acid group.(Liang & Kitts, 2016) Chlorogenic acid can chelate transition metals like Fe²⁺ to scavenge free radicals and break radical chains. Chlorogenic acid scavenges DPPH radicals, superoxide anions, hydroxyl radicals, and protect DNA from damage caused by oxidative stress.(Azam, Hadi, Khan, & Hadi, 2003) Chlorogenic acid also function as antibacterial. Chlorogenic acid is a phenolic bioactive compound that can cause physiological changes in bacterial cell membranes.(Kabir et al., 2014) Chlorogenic acid causes cell membrane rupture and release of the content inside the cellular. Chlorogenic acid changes in bacterial cell membrane permeability that will cause damage to bacterial nucleotides.(Lou et al., 2011)

Robusta coffee leaves with processed by JGTP method also contained caffeine. Caffeine prevents lipid oxidation by inhibiting the production of free radicals including hydroxyl radicals, peroxy radicals, and singlet oxygen.(Azam et al., 2003) Caffeine also acts as antibacterial agent. It can inhibit protein and DNA synthesis by inhibiting the merging of adenine and thymidine. Besides, caffeine also increases genotoxicity after DNA damage.(Pruthviraj et al., 2011)

The amount of chlorophyll in brewed Robusta coffee leaves was very low compared to the amount of chlorophyll in tea leaves. Chlorophyll acts as antioxidants, where it can prevent the Fenton reaction, by chelating Fe(II). Chlorophyll directly scavenges the hydroxyl free radicals generated from H₂O₂.(Hsu, Chao, Hu, & Yang, 2013) It can also act as antibacterial by increasing membrane permeability in bacteria.(Kustov et al., 2018) This will cause changes in membrane structure and accelerate intracellular leakage thereby destroying membrane integrity which will facilitate the entry of antibacterial.(Wu et al., 2016)

The results showed that Robusta coffee leaves with the JGTP method had a DPPH radical inhibition. Based on the results of this analysis, Robusta coffee leaves and brewed coffee leaves have high antioxidant activity, being able to capture free radicals by 50%. The results of this study are in line with research on green tea, where green tea has a DPPH radical inhibition more than 50%.

The results of antibacterial of brewed coffee leaves showed the inhibitory power of 5 g, 10 g, 15 g, and 20 g brewed Robusta coffee leaves against *Escherichia coli* and *Salmonella typhi* bacteria. *Escherichia coli* and *Salmonella typhi* belong to gram-negative bacteria. The greater the amount of leaves in the brewed process, the greater the diameter of the inhibition zone formed. The effectiveness of antibacterial substances will be influenced by the concentration of these substances. An increase in the concentration of the substance causes an increase in the content of active compounds such as flavonoids, caffeine and chlorogenic acid which functions as an antibacterial so that its ability to inhibit bacterial growth increases.(Mufti, Bahar, & Arisanti, 2017; Muslim & Dephinto, 2017) Previous study conducted on CTC brewed black tea with a comparison of tea weight and water volume (w/v) 1:50, 2:50, and 3:30 resulted in an area of 29.89 successive barriers; 33.64 and 45.02 mm².(Rohdiana et al., 2013)

Conclusion

Robusta coffee leaves in this research were processed by JGTP method with 75-second blanching. The best organoleptic based on color, aroma, and taste was 5 g of dried Robusta coffee leaves. Brewed Robusta coffee leaves processed by the Japanese Style Green tea Process (JGTP) have contain flavonoids 0.10 mg QE/g, chlorogenic acid 3.49 mg/g, caffeine 0.194 mg/g and high antibacterial activity in 20 g of brewed Robusta coffee leaves.

Conflict of Interest

No conflict of interest.

Acknowledgments

This research was supported by Research and Development (RPP) from the Faculty of Medicine, Universitas Diponegoro.

References

- Analysis., O. M. of. (2005). *Official Methods of Analysis*. (18th ed.). AOAC International: USA.
- Azam, S., Hadi, N., Khan, N. U., & Hadi, S. M. (2003). Antioxidant and Prooxidant Properties of Caffeine , Theobromine and Xanthine. *Med Sci Monit.*, 9(9), 325–331.
- Belay, A. (2011). Some biochemical compounds in coffee beans and methods developed for their analysis. *International Journal of the Physical Sciences*, 6(28), 6373–6378.
- Campa, C., Urban, L., Mondolot, L., Fabre, D., Roques, S., Lizzi, Y., ... Etienne, H. (2017). Juvenile Coffee Leaves Acclimated to Low Light Are Unable to Cope with a Moderate Light Increase. *Front Plant Sci.*, 8(1126), 1–16.
- Chacko, S. M., Thambi, P. T., Kuttan, R., & Nishigaki, I. (2010). Beneficial effects of green tea : A literature review. *Chin Med*, 5(15), 1–9.
- Chaturvedula, V. S. P., & Prakash, I. (2011). The aroma, taste, color and bioactive constituents of tea. *Journal of Medicinal Plants Research*, 5(11), 2110–2124.
- Chen, X. (2019). A Review on Coffee Leaves : Phytochemicals, Bioactivities and Applications. *Crit Rev Food Sci Nutr*, 59(6), 1008–1025. <https://doi.org/10.1080/10408398.2018.1546667>.
- Chen, X.-M., Ma, Z., & Kitts, D. D. (2018). Effects of processing method and age of leaves on phytochemical profiles and bioactivity of coffee leaves. *Food Chemistry*, 249, 143–153. <https://doi.org/10.1016/j.foodchem.2017.12.073>
- Elias, R. J., Kellerby, S. S., & Decker, E. A. (2008). Antioxidant Activity of Proteins and Peptides. *Critical Reviews in Food Science and Nutrition*, 48(5), 430–441.
- Farhadi, F., Khameneh, B., Iranshahi, M., & Iranshahi, M. (2018). Antibacterial activity of flavonoids and their structure – activity relationship : An update review. *Phyther Res*, 1–28.
- Gokul G, & Lakshmanan, R. (2016). Effect Of Chlorhexidie Mouthwash On Taste Alteration. *Asian J Pharm Clin Res*, 9(1), 102–104.
- Hamid, A. A., Usman, L. A., Aiyelaagbe, O., & Ameen, M. O. (2010). Antioxidants : Its Medicinal and Pharmacological Applications. *African J Pure Appl Chem*, 4(8), 142–151.
- Hasanah, M., Maharani, B., & Munarsih, E. (2017). Daya Antioksidan Ekstrak Dan Fraksi Daun Kopi Robusta (*Coffea robusta*) terhadap Pereaksi DPPH (2,2-difenil-1-pikrilhidrazil) (Antioxidant of Extract and Fraction *Coffea robusta* Leaves with Diphenylpicrylhydrazyl (DPPH) Method. *Indonesian Journal of Pharmaceutical Science and Technology*, 4(2), 42–49.
- Ho, C.-T., Zheng, X., & Li, S. (2015). Tea Aroma Formation. *Food Science and Human Wellness*, 4(1), 9–27.
- Hsu, C.-Y., Chao, P.-Y., Hu, S.-P., & Yang, C.-M. (2013). The Antioxidant and Free Radical Scavenging Activities of Chlorophylls and Pheophytins. *Food and Nutrition Sciences*, 4, 1–18.
- Irina, I., & Mohamed, G. (2012). Biological Activities and Effects of Food Processing on Flavonoids as Phenolic Antioxidants. In *Advances in Applied Biotechnology*. (pp. 112–114). Croatia.
- Kabir, F., Katayama, S., Tanji, N., & Nakamura, S. (2014). Antimicrobial Effects of Chlorogenic Acid and Related Compounds. *Journal of the Korean Society for Applied Biological Chemistry*, 57(3), 359–365.
- Kanisius, A. A. (1988). *Budidaya Tanaman Kopi* (Coffee Cultivation) (16th ed.). Yogyakarta: Kanisius.
- Karabiber, E., Yilmaz, E., & Zorba, N. N. D. (2018). antimicrobial and functional properties of the proteins

- extracted from lemon, orange and grapefruit seeds press meals. *Quality Assurance and Safety of Crops & Foods*, 10(2), 1–10.
- Keast, R. S. J., & Roper, J. (2007). A Complex Relationship Among Chemical Concentration, Detection Threshold, and Suprathreshold Intensity of Bitter Compounds. *Chem Senses*, 32(3), 245–253.
- Khotimah, K. (2014). Karakteristik Kimia Kopi Kawa dari Berbagai Umur Helai daun Kopi yang Diproses dengan Metode Berbeda (Chemical Characteristics of Kawa Coffee of Various Age Leaves Processed by Different Methods). *J Teknol Pertan*, 9(1), 40–45.
- Kustov, A. V., Belykh, D. V., Smirnova, N. L., Venediktov, E. A., Kudayarova, T. V., Kruchin, S. O., ... Berezin, D. B. (2018). Synthesis and investigation of water-soluble chlorophyll pigments for antimicrobial photodynamic therapy. *Dyes and Pigments*, 149, 553–559.
- Ky, C.-L., Louarn, J., Dussert, S., Guyot, B., Hamon, S., & Noirot, M. (2001). Caffeine, trigonelline, chlorogenic acids and sucrose diversity in wild *Coffea arabica* L. and *C. canephora* P. accessions. *Food Chemistry*, 75(2), 223–230.
- Li, X., Zhou, R., Xu, K., Xu, J., Jin, J., Fang, H., & He, Y. (2018). Rapid Determination of Chlorophyll and Pheophytin in Green Tea Using Fourier Transform. *Molecules*, 21, 1–13.
- Liang, N., & Kitts, D. D. (2016). Role of Chlorogenic Acids in Controlling Oxidative and Inflammatory Stress Conditions. *Nutrients*, 6, 16. <https://doi.org/10.3390/nu8010016>.
- Lou, Z., Wang, H., Zhu, S., Ma, C., & Wang, Z. (2011). Antibacterial Activity and Mechanism of Action of Chlorogenic Acid. *J Food Sci.*, 76(6), 398–403.
- Manach, C., Scalbert, A., Morand, C., Rémésy, C., & Jiménez, L. (2004). Polyphenols: Food Sources and Bioavailability. *The American Journal of Clinical Nutrition*, 79(5), 727–747.
- Martins, S. C. V., Araújo, W. L., Tohge, T., Fernie, A. R., & DaMatta, F. M. (2014). n High-Light-Acclimated Coffee Plants the Metabolic Machinery Is Adjusted to Avoid Oxidative Stress Rather than to Benefit from Extra Light Enhancement in Photosynthetic Yield. *PLoS ONE*, 9(4), 1–11.
- Mufti, N., Bahar, E., & Arisanti, D. (2017). Uji Daya Hambat Ekstrak Daun Sawo terhadap Bakteri *Escherichia coli* secara In Vitro (Inhibition Test of Inhibition Test of Sawo Leaf Extract Against *Escherichia coli* Bacteria in Vitro). *Jurnal Kesehatan Andalas*, 6(2), 289–294.
- Muslim, Z., & Dephinto, Y. (2017). Perbandingan Efektivitas Antimikroba Ekstrak Daun Kopi Robusta (*Coffea canephora*) Dengan Variasi Pengeringan Terhadap *Escherichia coli* (Comparison of Antimicrobial Effectiveness of Robusta (*Coffea canephora*) Leaf Extract with Drying Variations on *Escherichia coli*). *J Sains Dan Teknol Farm*, 19, 86–88.
- Naegele, E. (2016a). Determination of Caffeine in Coffee Products According to DIN 20481. *Agil Technol*, 1–6.
- Naegele, E. (2016b). Determination of Chlorogenic Acid in Coffee Products According to DIN 10767. *Agil Technol*, 1–8.
- Nayak, B., Liu, R. H., & Tang, J. (2015). Effect of Processing on Phenolic Antioxidants of Effect of Processing on Phenolic Antioxidants of Fruits, Vegetables, and Grains — A Review. *Crit Rev Food Sci Nutr.*, 55(7), 887–918.
- Nayeem, N., Denny, G., & Mehta, S. K. (2011). Comparative phytochemical analysis, antimicrobial and anti oxidant activity of the methanolic extracts of the leaves of *Coffea Arabica* and *Coffea Robusta*. *Der Pharmacia Lettre*, 3(1), 292–297.
- Pourcel, L., Routaboul, J.-M., Cheynier, V., Lepiniec, L., & Debeaujon, I. (2007). Flavonoid Oxidation in Plants : from Biochemical Properties to Physiological Functions. *Trends Plant Sci.*, 12(1), 29–36.
- Prawira-Atmaja, M. I., Shabri, Khomaini, H. S., Maulana, H., Harianto, S., & Rohdiana, D. (2018). Changes in chlorophyll and polyphenols content in *Camellia sinensis var. sinensis* at different stage of leaf maturity. *Earth Enviromental Sci*, 131, 1–7.
- Pruthviraj, P., Shital, K., Suchita, B., & Shilpa, K. (2011). Evaluation of Antibacterial Activity of Caffeine. *Int J Res Ayurveda Pharm*, 2(4), 1354–1357.
- Rhayanne T. M. Ramos, I. C. F. B., Ferreira, M. R. A., & Soares, L. A. L. (2017). Spectrophotometric Quantification of Flavonoids in Herbal Material, Crude Extract, and Fractions from Leaves of *Eugenia uniflora* Linn. *Pharmacognosy Res.*, 9(3), 253–260.
- Rohdiana, D., Arief, D. Z., & Budiman, A. (2013). Aktivitas penghambatan pertumbuhan bakteri *Escherichia coli* oleh berbagai jenis teh dan seduhannya. (Inhibitory activity of *Escherichia coli* by type of teas and its liquors). *Jurnal Penelitian Teh Dan Kina*, 16(1), 37–44.
- Sadino, A., Sahidin, I., & Wahyuni, W. (2018). Antibacterial Activity of *Polygonum pulchrum* Blume Ethanol Extract on *Staphylococcus aureus* and *Escherichia coli*. *Pharmacol Clin Pharm Res.*, 3, 26–32.
- Silva, L. A. L. da, Pezzini, B. R., & Soares, L. (2015). Spectrophotometric Determination of The Total Flavonoid Content in *Ocimum basilicum* L.(*Lamiaceae*) Leaves. *Pharmacogn Mag*, 11(41), 96–101.
- SNI. (2016). *SNI 3945:2016 Teh Hijau (Green Tea)*. Jakarta.
- Woldesenebet, A. (2005). Nutritional Composition, Phytochemical Screening, Processing Methods and Sensory Attributes of A Brew Made From Infusions of Matured Leaves of Arabica Coffee Tree Consumed In

- Sidama, Kambata and Harar Communities, Ethiopia. Addus Ababa University.
- Wu, Y., Bai, J., Zhong, K., Huang, Y., Qi, H., Jiang, Y., & Gao, H. (2016). Antibacterial Activity and Membrane-Disruptive Mechanism of 3-p-trans Coumaroyl-2-hydroxyquinic Acid, a Novel Phenolic Compound from Pine Needles of *Cedrus deodara*, against *Staphylococcus aureus*. *Molecules* *21*(8), 1–12.
- Zou, G., Xiao, Y., Wang, M., & Zhang, H. (2018). Detection of bitterness and astringency of green tea with different taste by electronic nose and tongue. *PLoS ONE*, *13*(12), 1–10.

Letter of Accepted

The screenshot displays an email client interface. On the left, a sidebar lists folders: 'Inbox' (187), 'Starred', 'Snoozed', 'Important', 'Sent', 'Drafts' (54), 'Categories', and 'More'. Below these are 'Labels' including '[imap]/Archive', '[imap]/Drafts', and 'Notes'. The main content area shows an email from 'ICHWB Submission and Review System' (conferences-noreply@ums.ac.id) received on 'Thu, Jan 23, 2020, 11:24 AM'. The subject is 'ICHWB2019 status your paper submission 14787 - Accepted'. The email body contains the following text:

Thu, 23/01/2020 - 11:24am
This is to inform you that your submission has been changed by the track-director.

The status of your submitted paper no 14787 is Accepted

After careful consideration by our internal editorial team and a group of expert reviewer, we feel that your manuscript "Bioactive components and antibacterial activity study in Robusta coffee leaves (coffea canephora)" has potential for publication. The manuscript, at this time, will be sent to Q3 scopus journal.

We will keep you informed of your article's progress.

Accepted
=====

Detail information of your paper and LETTER OF ACCEPTANCE can be found in your author account (login required).

For registration your presentation attendance, please visit <http://iseth.ums.ac.id/2019/reg19>



INTERNATIONAL JOURNAL OF PHARMACEUTICAL RESEARCH

A Step Towards Excellence
Published by : Advanced Scientific Research

ISSN
0975-2366

Home
About Us
Editorial Board
Instruction to Authors
Current Issue
Article In Press
Table Of Contents

CURRENT ISSUE

volume 15, Issue 1, Jan - Mar, 2023

ARTICLE IN PRESS

No Data found.

ADOBE READER

(Require Adobe Acrobat Reader to open. If you don't have Adobe Acrobat Reader)



[Click here to Download](#)

IJPR 9[3] JULY - SEPTEMBER 2017 SPECIAL ISSUE

July - September 9[3] 2017

[Click to download](#)

Volume 12, issue 3, July - Sept, 2020

REVIEW

The effect of Sida acuta on bacterial enzymes in azoxymethane-induced experimental colon cancer
PERUMAL SAMY MUNNESWARI, SRI RASHMY MADATHIL, KANNAPPAN POORNIMA

Nanomedicine as an Emerging Treatment Approach for Many Cancers: An Updated Review
SALIM M.

Metabolic Syndrome Parameters in Psychiatric Disease: Current Knowledge and Future Perspectives
SOMAYEH DEHKOHNEH, SARA AZADI, MORAD RASOULI-AZAD, JAFAR JALILIAN, AZAM GHASEMINEGHAD, NEDA VAHED, FARKHONDEH RAZZAGHI, LEILA JAFARI, SAMIRA ABBASZADEH-MASHKANI

Oxidative Stress, Glycemic Control, Inflammation, and Lipid Profiles in Substance Abuse Disorders: Current Knowledge and Future Perspectives
SOMAYEH DEHKOHNEH, SARA AZADI, MORAD RASOULI-AZAD, JAFAR JALILIAN, AZAM GHASEMINEGHAD, NEDA VAHED, FARKHONDEH RAZZAGHI, AMINEH SADAT ZAHIRY POUR, I FII A. JAFARI, SAMIRA ABBASZADEH-MASHKANI

Compared panoramic radiography and CBCT on the surgical technique of third molar removal: A systematic review and meta-analysis
NARIMAN NIKPARTO, REZA JALILIAN, SAEED HASANI MEHRABAN

Compared the effects of skeletal anchored maxillary protraction Vs dental anchored maxillary protraction in children with class iii malocclusion: A systematic review and meta-analysis
ALIREZA CHAMANI, PARISA DOROUDGAR, KHANMIRZA BADRI

Outcome measures (proms) of implant-supported, relative to tooth-supported fixed dental prostheses: A systematic review and meta-analysis
AISOODA AFSHARI, FATEMEH MOOSAZADE HASSAN KHADEH, MESHKAT NAEIMI DARESTANI, MEHRAB TAKHTDAR

Association of dietary inflammatory index with visceral adiposity index among obese female adolescents
SUKMA UTAMI, FILLAH FITHRA DIENY, ARYU CANDRA KUSUMASTUTI, A FAHMY ARIF TSANI

Bioactive components and antibacterial activity in Robusta coffee leaves (Coffea canephora)
GEMALA ANJANI, NURMASARI WIDYASTUTI, ZULFATUL MASRUROH, RIZKA AYU DWI YULIANA, VITA GUSTIN ALMIRA, A. FAHMY ARIF TSANI, CHOIRUN NISSA, M IQBAL PRAWIRA-ATMAJA

Production and characterization of polyhydroxybutyrate (PHB) from marine invertebrates
N. EZHILRAJ, S. CHIPPY RAJ, S. MARY JOSEPHINE PUNITHA

Thyroid Abnormalities Associated with Complete Blood Count Parameter
JONGKY PRAJITNO, SOEBAGUO ADI SOELISTUO, HERMAWAN SUSANTO, AGUNG PRANOTO

Serum Bicarbonate Level And Nutritional Status In Non-Diabetic And Non-Dialysis Stage 5 Chronic Kidney Disease Patients
EKY RIYADI, NUNUK MARDIANA, WIDODO



Embase[®]
A DIVISION OF THE AMERICAN CHEMICAL SOCIETY



ONLINE SUBMISSION

[Click here for Online Submission](#)

ARTICLE IN PRESS

No Data found.

ADOBE READER

(Require Adobe Acrobat Reader to open. If you don't have Adobe Acrobat Reader)



[Click here to Download](#)

IJPR 9[3] JULY - SEPTEMBER 2017 SPECIAL ISSUE

July - September 9[3] 2017

[Click to download](#)

Article Detail

Bioactive components and antibacterial activity in Robusta coffee leaves (Coffea canephora)

Author: GEMALA ANJANI, NURMASARI WIDYASTUTI, ZULFATUL MASRUROH, RIZKA AYU DWI YULIANA, VITA GUSTIN ALMIRA, A. FAHMY ARIF TSANI, CHOIRUN NISSA, M IQBAL PRAWIRA-ATMAJA

Abstract: Purpose: To analyze bioactive components and antibacterial activity in brewed Robusta coffee leaves that processed by Japanese Style Green Tea Process (JGTP). Methodology: This was an experimental research with the sample of Robusta coffee leaves and brewed Robusta coffee leaves. The coffee leaves were brewed in water at 70°C during 10 minutes. The analysis of nutrition contents by proximate method, total flavonoids by colorimeter method, chlorogenic acid and caffeine by HPLC method, antioxidant activity by DPPH method, total chlorophyll using the amon formula and antibacterial activity by disk diffusion method in 5, 10, 15, and 20 g. Results: There were significant differences in color, taste, and acceptability of brewed Robusta coffee leaves. The best acceptance of brewed Robusta coffee leaves was 5 g of dried coffee leaves. Robusta coffee leaves contain 3.1% water, 6.03% ash, 17.76% protein, 2.37% fat, and 70.83% carbohydrate. The content of flavonoids, chlorogenic acid and caffeine in Robusta coffee leaves were 0.81 mgQE/g, 16.92 mg/g, 0.46 mg while in the brewed were 0.10 mgQE/g, 3.49 mg/g, 0.19 mg/g, respectively. The inhibition of free radical in Robusta coffee leaves were 90.2 - 92.7%, while in the brewed were 83.3 - 86.1%. Total chlorophyll of dried leaves were 63.62 - 69.39 mg/L and the brewed were 2.08 - 6.01 mg/L. Antibacterial activity Escherichia coli 7.65 - 9.89 mm and Salmonella typhi 7.47- 8.70 mm inhibition zone. Applications/Originality/Value: The role of various bioactive components in Robusta coffee leaves on antioxidant and antibacterial activity.

Keyword: JGTP

Download: [Request For Article](#)



ONLINE SUBMISSION

[Click here for Online Submission](#)

USER LOGIN

Author Reviewer

Editor Subscriber

Username:

Bioactive components and antibacterial activity in Robusta coffee leaves (*coffea canephora*)

GEMALA ANJANI^{1,2}, * NURMASARI WIDYASTUTI^{1,2}, ZULFATUL MASRUROH¹, RIZKA AYU DWI YULIANA¹, VITA GUSTIN ALMIRA¹, A. FAHMY ARIF TSANI^{1,2}, CHOIRUN NISSA^{1,2}, M IQBAL PRAWIRA-ATMAJA³

¹ Department of Nutrition Science, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

² Center of Nutrition Research (CENURE), Universitas Diponegoro, Semarang, Indonesia

³ Research Institute for Tea and Cinchona, Bandung, Indonesia

* Corresponding Author

Email ID: gemaanjani@gmail.com

Received: 12.03.20, Revised: 13.04.20, Accepted: 18.05.20

ABSTRACT

Purpose: To analyze bioactive components and antibacterial activity in brewed Robusta coffee leaves that processed by *Japanese Style Green Tea Process* (JGTP).

Methodology: This was an experimental research with the sample of Robusta coffee leaves and brewed Robusta coffee leaves. The coffee leaves were brewed in water at 70°C during 10 minutes. The analysis of nutrition contents by proximate method, total flavonoids by calorimeter method, chlorogenic acid and caffeine by HPLC method, antioxidant activity by DPPH method, total chlorophyll using the arnon formula and antibacterial activity by disk diffusion method in 5, 10, 15, and 20 g.

Results: There were significant differences in color, taste, and acceptability of brewed Robusta coffee leaves. The best acceptance of brewed Robusta coffee leaves was 5 g of dried coffee leaves. Robusta coffee leaves contain 3.1% water, 6.03% ash, 17.76% protein, 2.37% fat, and 70.83% carbohydrate. The content of flavonoids, chlorogenic acid and caffeine in Robusta coffee leaves were 0.81 mgQE/g, 16.92 mg/g, 0.46 mg while in the brewed were 0.10 mgQE/g, 3.49 mg/g, 0.19 mg/g, respectively. The inhibition of free radical in Robusta coffee leaves were 90.2 - 92.7%, while in the brewed were 83.3 - 86.1%. Total chlorophyll of dried leaves were 63.62 -69.39 mg/L and the brewed were 2.08 - 6.01 mg/L. Antibacterial activity *Escherichia coli* 7.65 – 9.89 mm and *Salmonella tiphy* 7.47-8.70 mm inhibition zone.

Applications/Originality/Value: The role of various bioactive components in Robusta coffee leaves on antioxidant and antibacterial activity.

Keywords: JGTP

INTRODUCTION

Coffee plant (*Coffea sp.*) is included in the family *Rubiaceae*. In general, there are three types of coffee, including Arabica coffee (*Coffea arabica*), Robusta coffee (*Coffea canephora*), and Liberica coffee (*Coffea liberica*). The most widely consumed coffee in Indonesia is Robusta coffee. Robusta coffee can be easily grown at lower altitude of less than 1000 meters above sea level and it resilient against disease and pests.(Kanisius, 1988)

The results of research on Arabica coffee leaves have nutritional content based on proximate components. Proximate components of Arabica coffee leaves from the Ethiopian region include 5.5-7.8% moisture, 8.8-12.4% ash content, 14.4-19.0% protein, 4.5-12.5% fat, and 51.0-63.9% carbohydrate.(Woldesenebet, 2005) Robusta coffee leaves contain phytochemicals that vary depending on leaf development.(X.-M. Chen, Ma, & Kitts, 2018) The younger the coffee leaves, the higher the chlorogenic acid content even 10 times

higher than the old coffee leaves.(Campa et al., 2017) The content of chlorogenic acid in Robusta coffee is higher compared to Arabica coffee, respectively 11.3% and 4.1%.(Ky et al., 2001) The caffeine content in young Robusta coffee leaves (0.29-0.5%) is lower than the caffeine content in its fruit (1.6-2.4%).(Khotimah, 2014)

Robusta coffee leaves also contain phenolic acid and total flavonoids, respectively 27.04 µg/g and 10.90 µg/g higher than Arabica coffee leaves respectively 21.80 µg/g and 8.08 µg/g.(Nayeem, Denny, & Mehta, 2011) Flavonoids as natural antioxidants can prevent the occurrence of diseases, such as cancer, metabolic syndrome, diabetes mellitus, and atherosclerosis which associated to oxidative stress.(X. Chen, 2019; Manach, Scalbert, Morand, Rémésy, & Jiménez, 2004)

The content of caffeine, flavonoids, and chlorogenic acid of coffee leaves can also

function as the prevention of diarrhea. Caffeine can inhibit protein and DNA synthesis by inhibiting the merging of adenine and thymidine. Caffeine also increases genotoxicity after DNA damage.(Pruthviraj, Shital, Suchita, & Shilpa, 2011) Flavonoids as an antibacterial by inhibiting nucleic acid synthesis, inhibiting energy metabolism, and changes in cytoplasmic membrane function.(Farhadi, Khameneh, Iranshahi, & Iransahy, 2018) In addition to flavonoids, there are chlorogenic acid contents that function as antibacterial. Chlorogenic acid is a phenolic bioactive compound causes physiological changes in bacterial cell membranes that encourage damage to bacterial nucleotides.(Kabir, Katayama, Tanji, & Nakamura, 2014; Lou, Wang, Zhu, Ma, & Wang, 2011)

The processing of coffee leaves to obtain optimal antibacterial is done by minimizing chemical and enzymatic reactions to prevent damage in phytochemical content. The process can be obtained using green tea processing.(X.-M. Chen et al., 2018) The processing of green tea is able to produce phenolic components 6 times higher than black tea. Green tea processing aims to maintain the content of natural polyphenols which have beneficial effects on health. (Chacko, Thambi, Kuttan, & Nishigaki, 2010). Therefore, the aim of this research was to examine bioactive components, including caffeine, chlorogenic acid, total flavonoids, antioxidant activity, total chlorophyll, and nutritional contents and antibacterial activity in Robusta coffee leaves by green tea processing.

MATERIALS AND METHODS

The samples used *Robusta canephora* Pierre ex Froehner from Gambung village. Coffee leaves were processed using the Japanese Style Green Tea Process (JGTP) method at the Tea Quality Processing and Testing Laboratory, Tea and Quinine Research Center (PPTK) Gambung, Bandung Regency, West Java, Indonesia. Preliminary research was conducted to obtain the JGTP procedure at the withering stage, including 2 minutes steaming method and 45 s, 60 s, and 75 s of blanching. The best withering method was chosen based on the results of the best organoleptic tests conducted by trained panelists. The best withering result that will be used in the processing of coffee leaves was the blanching method for 75 s.

This study was an experimental study with one factor completely randomized design (CRD) with 2 samples, dried Robusta coffee leaves and brewed Robusta coffee leaves. Dried Robusta coffee leaves were brewed with 200 ml of water

in 70°C for 10 minutes. Samples were analyzed in duplicate with three replications.

Processing Robusta Coffee Leaves with JGTP Method

JGTP method was carried out through 5 steps, picking; blanching; cooling, crushing, and drying. Coffee leaves were picked from 2nd, 3rd, and 4th leaf of each branch of coffee plants, sorted, then carried out the blanching process for 75 s. Leaves were soaked in water and put in seducer for \pm 15 minutes while separating the leaves from its midrib. The next process was crushing the leaves three times using a crushing-tearing-curling machine (CTC) (CNC, Sri Lanka). The last process was the drying process. The drying process was 4-5 hours long by a rack drier machine (CNC, Sri Lanka) at 80°C.

Proximate Analysis

Proximate analyses using dried Robusta coffee leaves samples processed by JGTP to analyze the moisture, ash, fat, protein, and carbohydrate contents with AOAC 2005 method.(Analysis., 2005)

Total Flavonoid Analysis

5 grams of dried Robusta coffee leaves in 200 ml of methanol were homogenized with a magnetic stirrer for \pm 30 minutes. The solution was taken 60 ml for centrifugation, then coffee leaves produced supernatant in methanol. Meanwhile, 5 grams of dried Robusta coffee leaves were brewed in 200 ml of hot water (70°C) for 10 minutes and stirred. After being dissolved in water, 60 ml was taken to be centrifuged, coffee leaves supernatant will be produced in brewed water. Afterwards, two samples were analyzed.

Total flavonoid analysis using aluminum chloride (AlCl₃) calorimetry method with absorbance measurement using UV-VIS spectrophotometer. 1 ml of sample solution was mixed with 9 ml of methanol. Pipette 2.8 ml sample solution; 0.4 ml AlCl₃ 5%; 6.8 ml of 5% acetic acid. Incubation for 30 minutes, measure the absorbance (λ 428 nm). Calibration curves used the standard quercetin in methanol (0.005 mg / ml) with varying concentrations of 0 μ g/ml, 5 μ g/ml, 10 μ g/ml, 20 μ g/ml, 40 μ g/ml, and 60 μ g/ml. Standard was carried out using the same procedures as samples. (Rhayanne T. M. Ramos, Ferreira, & Soares, 2017; Silva, Pezzini, & Soares, 2015)

Analysis of Chlorogenic Acid and Caffeine

Analysis of chlorogenic acid using 20 g of dried coffee leaves extracted with 150 mL methanol:water (50:50 v/v) in soxhlet. After 4-5 cycles, soxhletation was stopped and cooled

down at room temperature. The extract was transferred into a 200 mL flask and the volume was adjusted to 200 mL with a mixture of methanol-water. Meanwhile, caffeine analysis using 10-20 g of coffee leaves were added to 5 g of MgO and put into 200 mL of water. Furthermore, the samples were heated in a water bath at 90°C and sterilized for 20 minutes. Then the extracts were filtered with 0.45µL, 25 mm cellulose filter cartridge. The filtrates were injected into the HPLC.(Naegele, 2016a, 2016b)

Total Chlorophyll Test

The total chlorophyll test was carried out by 5, 10, 15 and 20 grams of brewed Robusta coffee leaves compared to the samples dissolved with 80% acetone. Total chlorophyll test was conducted using 663 nm and 645 nm wave spectrophotometry. Chlorophyll levels were calculated using the Arnon formula:(Prawira-Atmaja et al., 2018)

Chlorophyll a = $(12.7 \times A663) - (2.69 \times A645)$

Chlorophyll b = $(22.9 \times A645) - (4.68 \times A663)$

Total chlorophyll = $(20.2 \times A645) + (8.02 \times A663)$

Analysis of Antioxidant Activity

Sample preparation was performed as a total flavonoid test. Antioxidant activity was analyzed using the DPPH method (1,1-diphenyl-2-picrylhydrazyl). A DPPH 0.4 mM solution was prepared by 0.00394 g of DPPH powder and dissolved in absolute methanol to 25 ml. The test was done by piping 0.1 ml of samples and sufficient volume with absolute methanol into a 5 ml measuring flask, then vortex. Measurement of antioxidant activity was done by adding 1 ml of DPPH 0.4 mM solution, then vortex for ± 1 minute. Incubate the solutions for 30 minutes in a dark place, measured the absorbance by UV-VIS spectrophotometry ($\lambda 517$ nm). A blank solution was obtained by piping 5 ml absolute methanol and 1 ml DPPH solution, left for 30 minutes in a dark place and measuring the absorbance as with the samples.(Hasanah, Maharani, & Munarsih, 2017)

Antibacterial Activity Test

Antibacterial activity test was carried out by the disk diffusion Kirby-Bauer method.(Rohdiana, Arief, & Budiman, 2013) This method was used to see the bacterial growth inhibition response based on the diameter of the clear zone of *E. coli* and *S. typhi*. Antibacterial activity test began by dissolving nutrients with aquades. The media stock and the equipment used were put into a

121°C autoclave for 20 minutes. The next step was making a suspension by taking an ose of pure bacteria and implanting it in a solution of NaCl. Next, applied the suspension to the hard agar using a sterile cotton swab. Blank discs that have been soaked in 5, 10, 15, and 20 grams of brewed Robusta coffee leaves for 20 minutes were placed on hard agar and incubated at 37°C for 24-hour incubation periods. Classification of bacterial growth inhibition was observed by clear zone diameter consists of 4 groups: weak response (diameter ≤ 5 mm), moderate (diameter 5-10 mm), strong (diameter 10-20 mm), and very strong (diameter ≥ 20 mm).(Sadino, Sahidin, & Wahyuni, 2018)

Organoleptic Test

Organoleptic test was carried out by 30 semi-trained panelists of Nutrition Sciences students with variations in weight of 5, 10, 15, and 20 grams of dried coffee leaves brewed with water at 70°C. The assessment of quality attributes of brewed coffee leaves used 4 favorite scales, 1 (dislike very much), 2 (dislike slightly), 3 (like slightly), and 4 (like very much).

STATISTIC ANALYSIS

Statistical analysis using univariate and bivariate analysis. If the data is normally distributed, use the paired t test, whereas if the data is normally distributed, use the Wilcoxon test. Analysis of antioxidant activity using the One Way Anova statistical test if normally distributed, and using the Kruskal Wallis test if the distribution is not normal. The analysis was performed with a 95% confidence level with p value 0.05 and $\alpha = 0.05$.

RESULTS

Processing Robusta Coffee Leaves with JGTP Method

Based on the appearance and color aspects, brewed coffee leaves with the blanching method had an average value from 7.5 to 8 (1-10) while the steaming method got a value of 4 (1-10) as seen in Figure 1. The taste of brewed using the blanching method had an average value from 34 to 35 (20-49) and obtaining a value of 27 for brewed by steaming method. The aroma of brewed and pulp of coffee leaves had a value of 5-5.50 (1-10) for the method of blanching water and steaming. From the aspect of the aroma of coffee grounds pulp obtained a value of 9 for each blanching method and 4 for the steaming method (1-10). Based on organoleptic test, the 75-second blanching method was a processing method that had the highest average score.




Fig.1: Selection of Withering Methods

Nutrient Content

The material used in this study was Robusta coffee leaves which have been dried through the Japanese Style Green Tea Process (JGTP) method as seen in Table 1 .

Table 1: Nutrient Content of Dried Robusta Coffee Leaves

Proximate Component	Mean±SD (%)
Moisture	3.1±0.00
Ash	6.03±0.90
Protein	17.76±2.06
Fat	2.37±0.25
Carbohydrate	70.83±2.83

Note : SD = standard deviation

Bioactive Components (total flavonoids, chlorogenic acid, caffeine)

The results of the analysis showed that the content of bioactive components, as total flavonoids, chlorogenic acid and caffeine in Robusta coffee leaves as seen in Table.

Table 2: The Content of Bioactive Components

Treatment	Total flavonoids (mg QE/g)	Chlorogenic acid (mg/g)	Caffeine (mg/g)
	Mean±SD	Mean±SD	Mean±SD
Dried coffee leaves	0.81 ± 0.02	16.58 ± 0.48	0.456 ± 0.08
5 g of brewed coffee leaves	0.10± 0.01	3.49 ± 1.62	0.194 ± 0.00

Note: 5 g of dried coffee leaves were brewed with 200 mL of water at 70°C for 10 minutes

Total Chlorophyll

The results of chlorophyll analysis showed a significant difference in the groups of brewed coffee leaves (p = 0.000). While the results of chlorophyll analysis showed significant differences in the coffee leaves group (p = 0.014) and brewed coffee leaves (p = 0.000). The results of total chlorophyll analysis showed significant

differences in each coffee leaves group (p = 0.006) and brewed coffee leaves group (p = 0.000). There were significant differences in chlorophyll a, chlorophyll b and total chlorophyll between coffee leaves and brewed coffee leaves (p = 0,000) as seen in Antioxidant Activity. The results of the analysis showed that there were no significant differences in the Robusta coffee

leaves group (p = 0.206) and brewed (p =

0.131) as seen in Table 3.

The results of the analysis showed that there were no significant differences in the Robusta coffee leaves group (p = 0.206) and brewed (p = 0.131) as seen in Table 3.

Table 2.

Antioxidant Activity

Table 2: Total Chlorophyll

Samples	Chlorophyll a		Chlorophyll b		Total Chlorophyll	
	Dried coffee leaves Mean ± SD	Brewed coffee leaves Mean ± SD	Dried coffee leaves Mean ± SD	Brewed coffee leaves Mean ± SD	Dried coffee leaves Mean ± SD	Brewed coffee leaves Mean ± SD
5 g	23.66±0.35 ^a	0.74±0.04 ^a	39.98±2.37 ^a	1.35±0.11 ^a	63.62±2.01 ^a	2.08±0.14 ^a
10 g	23.75±0.17 ^a	1.16±0.09 ^b	42.65±0.63 ^{ab}	2.19±0.13 ^b	66.37±0.67 ^{ab}	3.35±0.22 ^b
15 g	23.97±0.39 ^a	1.73±0.31 ^c	43.17±0.61 ^{ab}	3.19±0.44 ^c	67.13±0.99 ^{ab}	4.91±0.75 ^c
20 g	24.66±0.73 ^a	2.12±0.02 ^c	44.76±0.80 ^b	3.91±0.52 ^d	69.40±1.45 ^b	6.01±0.09 ^d
P ^x	0.098	0.000*	0.014*	0.000*	0.006*	0.000*

Note: One-way-anova.test. Dried coffee leaves were brewed with 200 mL of water at 70°C for 10 minutes.. *Significant (p < 0.05). Numbers followed by superscript letters (a,b,c,d) differ to show significant differences.

Table 3: Antioxidant Activity with DPPH Method

Treatment	Inhibition (%)		p
	Dried coffee leaves Mean ± SD(%)	Brewed coffee leaves Mean ± SD (%)	
5 gram	92.7 ± 0.61	84.9 ± 0.23	0.109 ^p
10 gram	91.8 ± 0.76	86.1 ± 0.85	0.019 ^q
15 gram	92.0 ± 2.00	85.6 ± 1.20	0.021 ^q
20 gram	90.2 ± 1.36	83.3 ± 1.47	0.000 ^q
p	0,206 ^z	0,131 ^y	

Note: Dried coffee leaves were brewed with 200 mL of water at 70°C for 10 minutes. *Significant (p < 0,05). ^p Wilcoxon test, ^q Paired t test, ^z One way Anova, ^y Kruskal Wallis test

Antibacterial Activity Test

The results of the antibacterial activity analysis in *Escherichia coli* and *Salmonella typhi* showed

significant differences in the groups of brewed coffee leaves (p = 000) as seen in Table 4.

Table 4: Antibacterial Activity

Samples	<i>Escherichia coli</i>		<i>Salmonella typhi</i>	
	Mean ± SD	Inhibition	Mean ± SD	Inhibition
5 g	7.65±0.10 ^a	Medium	7.47±0.28 ^a	Medium
10 g	8.47±0.15 ^b	Medium	8.20±0.05 ^b	Medium
15 g	9.17±0.15 ^c	Medium	8.55±0.26 ^b	Medium
20 g	9.89±0.26 ^d	Medium	8.70±0.19 ^c	Medium
p	0,000*		0,000*	

*One Way Anova test. *Significant (p < 0.05). Numbers followed by superscript letters (a,b,c,d) differ to show significant differences.

Acceptability of Brewed Coffee Leaves

There was a significant difference in the color and Taste of brewed coffee leaves (p = 0,000) and

did not show a significant difference in the aroma of brewed coffee leaves (p = 0.392) as seen in

Table 5.

Table 5: Acceptability of Brewed Coffee Leaves

Treatment	Color		Aroma		Taste	
	Median	Note	Median	Note	Median	Note
5 g	3(1-4) ^a	Like slightly	3(1-4) ^a	Like slightly	3(2-4) ^a	Like slightly
10 g	3(2-4) ^{ab}	Like slightly	3(2-4) ^a	Like slightly	2(1-3) ^a	Dislike slightly
15 g	2,5(1-4) ^{bc}	Dislike slightly	3(1-4) ^a	Like slightly	1(1-4) ^b	Dislike very much
20 g	2(1-4) ^c	Dislike slightly	3(1-4) ^a	Like slightly	1(1-2) ^c	Dislike very much
<i>p</i>	0,000*		0,392		0,000*	

Note: Kruskal Wallis test. *Significant ($p < 0.05$). Numbers followed by superscript letters (a,b,c,d) differ to show significant differences.

DISCUSSION

In this study, Robusta coffee leaves were processed by the Japanese Style Green Tea Process (JGTP). JGTP method is one of the tea processing without fermentation process. Previous study has explained that the JGTP method in Arabica coffee leaves has higher phytochemicals concentration compared to other tea-making methods.(X.-M. Chen et al., 2018). The blanching process was used at the withering stage. Blanching can deactivate the enzyme polyphenol oxidase (PPO) which can cause oxidation.(Pourcel, Routaboul, Cheynier, Lepiniec, & Debeaujon, 2007)

Oxidation is a degenerative process triggered by the presence of free radicals which causes rancidity and decreasing components in food. The oxidation process causes the loss of phenolic components of food.(X.-M. Chen et al., 2018) The blanching process increases the polyphenol content during extraction.(Nayak, Liu, & Tang, 2015) In addition, the processing of green tea maintain the composition of chlorophyll including chlorophyll a (Chl a) and chlorophyll b (Chl b) that acts as an antibacterial.(Li et al., 2018)

Based on acceptability of brewed coffee leaves, the best acceptance was 5 g of brewed coffee leaves. The more the amount of leaves that are brewed, it will produce a thick yellowish-green color. Coffee leaves products using the JGTP method do not undergo oxidation of polyphenol compounds so that the resulting color was greenish or yellowish-green. Green is the main color determined by the chlorophyll content. Chlorophyll a on the leaves will produce a dark green color and chlorophyll b will produce a yellowish-green color. In addition to the chlorophyll content, flavonol content also plays a role in producing a yellowish color on the coffee leaves by the JGTP method.(Chaturvedula & Prakash, 2011)

Based on the aroma described by panelists, the aroma of brewed coffee leaves such as the aroma of fresh leaves. The more the amount of leaves in brewed, the smell will be thicker and increasingly disliked by panelists. The aroma of the leaves is caused by the high compound of catechins.(Ho, Zheng, & Li, 2015) The taste of brewed coffee

leaves was bitter. The bitter taste is influenced by the presence of chlorogenic acid and caffeine.(Belay, 2011; Gokul G & Lakshmanan, 2016) Caffeine is a natural compound in food and drinks that contributes to affect the bitter taste. Some studies show that the main taste of caffeine is bitter. Caffeine generally amounts to 2-5% of the dry weight of tea. Caffeine will cause a bitter taste on the tongue depending on the level of concentration.(Keast & Roper, 2007; Zou, Xiao, Wang, & Zhang, 2018) The study explained that the bitter taste began to emerge when the caffeine content was 75-155 mg/L. Chlorogenic acid also plays a role in acidity and bitterness.(Belay, 2011)

The moisture and ash content in dried Robusta coffee leaves processed by the JGTP method met the Indonesian National Standard (SNI) green tea requirements, where the moisture of green tea is set to a maximum of 8% and ash content of 4-8%.(SNI, 2016) The moisture in accordance with SNI can prevents the growth of bacteria and fungi, and prevent oxidation process. Protein content in dried Robusta coffee leaves with JGTP method included in the range of green tea, where protein content in green tea about 15-20%. Protein has the potential as an additional antioxidant in food because it can inhibit fat oxidation through several pathways including inactivation of Reactive Oxygen Species (ROS) through antioxidant enzymes (superoxide dismutase (SOD) and catalase), scavenge free radicals, chelating prooxidative transition metals, reducing hydroperoxide, and reducing hydroperoxides, and enzymatically removes specific oxidants.(Elias, Kellerby, & Decker, 2008) Protein also acts as an antimicrobial. Previous study explained that protein extracted from lemon, orange and grapefruit seeds could inhibit the growth of some pathogenic bacteria.(Karabiber, Yilmaz, & Zorba, 2018)

The fat content in Robusta coffee leaves processed by the JGTP method was lower than in Arabica coffee leaves with the same method. Based on previous research, Arabica coffee leaves have higher fat content about 4.5-12.5%. Meanwhile, the carbohydrate content in Robusta coffee leaves was higher than in Arabica coffee

leaves. In previous study, carbohydrate content of Arabica coffee leaves was about 51.99-63.93%. (Woldesenebet, 2005)

In this study, Robusta coffee leaves processed by JGTP method have bioactive components of flavonoids, chlorogenic acid, caffeine, and chlorophyll. Previous study has shown that coffee leaves contain flavonoids such as quercetin, rutin, and kaempferol. (Martins, Araújo, Tohge, Fernie, & DaMatta, 2014) Flavonoids can act as antioxidants to scavenge free radicals, prevent the formation of reactive oxygen species (ROS), and increase endogenous antioxidants (superoxide dismutase, catalase, and glutathione peroxidase). The mechanism of flavonoids in preventing damage by free radicals is by giving hydrogen atoms to free radicals to form stable flavonoxyl radicals (flavonoids (O•)). In addition, flavonoids can protect the phospholipid membrane by giving one of the hydrogen ions to the radical lipid peroxyl, so that radical reactions can be stopped. (Hamid, Usman, Aiyelaagbe, & Ameen, 2010; Irina & Mohamed, 2012) Flavonoids can also function as antibacterial by inhibiting nucleic acid synthesis, inhibiting energy metabolism, changes in cytoplasmic membrane function, reducing cell adhesion and biofilm formation and causing damage to the cytoplasmic membrane. (Farhadi et al., 2018)

Chlorogenic acid is an ester formed from hydroxycinnamic acid (caffeic acid, ferulic acid, p-kumaric acid) and quinic acid and into the phenolic acid group. (Liang & Kitts, 2016) Chlorogenic acid can chelate transition metals like Fe²⁺ to scavenge free radicals and break radical chains. Chlorogenic acid scavenges DPPH radicals, superoxide anions, hydroxyl radicals, and protect DNA from damage caused by oxidative stress. (Azam, Hadi, Khan, & Hadi, 2003) Chlorogenic acid also functions as antibacterial. Chlorogenic acid is a phenolic bioactive compound that can cause physiological changes in bacterial cell membranes. (Kabir et al., 2014) Chlorogenic acid causes cell membrane rupture and release of the content inside the cell. Chlorogenic acid changes in bacterial cell membrane permeability that will cause damage to bacterial nucleotides. (Lou et al., 2011)

Robusta coffee leaves with processed by JGTP method also contained caffeine. Caffeine prevents lipid oxidation by inhibiting the production of free radicals including hydroxyl radicals, peroxyl radicals, and singlet oxygen. (Azam et al., 2003) Caffeine also acts as antibacterial agent. It can inhibit protein and DNA synthesis by inhibiting the merging of adenine and thymidine. Besides, caffeine also

increases genotoxicity after DNA damage. (Pruthviraj et al., 2011)

The amount of chlorophyll in brewed Robusta coffee leaves was very low compared to the amount of chlorophyll in tea leaves. Chlorophyll acts as antioxidants, where it can prevent the Fenton reaction, by chelating Fe(II). Chlorophyll directly scavenges the hydroxyl free radicals generated from H₂O₂. (Hsu, Chao, Hu, & Yang, 2013) It can also act as antibacterial by increasing membrane permeability in bacteria. (Kustov et al., 2018) This will cause changes in membrane structure and accelerate intracellular leakage thereby destroying membrane integrity which will facilitate the entry of antibacterial. (Wu et al., 2016)

The results showed that Robusta coffee leaves with the JGTP method had a DPPH radical inhibition. Based on the results of this analysis, Robusta coffee leaves and brewed coffee leaves have high antioxidant activity, being able to capture free radicals by 50%. The results of this study are in line with research on green tea, where green tea has a DPPH radical inhibition more than 50%.

The results of antibacterial of brewed coffee leaves showed the inhibitory power of 5 g, 10 g, 15 g, and 20 g brewed Robusta coffee leaves against *Escherichia coli* and *Salmonella typhi* bacteria. *Escherichia coli* and *Salmonella typhi* belong to gram-negative bacteria. The greater the amount of leaves in the brewed process, the greater the diameter of the inhibition zone formed. The effectiveness of antibacterial substances will be influenced by the concentration of these substances. An increase in the concentration of the substance causes an increase in the content of active compounds such as flavonoids, caffeine and chlorogenic acid which functions as an antibacterial so that its ability to inhibit bacterial growth increases. (Mufti, Bahar, & Arisanti, 2017; Muslim & Dephinto, 2017) Previous study conducted on CTC brewed black tea with a comparison of tea weight and water volume (w/v) 1:50, 2:50, and 3:30 resulted in an area of 29.89 successive barriers; 33.64 and 45.02 mm². (Rohdiana et al., 2013)

CONCLUSION

Robusta coffee leaves in this research were processed by JGTP method with 75-second blanching. The best organoleptic based on color, aroma, and taste was 5 g of dried Robusta coffee leaves. Brewed Robusta coffee leaves processed by the Japanese Style Green tea Process (JGTP) have contained flavonoids 0.10 mg QE/g, chlorogenic acid 3.49 mg/g, caffeine 0.194 mg/g and high antibacterial activity in 20 g of brewed Robusta coffee leaves.

CONFLICT OF INTEREST

No conflict of interest.

ACKNOWLEDGMENTS

This research was supported by Research and Development (RPP) from the Faculty of Medicine, Universitas Diponegoro.

REFERENCES

1. Analysis., O. M. of. (2005). *Official Methods of Analysis*. (18th ed.). AOAC International: USA.
2. Azam, S., Hadi, N., Khan, N. U., & Hadi, S. M. (2003). Antioxidant and Prooxidant Properties of Caffeine, Theobromine and Xanthine. *Med Sci Monit.*, 9(9), 325–331.
3. Belay, A. (2011). Some biochemical compounds in coffee beans and methods developed for their analysis. *International Journal of the Physical Sciences*, 6(28), 6373–6378.
4. Campa, C., Urban, L., Mondolot, L., Fabre, D., Roques, S., Lizzi, Y., ... Etienne, H. (2017). Juvenile Coffee Leaves Acclimated to Low Light Are Unable to Cope with a Moderate Light Increase. *Front Plant Sci.*, 8(1126), 1–16.
5. Chacko, S. M., Thambi, P. T., Kuttan, R., & Nishigaki, I. (2010). Beneficial effects of green tea: A literature review. *Chin Med*, 5(15), 1–9.
6. Chaturvedula, V. S. P., & Prakash, I. (2011). The aroma, taste, color and bioactive constituents of tea. *Journal of Medicinal Plants Research*, 5(11), 2110–2124.
7. Chen, X. (2019). A Review on Coffee Leaves: Phytochemicals, Bioactivities and Applications. *Crit Rev Food Sci Nutr*, 59(6), 1008–1025. <https://doi.org/10.1080/10408398.2018.1546667>
8. Chen, X.-M., Ma, Z., & Kitts, D. D. (2018). Effects of processing method and age of leaves on phytochemical profiles and bioactivity of coffee leaves. *Food Chemistry*, 249, 143–153. <https://doi.org/10.1016/j.foodchem.2017.12.073>
9. Elias, R. J., Kellerby, S. S., & Decker, E. A. (2008). Antioxidant Activity of Proteins and Peptides. *Critical Reviews in Food Science and Nutrition*, 48(5), 430–441.
10. Farhadi, F., Khameneh, B., Iranshahi, M., & Iranshahi, M. (2018). Antibacterial activity of flavonoids and their structure – activity relationship: An update review. *Phyther Res*, 1–28.
11. Gokul G, & Lakshmanan, R. (2016). Effect Of Chlorhexidie Mouthwash On Taste Alteration. *Asian J Pharm Clin Res*, 9(1), 102–104.
12. Hamid, A. A., Usman, L. A., Aiyelaagbe, O., & Ameen, M. O. (2010). Antioxidants: Its Medicinal and Pharmacological Applications. *African J Pure Appl Chem*, 4(8), 142–151.
13. Hasanah, M., Maharani, B., & Munarsih, E. (2017). Daya Antioksidan Ekstrak Dan Fraksi Daun Kopi Robusta (*Coffea robusta*) terhadap Pereaksi DPPH (2,2-difenil-1-pikrilhidrazil) (Antioxidant of Extract and Fraction *Coffea robusta* Leaves with Diphenylpicrylhydrazyl (DPPH) Method. *Indonesian Journal of Pharmaceutical Science and Technology*, 4(2), 42–49.
14. Ho, C.-T., Zheng, X., & Li, S. (2015). Tea Aroma Formation. *Food Science and Human Wellness*, 4(1), 9–27.
15. Hsu, C.-Y., Chao, P.-Y., Hu, S.-P., & Yang, C.-M. (2013). The Antioxidant and Free Radical Scavenging Activities of Chlorophylls and Pheophytins. *Food and Nutrition Sciences*, 4, 1–18.
16. Irina, I., & Mohamed, G. (2012). Biological Activities and Effects of Food Processing on Flavonoids as Phenolic Antioxidants. In *Advances in Applied Biotechnology*. (pp. 112–114). Croatia.
17. Kabir, F., Katayama, S., Tanji, N., & Nakamura, S. (2014). Antimicrobial Effects of Chlorogenic Acid and Related Compounds. *Journal of the Korean Society for Applied Biological Chemistry*, 57(3), 359–365.
18. Kanisius, A. A. (1988). *Budidaya Tanaman Kopi (Coffee Cultivation)* (16th ed.). Yogyakarta: Kanisius.
19. Karabiber, E., Yilmaz, E., & Zorba, N. N. D. (2018). antimicrobial and functional properties of the proteins extracted from lemon, orange and grapefruit seeds press meals. *Quality Assurance and Safety of Crops & Foods*, 10(2), 1–10.
20. Keast, R. S. J., & Roper, J. (2007). A Complex Relationship Among Chemical Concentration, Detection Threshold, and Suprathreshold Intensity of Bitter Compounds. *Chem Senses*, 32(3), 245–253.
21. Khotimah, K. (2014). Karakteristik Kimia Kopi Kawa dari Berbagai Umur Helai daun Kopi yang Diproses dengan Metode Berbeda (Chemical Characteristics of Kawa Coffee of Various Age Leaves Processed by Different Methods). *J Teknol Pertan*, 9(1), 40–45.
22. Kustov, A. V., Belykh, D. V., Smirnova, N. L., Venediktov, E. A., Kudayarova, T. V., Kruchin, S. O., ... Berezin, D. B. (2018). Synthesis and investigation of water-soluble chlorophyll pigments for antimicrobial photodynamic therapy. *Dyes and Pigments*, 149, 553–559.
23. Ky, C.-L., Louarn, J., Dussert, S., Guyot, B., Hamon, S., & Noirot, M. (2001). Caffeine, trigonelline, chlorogenic acids and sucrose diversity in wild *Coffea arabica* L. and *C. canephora* P. accessions. *Food Chemistry*, 75(2), 223–230.
24. Li, X., Zhou, R., Xu, K., Xu, J., Jin, J., Fang, H., & He, Y. (2018). Rapid Determination of Chlorophyll and Pheophytin in Green Tea Using

- Fourier Transform. *Molecules*, 21, 1–13.
25. Liang, N., & Kitts, D. D. (2016). Role of Chlorogenic Acids in Controlling Oxidative and Inflammatory Stress Conditions. *Nutrients*, 6, 16. <https://doi.org/10.3390/nu8010016>.
 26. Lou, Z., Wang, H., Zhu, S., Ma, C., & Wang, Z. (2011). Antibacterial Activity and Mechanism of Action of Chlorogenic Acid. *J Food Sci.*, 76(6), 398–403.
 27. Manach, C., Scalbert, A., Morand, C., Rémésy, C., & Jiménez, L. (2004). Polyphenols: Food Sources and Bioavailability. *The American Journal of Clinical Nutrition*, 79(5), 727–747.
 28. Martins, S. C. V., Araújo, W. L., Tohge, T., Fernie, A. R., & DaMatta, F. M. (2014). n High-Light-Acclimated Coffee Plants the Metabolic Machinery Is Adjusted to Avoid Oxidative Stress Rather than to Benefit from Extra Light Enhancement in Photosynthetic Yield. *PLoS ONE*, 9(4), 1–11.
 29. Mufti, N., Bahar, E., & Arisanti, D. (2017). Uji Daya Hambat Ekstrak Daun Sawo terhadap Bakteri *Escherichia coli* secara In Vitro (Inhibition Test of Inhibition Test of Sawo Leaf Extract Against *Escherichia coli* Bacteria in Vitro). *Jurnal Kesehatan Andalas*, 6(2), 289–294.
 30. Muslim, Z., & Dephinto, Y. (2017). Perbandingan Efektivitas Antimikroba Ekstrak Daun Kopi Robusta (*Coffea canephora*) Dengan Variasi Pengeringan Terhadap *Escherichia coli* (Comparison of Antimicrobial Effectiveness of Robusta (*Coffea canephora*) Leaf Extract with Drying Variations on *Escherichia coli*). *J Sains Dan Teknol Farm*, 19, 86–88.
 31. Naegele, E. (2016a). Determination of Caffeine in Coffee Products According to DIN 20481. *Agil Technol*, 1–6.
 32. Naegele, E. (2016b). Determination of Chlorogenic Acid in Coffee Products According to DIN 10767. *Agil Technol*, 1–8.
 33. Nayak, B., Liu, R. H., & Tang, J. (2015). Effect of Processing on Phenolic Antioxidants of Effect of Processing on Phenolic Antioxidants of Fruits, Vegetables, and Grains — A Review. *Crit Rev Food Sci Nutr.*, 55(7), 887–918.
 34. Nayeem, N., Denny, G., & Mehta, S. K. (2011). Comparative phytochemical analysis, antimicrobial and anti oxidant activity of the methanolic extracts of the leaves of *Coffea Arabica* and *Coffea Robusta*. *Der Pharmacia Lettre*, 3(1), 292–297.
 35. Pourcel, L., Routaboul, J.-M., Cheynier, V., Lepiniec, L., & Debeaujon, I. (2007). Flavonoid Oxidation in Plants: from Biochemical Properties to Physiological Functions. *Trends Plant Sci.*, 12(1), 29–36.
 36. Prawira-Atmaja, M. I., Shabri, Khomaini, H. S., Maulana, H., Harianto, S., & Rohdiana, D. (2018). Changes in chlorophyll and polyphenols content in *Camellia sinensis var. sinensis* at different stage of leaf maturity. *Earth Environmental Sci*, 131, 1–7.
 37. Pruthviraj, P., Shital, K., Suchita, B., & Shilpa, K. (2011). Evaluation of Antibacterial Activity of Caffeine. *Int J Res Ayurveda Pharm*, 2(4), 1354–1357.
 38. Rhayanne T. M. Ramos, I. C. F. B., Ferreira, M. R. A., & Soares, L. A. L. (2017). Spectrophotometric Quantification of Flavonoids in Herbal Material, Crude Extract, and Fractions from Leaves of *Eugenia uniflora* Linn. *Pharmacognosy Res.*, 9(3), 253–260.
 39. Rohdiana, D., Arief, D. Z., & Budiman, A. (2013). Aktivitas penghambatan pertumbuhan bakteri *Escherichia coli* oleh berbagai jenis teh dan seduhannya. (Inhibitory activity of *Escherichia coli* by type of teas and its liquors). *Jurnal Penelitian Teh Dan Kina*, 16(1), 37–44.
 40. Sadino, A., Sahidin, I., & Wahyuni, W. (2018). Antibacterial Activity of Polygonum pulchrum Blume Ethanol Extract on *Staphylococcus aureus* and *Escherichia coli*. *Pharmacol Clin Pharm Res.*, 3, 26–32.
 41. Silva, L. A. L. da, Pezzini, B. R., & Soares, L. (2015). Spectrophotometric Determination of The Total Flavonoid Content in *Ocimum basilicum* L.(Lamiaceae) Leaves. *Pharmacogn Mag*, 11(41), 96–101.
 42. SNI. (2016). *SNI 3945:2016 Teh Hijau (Green Tea)*. Jakarta.
 43. Woldesenebet, A. (2005). Nutritional Composition, Phytochemical Screening, Processing Methods and Sensory Attributes of A Brew Made From Infusions of Matured Leaves of Arabica Coffee Tree Consumed In Sidama, Kambata and Harar Communities, Ethiopia. Addis Ababa University.
 44. Wu, Y., Bai, J., Zhong, K., Huang, Y., Qi, H., Jiang, Y., & Gao, H. (2016). Antibacterial Activity and Membrane-Disruptive Mechanism of 3-p-trans Coumaroyl-2-hydroxyquinic Acid, a Novel Phenolic Compound from Pine Needles of *Cedrus deodara*, against *Staphylococcus aureus*. *Molecules*, 21(8), 1–12.
 45. Zou, G., Xiao, Y., Wang, M., & Zhang, H. (2018). Detection of bitterness and astringency of green tea with different taste by electronic nose and tongue. *PLoS ONE*, 13(12), 1–10.