

# Gestation Length and Litter Size of New Zealand White Grade Rabbit

*by Dela Ayu Lestari*

---

**Submission date:** 14-Mar-2023 10:06AM (UTC+0700)

**Submission ID:** 2036695630

**File name:** 16003-Article\_Text-41273-47365-10-20211021.pdf (244.77K)

**Word count:** 2402

**Character count:** 11725

## Gestation Length and Litter Size of New Zealand White Grade Rabbit

A. Setiaji\*, D. A. Lestari, S. Sutopo, Y. S. Ondho and E. Kurnianto

Departement of Animal Science, Faculty of Animal and Agricultural Sciences,  
Universitas Diponegoro, Tembalang Campus,  
Semarang, 50275, Central Java, Indonesia  
Corresponding Author: [asepsetiaji@lecturer.undip.ac.id](mailto:asepsetiaji@lecturer.undip.ac.id)

9

### ABSTRACT

The purpose of this study was to examine performances and factors that influence the gestation length (GL) and litter size (LS) of New Zealand White Grade (NZW Grade) rabbits. Reproduction records of NZW Grade rabbits were taken from April 2020 to March 2021. Data analysis was carried out using the general linear model (GLM) procedure from Statistical Analysis System (SAS) University Edition V.6p.2. software. Farm showed a significant effect ( $P < 0.001$ ) on LS and significant season ( $P < 0.05$ ) on GL and LS. GL in the rainy season (31.56) was shorter than the GL in the summer (32.26), but the LS in the rainy season (7.06) was greater than the LS in the rainy season (6.37). Maintenance management should be improved to reduce the impact of the season on GL and LS NZW Grade rabbits.

**Keywords:** General linear model; gestation length; litter size; New Zealand White Grade rabbit; reproduction records.

### INTRODUCTION

The New Zealand White Grade Rabbit (NZW Grade) is a broiler-type rabbit developed by most breeders in Central Java. The NZW Grade is the result of a cross between a New Zealand White rabbit imported from the United States in the last ten years with a rabbits that have been adapted to the tropical climate. NZW Grade rabbit have fast growth rates, good carcass quality, and well adapted in tropical environments. According to these performances NZW Grade have potential for meat producing livestock.

Reproductive performance is important in determining the success of rabbits breeding. The breed of rabbits should have a high fertility and birth rates and good ability to raise their offsprings (Widitania et al., 2016). Reproductive traits that used in assessing the reproductive ability of female rabbits were gestation length, litter size, and mortality of offspring.

Litter size (LS) is a reflection of the fertility of female rabbits and their ability to take care of their offspring (McNitt et al., 2013). The average LS of rabbits in have been reported in Indonesia was range from 7.2 to 8.1 (Brahmantiyo et al., 2017). According to Lebas et al. (1997), LS will increase in the second parity by 10% -20%, then will decrease in the third parity and remain in the fourth parity further decrease in the fifth parity; and so on. LS was influenced by several factors including the breeds, mating management, the fisiologis condition of the female feed quality during pregnancy period and

Low quality of sperm tend to produce fewer offspring (Utami et al., 2019).

Gestation length (GL) of rabbits is influenced by several factors such as breeds (McNitt and Moody, 1991), climate, and weather (Ferraz et al., 1991; Farghaly, 1996). The average GL in rabbits is 32 days, ranging from 27-35 days (Ehiobu et al., 1997). McNitt et al. (1997) reported that longer GL of rabbit in summer was follow with fewer number of LS. This study aimed to examine performance and factors that influence GL and LS of NZW Grade rabbits.

### MATERIAL AND METHODS

#### Data collection

Reproduction records of female NZW Grade rabbits were obtained from 8 farms in Semarang and Kendal Regencies, Central Java Province. Data collection was carried out from April 2020 to March 2021. A total of 182 data were obtained from 74 female rabbits from first to fifth parity. The parameters observed were GL and LS. GL is period from mating to partus, while LS is the number of offspring birth in one parity.

#### Statistical analysis

Data analysis was performed using Statistical Analysis System (SAS) University Edition V.6p.2. software. General Linear Model (GLM) procedur (SAS Institute Inc., Cary, NC, USA) was used to determine the effect of farm, season, and parity as well as the interaction between them for the GL and LS.

Statistical models was used as follow:

$$y_{ijk} = \mu + S_j + P_k + e_{ijk}$$

where:  $y_{ijk}$  is the observed value of a dependent variable (GL/LS), the  $\mu$ , the overall mean,  $S_j$ , the  $j$ th effect of the season,  $P_k$ , the  $k$ th effect of parity and  $e_{ijk}$  is the random residual of  $y_{ijk}$ .

## RESULTS AND DISCUSSION

The results (Table 1) significant factors on GL presented in table 1. The season had a significant effect ( $P < 0.001$ ) on GL, while farm and parity had no statistical effect on GL. The interaction between these factors did not affect on GL and (Table 2) so there is no discussion about the interaction. The results of the analysis were in agreement with the studies reported by Boyd and Bray (1989); Ferraz et al. (1991) and Farghaly, (1996) who stated that the

reproduction of rabbits was influenced by the seasons which has an impact on gestation and lactation of female rabbit. Lebas et al. (1986), reported that season is one of the external factors that play a role in the physiological and reproductive conditions of female rabbits. Parity has no significant effect on GL but GL will be longer in older female rabbits (Tuma et al., 2010). The Least-squares means analysis of GL in the rainy season is 31.56 weeks, in the summer is 32.26 (Table 3). These results are in accordance with McNitt et al. (1997) reported that GL of rabbit in the summer is longer than which in the winter. Fayeye and Ayorinde, (2010) reported different results of local rabbits in Nigeria, where; GL of rabbits in summer (32.03) was shorter than that in the rainy season (32.62). Overall, the GL of the NZW Grade rabbit in this study was longer than the GL of the New Zealand White Purebred Rabbit (30.4) by Ghosh et al. (2003) and (31.9) by Fadare and Fatoba (2018).

Table 1. Significance of factors affecting gestation length and litter size

Factor	db	Gestation length		Litter size	
		F Value	Pr > F	F Value	Pr > F
Farm	7	1,02	0,4173	3,94	0,0005
Season	1	3,95	0,0484	6,27	0,0132
Parity	4	0,06	0,9935	of 1.52	0,1974

Description : db = degrees of freedom; Pr = probability

Farm and season were significant ( $P < 0.05$ ) on LS, while parity was not statistically significant on LS in NZW Grade rabbits. This result shows that besides being influenced by genetics, LS is strongly influenced by farm management (Table 2). Farm managements are including breeding, feeding, and housing management. Eiben et al. (2001) reported that female rabbits have been fed a controlled diet showed good reproductive performance, large

number of LS and high milk production. Mating management also affected on LS, female rabbits mated during standing heat, with characteristics swollen and red vulva was increased the conception rate and number of LS (Maertens, 1998; Szendrő et al., 2006). Rebollar et al. (2009) reported that female rabbits reared with semi-intensive systems showed better reproductive performance than in the intensive systems.

Table 2. Least squares means for gestation length and litter size on the different farm

Farm	N	GL	LS
1	38	31,89	5,36
2	24	31,85	6,89
3	37	31,31	5,96
4	11	31,85	6,01
5	17	32,63	7,11
6	15	32,19	5,07
7	26	31,29	4,88
8	14	32,34	7,30

Description : N = number of samples; GL = gestation length ; LS = litter size

Table 3. Least squares means for GL and LS on different season and parity

Factor	Category	N	GL	LS
Season	rain	104	31,56	7,06
	dry	78	32,26	6,37
Parity	1	72	31,97	6.69
	2	56	31,95	5,82
	3	31	31,76	6.11
	4	19	32,00	5,83
	5	4	31,87	5,90

Description : N = number of samples; ; GL = gestation length; LS = litter size

The Least-squares means of LS in rainy season was 7.06 while in the summer was 6.37 (Table 3). These results are in accordance with Ayyat et al. (1995) that reported number of LS in summer was fewer number of LS in the winter. Fayeye and Ayorinde, (2010) reported different results where number of LS in the summer (5,12) was larger than number of LS in the rainy season (4,4) for local rabbits in Nigeria. The LS of the NZW Grade rabbits in this study was fewer than the LS of the New Zealand White, Hyla, and Hycle Rabbits (ranged from 7.2 to 8.1) (Brahmantiyo et al., 2017).

## CONCLUSION

Reproductive management of rabbits was difficult to formulated. The results of this study can be considered, among others: farm, feeding, and mating managements should be standardized to improve the reproductive performance of female NZW Grade rabbits. Hausing manipulation such as lighting and temperature can could be done as an effort to reduce the effect of season on the reproductive performance of female rabbits.

## ACKNOWLEDGMENT

This research was funded by Hibah Dana Penelitian Universitas Diponegoro Semarang with Contract No. 36/UN7.5.5.2/PP/2021 dated March 1, 2021.

## REFERENCES

- Ayyat, M.S., Marai, I.F.M. and Gh.A. El-Sayiad. 1995. Genetic and non-genetic factors affecting milk production and preweaning litter traits of New Zealand White does under egyptian conditions. world rabbit science 3:119-124. DOI: <https://doi.org/10.4995/wrs.1995.250>
- Brahmantiyo, B., Raharjo Y.C. dan L.H. Prasetyo. 2017. Performa produksi kelinci HyCole, New Zealand White dan persilangan timbal baliknya. Jurnal Ilmu Ternak dan Veteriner. 22(1):16-23. DOI:10.14334/jitv.v22i1.1590
- Boyd, I.L. and C.J. Bray. 1989. Nutritional ecology of the wild rabbit an input to timing of reproduction. Proceeding of Nutrition Society. 48:81-91. DOI: 10.1079/pns19890012
- Ehiobu, N.G., Utim A. and S.D. Gwaza. 1997. Some observations on reproductive performance of rabbits obtained in semi-humid tropical conditions in Nigeria. World Rabbit Science. 5:47-49. DOI: <https://doi.org/10.4995/wrs.1997.317>
- Eiben, Cs., Kustos, K., Kenessey, Á., Virág Gy. and Zs. Szendrő. 2001. Effect of different feed restriction during rearing on production performance in rabbit does. World Rabbit Science. 9:9- 14. DOI:10.4995/wrs.2001.440
- Fadare, A.O. and T.J. Fatoba. 2018. Reproductive performance of four breeds of rabbit in the humid tropics. Livestock Research for Rural Development. Volume. 30: article 114. <http://www.lrrd.org/lrrd30/7/delod30114.html>
- Farghaly, H.M. 1996. Analysis of incidence of pre and post mature gestations in rabbit populations. Proceeding 6<sup>th</sup> World Rabbit Congress 9-12 July 1996. Toulouse, France. p 273-277. <http://www.publications.zu.edu.eg/Pages/PubShow.aspx?ID=4127&&pubID=2pdf>

- Fayeye, T.R. and K.L. Ayorinde. 2010. Effects of season, generation, number of mating, parity and doe number of teat on doe and litter birth characteristics in domestic rabbit. *Research Journal of Animal and Veterinary Sciences*, 5: 6-9. <http://www.aensiweb.net/AENSIWEB/rjavs/rjavs/2010/6-9.pdf>
- Ferraz, J.B.S., Johnson R.K. and J.P. Eler. 1991. Effect of inbreeding on reproduction traits of Californian and New Zealand White rabbits. *Journal of applied Rabbit Research*. 15:211-217.
- Ghosh S.K., Das A., Bujarbaruah K.M., Das Asit., Dhiman K.R. and N.P. Singh. 2008. Effect of breed and season on rabbit production under subtropical climate. *World Rabbit Science*. 16:29-33. DOI: <https://doi.org/10.4995/wrs.2008.638>
- Lebas, F., P. Coudert, R. Rouvier and H.D. Rochambeau. 1986. The rabbit husbandry, health and production. Food and Agriculture Organization of The United Nation, Rome. Italy. [https://www.researchgate.net/publication/38977233\\_The\\_Rabbit\\_Husbandry\\_Health\\_and\\_Production](https://www.researchgate.net/publication/38977233_The_Rabbit_Husbandry_Health_and_Production)
- Lebas, F., P. Coudert, H. De Rochambeau, and T.G. Thebault. 1997. The Rabbit, Husbandry, Health and Production, New Revised Version. Food and Agriculture Organization of The United Nation, Rome. [https://www.doc-developpement-durable.org/file/Elevages/Lapins/The%20Rabbit\\_Husbandry\\_health%20and%20protection\\_FAO.pdf](https://www.doc-developpement-durable.org/file/Elevages/Lapins/The%20Rabbit_Husbandry_health%20and%20protection_FAO.pdf)
- McNitt, J. I. and G.L. Moody. 1991. Gestation length of four medium breeds of rabbits in Louisiana. *Journal of applied Rabbit Research*. 14:80-82.
- McNitt, J. I., Mellad K.E., Simon G., Negatu Z. and S.D. Lukefahr. 1997. Efficacy of prostaglandin F<sub>2a</sub> and its analogs in enhancing reproductive efficiency of doe rabbits. *World Rabbit Science*. 5:155-159. DOI: <https://doi.org/10.4995/wrs.1997.336>
- McNitt, J. I., Lukefahr, S. D., Cheeke, P. R. and N.M. Patton. 2013. *Rabbit production*, 9th Ed. CABI, Wallingford. U.K.
- Rebollar, P.G., Pérez-Cabal, M.A., Pereda, N., Lorenzo, P.L., Arias-Álvarez, P. and P. García-Rebollar. 2009. Effects of parity order and reproductive management on the efficiency of rabbit productive systems. *Livestock Science*, 121:227-233. <https://doi.org/10.1016/j.livsci.2008.06.018>
- SAS. 2011. *SAS/STAT User's Guide* (Release 9.3). SAS Inst. Inc., Cary NC, USA. Available at <https://citeseerx.ist.psu.edu/viewdoc/download?doi=10.1.1.226.6407&rep=rep1&type=pdf> Accessed January 2021.
- Tůma, J., E. Tůmová, E. and V. Valášek. 2010. The effect of season and parity order on fertility of rabbit does and kit growth. *Czech Journal of Animal Science*. 55:330-336. <https://doi.org/10.17221/317/2009-CJAS>
- Utami, P., Samsudewa, D. dan C.M.S. Lestari. 2019. Pengaruh perbedaan sistem perkawinan terhadap lama bunting dan litter size Kelinci New Zealand White. <https://doi.org/10.31186/jspi.id.14.1.70-74>
- Widitania, S. Ondho, Y.S. dan C.M.S. Lestari. 2016. Korelasi antara bobot badan induk dengan litter size, bobot lahir dan mortalitas anak kelinci New Zealand White. *Jurnal Ilmu-ilmu Peternakan*. 26(2):42-48. <http://dx.doi.org/10.21776/ub.jiip.2016.026.02.6>

# Gestation Length and Litter Size of New Zealand White Grade Rabbit

## ORIGINALITY REPORT

14%

SIMILARITY INDEX

11%

INTERNET SOURCES

7%

PUBLICATIONS

0%

STUDENT PAPERS

## PRIMARY SOURCES

- |   |   |    |
|---|---|----|
| 1 | <a href="http://publikasi.polije.ac.id">publikasi.polije.ac.id</a><br>Internet Source   | 2% |
| 2 | <a href="http://riunet.upv.es">riunet.upv.es</a><br>Internet Source   | 1% |
| 3 | <a href="http://www.reproduction-online.org">www.reproduction-online.org</a><br>Internet Source   | 1% |
| 4 | Samah Abdel-Rahman, Gamal Ashour, Noha Abd El-Azeem, Safaa Barakat, Dessouki M, Mehaisen G.. "PROTECTIVE EFFECT OF N-ACEETYL CYSTEINE AGAINST OXIDATIVE STRESS IN RELATION TO SEMEN CHARACTERISTICS IN NEW ZEALAND WHITE RABBITS", Egyptian Journal of Animal Production, 2022<br>Publication | 1% |
| 5 | <a href="http://repository.ub.ac.id">repository.ub.ac.id</a><br>Internet Source   | 1% |
| 6 | alaa desouky. "Influence of Genotypes, Season of Birth, Parity Order and the Interactions between Them on Litter Traits   | 1% |

and Body Weight Measurements of Rabbits",  
Annals of Agricultural Science, Moshtohor,  
2021

Publication

---

7	<a href="https://repository.stikim.ac.id">repository.stikim.ac.id</a> Internet Source	1 %
8	<a href="https://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a> Internet Source	1 %
9	<a href="https://era.ed.ac.uk">era.ed.ac.uk</a> Internet Source	1 %
10	<a href="https://ijlr.org">ijlr.org</a> Internet Source	1 %
11	<a href="https://erl.ucc.edu.gh:8080">erl.ucc.edu.gh:8080</a> Internet Source	1 %
12	<a href="https://hdl.handle.net">hdl.handle.net</a> Internet Source	1 %
13	<a href="https://www.mdpi.com">www.mdpi.com</a> Internet Source	1 %
14	I. F. M. Marai, A. A. M Habeeb, A. E. Gad. "Reproductive traits of female rabbits as affected by heat stress and lighting regime under subtropical conditions of Egypt", Animal Science, 2016 Publication	1 %
15	<a href="https://revista-agroproductividad.org">revista-agroproductividad.org</a> Internet Source	1 %

---



---

16

safaa barakat. "GROWTH PERFORMANCE AND  
PHYSIOLOGICAL RESPONSES OF GROWING  
RABBITS SUPPLEMENTED WITH GREEN  
COFFEE EXTRACT", Egyptian Poultry Science  
Journal, 2021

Publication

1 %

---

17

[www.biobasedeconomy.eu](http://www.biobasedeconomy.eu)

Internet Source

1 %

---

---

Exclude quotes      On

Exclude matches      < 1%

Exclude bibliography      On