

SERTIFIKAT

Kementerian Riset dan Teknologi/
Badan Riset dan Inovasi Nasional



Petikan dari Keputusan Menteri Riset dan Teknologi/
Kepala Badan Riset dan Inovasi Nasional
Nomor 200/M/KPT/2020
Peringkat Akreditasi Jurnal Ilmiah Periode III Tahun 2020
Nama Jurnal Ilmiah
Bulletin of the Marine Geology
E-ISSN: 25278843

Penerbit: Pusat Penelitian dan Pengembangan Geologi Kelautan – ESDM

Ditetapkan sebagai Jurnal Ilmiah

TERAKREDITASI PERINGKAT 2

Akreditasi Berlaku selama 5 (lima) Tahun, yaitu
Volume 35 Nomor 1 Tahun 2020 sampai Volume 39 Nomor 2 Tahun 2024

Jakarta, 23 Desember 2020

Menteri Riset dan Teknologi/
Kepala Badan Riset dan Inovasi Nasional
Republik Indonesia,




Bambang P. S. Brodjonegoro

ISSN 1410-6175

BULLETIN OF THE MARINE GEOLOGY

VOLUME 36. NUMBER 1 . JUNE 2021



BULL MGI

VOL. 36

NO.1

PAGE 1-56

BANDUNG, JUNE 2021

ISSN 1410-6175

Accredited : RISTEKDIKTI No. 21/E/KPT/2018



MARINE GEOLOGICAL INSTITUTE

RESEARCH AND DEVELOPMENT AGENCIES FOR ENERGY AND MINERAL RESOURCES
MINISTRY OF ENERGY AND MINERAL RESOURCES

PUSAT PENELITIAN DAN PENGEMBANGAN GEOLOGI KELAUTAN

BADAN PENELITIAN DAN PENGEMBANGAN ENERGI DAN SUMBER DAYA MINERAL
KEMENTERIAN ENERGI DAN SUMBER DAYA MINERAL

BULLETIN OF THE MARINE GEOLOGY

Vol. 36, No. 1, June 2021

INSURED EDITOR

Director of Marine Geological Institute

CHIEF OF EDITORIAL BOARD

Prof. Dr. Ir. Hananto Kurnio, M. Sc.

VICE CHIEF OF EDITORIAL BOARD

Dr. Luli Gustiantini, S.T., M.T.

EDITORIAL BOARDS

Dr. Ir. Susilohadi (*Marine Geological Institute of Indonesia*)
Dr. Ir. Noor Cahyo Dwi Aryanto, M.T. (*Marine Geological Institute of Indonesia*)
Imam Setiadi, S.Si., M.T. (*Marine Geological Institute of Indonesia*)
Ir. Immaculata Christiana, MT. (*Marine Geological Institute of Indonesia*)
Ir. Mustafa Hanafi, M.Si. (*Marine Geological Institute of Indonesia*)
Ir. Joni Widodo, M.Si. (*Marine Geological Institute of Indonesia*)
Tumpal Bernhard Nainggolan, S.T., M.T. (*Marine Geological Institute of Indonesia*)
Dra. Ai Yuningsih (*Marine Geological Institute of Indonesia*)
Nazar Nurdin, S.T., M.T. (*Marine Geological Institute of Indonesia*)
Yulinar Firdaus, S.Si., M.T. (*Marine Geological Institute of Indonesia*)
Siti Marina, S.T., M.Phil. (*Marine Geological Institute of Indonesia*)
Drs. Judy Muliawan Eddy (*Marine Geological Institute of Indonesia*)

SCIENTIFIC REVIEWERS

Dr. Eng. Totok Suprijo, (*Institute of Technology Bandung*)
Dr. Eddy Hartantyo, M.Si. (*Gadjah Mada University*)
Dr. Hermes Panggabean, M.Sc. (*Geological Agency*)
Dr. Muhammad Ma'ruf Mukti (*Research Center for Geotechnology*)
Dr. Ir. Baskoro Rohadi, M.T. (*Diponegoro University*)

PUBLISHER BOARDS

Adithya Kandiawan S.P., M.M.
Slamet
Swasty Aninda Piranti, ST.
Dwinanda Pratya Annisa M., S.pd.
Dery Rochiman, A.Md.

GRAPHIC DESIGN

Arif Suprayitno, S.Kom., M.Kom.

For communication of this publications, please contact :

MARINE GEOLOGICAL INSTITUTE

Dr. Junjunan 236, Bandung-40174, Indonesia
Telephone : +62-22-6032020, 6032201, Fax : +62-22- 6017887
E-mail : ejournal.p3gl@gmail.com

BULLETIN OF THE MARINE GEOLOGY

Vol. 36, No. 1, June 2021

CONTENTS

QFL And Litho Facies: Predicting Reservoir Quality Of The Middle Miocene Deep-Water Facies At Kutei And North Makassar Basins

Kuntadi Nugrahanto, Ildrem Syafri, Budi Muljana-----1-14

Distribution Of Subsurface Sedimentation As A Potential Mineral Placer Deposits In South Bintan Island Waters

Muhammad Zulfikar, Nazar Nurdin, Noor Cahyo Dwi Ariyanto, Ildrem Syafri, Budi Mulyana, Andi Agus Nur

----- 15-26

Assessment Of Potential Marine Current Energy In The Straits Of The Lesser Sunda Islands

Ai Yuningsih, Mario D. Saputra ----- 27-36

The Seawater And Freshwater Influence On Expansivity Behaviors Of Clay Minerals

Dian Agus Widiarso, Nurakhmi Qadaryati, Wiyatno Haryanto ----- 37-44

PETROPHYSICAL ANALYSIS AND SEISMIC STRATIGRAPHY INTERPRETATION TO DETERMINE HYDROCARBON RESERVOIR IN TARAkan BASIN, BUNYU ISLAND WATERS

Daffa Dzakwan Shiddiq, Eleonora Agustine, Tumpal Bernhard Nainggolan, Imam Setiadi, Shaska Zulivandama ----- 45-56

THE SEAWATER AND FRESHWATER INFLUENCE ON EXPANSIVITY BEHAVIORS OF CLAY MINERALS

PENGARUH AIRLAUT DAN AIRTAWAR TERHADAP DAYA KEMBANG SUSUT MINERAL LEMPUNG

Dian Agus Widiarso¹, Nurakhmi Qadaryati¹, Wiyatno Haryanto²

¹Department of Geological Engineering, Universitas Diponegoro, Jl. Prof. Soedharto, SH., Tembalang, Semarang, Indonesia

²Indonesian Society of Geotechnical Engineering, Jakarta, Indonesia

Corresponding author: dianagus@lecturer.undip.ac.id

(Received 07 July 2021; in revised from 22 July 2021; accepted 22 September 2021)

ABSTRACT: Semarang City subsurface geology is characterized by an intercalating of loam-silt and clayey units. The behavior of clay materials are their expansivity potency of their volume when in contact with water. Some problems in the presence of tidal flooding appears when the seawater ingress to the shoreland that causes severe damage to infrastructures. This research attempts to reveal the influence of both seawater and freshwater on the expansivity behaviors of the clayey materials based on its mineral composition. In order to observe expansivity potency of clay minerals, 28 samples from four drill hole samples were soaked in seawater and freshwater for twenty-four hours. The initial volume of each samples were then compared with the samples volume after being soaked. X-Ray Diffraction (XRD) analysis was also conducted on selected samples to determine their mineral composition. The ANOVA test was then introduced to distinguish the influence of certain mineral types and composition to the clay expansivity behaviors. Confidence level and alpha (α) are about ninety-five percent and about five percent, respectively. The result of this research has proven that montmorillonite is clay mineral type that is most affected by expansivity behaviors when immersed in seawater and freshwater compared to kaolinite and illite minerals.

Keywords: Clay minerals, montmorillonite, expansivity, seawater, freshwater, tidal flooding.

ABSTRAK: Kota Semarang tersusun dari satuan lanau dan lempung. Mineral lempung memiliki sifat kembang susut dari segi volumenya ketika mengalami kontak dengan air. Permasalahan yang timbul akibat terjadinya peristiwa banjir pasang di perkotaan, adalah masuknya air laut menuju ke pantai dan daratan yang dapat merusak infrastruktur. Penelitian ini bertujuan untuk menunjukkan dampak yang ditimbulkan oleh air tawar maupun air laut terhadap kembang susut mineral lempung berdasarkan komposisi mineralnya. Untuk mengamati perubahan volume dalam mineral lempung, sebanyak 28 sampel dari empat sampel sumur bor direndam dalam air tawar dan air laut selama 24 jam. Kemudian volume awal setiap sampel dibandingkan dengan volume sampel sesudah dilakukan perendaman. Uji X-Ray Diffraction (XRD) juga dilakukan pada sampel pilihan untuk mengetahui komposisi mineral sampel tersebut. Uji ANOVA dengan tingkat kepercayaan 95% dan alpha (α) 5% diterapkan untuk mengetahui pengaruh tipe dan komposisi mineral lempung tertentu terhadap kembang susut yang terjadi. Penelitian ini berhasil membuktikan bahwa montmorillonit merupakan jenis mineral lempung yang paling terpengaruh oleh sifat kembang susut di daerah penelitian ketika direndam dengan air tawar dan air laut, dibandingkan dengan kaolinit dan illit.

Kata Kunci: Mineral lempung, montmorillonit, kembang susut, air laut, air tawar, banjir pasang

QFL AND LITHO FACIES: PREDICTING RESERVOIR QUALITY OF THE MIDDLE MIOCENE DEEP-WATER FACIES AT KUTEI AND NORTH MAKASSAR BASINS

QFL DAN FASIES BATUAN: PREDIKSI KUALITAS RESERVOIR FASIES LAUT DALAM BERUMUR MIOSEN TENGAH DI CEKUNGAN KUTEI DAN MAKASSAR UTARA

Kuntadi Nugrahanto^{1,2}, Ildrem Syafri¹, and Budi Muljana¹

¹ Postgraduate Program, Faculty of Geological Engineering, the University of Padjadjaran

² Upstream Subholding Pertamina Hulu Energi (PHE)

Corresponding author: kuntadi19001@mail.unpad.ac.id; kuntadi.nugrahanto@pertamina.com

(Received 21 June 2021; in revised form 16 June 2021; accepted 08 July 2021)

ABSTRACT: As it is widely known the oil and gas wellbores offshore Kutei and North Makassar have not optimally penetrated the objective strata, which is the Middle Miocene's deep-water reservoirs. Therefore, evaluating the quality of these reservoirs with onshore dataset then comparing them with the proven Late Miocene's deep-water producing reservoirs had been very fundamental. The study focuses on the assessment of QFL (Quartz, Feldspar and Lithic fragments) and sandstones litho-facies based on the rock samples from conventional-core and side-wall core, and well-logs data from forty wells onshore and offshore. These rock samples are bounded by the key biostratigraphy intervals of M40M33, M45M40, M50M45 (Middle Miocene), and M65M50, M66M65, M70M66, M80M70 (Late Miocene). Subdivisions of the reservoirs considered the sandstone litho facies, NTG ratio, sorting, and grain size, to come up with five groups in the Middle Miocene deltaic facies: FLU_SX, DC_SX, DC_SM, DC_SM, and DF_SC; and four groups in the Late Miocene deep-water facies: SSWS, MSWS, SSPS, and MSPS. Core-based porosity and permeability further explain the relationship between the reservoir quality with the sandstones' composition and litho facies, and concluded that high-energy depositional system is mainly associated with the FLU_SX, DC_SX, SSWS and MSWS being the reservoir with best quality. Oppositely, the DF_SC, SSPS, and MSPS are classified the reservoir with worst to none quality. A cross plot between core-based porosity and maximum burial depth is able to postulate the relational trend of decreasing reservoir quality with deeper depth.

Keywords: Middle Miocene, deep water, QFL, litho facies, Kutei, North Makassar

ABSTRAK: Telah umum diketahui bahwa sumur bor minyak dan gas di lepas pantai Kutei dan Makassar Utara belum secara optimal menembus reservoir fasies laut dalam berumur Miosen Tengah, yang merupakan obyekti utama dalam penelitian ini. Oleh karenanya penulis melakukan studi banding antara reservoir obyektif yang umumnya berlokasi di daratan dengan reservoir berumur Miosen Atas yang telah terbukti memproduksi hasil migas di lepas pantai laut dalam. Fokus studi ini adalah melakukan pengkajian komposisi QFL (fragmen Kuarsa, Felsfar dan Litik) dan fasies batuan batupasir berdasarkan sampel batuan yang diambil dari data inti batuan (core) dan data dinding bor (side-wall core), serta data log elektrik yang berasal dari sekitar empat puluh buah sumur. Sampel batuan ini dibatasi oleh marker biostratigrafi: M40M33, M45M40, M50M45 (Miosen Tengah), dan M65M50, M66M65, M70M66, M80M70 (Miosen Akhir). Fasies reservoir dapat dikenali berdasarkan fasies batuan batupasir; rasio NTG, sortasi, dan ukuran butir terutama pada fasies delta berumur Miosen Tengah: FLU_SX, DC_SX, DC_SM, DC_SM, dan DF_SC, dan untuk fasies laut dalam Miosen Akhir: SSWS, MSWS, SSPS, dan MSPS. Selanjutnya porositas dan permeabilitas yang diukur pada batuan inti bor tersebut dapat digunakan untuk menerangkan hubungan antara kualitas reservoir dengan komposisi batupasir maupun fasies batuan. Sistem pengendapan berenergi tinggi terutama yang berhubungan dengan fasies batuan FLU_SX, DC_SX, SSWS dan MSWS merupakan reservoir dengan kualitas terbaik; dan fasies batuan DF_SC, SSPS, dan MSPS kebalikannya. Hubungan antara porositas berbasis data inti batuan dengan kedalaman (depth of burial) secara umum dapat menggambarkan pola penurunan kualitas reservoir seiring bertambahnya kedalaman.

Kata Kunci: Miosen Tengah, laut dalam, QFL, fasies batuan, Kutai, Makassar Utara

ASSESSMENT OF POTENTIAL MARINE CURRENT ENERGY IN THE STRAITS OF THE LESSER SUNDA ISLANDS

KAJIAN POTENSI ENERGI ARUS LAUT DI SELAT-SELAT KEPULAUAN SUNDA KECIL

Ai Yuningsih¹ and Mario D. Saputra¹

¹Marine Geological Institute of Indonesia

Jl. Dr. Djunjunan No.236 Bandung, Indonesia

Corresponding author: ai.yuningsih@esdm.go.id

(Received 11 May 2021; in revised from 28 May 2021; accepted 7 September 2021)

ABSTRACT: The Lesser Sunda Islands extend from Bali to Timor and consist of two geologically distinct parts formed by a subduction system of oceanic crust along the Java-Timor Trench. The northern part which includes Bali, Lombok, Sumbawa, Flores, Wetar, Pantar and Alor, is volcanic in origin; whilst the southern part is non-volcanic, encompassing the islands of Sumba, Timor and Rote. The straits along the Lesser Sunda Islands are formed as a result of very complex geological processes and tectonics in this area. These straits are the most important cross-sections in the southern part of the Indonesian Throughflow (ITF), functioning as outlets for the mass flows of seawater from the Pacific Ocean to the Indian Ocean through the Flores and the Savu Seas. In these straits, relatively high current speeds are occurred, not only caused by the ITF but also due to its geometry, the influence of tidal flow, and monsoonal currents.

Site study and ocean current measurement were conducted by using an echosounder, a pair of Acoustic Doppler Current Profilers (ADCP), and other supporting equipment. In general, the average of most ocean current speeds is less than 1.5 m/s with a duration flow of 8 -12 hours a day, and the maximum speed reaches up to 3 m/s. The tidal types in almost all the straits are mixed semidiurnal tides, in which two high waters and two low waters occur twice a day, with the high and low tides differ in height.

The Lesser Sunda Straits were selected as the potential sites for ocean current power plant because their current speeds are relatively high and their characteristics are more predictable compared with other straits from other regions. Based on the results of bathymetry survey and current characteristics from the deployed ADCP at a fixed (stationary) location on the seabed, the best location for the current power turbines is at the depth of 15-30 m where the seabed gently sloping.

Keywords: Ocean Current, Sunda Lesser, Renewable Energy

ABSTRAK: Kepulauan Sunda Kecil membentang dari Bali ke Timor yang terbagi menjadi dua bagian yang berbeda secara geologis dan terbentuk karena subduksi kerak samudera di sepanjang Palung Jawa-Timor. Bagian utara Kepulauan Sunda Kecil, yang meliputi Bali, Lombok, Sumbawa, Flores, Wetar, Pantar dan Alor, merupakan kepulauan busur vulkanik, sedangkan pulau-pulau di bagian selatannya adalah pulau non-vulkanik yang mencakup Sumba, Timor dan Rote. Selat-selat di sepanjang Kepulauan Sunda Kecil adalah bentukan dari proses geologi dan tektonik yang kompleks yang terjadi di daerah ini. Selat-selat tersebut menjadi perpotongan yang paling penting pada bagian selatan Arus Lintas Indonesia (Arlindo) dan berfungsi sebagai alur keluar aliran massa air laut dari Samudera Pasifik yang melintas melalui Laut Flores dan Laut Sawu hingga ke Samudra Hindia. Kecepatan arus di selat-selat tersebut relatif tinggi. Hal ini tidak hanya disebabkan oleh adanya Arlindo, tetapi juga karena adanya pengaruh geometri, siklus pasang surut dan arus Muson.

Studi lokasi dan pengukuran arus laut telah dilakukan dengan menggunakan alat perum gema, sepasang Acoustic Doppler Current Profiler (ADCP) dan peralatan pendukung lainnya. Secara umum kecepatan arus laut rata-rata kurang dari 1.5 m/detik dengan durasi aliran 8 -12 jam per hari, dan kecepatan maksimum bisa mencapai lebih dari 3 m/detik. Jenis pasang surut di hampir semua selat perairan Sunda Kecil adalah pasang surut campuran harian ganda, yaitu dalam satu hari terjadi dua kali pasang naik dan dua kali pasang surut yang berbeda tinggi.

The Seawater And Freshwater Influence On Expansivity Behaviors Of Clay Minerals

by Dian Agus Widiarso

Submission date: 05-Apr-2023 12:48PM (UTC+0700)

Submission ID: 2056371198

File name: 5_The_sea_water_and_fresh_water_influence.pdf (540.94K)

Word count: 3514

Character count: 18920

THE SEAWATER AND FRESHWATER INFLUENCE ON EXPANSIVITY BEHAVIORS OF CLAY MINERALS

PENGARUH AIRLAUT DAN AIRTAWAR TERHADAP DAYA KEMBANG SUSUT MINERAL LEMPUNG

Dian Agus Widiarso¹, Nurakhmi Qadaryati¹, Wiyatno Haryanto²

¹Department of Geological Engineering, Universitas Diponegoro, Jl. Prof. Soedharto, SH., Tembalang, Semarang, Indonesia

²Indonesian Society of Geotechnical Engineering, Jakarta, Indonesia

Corresponding author: dianagus@lecturer.undip.ac.id

(Received 07 July 2021; in revised form 22 July 2021; accepted 22 September 2021)

ABSTRACT : Semarang City subsurface geology is characterized by an intercalating of loam-silt and clayey units. The behavior of clay materials are their expansivity potency of their volume when in contact with water. Some problems in the presence of tidal flooding appears when the seawater ingress to the shoreland that causes severe damage to infrastructures. This research attempts to reveal the influence of both seawater and freshwater on the expansivity behaviors of the clayey materials based on its mineral composition. In order to observe expansivity potency of clay minerals, 28 samples from four drill hole samples were soaked in seawater and freshwater for twenty-four hours. The initial volume of each samples were then compared with the samples volume after being soaked. X-Ray Diffraction (XRD) analysis was also conducted on selected samples to determine their mineral composition. The ANOVA test was then introduced to distinguish the influence of certain mineral types and composition to the clay expansivity behaviors. Confidence level and alpha (α) are about ninety-five percent and about five percent, respectively. The result of this research has proven that montmorillonite is clay mineral type that is most affected by expansivity behaviors when immersed in seawater and freshwater compared to kaolinite and illite minerals.

Keywords: Clay minerals, montmorillonite, expansivity, seawater, freshwater, tidal flooding.

ABSTRAK: Kota Semarang tersusun dari satuan lanau dan lempung. Mineral lempung memiliki sifat kembang susut dari segi volumenya ketika mengalami kontak dengan air. Permasalahan yang timbul akibat terjadinya peristiwa banjir pasang di perkotaan, adalah masuknya air laut menuju ke pantai dan daratan yang dapat merusak infrastruktur. Penelitian ini bertujuan untuk menunjukkan dampak yang ditimbulkan oleh air tawar maupun air laut terhadap kembang susut mineral lempung berdasarkan komposisi mineralnya. Untuk mengamati perubahan volume dalam mineral lempung, sebanyak 28 sampel dari empat sampel sumur bor direndam dalam air tawar dan air laut selama 24 jam. Kemudian volume awal setiap sampel dibandingkan dengan volume sampel sesudah dilakukan perendaman. Uji X-Ray Diffraction (XRD) juga dilakukan pada sampel pilihan untuk mengetahui komposisi mineral sampel tersebut. Uji ANOVA dengan tingkat kepercayaan 95% dan alpha (α) 5% diterapkan untuk mengetahui pengaruh tipe dan komposisi mineral lempung tertentu terhadap kembang susut yang terjadi. Penelitian ini berhasil membuktikan bahwa montmorilonit merupakan jenis mineral lempung yang paling terpengaruh oleh sifat kembang susut di daerah penelitian ketika direndam dengan air tawar dan air laut, dibandingkan dengan kaolinit dan ilit.

Kata Kunci: Mineral lempung, montmorilonit, kembang susut, air laut, air tawar, banjir pasang

INTRODUCTION

In general, a clayey material is an important part of the soil, which mostly involves chemical weathering of rock-forming minerals and will change in some engineering behaviors when it contacts seawater (Aksoy et al., 2008). Some research on mineralogical and geotechnical approaches in coastal areas with a predominance of clay lithology has been performed widely in several countries. For example Veerasingam et Al.(2014) studied mineralogy and geotechnic for a construction planning in Lianyungang – a region of China that requires some investigative data regarding the origin of rocks, and some geotechnical studies of the clay minerals onsite by Liu et al. (2011). Similar case from the Changi project in Singapore within Quaternary sediment, which are composed of soft marine clay recognized as problematic soil for geotechnical engineering purposes (Bo et al., 2015). The major decision in construction process involves selecting a suitable site with the best soil conditions (Manimaran et al., 2019).

Based on geological setting, Semarang City is a part of the Holocene sediment which is composed of a river, flood-plain, swamp, tidal, and coastal deposits (Thanden et al., 1996). The sediment is characterized by intercalating of a dominant loam-silt and clay units and constituted normally consolidated clays with a soft and very soft consistency (Widiarso et al., 2019). This city is the capital city of Central Java Province, located in a large coastal deposit area with significant environmental issue. One problem such as tidal flooding in which the seawater ingresses frequently to the shoreland area during a high tide period (Wahyudi, 2007), causes serious damage in infrastructures and residential areas (Marfai and King, 2008; Muslim et al., 2019). The quality of the circumference decreases due to being influenced by tidal flooding (Sulaksana et al., 2019). One of the most important goals of disaster management teams is to protect the assets and infrastructures of the community from natural disasters such as floods (Darani and Bashiri, 2018).

Based on previous research, it was explained that claystone which has montmorillonite mineral composition more than 72% may indicate a reduction in the expansivity of the clay material when mixed with seawater, conversely, it exhibits less expansive when mixed with freshwater (Elmashad and Ata, 2014). Supporting that study result, other studies explained that the clay minerals such as kaolinite, chlorite, and mixing of other clay minerals have a lower reaction (low reactivity) than montmorillonite mineral when mixed with seawater (Aksoy et al, 2008). Therefore, the authors intend to present this research to study the influence of seawater and freshwater on the expansivity behaviors of clay materials based on particular mineral composition.

Clay Minerals Overview

Soil is defined as a constituent mineral with or without organic material left over from plants and fauna that are weathered, structured, and textured. It is useful as a construction material in various kinds of civil engineering works. As addition, the land also functions as a support for building foundation (Das, 2008). Based on the cohesivity, soils can be distinguished as cohesive and non-cohesive soils, fine-grained or coarse-grained soils (Bowles, 2015).

The abundance of clay minerals in the soils varies greatly, influenced by various things, including the type of origin rock, weathering, and diagenesis process, which cause variations both vertically and laterally. The most influential factor that determines soil types is the particle size distribution (Kaream et al., 2020). Clay materials with particle sizes smaller than 0.002-mm can be divided into three subgroups of clay minerals, are kaolinite, illite, and montmorillonite (Dunn et al., 1980).

Tidal Flood

Tidal flood is a flood that inundating the lowland areas on the coast, including estuaries and deltas, resulting in salty/brackish groundwater (Marfai, 2004). The underlying problem of tidal flooding affecting shoreland and coastal area is that the land level is lower than the high tide level. Furthermore, human activities include excessive groundwater extraction, dredging on the shipping lines, coastal reclamation, etc., are some factors contribute the tidal flooding.

METHODS (AND MATERIALS)

The research area is in Semarang City, especially on the alluvium and flood plains as referring to the Geological Map of Magelang and Semarang, Java (Thanden et al., 1996). The soil mechanics laboratory tests are performed on 28 clay soil samples, selected from 4 (four) drill hole samples (BM01, BM03, BM04, and BM05 in Figure 1). The depths of the drill hole samples are 60m, 40m, 30m, and 30m, respectively. The samples were then soaked d in seawater and freshwater for twenty-four hours. The samples volume were measured prior and after the soaking test. During the time, we observed the changes in their volumetric parameters. The difference of the lithologic volume is a product of the reduction between a final volume and the initial volume. The freshwater and seawater used in the soaking tests have been analyzed in terms of chemical and physical properties in Groundwater Utilization Sector, Geological Agency, Ministry of Energy and Mineral Resources, Bandung. (Table 1).

Table 1. The results of chemical analysis of fresh water and seawater sample

Type of Water	Indicators (mg/L)					
	Ca ²⁺	Mg ²⁺	K ⁺	Na ⁺	Cl ⁻	SO ₄ ²⁻
Seawater	315	749,5	248,5	7293,5	13393,9	246,5
Freshwater	25,4	6,4	6	19,1	14,4	1

X-Ray Diffraction (XRD) tests were performed on the similar samples as the soaking test. The purpose of this analysis is to determine the clay mineral type of the samples. The results were then processed by using X-pert Highscore and Siroquant software. The software is used to obtain the quantitative data on the diffraction pattern of a sample. ANOVA was then performed to observe the effect of mineral types and composition on the clay mineral expansivity. It is also used as an analytical tool to test the hypothesis by assessing whether there are any differences among the groups. Hypothesis testing is performed at a ninety-five percent confidence level and alpha (α) five percent.

RESULTS AND DISCUSSION

The result of the soaking test is shown in Table 2, it is observed that 3 samples were broken after soaking in both seawater and freshwater (BM 05-2, BM 05-5, BM 05-6). Therefore, these three samples were not taken into account for statistical ANOVA test. In addition to these three samples, one sample BM 05-3 was also broken after soaking in freshwater. Meanwhile, the result of XRD (X-ray diffraction) analysis in the clay mineral compositions is demonstrated in Table 3.




Figure 1. Geotechnical Drilling Map

Table 2. Test parameters of soaking

No	Code	Depth	Soaking in Sea Water		Soaking in Fresh Water	
			Initial Volume (cm ³)	End Volume (cm ³)	Initial Volume (cm ³)	End Volume (cm ³)
1	BM 01-1	5 - 6	166	263	133	120
2	BM 01-2	11 - 12	113	140	113	148
3	BM 01-3	15 - 16	113	159	113	135
4	BM 01-4	23 - 24	113	119	107	159
5	BM 01-5	25 - 26	90	453	90	509
6	BM 01-6	31 - 32	73	145	84	428
7	BM 01-7	41 - 42	113	127	113	137
8	BM 01-8	51 - 52	106	133	106	136
9	BM 01-9	59 - 60	129	154	129	145
10	BM 03-1	5 - 6	113	176	113	206

11	BM 03-2	11 - 12	73	122	113	161
12	BM 03-3	15 - 16	109	168	113	183
13	BM 03-4	21 - 22	107	139	140	190
14	BM 03-5	25 - 26	127	468	127	171
15	BM 03-6	29 - 30	113	144	106	146
16	BM 03-7	39 - 40	159	200	172	170
17	BM 04-1	5 - 6	113	146	113	176
18	BM 04-2	11 - 12	99	129	99	154
19	BM 04-3	15 - 16	109	134	109	166
20	BM 04-4	21 - 22	95	117	79	76
21	BM 04-5	25 - 26	95	142	79	956
22	BM 04-6	29 - 30	113	662	113	708
23	BM 05-1	5 - 6	113	108	55	163
24	BM 05-2	11 - 12	104	broken	104	broken
25	BM 05-3	15 - 16	132	190	112	broken
26	BM 05-4	21 - 22	119	468	79	179
27	BM 05-5	25 - 26	119	broken	119	broken
28	BM 05-6	25,5 - 26	98	broken	95	broken

Statistical Test on Soaking in Seawater

The independent variables in the statistical test with ANOVA were the mineral content in the sample, which are kaolinite, montmorillonite, and illite. Statistical tests were carried out by observing changes in the sample volume before and after soaking without considering the soil water content. The coefficient of significance value (α) is

determined at 0.05. The significance or probability column of the ANOVA test results indicate the level of significance. The significance value (sig value) is used as a cut-off model to determine if the independent variable affects or does not affect the dependent variable. The independent variable is considered to affect the dependent variable if the sig value is lower than 0.05.

Table 3. Percentage of identified mineral from XRD analysis

No	Code	Depth	Identified minerals (%)			
			Quartz	Kaolinite	Illite	Montmorillonite
1	BM 01-1	5 - 6	54.97	40.56	4.47	0
2	BM 01-2	11 - 12	57.33	35.09	7.58	0
3	BM 01-3	15 - 16	59.69	39.54	0	0.76
4	BM 01-4	23 - 24	42.35	48.53	9.12	0
5	BM 01-5	25 - 26	29.48	43.98	0	26.54
6	BM 01-6	31 - 32	28.47	40.95	0	30.58
7	BM 01-7	41 - 42	35.71	0	63.77	0.52
8	BM 01-8	51 - 52	40.48	13.42	33.55	12.54
9	BM 01-9	59 - 60	28.54	44.23	0	27.23
10	BM 03-1	5 - 6	47.26	20.15	31.70	0.89
11	BM 03-2	11 - 12	48.36	17.41	33.18	1.04
12	BM 03-3	15 - 16	47.98	22.34	28.34	1.35
13	BM 03-4	21 - 22	43.62	33.87	0	22.51
14	BM 03-5	25 - 26	27.03	0	60.36	12.61
15	BM 03-6	29 - 30	23.82	47.83	0	28.35
16	BM 03-7	39 - 40	28.06	44.60	0	27.34
17	BM 04-1	5 - 6	72.61	0	26.11	1.27

18	BM 04-2	11 - 12	64.78	28.85	0	6.37
19	BM 04-3	15 - 16	45.72	29.01	22.06	3.21
20	BM 04-4	21 - 22	50.19	49.05	0	0.76
21	BM 04-5	25 - 26	33.84	0	59.44	6.72
22	BM 04-6	29 - 30	28.04	38.10	0	33.86
23	BM 05-1	5 - 6	60.04	34.40	0	5.56
24	BM 05-2	11 - 12	48.12	25.29	24.71	1.88
25	BM 05-3	15 - 16	94.37	0	0	5.63
26	BM 05-4	21 - 22	59.62	0	0	40.38
27	BM 05-5	25 - 26	12.80	0	87.20	0
28	BM 05-6	25,5 - 26	63.55	35.81	0	0.65

Based on Table 4, it shows that the sig value is 0.024, smaller than 0.05 (α), suggests that the model passed and can be used to specify the influence of the mineral contents in samples on sample volume changes after soaked in seawater. The probability value (significance) is below 0.05, indicates all independent variables influence the dependent variable and vice versa.

Tabel 4. ANOVA test result on soaking in seawater

Model	df	Mean square	F	Sig.
Regression	3	48961.1	3.897	0.024 ^b
Residual	20	12563.6		
Total	23			

a. Dependent variable: after soaked in seawater

b. Predictors: (constant), montmorillonite, kaolinite, and illite.

Afterwards, the F-count value (F in table 4) is then compared to the F-table parameter. The F-table is obtained by calculating the number of df (degree of freedom). The df of multiple regression consists of df1 and df2 with the following formula:

$$df1 = k - 1 \dots \dots \dots (1)$$

$$df2 = n - k \dots \dots \dots (2)$$

under the following condition:

n = number of samples

k = number of independent variables

For this statistical tests, 25 samples were analyzed. As has been mentioned before, from total 28 samples, 3 samples (BM 05-2, BM 05-5, and BM 05-6) were broken after being soaked in seawater, therefore were not considered for the statistical test. The soaking test in seawater has k value = 3 and n = 25. Hence, the values of df1 = 2 and df2 = 22 are obtained. Based on F distribution table at 0.05 sig level, the F-table value is 3.443 (Nuryadi et al., 2017), which is smaller than F-count (3.897). Therefore, it can be concluded that all independent variables influence the dependent variable. The test results of the coefficient value on the soaking in seawater is displayed in Table 5.

In Table 5, t value is the benchmark to see the influence of variable (partial) by comparing their each t value to the t-table value. When the value of t-count is greater than the t-table value, denotes the variable has an impact. By calculating the number of df (degree of freedom), t-table is obtained with the following formula:

$$df = n - (k - 1) \dots \dots \dots (3)$$

under the following condition:

n = number of samples

k = number of independent variables

The calculation of df (degree of freedom) = 25 - (3 - 1) = 23.

Based on t-table value for multiple regression df 23 at sig 0.05 is 1.71387 (Nuryadi et al., 2017). T-count value of montmorillonite mineral reveals a value of 3.303 (Table 5), which is greater than t-table value. In contrast, t-count value of kaolinite and illite are -0.375 and 0.568, respectively, which are lower than t-table (1.71387). It is concluded that montmorillonite mineral has the most influence on the changes in sample volumetric when it is soaked in seawater compared to kaolinite and illite minerals.

The sig column in Table 5 represents the significance level of each independent variable to determine whether the variables influence the dependent variable or not. When sig value is smaller than 0.05 (α), signify that there is significant impact, in contrast, when it is greater than 0.05 (α), then there is no influence at all. The result reveals that only sig value of montmorillonite mineral has a lower value than 0.05 (0.004). So, montmorillonite mineral has the most influence on changes in sample volume compared to kaolinite and illite minerals when soaked in seawater.

Statistical Test on Soaking in Freshwater

The data is processed using multiple regression analysis methods with the dependent variable is the percentage of difference between the initial volume with the sample final volume, after soaked in freshwater for

Table 5. Coefficient test result of soaked in seawater

Model	t	Sig.
(Constant)	1.503	0.148
Kaolinite	-0.375	0.712
Illite	0.568	0.576
Montmorillonite	3.303	0.004

twenty-four hours. The ANOVA test result shown in Table 6 with the sig value of about 0.249.

Table 6. ANOVA test result on soaked in freshwater

Model	df	Mean square	F	Sig.
Regression	3	93014.1	1.484	0.249 ^b
Residual	20	62688.1		
Total	23			

Dependent variable: after soaked in freshwater

Predictors: (Constant), montmorillonite, kaolinite, and illite

The value is greater than 0.05 (α), so the model is considered failed, and cannot be used to determine the influence of sample's mineral content in the changes of sample volume after being soaked in freshwater. The probability value (significance) is above 0.05, suggests not all independent variables have an affect on the dependent variable and vice versa.

Only 24 out of 28 samples were analyzed in the statistical tests after being soaked in freshwater because four samples (BM 05-2, BM 05-3, BM 05-5, BM 05 -6) were broken after being soaked. The soaking test in freshwater has a value of $k = 4$ and a value of $n = 24$, so that the values of $df_1 = 3$ and $df_2 = 21$ are obtained. Based on the F distribution table at the 0.05 sig level, the F-table value is 3.072 (Nuryadi et al., 2017), greater than the F-count amounting to 1.484 (Table 6). It can be concluded that not all independent variables have an influence on the dependent variable.

Table 7. Coefficient test result of soaked in freshwater

Model	t	Sig.
(Constant)	-0.012	0.990
Kaolinite	0.364	0.719
Illite	1.304	0.207
Montmorillonite	1.837	0.081

The df value for the soaking test in freshwater is = $24 - (4 - 1) = 21$. Based on the t-table value for the multiple regression df 21 at sig 0.05 is 1.72074 (Nuryadi, et al., 2017). The t-count value of the montmorillonite mineral in Table 7 shows a value of 1.837, so the montmorillonite mineral has the most influence in the changes of sample volume compared to the kaolinite and illite minerals when soaked in freshwater.

Based on a comparison of the ANOVA difference test for soaking of the clay minerals in seawater and freshwater, it is obtained that the F value is 3.897 with a sig of 0.024, while the results of soaking in freshwater, the F value is 1.484 with sig value 0.24. Hence, it is suggested that seawater has a more significant influence on the expansion process of the clay minerals compared to freshwater.

Montmorillonite is a mineral formed by two sheets of silica and one sheet of aluminium. The octahedral sheet is located between two silica sheets with the ends of the tetrahedra. Tetrahedra is mixed with hydroxyl of the octahedral sheet to form an aluminium single layer by magnesium. Due to the weak Van der Waals bonding forces (bonds due to changes in the number of electrons at any time in one part of the atomic nucleus) between the ends of the silica sheet and the lack of negative charge in the octahedral sheet, water (H_2O) and moving ions can enter and separate the layers (Ariesnawan, 2015). This, results in high swelling properties. The ionic content of seawater which is higher than that in fresh water, induces the swelling properties of mineral montmorillonite to be more developed when exposed to seawater than when exposed to fresh water.

CONCLUSIONS

It is proven that montmorillonite mineral is the most influential in changing the volumetric expansivity of soil when the soil is soaked in seawater and freshwater compared to kaolinite and illite minerals. The montmorillonite minerals have more effect on volumetric changes when the clay minerals are immersed in seawater than when they are immersed in freshwater. This study provides an understanding that it is also necessary to have a good knowledge regarding the composition of clay minerals in the planning, especially when the land area is dominated by clay material and affected by tidal flooding. Engineering geological planning should not only observe the physical properties of lithology, they must consider the chemical properties of lithology as well.

ACKNOWLEDGEMENTS

This paper is a part of Research Development and Implementation Research (RPP) Fund Sources Apart from the Diponegoro University State Budget for the 2021 Fiscal Year. The authors are grateful to all those who have supported and helped during the research process.

REFERENCES

- Aksoy, Y. Y., Kaya, A., Oren, A. H., 2008. Seawater effect on consistency limits and compressibility characteristics of clays. *Engineering Geology*, 102: 54-61.
- Ariesnawan, R. A., 2015. The Mechanic And Dynamic Characteristic of Tuban Clay Shale Due To Changes of Water Content. Thesis. Institut Teknologi Sepuluh Nopember Surabaya.
- Bo, M. W., Arulrajah, A., Sukma, P., Horpibulsuk, S., 2015. Mineralogy and Geotechnical Properties of Singapore Marine Clay at Changi. *Soils and Foundation*, 55: 600- 613.
- Bowles, J. E., 2015. *Physical and Geotechnical Properties of Soils*. McGraw-Hill International Book Company.
- Darani, S. K., Bashiri, M., 2018. A Multi-district Asset Protection Problem with Time Windows for Disaster Management. *International Journal of Engineering*, 31(11):1929-1934.
- Das, B. M., 2008. *Advanced Soil Mechanics, Third Edition*. Taylor & Francis Group.
- Dunn, I. S., Anderson, I. R., dan Kiefer, F. W., 1980. *Basics of Geotechnical Analysis, First Edition*. IKIP Semarang Press.
- Elmashad, M. E., and Ata, A. A., 2014. Effect Of Seawater On Consistency, Infiltration Rate And Swelling Characteristics Of Montmorillonite Clay. *Housing and Building National Research Center*, 12: 175-180.
- Kaream, K. W. A, Fattah, M. Y., Khaled, Z. S. M., 2020. Assessment of Changes in Shear Strength Parameters for Soils below Circular Machine Foundation. *International Journal of Engineering*, 33(8):1491-1498.
- Liu, S. Y., Shao, G. H., Du, Y. J., Cai, G. J., 2011. Depositional and Geotechnical properties of Marine Clays in Lianyungang, China. *Engineering Geology*, 121:66-74.
- Manimaran, A., Seenu, S., Ravichandra, P. T. N., 2019. Stimulation Behaviour Study on Clay Treated with Ground Granulated Blast Slag and Groundnutshell Ash. *International Journal of Engineering*, 32(5):673-678.
- Marfai, M. A., 2004. Tidal Flood Hazards Assessment: Modelling in Raster GIS, Case in Western Part of Semarang Coastal Area. *Indonesian Journal of Geography*, 36(1):25-28.
- Marfai, M. A. and King, L., 2008. Potential Vulnerability Implications if Coastal inundation Due Sea Level Rise for The Coastal Zone Of Semarang City, Indonesia. *Journal Environmental Geology*, 54:1235-1245.
- Muslim, D, Haerani E., Muslim F. N. and Muslim G. O., 2019. Toward the Safe Live-able Built Environment around Ciletuh-Palabuhanratu Geopark Area in Sukabumi Regency, Indonesia. *IOP Conf. Series: Earth and Environmental Science*, 248: 012036. doi:10.1088/1755-1315/248/1/012036.
- Nuryadi, Astuti, T. D., Utami, E. S., Budiantara, M. 2017. *Dasar-dasar Statistik Penelitian*. Sibuku Media. Yogyakarta.
- Sulaksana, N., Iskandarsyah, T. Y.W.M., Rifai, A., Rendra, P.P.R. and Sulastri, M., 2019 Prospective Zone Area for Agriculture and Residential Based on Geological Disaster Potentials in South Bandung Region. *Journal of Geological Sciences and Applied Geology*, 3(1): 26-37.
- Thanden, R. E., Sumadirdja, H., Richards, P. W., dan Sutisna, K., 1996. *Peta Geologi Lembar Magelang dan Semarang, Jawa skala 1:100.000*. Pusat Survey Geologi.
- Veerasingam, S., Venkatachalapathy, R., Ramkumar, T., 2014. Distribution of Clay Minerals in Marine Sediments of Chennai, Bay of Bengal, India: Indicators of Sediment Sources and Transport Processes. *International Journal of Sediment Research*, 29:11-23.
- Wahyudi, 2007. Tingkat Pengaruh Elevasi Pasang Laut Terhadap Banjir dan Rob di Kawasan Kaligawe Semarang. *Jurnal dalam Riptek*, 1(1):27-34.
- Widiarso, D. A., Haryanto, W., Muslim, D., Zakaria, Z. and Iskandarsyah, T. Y., 2019. Potential Consolidation Settlement Due to Load Stresses of Building Structures. *International Journal of GEOMATE*, 17(60):204-210.

The Seawater And Freshwater Influence On Expansivity Behaviors Of Clay Minerals

ORIGINALITY REPORT

4%

SIMILARITY INDEX

3%

INTERNET SOURCES

2%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|------|
| 1 | turkiyeklinikleri.com
Internet Source | 1 % |
| 2 | jurnal.stie-aas.ac.id
Internet Source | 1 % |
| 3 | "Geotechnical Engineering and Sustainable Construction", Springer Science and Business Media LLC, 2022
Publication | <1 % |
| 4 | Hadiratul Kudsiah, Suwarni, Sri Wahyuni Rahim, Dwi Fajriyati Inaku, Muhammad Ahsin Rifa'i, Sri Haslina. "Growth patterns and condition factors of the bungo fish, Glossogobius giuris (Buchanan, 1822) in Sidenreng Lake waters, Sidenreng Rappang Regency, South Sulawesi", IOP Conference Series: Earth and Environmental Science, 2022
Publication | <1 % |
| 5 | www.tribology.rs
Internet Source | <1 % |

6

garuda.kemdikbud.go.id

Internet Source

<1 %

7

teachers.bcps.org

Internet Source

<1 %

8

Subramanian VEERASINGAM, Ramdoss VENKATACHALAPATHY, Thirunavukkarasu RAMKUMAR. "Distribution of clay minerals in marine sediments off Chennai, Bay of Bengal, India: Indicators of sediment sources and transport processes", International Journal of Sediment Research, 2014

Publication

<1 %

9

Wolfgang Lieske, Wiebke Baille, Joyce Schmatz, Stephan Kaufhold, Reiner Dohrmann. "Characterisation of natural and remoulded Onsøy clay with focus on the influence of mica", Engineering Geology, 2021

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

The Seawater And Freshwater Influence On Expansivity Behaviors Of Clay Minerals

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8