

The Role of Zinc Intake in Serotonin and Cortisol Level In Patient with Depression

by Alifiati Fitrikasari

Submission date: 13-Apr-2023 09:56AM (UTC+0700)

Submission ID: 2063103616

File name: e_in_Serotonin_and_Cortisol_Level_In_Patient_with_Depression.pdf (280.32K)

Word count: 4290

Character count: 23220

JOURNAL OF BIOMEDICINE AND TRANSLATIONAL RESEARCH

Available online at JGTR website: <https://jbr.fk.unidip.ac.id>

Copyright © 2022 by Faculty of Medicine Universitas Diponegoro and Indonesian Society of Human Genetics

Original Research Article

The Role of Zinc Intake in Serotonin and Cortisol Level In Patient with Depression

Tanjung Ayu Sumekar^{1,2*}, Innawati Jusup¹, Natalia Dewi Wardani¹, Titis Hadiati¹, Mohammad Sulchan¹, Alifiaty Fitrikasari¹

¹Department of Psychiatry, Faculty of Medicine, Universitas Diponegoro, Indonesia

²Department of Physiology Faculty of Medicine, Universitas Diponegoro, Indonesia

³Department of Nutrition, Faculty of Medicine, Universitas Diponegoro, Indonesia

Article Info

History

Received: 07 Mar 2022

Accepted: 25 Apr 2022

Available: 28 Apr 2022

Abstract

Background: Low zinc levels affect the relationship between the glutamatergic and serotonergic systems in major depressive disorders that cause stress and inflammation. Decreased zinc in the hippocampus can activate the HPA axis associated with increased in cortisol. Several studies documented the relationship between zinc and clinical depression. However, further research including biological measurements is needed to support these studies.

Objective: To analyze the correlation between zinc intake with serotonin and cortisol serum in patients with depression

Methods: This was an observational study with a cross sectional design. Subjects were patients with depression who admitted to Dr. Kariadi Hospital, Tugurejo Hospital, Diponegoro National Hospital and Permata Medika Hospital. The level of depression was determined using Beck Depression Inventory-II (BDI-II). The food frequency questionnaire (FFQ) was used to assess daily zinc intake. The levels of serum serotonin and cortisol were measured using the ELISA technique.

Results: Of the 53 subjects, there was a significant correlation between zinc intake with serotonin serum level ($p=0.038$); however, there was no significant correlation between zinc intake with cortisol serum level ($p=0.845$).

Conclusion: High zinc intake is correlated with serotonin level in patient with depression

Keywords: cortisol; depression; FFQ; serotonin; zinc intake

Permalink/ DOI: <https://doi.org/10.14710/jbr.v8i1.13808>

INTRODUCTION

Depression is a mental disorder associated with decreased productivity, poor psychosocial outcomes and decreased quality of life and wellbeing. According to the World Health Organization (WHO), the total number of people living with depression in the world is about 350 million. Nearly half of these people live in South-East Asia and Western Pacific.¹ In Indonesia, with a variety of biological, psychological and social factors with diversity in the population, the total estimated number of cases of mental disorders increase. This may result burden and decrease in human productivity. Data of Basic Health Research of Ministry of Health of the Republic of Indonesia in 2018 showed the prevalence of depression among the population of aged 15 years and

above was about 14 million people or 6.1%, with a prevalence of 4.7% in Central Java.² Although there are many effective treatments for depression, pharmacotherapy is usually costly and has potential side effects and psychotherapy requires time and commitment. Take the widely promoted monoaminergic antidepressant, for example, more than 30% of the patients did not respond to this treatment.³ Therefore, there is a need to investigate alternative treatment or prevention strategies.

* Corresponding author:

E-mail: tanjung.sumekar@fk.unidip.ac.id

(Tanjung Ayu Sumekar)

Recent research has focused on the role of micronutrient in various mental health disorders including depression.^{4,5} Zinc is one of micronutrients essential for optimal function of the human body, especially the brain and neural structures, where it is found at the highest concentration in the hippocampus and amygdala regions of the brain.⁶

The essential trace ion zinc participates in numerous biological processes. Zinc regulates intracellular signal transduction and contributes to efficient synaptic transmission in the central nervous system.⁷ Zinc deficiency is a recognized global public health concern in developing countries and is also becoming a prevalent concern in the aging population of developed countries.⁸ Zinc deficiency is also associated with neuropsychiatric manifestation, including cognitive and behavioral changes and depression.⁹ A case-control study showed that serum zinc levels in patients with depression were significantly lower than in normal controls.¹⁰ Furthermore, a meta-analysis study concluded that zinc supplementation may improve depression in individuals treated with antidepressant.¹¹

The interaction of zinc and the serotonergic receptor 5-hydroxytryptamine 1A (5-HT_{1A}) in a mouse test with reduced antidepressant administration showed that zinc modulates serotonergic receptors that can improve depression.¹² Serotonin is involved in various central nervous system functions to regulate pain, sleep, blood pressure, mood, and behaviour. The role of serotonergic transmission in depression is well known so that it is one of the mechanisms of action in anti-depressant drugs. Effect on serotonergic receptors is one of the explanations of the antidepressant-like properties of zinc, which are observed in both preclinical and clinical studies. Satake et al. extensively explored the pharmacological profile of zinc at the 5-HT_{1A} receptors using the agonist of 5-HT_{1A}, [3H]-8-OH-DPAT.¹³ The interaction of zinc with glutamate and the specific N-methyl-D-aspartate (NMDA) receptor is another mechanism in reducing symptoms of depression.¹⁴ Low zinc levels disrupt the relationship between the glutamatergic and serotonergic systems in major depressive disorders causing stress and inflammation.¹⁵ Some studies showed that increased glucocorticoids in stress and inflammation in the dysregulation of the Hypothalamic Pituitary Adrenal (HPA) axis produce symptoms of depression. Decreased zinc level in the hippocampus activates the HPA axis associated with increased cortisol.¹⁶ The role of dietary zinc intake in serotonin and cortisol levels has never been reported in depression patients. This study aimed to analyze the correlation between zinc intake with serotonin and cortisol serum in patient with depression.

MATERIALS AND METHODS

Study Design and Participants

This was an observational study with cross-sectional design. Subjects participated in this study were patients with diagnosis of depression under treatment based on the medical records who admitted to Psychiatry Clinic at Dr. Kariadi Hospital, Tugurejo Hospital, Diponegoro National Hospital, and Permata Medika Hospital between January - December 2019 with age ranged from 18 to 60 years old. Beck Depression Inventory - II (BDI-

II) was administered by authors who were psychiatrist to determine level of depression. Subjects signed a consent form that approved by our institutional review board No. 126/EC/KEPK-FK-UNDIP/VI/2020.

Measurements

Subjects filled the demographic questionnaires including sex, occupation, marital status, education, income, and history of psychosocial stressors. Subjects underwent a full medical evaluation, which were medical history, physical and psychiatric examination. Subjects who had diabetes mellitus, cardiovascular disease, renal disease, gastrointestinal disease, malignancy, smoking, pregnancy and breastfeeding were excluded. Body weight and height were measured to calculate the body mass index (BMI).

Zinc intake was calculated from a dietary intakes assessment using food frequency questionnaires (FFQ).¹⁷ FFQ was administered by trained physician and data conversion was carried by nutritionist. Zinc daily intake of each participant was obtained from the conversion of FFQ in gram. Zinc daily intake level was categorized as insufficient when the zinc intake was below 77% of daily requirement and normal when the zinc intake was above or the same as daily requirement.^{18,19}

Serum were obtained from peripheral blood vein at 8:00 - 9:30 am for each subject. The Sandwich ELISA method using Serotonin high sensitive ELISA kit, IBL International GmbH (Hamburg, Germany) and Microplate Reader Biorad 680 were applied to measure the serotonin level. Architect Cortisol Reagen (Illinois, USA), and Architect i2000SR were used to measure the serum cortisol level. Normal serum serotonin levels ranged from 101-283 ng/ml and normal serum cortisol level ranged from 10-20 g/dL.²⁰

44

Table 1 Demographic Characteristics of Subjects

Characteristics	Mean ± SD (min-max)
Age	35.91 ± 13.26 (19-60)
Height (cm)	156.87 ± 8.12 (140-176)
Weight (kg)	60.85 ± 9.32 (45-85)
BMI (kg/m ²)	24.77 ± 3.59 (17.4-34.4)
BDI-II	21.15 ± 11.41 (1-53)
Serum serotonin level	91.15 ± 63.00 (8.12-327.66)
Serum cortisol level	8.26 ± 3.65 (1.7-16.4)

SD = Standard Deviation; min = minimum; max = maximum
BMI = Body Mass Index; BDI-II = Beck Depression Inventory - II (BDI - II)

Table 2. Correlation between zinc intake and serum serotonin level

	Zinc Intake	
Serum serotonin level	r	0.285
	p	0.038*
Serum cortisol level	r	-0.027
	p	0.845

r = correlation coefficient; p = significant value

Table 3. Correlation between demographic variables and serum serotonin and cortisol levels

Variable	n	7 Serotonin	Cortisol
Sex		$p=0.185, r=0.185^*$	$p=0.001^*, r=-0.430^*$
Male	13	72.27 (8.12 -157.55)	10.7 (7.2 - 16.4)
Female	40	85.05 (17.48 -327.66)	7 (1.8 - 18.7)
Age		$p=0.119, r=-0.217^*$	$p=0.091, r=-0.235^*$
Nutritional status		$p=0.039^*, r=-0.284^*$	$p=0.930, r=0.012^*$
Underweight	1	90.08 (90.08 -90.08)	10.7 (10.7 -10.7)
Normal	19	86.06 (17.48 -327.66)	7.9 (2.1 -14.3)
Overweight	13	87.72 (25.07 -243.87)	8.7 (3.5 - 18.7)
Obesity	20	53.11 (8.12 -229.47)	8.2 (1.8 - 16.4)
BDI-II		$p=0.758, r=0.043^*$	$p=0.058, r=0.262^*$
Minimum	14	87.06 (17.48 -198.13)	7.4 (1.8 -11.5)
Mild	12	66.07 (38.93 -327.66)	7 (2.1 -12.8)
Moderate	11	77.51 (8.12 -243.87)	9.3 (4.4 -18.7)
Severe	15	76.98 (17.72 -254.26)	9.2 (3.3 -14.9)
Length of antidepressant therapy		$p=0.349, r=0.131^*$	$p=0.042^*, r=0.281$
< 1 Month	7	90.08 (41.20 -254.26)	11.9 (7.1 -14.9)
>1 Month	46	76.15 (8.12 -327.66)	8 (1.8 -18.7)
Stressor		$p=0.059, r=0.262^*$	$p=0.977, r=0.004^*$
Occupation	2	58.92 (8.12 -156.41)	9.9 (6.3 -16.4)
Psychosocial environment and Primary Support Group	8	58.82 (37.97 -254.26)	8 (1.8 -14.9)
Health condition	27	86.06 (10.91 -327.66)	8 (2.1 - 18.7)
Education	4	110.05 (84.05 -157.55)	7.4 (5.9 -9.2)
Financial	7	77.51 (40.33 -180.85)	9.5 (6.7 -13.5)
	2	95.79 (50.54 -141.05)	6.3 (2.2 -10.5)

* Significant ($p < 0.05$);

* Spearman's correlation

* Pearson correlation

Statistical Analysis

The normality test was carried out using the Shapiro-Wilk test. The correlation between zinc intake and serotonin and cortisol levels was analyzed. Pearson test was done to analyze the role of age in serotonin and cortisol level and the Spearman correlation test to analyze the role of, length of therapy, stressor, nutritional status, sex, and BDI-II score in serum serotonin and cortisol level. The p value is considered significant if $p \leq 0.05$.

RESULTS

Subjects participated in studies at Kariadi Hospital, Tugurejo Hospital, Diponegoro National Hospital and Permata Medika Hospital between January and December 2019. This study included 53 patients who met criteria with the mean age $35.91 \text{ years} \pm 13.26$ (19 to 60 years) with mean BDI-II score was 21.15 ± 11.41 (1 to 53), mean body weight $60.85 \text{ kg} \pm 9.32$ (45 to 85), mean height $156.87 \text{ cm} \pm 8.12$ (140 to 176), and mean body mass index 24.77 ± 3.59 (17.4 to 34.4). The characteristics of the study subjects are shown in Table 1. Serum serotonin levels were measured with levels ranging from 8.12 ng/mL to 327.66 ng/mL. (mean $91.15 \pm \text{SD } 63.00$). Cortisol serum levels ranging from 1.7 g/dL to 16.4 g/dL. (mean $8.26 \pm \text{SD } 3.65$).

Table 2 summarizes the correlation between zinc intake and serum serotonin and cortisol levels. It showed a significant positive correlation between zinc intake and serum serotonin level ($p=0.038$) but not with serum cortisol level ($p=0.0845$).

Table 3 shows the result of the bivariate analysis between demographic variables i.e., occupation, marital status, education, monthly income, length of antidepressant therapy, stressor, nutritional status, sex, age and BDI-II and serum serotonin and cortisol levels. From this analysis, we found a significant negative correlation between BMI and serum serotonin levels in the subjects. ($p=0.039$) with a weak level of coefficient correlation ($r=-0.284$). The higher the BMI, the lower the serum serotonin levels. There was a significant negative correlation between the length of therapy and serum cortisol levels of subjects ($p=0.042$) and there was a significant negative correlation between gender and serum cortisol levels ($p=0.001$).

DISCUSSION

This study showed a significant correlation between zinc intake and serum serotonin levels in patients with depression. A publication demonstrates differences in serum concentrations in patient with depression.¹⁰ In the etiology of depression there is a monoamine hypothesis which states that depression is associated with a decrease in the amount or function of serotonin in cortex and limbic system. In this study the measured zinc intake was obtained from interview using food frequency questionnaire similar to a study conducted by Li et al which assessed total zinc level using a 24-h recall survey.²¹ Serotonin levels play a role in the mechanism of action of antidepressants. The relationship between zinc intake and depression can be explained by various mechanisms. Zinc homeostasis in the brain is regulated by the food intake containing zinc. Zinc has a synergistic effect like SSRIs that can regulate 5-HT1A receptors that

can inhibit agonist and antagonist binding at synapses. Zinc can also affect serotonin neurotransmission, NMDAR activation and BDNF activity involved in depression.²² In addition, zinc also has anti-inflammatory and antioxidant properties that contribute to depression.²³

Zinc plays a significant role with respect to the stress response. Proper maintenance of zinc status can help stabilize serum cortisol levels over time and zinc intake has been shown to temporarily inhibit cortisol secretions.²⁴ However, prolonged stress will deplete zinc concentrations in the blood. Zinc deficiency has been demonstrated to increase plasma cortisol and pro-inflammatory cytokine of interleukin-6 (IL-6), IL-1 and nitric oxide levels. The mechanism of food consumption-induced cortisol stimulation is not known, it may be due to stimulation of pituitary ACTH secretion by amino acid products or protein digestion, release of gut messengers that secondarily release ACTH or release locally generated ACTH from the intestine. The results also indicated that there was no significant correlation between zinc intake and serum cortisol levels in patients with depression. This may be because cortisol is a biological marker of stress in general and not a specific marker for depression.²⁴ We found that almost all subjects had normal serum cortisol levels which was possible because the subjects were already on antidepressant treatment and were not currently exposed to stressful stimuli. Across multiple mental health-related measures, a polymorphism (5-HTTLPR) within the serotonin transporter gene promoter has been associated with differential psychological sensitivity to stressful experience. The short/short genotype of the 5-HTTLPR is associated with greater cortisol reactivity to social threat. When short/short individuals experience stressful life events, they might be at greater risk for the adverse psychological and physical health consequences associated with heightened cortisol exposure.²⁵

Although acute or chronic stress can be measured in physiological parameters such as heart rate, blood pressure, and various metabolic hormones, it is difficult to understand whether changes in levels of circulating stress mediators such as cortisol can reflect acute, chronic or diurnal stress.^{24,26} In the analysis of confounding variables on serotonin levels, it was found that serum serotonin levels were correlated by body mass index. Abnormal BMI is also related to the availability of serotonin transporters that disrupted in individual with obesity. This is consistent with a study assessed the correlation between plasma tryptophan levels and BMI with symptoms of depression.²⁷ Another study demonstrated that BMI and body fat mass were associated decreased serotonin production in children and adolescents with severe obesity. A chronic low-grade inflammatory state is a trait in obesity and assumed to be one of the underlying pathomechanism. This mechanism increase tryptophan catabolism via the kynurenine pathway regulated by nutritional and inflammatory signals and linked to decrease in serotonin production.^{28,29} This study also showed a significant negative correlation between gender and serum cortisol levels, which is consistent with previous studies about gender differences in cortisol responses and females may be at increased risk for depression because of HPA-axis

dysregulation.³⁰⁻³² Animal and human studies also suggested that female sex hormone influence the HPA axis and corticoid receptors in the hypothalamus and amygdala.^{33,34} Decreased serotonin levels is one of the pathophysiology mechanism of depression.³⁵ The limitation of our study was the subjects were patients with depression under treatment, however we analyzed the correlation between length of therapy with serotonin and cortisol level. We found that there was no significant correlation between length of therapy with serotonin but there was a significant correlation between length of therapy with cortisol level. Serum cortisol levels were influenced by the length of therapy. This can be because the use of long-term antidepressants can affect the HPA axis through changes in glucocorticoid and mineralocorticoid receptors so that serum cortisol levels will decrease. Changes in these receptors affect the effectiveness of therapy. Long term use of antidepressants for both SSRI and non-SSRI can reduce the intensity of HPA axis activity that eventually reduces serum cortisol levels.³⁶

CONCLUSION

From the study, we conclude that there was a significant correlation between zinc intake and serum serotonin levels, thus the higher the zinc intake the higher the serum serotonin levels. There was no significant correlation between zinc intake and serum cortisol levels. In addition, BMI has a significant correlation with serotonin levels. The length of antidepressant therapy and gender significantly correlated with serum cortisol levels. Further research is needed to assess other risk factors affecting serum serotonin and cortisol levels.

ACKNOWLEDGMENTS

This work was supported by Department of Psychiatry, Faculty of Medicine, Diponegoro University. We thank Dr. Kariadi Hospital, Diponegoro National Hospital, Tugurejo Hospital, Permata Medika Hospital and all participants and their families. We also thank Prof. Tri Indah Winarni, MD, PhD for the manuscript preparation. We thank to Hardian, MD, PhD, Nurhasanah, MD and Darmawati Ayu Indraswari, MD for their contribution with data processing.

REFERENCES

1. World Health Organization. *Depression and Other Common Mental Disorders: Global Health Estimates*. WHO Press; 2017.
2. Litbang Kementerian Kesehatan RI. *Riset Kesehatan Dasar*. Kemenkes RI; 2018.
3. Al-Harbi KS. Treatment-resistant depression: Therapeutic trends, challenges, and future directions. *Patient Prefer Adherence*. 2012;6:369-388. <https://doi.org/10.2147/PPA.S29716>
4. Opie RS, Itsiopoulos C, Parketa N, et al. Dietary recommendations for the prevention of depression. *Nutr Neurosci*. 2017;20(3):161-171. <https://doi.org/10.1179/1476830515Y.0000000043>
5. Marx W, Moseley G, Berk M, Jacka F. Nutritional psychiatry: The present state of the evidence. *Proc Nutr Soc*. 2017;76(4):427-436. <https://doi.org/10.1017/S0029665117002026>

6. Wang J, Um P, Dickerman BA, Liu J. Zinc, magnesium, selenium and depression: A review of the evidence, potential mechanisms and implications. *Nutrients*. 2018;10(5):1-19. <https://doi.org/10.3390/nu10050584>
7. Peetbury SD, Adlard PA. Zinc signal in brain diseases. *Int J Mol Sci*. 2017;18(12). <https://doi.org/10.3390/ijms18122506>
8. Maron E, Toru I, Hirvonen J, et al. Gender differences in brain serotonin transporter availability in panic disorder. *J Psychopharmacol*. 2011;25(7):952-959. <https://doi.org/10.1177/0269881110389207>
9. Sathyanarayana Rao T, Asha M, Ramesh B, Jagannatha Rao K. Understanding nutrition, depression and mental illnesses. *Indian J Psychiatry*. 2008;50(2):77. <https://doi.org/10.4103/0019-5545.42391>
10. Styczen K, Sowa-kuzma M, Siwek M, et al. The serum zinc concentration as a potential biological marker in patients with major depressive disorder. *Metab Brain Disord*. 2017;32:97-103. <https://doi.org/10.1007/s11011-016-9888-9>
11. Da Silva LEM, De Santana MLP, Costa PRDF, et al. Zinc supplementation combined with antidepressant drugs for treatment of patients with depression: A systematic review and meta-analysis. *Nutr Rev*. 2021;79(1):1-12. <https://doi.org/10.1093/nutrit/nuaa039>
12. Szweczyk B, Poleszak E, Wlaz P, et al. The involvement of serotonergic system in the antidepressant effect of zinc in the forced swim test. *Prog Neuro-Psychopharmacology Biol Psychiatry*. 2009;33(2):323-329. <https://doi.org/10.1016/j.pnpb.2008.12.011>
13. Satala G, Duszyńska B, Stachowicz K, et al. Concentration-Dependent Dual Mode of Zn Action at Serotonin 5-HT1A Receptors: In Vitro and In Vivo Studies. *Mol Neurobiol*. 2016;53(10):6869-6881. <https://doi.org/10.1007/s12035-015-9586-3>
14. Cowen P, Browning M. What has serotonin to do with depression? *World Psychiatry*. 2015;14(2(June)):11-13.
15. Bonaventura P, Benedetti G, Albarede F, Miossec P. Zinc and its role in immunity and inflammation. *Autoimmun Rev*. 2014;14(4):277-285. <https://doi.org/10.1016/j.autrev.2014.11.008>
16. Brandao-Neto J, De Mendonca B, Shuhama T, Marchini J, WP P, Tomero M. Zinc Acutely and Temporarily Inhibits Adrenal Cortisol Secretion in Humans: A Preliminary Report. *Biol Trace Elem Researcj*. 1990;24:83-89.
17. Fitri N, Jafar N, Indriyani R. Validation Study Semi-Quantitative Food Frequency Questionnaire with 24-hour Food Recall on Micronutrient Intake in Adolescent at Athirah High School Makassar. Published online 2013:1-13.
18. Menteri Kesehatan Republik Indonesia. Peraturan Menteri Kesehatan Republik Indonesia Nomor 28 Tahun 2019 tentang Angka Kecukupan Gizi yang Dianjurkan untuk Masyarakat Indonesia. Published online 2019:15-16. http://hukor.kemkes.go.id/uploads/produk_hukum/PMK_No_28_Th_2019_ttg_Angka_Kecukupan_Gizi_Yang_Dianjurkan_Untuk_Masyarakat_Indone sia.pdf
19. Gibson RS. *Principles of Nutritional Assessment*. 2nd ed. Oxford University Press; 2015.
20. Szeitz A, Bandiera SM. Analysis and measurement of serotonin. *Bioimed Chromatogr*. 2018;32(1). <https://doi.org/10.1002/bmc.4135>
21. Li Z, Wang W, Xin X, Song X, Zhang D. Association of total zinc, iron, copper and selenium intakes with depression in the US adults. *J Affect Disord*. 2018;228(November 2017):68-74. <https://doi.org/10.1016/j.jad.2017.12.004>
22. Doboszewska U, Wlaz P, Nowak G, Radziwon-Zaleska M, Cui R, Mlyniec K. Zinc in the Monoaminergic Theory of Depression: Its Relationship to Neural Plasticity. *Neural Plast*. 2017;2017. <https://doi.org/10.1155/2017/3682752>
23. Bonaventura P, Benedetti G, Albarede F, Miossec P. Zinc and its role in immunity and inflammation. *Autoimmun Rev*. 2015;14(4):277-285. <https://doi.org/10.1016/j.autrev.2014.11.008>
24. Lee DY, Kim E, Choi MH. Technical and clinical aspects of cortisol as a biochemical marker of chronic stress. *BMB Rep*. 2015;48(4):209-216. <https://doi.org/10.5483/BMBRep.2015.48.4.275>
25. Way BM, Taylor SE. The Serotonin Transporter Promoter Polymorphism Is Associated with Cortisol Response to Psychosocial Stress. *Biol Psychiatry*. 2010;67(5):487-492. <https://doi.org/10.1016/j.biopsych.2009.10.021>
26. Adam EK, Quinn ME, Tavemier R, McQuillan MT, Dahlke KA, Gilbert KE. Diurnal cortisol slopes and mental and physical health outcomes: A systematic review and meta-analysis. *Psychoneuroendocrinology*. 2017;83:25-41. <https://doi.org/10.1016/j.psycheneu.2017.05.018>
27. Nam SB, Kim K, Kim BS, et al. The Effect of Obesity on the Availabilities of Dopamine and Serotonin Transporters. *Sci Rep*. 2018;8(1):1-6. <https://doi.org/10.1038/s41598-018-22814-8>
28. Lischka J, Schanzer A, Baumgartner M, de Gier C, Greber-Platzer S, Zeyda M. Tryptophan Metabolism Is Associated with BMI and Adipose Tissue Mass and Linked to Metabolic Disease in Pediatric Obesity. *Nutrients*. 2022;14(2). <https://doi.org/10.3390/nu14020286>
29. Bernard A, May C Le, Dastugue A, et al. The tryptophan/kynurenine pathway: A novel cross-talk between nutritional obesity, bariatric surgery and taste of fat. *Nutrients*. 2021;13(4):1-15. <https://doi.org/10.3390/nu13041356>
30. Bouma EMC, Riese H, Ormel J, Verhulst FC, Oldehinkel AJ. Adolescents' cortisol responses to awakening and social stress: Effects of gender, menstrual phase and oral contraceptives. The TRAILS study. *Psychoneuroendocrinology*. 2009;34(6):884-893. <https://doi.org/10.1016/j.psycheneu.2009.01.003>
31. Kudielka BM, Kirschbaum C. Sex differences in HPA axis responses to stress: A review. *Biol Psychol*. 2005;69(1 SPEC. ISS.):113-132. <https://doi.org/10.1016/j.biopsycho.2004.11.009>

32. Oldehinkel AJ, Bouma EMC. Sensitivity to the depressogenic effect of stress and HPA-axis reactivity in adolescence: A review of gender differences. *Neurosci Biobehav Rev.* 2011;35(8):1757-1770. <https://doi.org/10.1016/j.neubiorev.2010.10.013>
 33. Bangasser DA, Curtis A, Reyes BAS, et al. Sex differences in corticotropin-releasing factor receptor signaling and trafficking: Potential role in female vulnerability to stress-related psychopathology. *Mol Psychiatry.* 2010;15(9):896-904. <https://doi.org/10.1038/mp.2010.66>
 34. Chrousos GP, Torpy DJ, Gold PW. Interactions between the hypothalamic-pituitary-adrenal axis and the female reproductive system: Clinical implications. *Ann Intern Med.* 1998;129(3):229-240. <https://doi.org/10.7326/0003-4819-129-3-199808010-00012>
 35. Cowen PJ, Browning M. What has serotonin to do with depression? *World Psychiatry.* 2015;14(2):158-160. <https://doi.org/10.1016/j.wpsyc.2014.12.007>
 36. Manthey L, Leeds C, Gilray EJ, et al. Antidepressant use and salivary cortisol in depressive and anxiety disorders. *Eur Neuropsychopharmacol.* 2011;21(9):691-699. <https://doi.org/10.1016/j.eurocnuro.2011.03.002>
-

The Role of Zinc Intake in Serotonin and Cortisol Level In Patient with Depression

ORIGINALITY REPORT

23%

SIMILARITY INDEX

20%

INTERNET SOURCES

17%

PUBLICATIONS

13%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|----|
| 1 | ri.conicet.gov.ar
Internet Source | 1% |
| 2 | Naintina Lisnawati, Nur Amin, Yanesti Nuravianda Lestari. "Sport drink containing maltodextrin to improve physical performance of Soccer athletes", AIP Publishing, 2023
Publication | 1% |
| 3 | www.termedia.pl
Internet Source | 1% |
| 4 | Jin-Sun Jun, Jun-Sang Sunwoo, Jung-Ick Byun, Jung-Won Shin, Tae-Joon Kim, Carlos H Schenck, Ki-Young Jung. "Emotional and Environmental Factors Aggravating Dream Enactment Behaviors in Patients with Isolated REM Sleep Behavior Disorder", Nature and Science of Sleep, 2022
Publication | 1% |
| 5 | "Original Research Abstract", European Heart Journal Supplements, 2019 | 1% |

6	Submitted to Universitas Diponegoro Student Paper	1 %
7	www.mdpi.com Internet Source	1 %
8	Submitted to University of North Carolina, Greensboro Student Paper	1 %
9	Submitted to Mount St. Marys College Student Paper	1 %
10	www.bsmiab.org Internet Source	1 %
11	www.cerascreen.co.uk Internet Source	1 %
12	hal.inrae.fr Internet Source	1 %
13	research.vumc.nl Internet Source	1 %
14	www.yumpu.com Internet Source	1 %
15	www.ncbi.nlm.nih.gov Internet Source	1 %
16	Submitted to Forum Perpustakaan Perguruan Tinggi Indonesia Jawa Timur II Student Paper	1 %

17	Miguel H. Vicco, Nazarena Pujato, Iván Bontempi, Luz Rodeles, Iván Marcipar, Oscar A. Bottasso. " β 1-Selective Adrenoceptor Antagonists Increase Plasma Levels of Anti-p2 β Antibodies and Decrease Cardiac Involvement in Chronic Progressive Chagas Heart Disease", Canadian Journal of Cardiology, 2014 Publication	1 %
18	hal.archives-ouvertes.fr Internet Source	1 %
19	repository.universitasbumigora.ac.id Internet Source	1 %
20	www.scielo.br Internet Source	<1 %
21	academic.oup.com Internet Source	<1 %
22	www.ryghealth.com Internet Source	<1 %
23	András Szeitz, Stelvio M. Bandiera. "Analysis and measurement of serotonin", Biomedical Chromatography, 2018 Publication	<1 %
24	Jiajia Wan, Wenxiu Duan, Kai Chen, Yanduo Tao, Jun Dang, Kehao Zeng, Yushu Ge, Jiang Wu, Dan Liu. "Selective and sensitive	<1 %

detection of Zn(II) ion using a simple peptide-based sensor", Sensors and Actuators B: Chemical, 2018

Publication

25

Submitted to Queen Mary and Westfield College

Student Paper

<1 %

26

medic.upm.edu.my

Internet Source

<1 %

27

www.frontiersin.org

Internet Source

<1 %

28

Submitted to Swinburne University of Technology

Student Paper

<1 %

29

espace.curtin.edu.au

Internet Source

<1 %

30

ejournal2.undip.ac.id

Internet Source

<1 %

31

Harmiyati Harmiyati, Ariawan Soejoenoes, Sri Wahyuni M, Kun Aristiati, Soeharyo Hadisaputro. "THE IMPACT OF HONEY ON CHANGE IN NUTRITIONAL STATUS IN CHILDREN WITH POOR NUTRITION", Belitung Nursing Journal, 2017

Publication

<1 %

32	Matthew J. Zdilla, Julia R. Saling, Leah D. Starkey. "Zinc sulfate taste acuity reflects dietary zinc intake in males", Clinical Nutrition ESPEN, 2016 Publication	<1 %
33	libstore.ugent.be Internet Source	<1 %
34	Submitted to Cardiff University Student Paper	<1 %
35	d-scholarship.pitt.edu Internet Source	<1 %
36	jneuroinflammation.biomedcentral.com Internet Source	<1 %
37	med-expert.com.ua Internet Source	<1 %
38	bapin-ismki.e-journal.id Internet Source	<1 %
39	Melita Stipancic, Walter Renner, Peter Schütz, Renata Dond. "Effects of Neuro-Linguistic Psychotherapy on psychological difficulties and perceived quality of life", Counselling and Psychotherapy Research, 2010 Publication	<1 %
40	Tao Li, Huihui Wang, Yaxin Zhang, Huiyuan Wang et al. "Comprehensive profiling of	<1 %

epigenetic modifications in fast-growing moso bamboo shoots", Plant Physiology, 2022

Publication

41

d.lib.msu.edu

Internet Source

<1 %

42

doi.org

Internet Source

<1 %

43

ouci.dntb.gov.ua

Internet Source

<1 %

44

www.discoveryjournals.org

Internet Source

<1 %

45

1library.net

Internet Source

<1 %

46

Hanmo Li, Andrew Bersten, Ubbo Wiersema, David Schembri, Elena Cavallaro, Dani-Louise Dixon, Shailesh Bihari. "Bolus intravenous 0.9% saline leads to interstitial permeability pulmonary edema in healthy volunteers", European Journal of Applied Physiology, 2021

Publication

<1 %

47

Meira Erawati, Nyoman Suci Widyastiti, Tri Indah Winarni, Edi Dharmana. "beta-Glucan Increases IFN-gamma and IL-12 Production of Peripheral Blood Mononuclear Cells with/without Induction of Mycobacterium

<1 %

tuberculosis Wild-type/Mutant DNA", The Indonesian Biomedical Journal, 2019

Publication

48	bmcinfectdis.biomedcentral.com Internet Source	<1 %
49	bmcp psychiatry.biomedcentral.com Internet Source	<1 %
50	ghrp.biomedcentral.com Internet Source	<1 %
51	nlistsp.inflibnet.ac.in Internet Source	<1 %
52	repositorio.unb.br Internet Source	<1 %
53	www.agrimaroc.org Internet Source	<1 %
54	www.dovepress.com Internet Source	<1 %
55	hdl.handle.net Internet Source	<1 %
56	Bjarni Sigurdsson, Sigurdur Pall Palsson, Olafur Aevansson, Maria Olafsdottir, Magnus Johannsson. "Saliva testosterone and cortisol in male depressive syndrome, a community study. The Sudurnesjamenn Study", Nordic Journal of Psychiatry, 2014 Publication	<1 %

57

Eirini Martinou, Irena Stefanova, Evangelia Iosif, Angeliki M. Angelidi. "Neurohormonal Changes in the Gut-Brain Axis and Underlying Neuroendocrine Mechanisms following Bariatric Surgery", *International Journal of Molecular Sciences*, 2022

Publication

<1 %

58

Roopa P Nayak, Ramesh Kumar Karri. "Evaluation of antidepressant properties of ethanolic extract of root of in Swiss albino mice", *Indian Journal of Pharmacy and Pharmacology*, 2022

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off

The Role of Zinc Intake in Serotonin and Cortisol Level In Patient with Depression

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
