

# Optimization of Eugenol Extraction from Clove Oil using Response Surface Methodology

*by* Bambang Cahyono

---

**Submission date:** 05-Mar-2020 08:47AM (UTC+0700)

**Submission ID:** 1269510355

**File name:** 53732-185059-1-SM.pdf (1.33M)

**Word count:** 3221

**Character count:** 16868

## Optimization of Eugenol Extraction from Clove Oil using Response Surface Methodology

Widayat<sup>1</sup>, Hadiyanto<sup>1</sup>, Bambang Cahyono<sup>2</sup> & Ngadiwiyanana<sup>2</sup>

<sup>1</sup> Department of Chemical Engineering, Diponegoro University, Indonesia

<sup>2</sup> Department of Chemistry, Diponegoro University, Indonesia

Correspondence: Hadiyanto, Department of Chemical Engineering, Diponegoro University, Indonesia. Tel: 62-247-460-058. E-mail: h.hadiyanto@undip.ac.id

Received: May 8, 2015

Accepted: June 29, 2015

Online Published: September 30, 2015

doi:10.5539/mas.v9n11p68

URL: <http://dx.doi.org/10.5539/mas.v9n11p68>

*The research is financed by MP3EI research grant 2013, Directorate General of Higher Education, Ministry of Education and Culture, Indonesia.*

### Abstract

The objective of this research was to obtain optimum condition of eugenol production from clove oil using a central composite design method. The main process occurred in the eugenol production was saponification and neutralization processes. In order to optimize these processes, the ratio of NaOH/clove oil and temperature were studied as design variables i.e. ratio of NaOH/clove oil=1:2.5-1:3.5 while temperature was varied between 40 and 60°C. The yield of eugenol was considered as the main response in of this experiment. The result showed that the optimum condition was achieved when the temperature and the ratio of NaOH/clove oil were 50°C and 2.75:1, respectively and the yield was 39.17%.

**Keywords:** eugenol, clove oil, saponification –neutralization process, yield of eugenol, and Response surface response methods (RSM)

### 1. Introduction

Since the production of clove oil has reached 250 tons in 2011, Batang regency has been considered as one of the largest producers of clove oil in Indonesia (Widayat et al, 2011). The clove oil is mostly extracted from part of clove plants (*Eugenia caryophyllata* Thunb) such as leaves, flower, and stem. The quality of cloves oil is determined by the content of phenol compound, especially eugenol (BSN, 2010) which is still considered as the main problem of clove oil in Indonesia. Currently, the clove oil still has high content of eugenol and it is higher than the National Indonesian Standard (SNI). It has been reported that most of clove oil at small medium entrepreneurship (SME) in Batang Regency have eugenol contents up to 80% (Widayat et al, 2014). Besides eugenol, the clove oil is also determined by  $\beta$ -Caryophyllene, an impurities that decrease the clove oil quality (Widayat et al, 2014).




Figure 1. The molecule structure of eugenol

However, besides it is resulted from side product of clove oil, eugenol has potential for industries. Eugenol or Phenol, 2-methoxy-4-(2-propenyl) (Figure 1) is the main component of clove oil and it is colourless, has spicy taste and special odour which therefore it was mostly used for fragrance and flavour industries. The eugenol is also mostly utilized in the perfume industries, flavor concentrates and in the pharmaceutical industries as an antiseptic and anesthetic drug. Oyedemi et al. (2009) reported that eugenol is efficient for the metabolic activity

4  
of bacteria *Listeria monocytogenes*, *Streptococcus pyogenes*, *Escherichia coli* and *Proteus vulgaris*. Furthermore, Cheng et al. (2008) has utilized this compound and sinamaldehyd as an anti-fungal for the type of fungus and *Laetiporus sulphureus*, which is considerable used as activity inhibitory. Shelly et al. (2010) used eugenol derivative such as methyl eugenol to effectively increase the fertility of insects. Sadeghian et. al., (2008) had utilized eugenol derivatives to inhibit the activity of the enzyme 15-lipogenase which is involved in many diseases such as asthma and lung cancer. The results showed that these compounds could inhibit the performance of the enzyme 15-lipogenase. Furthermore, Chami et al. (2004) also have tested eugenol as anti fungus *Candida albicans*. Vidhya and Devaraj (2011) experienced eugenol for inducing lung cancer, which showed inhibitory phenomena in lung cancer.

The price of clove oil is determined by its purity, the clove oil with purity under 70% will have price of Rp 120.000,00/ kg, while the clove oil with purity 98 % have price of 500.000/kg. Therefore, extraction and separation of eugenol from clove oil is an essential step to obtain such valuable product. According to USP (United States Pharmacopeia) standard, eugenol product must have purity higher than 98 %.

Another aspect for increasing eugenol contents is by implementation of chelating agent to reduce the darkness. The chelating agents that mostly used are as citric acid and sodium EDTA (Marwati et al, 2005). Commonly, this process can increase eugenol contents up to 80%. Another process for separation of eugenol from clove oil is by using saponification and distillation as well as fractionation. In this method, sodium hydroxide will be reacted with clove oil and then was neutralized by adding sulphuric acid or hydrochloride acid. The eugenol product then is separated using decantation and distillation process. This process could increase eugenol contents up to 82,6 % (minimum) (Anny, 2002; Sukarsono et al, 2003). However, the optimization is still required to estimate the associated variables in eugenol production. Therefore, the objective of this research was to conduct optimization of eugenol production from clove oil using neutralization process and fractional distillation by using Response Surface Methodology (RSM).

## 2. Method

### 2.1 Materials

The clove oil used in this study was obtained from essential oils cluster Batang District. The clove oil has eugenol contents of 80%.

The equipment used for saponification and distillation process is presented in Figure 2. The distillation was operated in vacuum pressure that obtained with vacuum pump. Distillation column used packing system.

### 2.2 Experimental Design

The experimental design used is Response Surface Methodology (RSM) as described in Table 1 with two variables: the operation temperature and the ratio of clove oil to sodium hydroxide (NaOH). Experiment process were performed on a laboratory scale and batch processes. The operating conditions used as follows:

#### a. Fixed Variables

- ✓ Clove oil volume (ml) : 150
- ✓ Materials : clove oil
- ✓ Operational time : 45 minute

#### b. Experimental variables:

- Temperature : (-1) : (40) and (+1) (60) °C
- Ratio NaOH/Clove oil (v/v) : (-1) (2,5) and (+1) (3,5)




Figure 2. Vacuum Distillation equipment

### 2.3 Procedure

The experiments began with oil density measurement and specific volume of clove leaf oil. Prior to experiment, clove oil was purified using citric acid (Widayat et al, 2014). A 150 ml of clove oil and NaOH were added under specified ratios of variables. The saponification reaction occurred for 45 minutes, and then followed by decantation for 1 day. The formed soaps were then neutralized with hydrochloride acid and were reacted for 5 minutes. This process was continued by decantation for 15 minutes. The eugenol product was purified by using distillation for 1.5 hours. The final product was collected and was measured for its volume and refractive index. The concentration of products was also analyzed by using GC and data was analyzed by using STATISTICA.

Table 1. Variable Central Composite Design for Experiment

Run	Block	X <sub>1</sub>	X <sub>2</sub>	Y
1	1	-1	-1	Y <sub>1</sub>
2	1	-1	+1	Y <sub>2</sub>
3	1	+1	-1	Y <sub>3</sub>
4	1	+1	+1	Y <sub>4</sub>
5	1	0	0	Y <sub>5</sub>
6	2	0	-1,424	Y <sub>6</sub>
7	2	0	+1,424	Y <sub>7</sub>
8	2	-1,424	0	Y <sub>8</sub>
9	2	+1,424	0	Y <sub>9</sub>
10	2	0	0	Y <sub>10</sub>

Where:

X<sub>1</sub> = Coded variables for ratio of oil to NaOH

X<sub>2</sub> = Coded variables for temperature

Y = Yield of eugenol (%)

### 2.4 Data Analysis

The responses of this experiment were density, viscosity and eugenol concentration. The eugenol concentration

was analyzed by using gas chromatography at Malang Polytechnic Laboratory. The liquid product was analysed by using gas chromatography (HP 5890, with HP 608 column) equipped with FID detector. The operating condition used helium as gas carries with flow rate of 20 ml/minute, temperature of 100-200 °C with temperature gradient 5°/minute and initial time 5 minute. Detector temperature was 275 °C. The product was analysed with internal standard methods. 6 µL eugenol standard solutions was added with internal standard (benzyl alcohol) 6 µL and 1 ml of solvent. The solution injected on GC equipment and conducted in triple runs. The response of detector was calculated with sample area divided internal standard area. The yield of eugenol was calculated according to Eq 1.

$$\text{Yield (\%)} = \frac{\text{weight of eugenol}}{\text{weight of clove oil}} \times 100 \quad (1)$$

### 3. Results

#### 3.1 Chromatography Analysis

Fig. 3 shows the chromatogram analysis of methanol(solvent), benzyl alcohol and eugenol with retention time of 3.39; 10.04 and 18.36 minute, respectively. According to this chromatogram, the eugenol contents in the product were varies between 75.10 to 96.00%. This result showed that the euogenol producton could achieve higher concentration of eugenol from clove oil. Figure 3a shows a chromatogram of eugenol and internal standard and Figure 3.b for sample product and both figures show similarities, although there was a small shift of retention time.




Figure 3. Chromatogram of eugenol (a). Eugenol and internal standards, (b). Sample with internal standard

### 3.2 Response Surface Methods

The results of statistic analysis was performed by developing mathematical model, t test, analysis of variance, pareto analysis as well as model validation. The polynomial equation of multi linear regression was proposed by following form:

$$Y = X_0 + \beta_1 X_1 + \beta_2 X_2 + \beta_{11} X_1^2 + \beta_{22} X_2^2 + \beta_{12} X_1 X_2 \quad (2)$$

Where:

Y = predicted response

$\beta_1, \beta_2$  = linear coefficient for 1, 2 variable

$\beta_{11}, \beta_{22}$  = squared term coefficient for 1,2 variable

$\beta_{12}$  = interactions variable coefficient

$X_1, X_2$  = non dimensional number of independent variables

The variable or coefficients were obtained by determining the multi linear regression by using STATISTICA software. Through this software, the coefficients were obtained and showed by Eq 3.

$$Y = 34.4639 - 0.5886 X_1 + 19.6363 X_1^2 - 5.6989 X_2 + 9.2652 X_2^2 + 3.2853 X_1 X_2 \quad (3)$$

This equation represents correlation between yield of eugenol and realted parmeters to obtain the yield. However, this variable of X1 and X2 are still in coded variable. The real values of condition process must be determined inversely.

In Eq 3, it is shown that the values of coefficient for X1 and X2 are negative. This leads to the fact that if the ratio of oil to NaOH (X1) and temperature (X2) increase then the yield of eugenol will decrease (for X1 and X2>0). In addition, the interaction between X1 and X2 and also quadratic variable are positive, which means that the increase of both variables will also increase of the eugenol yield. The complete statistical analysis of coefficient by using multivariate variables in Eq 3 is shown by Table 2.

Table 2. Regression analysis result

Parameters	Regression	Std.Err.	t(3)	p	-95.%	+95.%
Mean/Interc.	34.4639	11.0747	3.1119	0.0528	-0.7808	69.7085
Blocking	-5.5849	4.9528	-1.1276	0.3415	-21.3468	10.1770
X <sub>1</sub>	-0.5886	5.5374	-0.1063	0.9221	-18.2109	17.0337
X <sub>1</sub> <sup>2</sup>	19.6363	7.3252	2.6806	0.0750	-3.6759	42.9485
X <sub>2</sub>	-5.6989	5.5374	-1.0292	0.3791	-23.3212	11.9235
X <sub>2</sub> <sup>2</sup>	9.2652	7.3252	1.2648	0.2952	-14.0470	32.5773
X <sub>1</sub> X <sub>2</sub>	3.2853	7.8310	0.4195	0.7031	-21.6365	28.2070

Variance of mathematical model was analyzed and the results presented in Table 3. Table 3 can also be used to determine whether the independent variables simultaneously significant effect on the dependent variable. The degree of confidence was 0.05(Bos et al, 2005; Lazic, 2004)). The parameters have F value more than p for all parameters except for X1 quadratic variable (F value = 0.0113 p=0.9221) and interaction variable(F=0.1759 and p= 0.2953). The results of variance analysis support the t-test in Table 2 for X1 and X2 variable (single variable). It also shows that the mathematical model already in optimum condition (maximum / minimum) no stripes ascend and mathematical model can be used directly for optimization.

Table 3. Analysis of variance(ANOVA)

Parameter	SS	df	MS	F	p
<b>Blocks</b>	311.911	1	311.911	1.2716	0.3415
<b>X<sub>1</sub></b>	2.771	1	2.771	0.0113	0.9221
<b>X<sub>1</sub><sup>2</sup></b>	1762.671	1	1762.671	7.1858	0.0750
<b>X<sub>2</sub></b>	259.816	1	259.816	1.0592	0.3791
<b>X<sub>2</sub><sup>2</sup></b>	392.427	1	392.427	1.5998	0.2953
<b>X<sub>1</sub>X<sub>2</sub></b>	43.172	1	43.172	0.1759	0.7031
<b>Error</b>	735.896	3	245.299		
<b>Total SS</b>	3120.280	9			

Pareto diagram can be a help to assist determination of the most important parameter in the process(Lazic ,2004). Pareto diagram (Figure 4) show that a linear variable X2 has the smallest value. Therefore this variable can be neglected due to no having effect in the process. All histogram of variables don't cross the line  $p = 0.05$ . Pareto chart show quadratic variable of X22 which has a histogram near to line  $p = 0.05$ . This proofs that quadratic variable X22 has the most effect in the eugenol yield . This condition can be increased for obtaining the optimum condition.




Figure 4. Parreto Diagram

Figure 5 shows optimum condition in term of a surface response graph that consists of x and y axis as independent variable ( $X_1$  and  $X_2$ ) and z-axis as dependent variable or yield of eugenol (Y). Figure 5 has a minimum poin meaning that the process is already minimized.






Figure 5. Countour Graphics of optimization results material ratio and temperature




Figure 6. Model validations with experimental data

Mathematical model described by equations 3 was validated with experiments data (Figure 6). Figure 6 also shows that the mathematical model has lower significant regression value ( $R^2=0.764$ ). However, such value is still sufficient to obtain the optimum condition.

In Table 4, the critical value of dimensionless numbers for each variable. Critical dimensionless value obtained for X1 (ratio clove oil to sodium hydroxide) is -0.0109 and X2 (temperature) 0.3095. X1 and X2 critical value is substituted to equation 6 to obtained yield of eugenol. Yield of eugenol in minimum condition is 39.17%.



Table 4. Critical value results / optimization

	Solution: minimum		
	Observed	Critical	Observed
X <sub>1</sub>	-1.4142	-0.0109	1.4142
X <sub>2</sub>	-1.4142	0.3095	1.4142

#### 4. Discussion

The clove oil has been purified and separated from its eugenol contents. Eugenol which is one of indicator for quality of clove oil has been considered also as an important product for food industries. The result shows that 75-96% eugenol can be extracted from clove oil by using separation method of saponification, and distillation.

The Surface Response Method (SRM) has been employed for optimization and analysis of production and purification of eugenol from clove oil by using neutralization and fractionation distillation. The minimum yield was obtained as 39.17% under X<sub>1</sub> (ratio clove oil to sodium hydroxide) is -0.0109 (equal to 2.75:1) and X<sub>2</sub> (temperature) of 0.3095 (equal to 55°C). The regression coefficient for the mathematical model was achieved at R<sup>2</sup>= 0.7642.

The further research is required to increase the eugenol purity and yield. Conventional steam distillation is typically considered the best way to obtain essential oils including eugenol from clove oil. However, this method produces varying eugenol product qualities dependent upon the temperature, pressure and time used for distillation. An important point regarding steam distillation of eugenol oils is that the temperature involved in the process changes the molecular composition of the plant matter. However, in practice most essential oils including eugenol are obtained by distillation at rather elevated temperature, in order to optimize the yield. Other alternative were proposed to use process intensification through microwave, ultrasound or supercritical fluid extraction and distillation. However, their economical feasibility must also be considered. Further action is also required to investigate the eugenol risk to human in term of their use for food.

#### 5. Conclusion

This study showed the distillation process of eugenol out of clove oil. The distillation method give much better improvements in term of its yield and purity of products. The optimum condition was achieved when the temperature 55°C and the ratio of clove oil/NaOH was 2.75 :1.

Further improvement is required to increase the yield by using process intensification through microwave, ultrasound or supercritical technology.

#### Acknowledgement

The authors would like to express sincerely thanks to Higher Education Directorate, Ministry of Education and Culture for their support in providing MP3EI 2013 research grant.

#### References

- Anny, S. (2002). *Pengolahan Lanjut Minyak Atsiri dan Penggunaannya Dalam Negeri*. Workshop Nasional Minyak Atsiri 30 Oktober 2002, Dirjen Industri Kecil Dagang Menengah, Depperindag.
- Box, G. EP., Hunter, J. S., & Hunter, W. G. (2005). *Statistics for Experiments second edition*. John Wiley and Sons, New York.
- BSN (Badan Standarisasi Nasional). (2010). Standar Nasional Indonesia Minyak Daun Cengkeh. SNI 06-2387-2006 2006.
- Chami, N., Chami, F., Bennis, S., Trouillas, J., & Remmal, A. (2004). Antifungal treatment with carvacrol and eugenol of oral candidiasis in immunosuppressed rats. *Braz. J. Infect. Dis.*, 8(3), 217-226.
- Lazic, Z. (2004). *Statistics Experiments for Chemical engineering*. WILEY-VCH Verlag GmbH & Co. KGaA, Weinheim.
- Marwati, T., Rusli, M. S., Noor, E., & Mulyono, E. (2005). Peningkatan Mutu Minyak Daun Cengkeh Melalui Proses Pemurnian. *Jurnal Pasca Panen*, 2, 45-52.
- Oyedemi, S. O., Okoh, A. I., Mabinya, L.V., Pirochenva, G., & Afolayan, A. J. (2009). The proposed mechanism of bactericidal action of eugenol,  $\alpha$ -terpineol and  $\alpha$ -terpinene against listeria monocytogenes, Streptococcus pyogenes, Proteus vulgaris and Escherichia coli. *Afr. J. Biotechnol.*, 8(7), 1280-1286.
- Sadeghian, H., Seyedi, S. M., Saberi, M. R., Arghiania, Z., & Riazi, M. (2008). *Design and synthesis of eugenol*

- derivatives, as potent 15-lipoxygenase inhibitors. Bioorgan. Med. Chem., 16, 890-901.*
- Shelly, T. E., Edu, J., & McInnis, D. (2010). Pre-release consumption of methyl eugenol increases the mating competitiveness of sterile males of the oriental fruit fly, *Bactrocera dorsalis*, in large field enclosures. *J. Insect Sci., 10, 8.*
- Sukarsono, Dahroni, I., & Sucahyo, D. H. (2003). Kajian Pemisahan Komponen – komponen dari Minyak Cengkeh”. Yogyakarta Puslitbang Teknologi Maju BATAN.
- Vidhya, N. and S.N. Devaraj, (2011). *Induction of apoptosis by eugenol in human breast cancer cells.* Indian J. Exp. Biol., 49: 871-878.
- Widayat, Cahyono B., Hadiyanto & Ngadiwiyana. (2014). Improvement of Clove Oil Quality by Using Adsorption-distillation Process. *Research Journal of Applied Sciences, Engineering and Technology, 7(18), 3867-3871,*
- Widayat, Cahyono, B, Hadiyanto & Ngadiwiyono. (2011). Rancang Bangun dan Uji Alat Proses Peningkatan Minyak Cengkeh Pada Klaster Minyak Atsiri kabupaten Batang. *Jurnal Ilmu Lingkungan, 9(2), 57-61.*
- Widayat, Haidar, M. H., Nurdiana, L., Cahyono B., Ngadiwiyana, & Satriadi, H. (2013). Fractional Distillation of Clove Oil at Vacuum Pressure, Proceeding International Seminar 2013 Faculty of Engineering University Muhammadiyah Purwokerto.

#### **Copyrights**

Copyright for this article is retained by the author(s), with first publication rights granted to the journal.

This is an open-access article distributed under the terms and conditions of the Creative Commons Attribution license (<http://creativecommons.org/licenses/by/3.0/>).

# Optimization of Eugenol Extraction from Clove Oil using Response Surface Methodology

## ORIGINALITY REPORT

5%

SIMILARITY INDEX

3%

INTERNET SOURCES

2%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.ifrj.upm.edu.my">www.ifrj.upm.edu.my</a> Internet Source	<1%
2	Submitted to Asian Institute of Technology Student Paper	<1%
3	<a href="http://files.eric.ed.gov">files.eric.ed.gov</a> Internet Source	<1%
4	Submitted to Queen Margaret University College, Edinburgh Student Paper	<1%
5	<a href="http://es.scribd.com">es.scribd.com</a> Internet Source	<1%
6	Kate C. Blanco, Cristian J. B. de Lima, Rubens Monti, Joaquim Martins, Natalia S. Bernardi, Jonas Contiero. "Bacillus lehensis—an alkali-tolerant bacterium isolated from cassava starch wastewater: optimization of parameters for cyclodextrin glycosyltransferase production", <i>Annals of Microbiology</i> , 2011 Publication	<1%

7	<a href="http://mid.ism.ac.jp">mid.ism.ac.jp</a> Internet Source	<1%
8	T D Kusworo, Widayat, Budiyono, A A Siahaan, G K Iskandar, D P Utomo. "Development of Cellulose Acetate – PVP blend membrane and UV irradiation treatment to increase membrane selectivity for clove oil purification", Journal of Physics: Conference Series, 2019 Publication	<1%
9	<a href="http://phd.lib.uni-corvinus.hu">phd.lib.uni-corvinus.hu</a> Internet Source	<1%
10	<a href="http://www.academicjournals.org">www.academicjournals.org</a> Internet Source	<1%
11	Submitted to Flinders University Student Paper	<1%
12	Submitted to Liverpool John Moores University Student Paper	<1%
13	<a href="http://www.lifesciences.ic.ac.uk">www.lifesciences.ic.ac.uk</a> Internet Source	<1%

Exclude quotes  Off

Exclude matches  Off

Exclude bibliography  On

# Optimization of Eugenol Extraction from Clove Oil using Response Surface Methodology

---

## GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---