

The Differences of N-Acetyl- β - Glucosaminidase and β 2 Microglobulin levels in Patients with and without Early Diabetic Nephropathy

by Banundari Rachmawati

Submission date: 07-Apr-2023 03:36PM (UTC+0700)

Submission ID: 2058271513

File name: The Differences of N Acetyl beta Glucosaminidase and beta2.pdf (544.72K)

Word count: 4115

Character count: 21323

The Differences of N-Acetyl- β -Glucosaminidase and β 2 Microglobulin levels in Patients with and without Early Diabetic Nephropathy

Cynthia Citra, Edward Kurnia Setiawan Limijadi, Banundari Rachmawati

Department of Clinical Pathology, Faculty of Medicine, Diponegoro University/Dr. Kariadi Hospital, Semarang, Indonesia. E-mail: cynthiatjitra@gmail.com

ABSTRACT

Diabetic Nephropathy (DN) becomes the most serious microvascular complication, which is characterized by the presence of persistent albuminuria. N-acetyl- β -glucosaminidase is a dominant lysosomal enzyme in renal tubular epithelial cells. β 2 microglobulin is a low molecular weight protein produced by Major Histocompatibility Complex Class 1 (MHC-1) expressed in all nucleated cells. Both N-acetyl- β -glucosaminidase and β 2 microglobulin could be a new useful marker for early detection of DN. Analytic observational study with a cross-sectional approach was performed from May to July 2019 involving 27 non-diabetic patients (K1), 27 diabetic patients without DN (K2), and 27 diabetic patients with early DN (K3) at the Clinical Pathology Department of the Faculty of Medicine, Diponegoro University and Diabetic Clinic. Data presented in this study were age, gender, fasting blood glucose, blood pressure, and urine albumin creatinine ratio. N-acetyl- β -glucosaminidase levels between groups were analyzed using ANOVA, β 2 microglobulin levels between groups were analyzed using Kruskal-Wallis test, $p < 0.05$ was considered significant. This study showed that there were significant differences in NAG level and β 2M level among three groups consisting of non-DM patients, DM patients with and without early DN. It was suggested that both parameters can be used as alternative markers in diabetes mellitus with early DN.

Keywords: Diabetic nephropathy, N-acetyl- β -glucosaminidase, β 2 microglobulin, urine albumin creatinine ratio

INTRODUCTION

The prevalence of diabetes is increasing every year all over the world. Based on the data by the International of Diabetic Federation (IDF), the global prevalence rate of people with Diabetes Mellitus (DM) in 2014 was 8.3%. Indonesia was the 7th most populated country with DM in the world, which had 8.5 million people with DM.¹ The diabetic proportion based on Riset Kesehatan Dasar (RISKESDAS) in Indonesia was 10.9% with most of them at the age of 35-54 years old.²

Diabetic Nephropathy (DN) is one of the serious microvascular complications of diabetes. It is the main cause of chronic kidney disease to End-Stage Renal Disease (ESRD).³ Data of RISKESDAS in 2018 showed that the prevalence of the Indonesian population who had chronic kidney disease was 0.38%.² Diabetic nephropathy is a clinical syndrome characterized by persistent urine albumin (> 300 mg/g or > 200 μ /min), clinically at a minimum of two tests in a period of 3 to 6 months, followed by increased blood pressure and a decreased Glomerular Filtration Rate (GFR). Urine albumin in

diabetic patients is considered as a sensitive and specific marker of kidney damage to diagnose and give the treatment of early diabetic nephropathy.⁴ Diabetic nephropathy is diagnosed based on normal albumin level (< 30 mg/g creatinine) and microalbuminuria (30-300 mg/g creatinine).^{4,5}

Urine albumin is a parameter of endothelial dysfunction in which endothelial defense functions are lost and leakage of protein or albumin occurs. It can be predicted on early DN by measuring Albumin Creatinine Ratio (ACR).^{5,6} However, ACR has several limitations such as low sensitivity and large variability. In some cases, a decrease in GFR can occur with normal urinary albumin excretion. The high sensitivity (81.8%) and specificity (96.8%) of microalbuminuria is the reason why it is necessary to perform further studies by using a marker with higher sensitivity.^{7,8} Several markers of tubular and glomerular damage have been investigated for early DN markers.

N-acetyl- β -glucosaminidase (NAG) is a lysosomal enzyme found in proximal tubular epithelial cells and has a relatively high molecular weight. Therefore, it is not filtered through the glomerulus and is only

released into the urine after tubular damage. N-acetyl- β -glucosaminidase has a higher diagnostic level compared to albumin excretion in diabetic patients, even NAG level can increase in normoalbuminuria.^{7,8} Sheira *et al.* suggested that NAG could be used as a marker of early DN because urine NAG increased in tubulointerstitial damage.⁷

β 2 microglobulin is a low molecular weight protein produced by cells expressing Major Histocompatibility Complex Class 1 (MHC-1) and found in all nucleated cells. Because it is almost entirely filtered by the glomerulus, it can be used to determine the glomerular filtration rate.^{8,9} The precision and accuracy of GFR estimation with creatinine were dissatisfactory because it is influenced by an external factor such as the variation of muscle mass. β 2 microglobulin can be used as an additional marker to measure LFG.^{10,11} In addition, Abdullah *et al.* suggested that it also could be used as a marker of early DN.¹⁰

The researchers conducted a study to investigate the difference of NAG and β 2M levels in non-DM patients, type 2 DM patients without diabetic nephropathy, and type 2 DM patients with early diabetic nephropathy. This study aimed to gain information on alternative parameters as a marker of early DN.

METHODS

This was an observational analytic study with a cross-sectional design conducted between May to July 2019 by observing type 2 DM patients in Kagok Community Health Center, Semarang, and non-DM patients in the Department of Clinical Pathology, Medical Faculty of Diponegoro University. N-acetyl- β -glucosaminidase and β 2 microglobulin levels were measured using Enzyme-Linked Immunosorbent Assay (ELISA) method in GAKI Laboratory Medical Faculty Diponegoro University, whereas urine albumin (ACR) was tested on Sarana Medika Laboratory Semarang. Samples were taken consecutively. The research subjects were type 2 DM patients with and without diabetic nephropathy taken from Kagok Community Health Center, Semarang and non-DM patients taken from the Department of Clinical Pathology, Medical Faculty of Diponegoro University who were willing to participate in this study, had signed informed consent and met the inclusion and exclusion criteria. The characteristic of subjects (age, gender, blood pressure, fasting blood glucose, and urine albumin creatinine ratio) was collected from the patient's medical record.

The sample size in this study was 81 patients. Data were presented in median (minimum-maximum). They were divided into 3 groups; group I: 27 non-diabetic (healthy persons), group II: 27 DM patients without nephropathy, and group III: 27 DM patients with early DN. Inclusion criteria of non-DM patients in this study were males and females, aged 30-55 years old, fasting blood glucose < 110 mg/dL, normal body temperature, and normal blood pressure. Inclusion criteria of DM patients without nephropathy were males and females, aged 30-55 years old, fasting blood glucose > 110 mg/dL, normal body temperature and blood pressure (\leq 120/80 mmHg) with normoalbuminuria (ACR \leq 30 mg/g). Inclusion criteria of DM patients with early DN were males and females, aged 30-55 years old, fasting blood glucose > 110 mg/dL, normal body temperature, and normal blood pressure (\leq 120/80 mmHg) with microalbuminuria (ACR 30-300 mg/g). Exclusion criteria for three groups were patients with liver diseases, cardiovascular diseases, cancer, or smoking habit.

Data were processed using SPSS 15. Because the sample size was less than 50, the Shapiro-Wilk test was used to determine the data normality. Multivariate analysis was performed on each variable to determine sample characteristics. N-acetyl- β -glucosaminidase levels were firstly transformed due to abnormal distribution. However, because it remained abnormally distributed in spite of the transformation, the Kruskal-Wallis test was used. Contrastingly, due to normal distribution, β 2 microglobulin data were analyzed using the oneway ANOVA test. Both the statistical tests showed significant results, so the analysis was continued with the Post-Hoc test. $p < 0.05$ was significant.

The research has obtained ethical clearance from the Health Research Ethics Commission, Faculty of Medicine, Diponegoro University, Semarang with number 235/EC/FK-UNDIP/VI/2019.

RESULTS AND DISCUSSIONS

From the total of 90 subjects who took part in the study, 81 subjects met inclusion and exclusion criteria, consisting of 27 non-DM patients, 27 type 2 DM patients without nephropathy, and 27 type 2 DM patients with early diabetic nephropathy. However, 9 subjects dropped out because 3 subjects had a history of heart disease, 1 subject had a history of malignancy, 5 subjects had high blood pressure.

The characteristics of subjects were presented in the in mean \pm SD and median (min;max) (Table 1).

Table 1. General characteristics of research subjects

Variable	Group					
	Non-DM		DM without Nephropathy		DM with Early DN	
	Mean±SD	(n)%	Mean±SD	(n)%	Mean±SD	(n)%
Gender						
Male		9(33.3)		7(25.9)		9(33.3)
Female		18(66.7)		20(74.1)		18(66.7)
Age (year)	34.33±5.44		50.41±3.04		50.89±2.87	
Fasting blood glucose (mg/dL)	94.85±6.19		244,48±106,45		241,81±95.54	
Systolic (mmHg)	118,52±4.56		120.19±5.46		118,33±4.16	
Diastolic (mmHg)	78.15±3.96		80.00±5.55		78.52±4.56	
ACR (mg/g)	6.65±3.42		13.96±6.12		89.56±55.61	

Table 2. Differences in NAG and β 2M levels in non-DM, DM patients without nephropathy, and DM patients with early diabetic nephropathy

Group	N	NAG	p	β 2M	p
		Median (min-max) ng/mL		Mean ±SD ng/mL	
Non-DM	27	1.91 (0.56-4.38)	0.001 [†] *	2009.52±336.77	0.017 [§] *
DM without nephropathy	27	4.09 (0.89-13.5)		2190.85±212.07	
Non-DM	27	1.91(0.56-4.38)	<0.001 [†] *	2009.52±336.77	<0.001 [§] *
DM with early DN	27	6.88 (1.49-24.85)		2773.45±257.17	
DM without nephropathy	27	4.09 (0.89-13.5)	0.003 [†] *	2190.85±212.07	<0.001 [§] *
DM with early DN	27	6.88 (1.49-24.85)		2773.45±257.17	

NAG: N-acetyl- β -glucosaminidase, β 2M: β 2 microglobulin, *Significant ($p < 0.05$); [†] Post-Hoc Mann-Whitney. [§] Post-Hoc LSD

Table 1 showed that most of the subjects were female. The mean age of DM patients was higher than non-DM patients. The mean age of DM patients without nephropathy and early diabetic nephropathy was similar, which was 50 years old. Urine albumin levels of non-DM patients had a normal median (6.65 mg/g). Type 2 DM patients without nephropathy were included in the normoalbuminuria group because their median urine albumin level was 13.96 mg/g, while type 2 DM patients with early DN was included in the microalbuminuria group (30-300 mg/gr creatinine).

Due to the abnormal distribution of NAG levels in the three groups, data were then presented in the median (min-max). Contrastingly, due to the abnormal distribution of β 2M levels in the three groups, data were then presented in mean±SD. The analysis result of NAG and β 2M levels in the three groups can be seen in Table 2.

The results of the statistical test among all study groups were shown in Table 2. There were significant differences in N-acetyl- β -glucosaminidase levels between K1 and K2 ($p = 0.01$), K1 (non-DM and DM without nephropathy), and K3 ($p = < 0.01$), K2 and K3 ($p = 0.03$). β 2 microglobulin between K1 and K2 ($p = 0.02$), K1 and K3 ($p = < 0.01$), K2 and K3 ($p < 0.01$). N-acetyl- β -glucosaminidase and β 2 microglobulin levels were higher in DM patients compared to non-DM patients and increased higher in DM patients with early DN compared to non-DM patients and DM patients without nephropathy.

The percentage of female patients was greater than males in this study. The cause of this finding remains debatable among experts, but it was suspected that type 2 DM occurred in most females at the age of nearly half a century. This was supported by a study by Alotaibi *et al.*, which found four studies reported significantly higher prevalence

4
rates for type 2 DM patients in males than in females; one regional study from the Eastern province and two nationwide studies, conducted between 2004 and 2005 and between 2007 and 2009 reported significantly higher prevalence rates for type 2 DM patients among females than males.¹²

Non-DM subjects were younger compared to DM subjects with a mean age of 33 years. The mean age of DM patients with and without DN was almost similar, around 50 years. These results were similar to those in a study by Retnakaran *et al.*, which found that individuals aged ≥ 45 years had a greater risk of developing DM compared with individuals aged <45 years, and in accordance with several epidemiological studies, which suggested that the susceptibility level of type 2 DM was proportional with increasing age.¹³ According to RISKESDAS in 2018, the incidence of type DM was greatest at the age 35-54 years (10.9%).²

Albumin creatinine ratio in non-DM and type 2 DM subjects without nephropathy were normal (ACR levels < 30 mg/g) of 5 and 12, respectively. It is an early predictor of DN complications assessed by measuring ACR.⁵ DM patients with early DN was included in the microalbuminuria group (ACR levels of 30-300 mg/g). Urine albumin in DM patients is considered the best predictor of kidney damage in assisting early diagnosis of DN. It is a parameter of endothelial dysfunction in which endothelial defense functions are lost and leakage of protein or albumin occurs.^{5,14}

The median blood glucose level in non-DM patients was within the normal range of ≤ 126 mg/dL. The median blood glucose level in type 2 DM patients without nephropathy and with early DN was ≥ 126 mg/dL. The mean fasting blood glucose of type 2 DM patients without nephropathy was higher than type 2 DM with DN patients. This might be influenced by the duration of fasting time, oral antidiabetic drug doses, and medication compliance.

The statistical test result of NAG levels in non-DM patients was significantly lower than type 2 DM patients without nephropathy ($p=0.001$). According to a study by Hong *et al.*, NAG is a lysosomal enzyme, which is normally excreted in small amounts in urine through physiological exocytosis in proximal tubular cells, which was significantly higher in type 2 DM patients compared with healthy subjects.^{8,15} Kaufmann *et al.* also found that there was a positive relationship between urine NAG and creatinine clearance in type 2 DM patients. The increase in urine NAG was proportional to a decreased kidney function in the early stages of DN development.¹⁶

The increase in urine NAG levels was due to the increased activity of proximal tubular cell lysosomes because of injury of tubular cells.^{15,17} This study was in accordance with previous studies, including a study by Siddiqui *et al.*, which reported that Cystatin C, NGAL, and NAG biomarkers were significantly higher in Diabetic kidney disease.¹⁸

The NAG levels in non-DM patients were significantly lower than type 2 DM patients with early DN ($p<0.001$). Increased level of NAG portrayed the level of kidney tubules damage. Sheira *et al.* stated that an increase in NAG observed in three groups of patients with albuminuria was directly proportional to the severity of kidney disorders.⁷

Microalbuminuria is a clinical marker of diabetic nephropathy, but its clinical manifestation can only be found in stage three of the five stages of DN development.^{8,17} Lysosomal enzyme from the tubular cell injury is secreted into the urine during the development of diabetic nephropathy. The activity of lysosomal enzymes shows an increase in the development of DN earlier than microalbuminuria.^{7,17}

The NAG level of type 2 DM patients without nephropathy was significantly lower than type 2 DM patients with early DN ($p=0.003$). Increased NAG through the process of exocytosis by proximal tubular cells triggered proteinuria. This showed the relationship between urine NAG level and the severity of proteinuria.¹⁸

According to a previous study, the detection of microalbuminuria in DM patients showed glomerular involvement in early kidney damage.¹⁵ Recent studies showed tubular involvement in DN complications, as demonstrated by the presence of renal tubular protein and enzyme in urine. Tubular involvement preceded the glomerular because many of this protein and enzyme tubular could be detected even before microalbuminuria occurred. At the beginning of glomerular damage, increased albumin leakage could be absorbed by the tubules if its function was still normal that albuminuria wouldn't occur.^{14,19}

The mean of $\beta 2M$ levels in non-DM patients was lower than type 2 DM patients without nephropathy. The $\beta 2M$ level increased in accordance with the degree of renal dysfunction.⁹ The median of $\beta 2M$ levels in non-DM patients and type 2 DM patients without nephropathy was 2085,9 and 2161,5, respectively. The normal median of $\beta 2M$ levels indicated the absence of kidney dysfunction. The $\beta 2M$ measurement was used in this study to assess kidney function.¹⁰ Many studies had analyzed the ability of $\beta 2M$ to assess renal function and its levels

were found superior to creatinine serum because the decreasing of GFR just starts in stage 3 of diabetic nephropathy.¹¹

There were significant differences in β 2M levels in non-DM patients and type 2 DM patients without nephropathy ($p=0.017$). This might be due to increased serum β 2M level in inflammatory conditions. In type 2 DM patients, chronic hyperglycemia-induced the formation of AGE. Increased expression of AGE in the RAGE receptor in endothelial cells triggered oxidative stress and induced the production of proinflammatory cytokines.^{10,11} Valdet *et al.* stated that hyperglycemia significantly increased inflammatory markers such as TNF- α and IL-6, which played a role in the pathogenesis of type 2 DM and its complications such as atherosclerosis and neuropathy.²⁰ Vaia *et al.* also stated that hyperglycemia played a significant role in the production of inflammatory cytokines.²¹

The mean of β 2M level in non-DM patients was significantly lower than type 2 DM patients with early DN ($p<0.001$). Serum β 2M can be used as a marker of chronic kidney disease associated with kidney dysfunction, especially in glomerular disorders. Measurement of the GFR index was considered less sensitive in the early stages of the disease because it could be normal.²⁰ Colombo *et al.* reported that β 2M concentrations increased significantly in the normoalbuminuria and microalbuminuria groups compared with controls.²² Many studies identified the potential of β 2M as a marker for diabetic renal disease and end-stage renal disease.^{20,22}

The mean of β 2M levels in type 2 DM patients without nephropathy was significantly lower than type 2 DM patients with early DN ($p <0.001$). DM patients with early DN had a mean β 2M above the normal value of 1.100-2.600 ng/mL. The increase of β 2M occurred in patients with renal dysfunction.

This study was in accordance with several previous studies, which reported that the microalbuminuria group had an increase in β 2M compared to the normoalbuminuria group. Colombo *et al.* stated that β 2M level was higher in DM patients with microalbuminuria compared to patients without microalbuminuria and the control group.²²

This study did not consider the duration of DM, which might influence NAG and β 2M. Another study, which uses a larger sample size and considers other risk factors of diabetic nephropathy, which may affect study variables should be performed. A diagnostic study was also considered necessary to be conducted.

CONCLUSIONS AND SUGGESTIONS

This study showed that there were significant differences in NAG and β 2M levels in three groups, which each consisted of non-DM patients, DM patients with and without early DN. This research suggested that both parameters can be useful to be alternative markers in early DN.

REFERENCES

1. American Diabetes Association (ADA). Standard of medical care in diabetes. *Diabetes Care*, 2017; 40(3): s4-s128.
2. Badan Penelitian dan Pengembangan Kesehatan Kementerian Kesehatan Republik Indonesia. *Riskesmas* 2018. Jakarta, Kementerian Kesehatan Republik Indonesia. 2018; 123-143
3. Qi C, Mao X, Zhang Z, Wu H. Classification and differential diagnosis of diabetic nephropathy. *Hindawi Journal of Diabetes Research*, 2017; 2017: articleID 8637138.
4. Lin CY, Chang YH, Yang SY, Wu KD, Chu TS. Update of pathophysiology and management of diabetic kidney disease. *Journal of the Formosan Medical Association*, 2018; 117(8): 662-675.
5. Wei C, Linlin M. Significance of joint detection of urinary proteins with low molecular weight as prognostic indicators for early diagnosis of diabetic nephropathy. *Int J Clin Exp Med*, 2016; 9(6): 1877-82.
6. Marketou N, Gantenbein CK, Marketos N, Chrousos G, Papassotiropoulos I. Biomarkers of diabetic nephropathy: A 2017 update. *Critical Reviews in Clinical Laboratory Sciences*, 2017; 54(5): 326-342.
7. Sheira G, Noreldin N, Tamer A, Saad M. Urinary biomarker N-acetyl- β -D-glucosaminidase can predict severity of renal damage in diabetic nephropathy. *J Diab Metab Disord*, 2015; 14(4): 33-6.
8. Hong N, Lee M, Park S, Lee HY, Jin MS, Hyeon K. Elevated urinary N-acetyl- β -D-glucosaminidase with high glycoalbumin to haemoglobin A1c ratio in diabetes patients with early diabetic kidney disease. *Scientific Reports*, 2018; 8(3): 6710-15.
9. Inker L, Tighiouart H, Coresh J, Foster M, Anderson A, Beck GJ, Contreras G. GFR estimation using β -trace protein and β 2-microglobulin in CKD. *Am J Kidney Dis*, 2016; 67(1): 40-8.
10. Abdullah, Kadam P, Yachha M, Srivastava G, Pillai A, Pandita A. Urinary beta-2 microglobulin as an early predictive biomarker of acute kidney injury in neonates with perinatal asphyxia. *Eur J Pediatr* 2022; 181(1): 281-286.
11. Barton K, Kakajiwala A, Dietzen D, Goss CW, Gu H, Dhamidharka V. Using the newer kidney disease: Improving global outcomes criteria, beta-2-microglobulin levels associate with severity of acute kidney injury. *Clinical Kidney Journal*, 2018; 1-6.
12. Alotaibi A, Perry L, Gholizadeh L, Ganmi A. Incidence and prevalence rates of diabetes mellitus in Saudi

- Arabia: An overview. *Journal of epidemiology and Global Health*, 2017; 211-18.
13. Nanayakkara N, Ranasinha S, Gadowski AM, Davis W, Flack JR, Wischer N, Andrikopoulos S, Zoungas S. Age-related differences in glycaemic control, cardiovascular disease risk factors and treatment in patients with type 2 diabetes: A cross-sectional study from the Australian National Diabetes Audit. *BMJ Open*, 2017; 8(8): e020677.
 14. Persson F, Rossing P. Diagnosis of diabetic kidney disease: state of the art and future perspective. *Kidney International Supplements*, 2018; 8: 2–7.
 15. Zeni L, Norden AG, Cancarini G, Robert J. A more tubulocentric view of diabetic kidney disease. *J Nephrol*, 2017; 30: 701–717.
 16. Kaufmann M, Schlossbauer M, Hubauer U, Stadler S, Fischer M, *et al.* N-acetyl-beta-D-glucosaminidase: A potential biomarker for early detection of acute kidney injury in acute chest pain. *Nephrology*, 2019; 1–9.
 17. Gluhovschi C, Gluhovschi G, Timar B. Urinary biomarkers in the assessment of early diabetic nephropathy. *J Diabetes Research*, 2016; 46(26): 1-13.
 18. Siddiqui K, Al-Malki B, George TP, Nawaz SS, Rubeaan KA. Urinary N-acetyl-beta-D-glucosaminidase (NAG) with Neutrophil Gelatinase-Associated Lipocalin (NGAL) improves the diagnostic value for proximal tubule damage in diabetic kidney disease. *Biotech*, 2019; 9(3): 66.
 19. Omozee EB, Okaka EI, Edo AE, Obika LF. Urinary N-acetyl-beta-d- glucosaminidase levels in diabetic adults. *Journal of Laboratory Physicians*, 2019; 11(1): 1-4.
 20. Valdet T, Rame S, Miftari R, Haxhibeqiri V. Association of beta-2 microglobulin with inflammation and dislipidemia in high-flux membrane hemodialysis patients. *Med Arch*, 2016; 70(5): 348-50.
 21. Vaia D, Despina K. The relationship between glycemic control, beta2-microglobulin and inflammation in patients on maintenance dialysis treatment. *J Diabetes Metab Disord*, 2015; 14(2): 34-40.
 22. Colombo M, Helen C, Farran B, Hees S, Groop L, *et al.* Serum kidney injury molecule 1 and β 2-microglobulin perform as well as larger biomarker panels for prediction of rapid decline in renal function in type 2 diabetes. *J Diabetologia*, 2019; 62(1): 156–68.

The Differences of N-Acetyl- β -Glucosaminidase and β 2 Microglobulin levels in Patients with and without Early Diabetic Nephropathy

ORIGINALITY REPORT

20%

SIMILARITY INDEX

12%

INTERNET SOURCES

14%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	ejmcm.com Internet Source	4%
2	Kalia, K.. "Non-enzymatic glycosylation of immunoglobulins in diabetic nephropathy", <i>Clinica Chimica Acta</i> , 200409 Publication	1%
3	link.springer.com Internet Source	1%
4	Abdulellah Alotaibi, Lin Perry, Leila Gholizadeh, Ali Al-Ganmi. "Incidence and prevalence rates of diabetes mellitus in Saudi Arabia: An overview", <i>Journal of Epidemiology and Global Health</i> , 2017 Publication	1%
5	Submitted to Universitas Diponegoro Student Paper	1%
6	<i>Clinical Nephrotoxins</i> , 2003. Publication	1%

7	C. Bazzi. "Urinary N-acetyl-beta-glucosaminidase excretion is a marker of tubular cell dysfunction and a predictor of outcome in primary glomerulonephritis", <i>Nephrology Dialysis Transplantation</i> , 2002 Publication	1 %
8	www.science.gov Internet Source	1 %
9	Vuthi Khanijou, Neda Zafari, Melinda T. Coughlan, Richard J. MacIsaac, Elif I. Ekinci. "Review of potential biomarkers of inflammation and kidney injury in diabetic kidney disease", <i>Diabetes/Metabolism Research and Reviews</i> , 2022 Publication	1 %
10	downloads.hindawi.com Internet Source	1 %
11	www.frontiersin.org Internet Source	1 %
12	dagensdiabetes.se Internet Source	<1 %
13	Submitted to iHeed Student Paper	<1 %
14	Monika Sharma, Varsha Pandey, Chandra Kant Sharma. "Advanced Approaches Based on Expression of Different Genes and	<1 %

Biomarkers for the Diagnosis and Treatment of Renal Disorders", Critical Reviews in Eukaryotic Gene Expression, 2019

Publication

15

Christos P. Argyropoulos, Shan Shan Chen, Yue-Harn Ng, Maria-Eleni Roumelioti et al. "Rediscovering Beta-2 Microglobulin As a Biomarker across the Spectrum of Kidney Diseases", Frontiers in Medicine, 2017

Publication

<1 %

16

Ji Zhou, Rui Peng, Tianju Li, Xiaomei Luo, Huimin Peng, He Zha, Pin Yin, Li Wen, Zheng Zhang. "A potentially functional polymorphism in the regulatory region of let-7a-2 is associated with an increased risk for diabetic nephropathy", Gene, 2013

Publication

<1 %

17

Nina Vodošek Hojs, Sebastjan Bevc, Robert Ekart, Nejc Piko, Tadej Petreski, Radovan Hojs. "Mineralocorticoid Receptor Antagonists in Diabetic Kidney Disease", Pharmaceuticals, 2021

Publication

<1 %

18

Joana Barbosa, Juliana Faria, Fernanda Garcez, Sandra Leal et al. "Repeated Administration of Clinical Doses of Tramadol and Tapentadol Causes Hepato- and

<1 %

Nephrotoxic Effects in Wistar Rats", Pharmaceuticals, 2020

Publication

19

Submitted to Writtle Agricultural College

Student Paper

<1 %

20

Fernández, José A., Peter Bankhead, Huiyu Zhou, J. Graham McGeown, and Tim M. Curtis. "Automated Detection and Measurement of Isolated Retinal Arterioles by a Combination of Edge Enhancement and Cost Analysis", PLoS ONE, 2014.

Publication

<1 %

21

academic.oup.com

Internet Source

<1 %

22

jhoonline.biomedcentral.com

Internet Source

<1 %

23

worldwidescience.org

Internet Source

<1 %

24

Temesgen Fiseha. "Urinary biomarkers for early diabetic nephropathy in type 2 diabetic patients", Biomarker Research, 2015

Publication

<1 %

25

repositorio.ufmg.br

Internet Source

<1 %

26

www.isainsmedis.id

Internet Source

<1 %

27

Gulnaz Bano, Mohammad Tarique Imam, Ram Bajpai, Ghada Alem, Varun Kumar Kashyap, Anwar Habib, Abul Kalam Najmi. "Expression of Angiopoetin-Like Protein-4 and Kidney Injury Molecule-1 as Preliminary Diagnostic Markers for Diabetes-Related Kidney Disease: A Single Center-Based Cross-Sectional Study", *Journal of Personalized Medicine*, 2023

Publication

<1 %

28

Quang Thuan Huynh, Nguyet Minh Pham, Diep Thao Pham, Minh Thi Hoang et al. "

Evaluation of Urinary Neutrophil Gelatinase Associated Lipocalin and Kidney Injury Molecule-1 as Diagnostic Markers for Early Nephropathy in Patients with Type 2 Diabetes Mellitus

", *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 2020

Publication

<1 %

29

Yun-Ting Liu, Li-Ying Lin, Chi-Wei Tuan, Chun-Ying Yang, Pei-Ling Tang. "Analyzing the Association HbA1c control by Depression, social participation and Utilizing Self-management Questionnaire", *Diabetes Research and Clinical Practice*, 2019

Publication

<1 %

30

Internet Source

<1 %

31

krcp-ksn.org
Internet Source

<1 %

32

www.fda.gov
Internet Source

<1 %

33

www.jocmr.org
Internet Source

<1 %

34

www.wjgnet.com
Internet Source

<1 %

35

Jia Chen, Wei-Wei Zhang, Ke-Hong Chen, Li-Rong Lin et al. "Urinary DcR2 is a novel biomarker for tubulointerstitial injury in patients with diabetic nephropathy", American Journal of Physiology-Renal Physiology, 2017
Publication

<1 %

36

Samuel N. Heyman, Itamar Raz, Jamie P. Dwyer, Roni Weinberg Sibony, Julia B. Lewis, Zaid Abassi. "Diabetic Proteinuria Revisited: Updated Physiologic Perspectives", Cells, 2022
Publication

<1 %

37

Samuel Uwaezuoke. "The role of novel biomarkers in predicting diabetic nephropathy: a review", International Journal

<1 %

of Nephrology and Renovascular Disease, 2017

Publication

38

C. Bazzi. "Urinary N-acetyl-beta-glucosaminidase excretion is a marker of tubular cell dysfunction and a predictor of outcome in primary glomerulonephritis", Nephrology Dialysis Transplantation, 11/01/2002

Publication

<1 %

39

Yongzhang Qin, Shuang Zhang, Xiaofang Shen, Shunming Zhang et al. "Evaluation of urinary biomarkers for prediction of diabetic kidney disease: a propensity score matching analysis", Therapeutic Advances in Endocrinology and Metabolism, 2019

Publication

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

The Differences of N-Acetyl- β -Glucosaminidase and β 2 Microglobulin levels in Patients with and without Early Diabetic Nephropathy

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6
