

The appearance of the humpback whale (*Megaptera novaeangliae*) during the rainy season in the Alor Waters, Indonesia.

by Max Rudolf Muskananfolo


Submission date: 30-May-2022 07:34PM (UTC+0700)

Submission ID: 1847178531

File name: ngliae_during_the_rainy_season_in_the_Alор_Waters,_Indonesia.pdf (1.24M)

Word count: 5309

Character count: 28232



The appearance of the humpback whale (*Megaptera novaeangliae*) during the rainy season in the Alor Waters, Indonesia

¹Jahved F. Maro, ²Agus Hartoko, ²Sutrisno Anggoro, ²Max R. Muskananfolo, ³Lumban N. L. Toruan, ⁴Micael Dakahamapu, ⁵Erick Nugraha

¹ Faculty of Fisheries and Marine Science, Coastal Resources Management Doctoral Program, Diponegoro University, Semarang, Indonesia; ² Faculty of Fisheries and Marine Science, Coastal Resources Management, Diponegoro University, Semarang, Indonesia; ³ Faculty of Fisheries and Marine Science, Nusa Cendana University, East Nusa Tenggara, Indonesia; ⁴ Environmental Practitioners, East Nusa Tenggara, Indonesia; ⁵ Faculty of Fishing Technology, Jakarta Technical University of Fisheries, South Jakarta, Indonesia.

Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. The appearance of humpback whale (*Megaptera novaeangliae*) in Alor Waters, Indonesia, located in the Eastern Indian Ocean, is one of the new findings in the marine protected area of Pantar Strait and the surrounding sea of Alor regency. Since 2015, when the waters of Alor regency have been designated as a marine protected area, *M. novaeangliae* specimens have never been found in these waters. The research aimed to track their trajectory, based on bio-oceanography predictions, in the marine protected area of Pantar Strait and its surrounding sea. This research used a field data survey, and database sources like the Copernicus Marine Service Data, Argo Floats and GEBCO data. The field survey was conducted from March to December 2020. The results of the research showed that around 15 *M. novaeangliae* specimens crossed the Alor waters on December 1st 2020 in the afternoon, from 05.00 to 05.45 pm. They were jumping above the water surface for about 3 seconds and diving into the water, then they were reappearing at the water surface for about 5 minutes and jumping again for about 5 seconds. The coordinates of the appearance were -8.123086° latitude and 124.063298° longitude. The average physical factors of the waters were: a sea surface temperature ranging from 25 to 31°C, a chlorophyll content ranging between 0.2 and 0.8 mm m⁻³, a waters depth of >50 m, and a speed of the water surface currents of 0.2 to 0.6 m s⁻¹. The type of dominant zooplankton found is the crustacean from the phylum Arthropoda, with a density of 1,957 individuals L⁻¹. These findings will inform the policies concerning the *M. novaeangliae* protection in the Alor waters.

Key Words: oceanographic variables, remote sensing, Pantar Strait, mammals.

Introduction. Indonesian waters are inhabited by 31 species of cetaceans (whale, porpoise, dolphin), twelve of which being whales and the others being dolphins and dugongs (*Dugong dugong*) (Rosas et al 2012). Both resident and migrant, these species are distributed throughout the coastal waters, towards the deep sea zone (Salim 2011). Several species of cetaceans are true migratory, using the Eastern Indonesian waters as a migration route from the Indian Ocean to the Pacific Ocean and vice-versa, crossing the waters of the Komodo Islands, Solor-Lembata (East Nusa Tenggara), Banda Sea (Moluccas), Southeast Sulawesi, North Sulawesi and Papua (Sorong and Fakfak) (Salim 2011). The waters of Eastern Indonesia, particularly in several inter-island deep canals, are assumed to serve as the entry point for the migration routes of marine mammals (cetaceans), such as whales and dolphins.

Nowadays, the protection of marine mammals is a priority for the marine biology research, being based on the study of migratory and distribution patterns, especially in the cetacean's case (Dréo et al 2019). The decrease of the population of cetaceans is due to the influence of human activities, resulting in pollution and environmental destruction

6
(Bejder et al 2019). The humpback whale (*Megaptera novaeangliae*) is a cosmopolitan species that exists in all major ocean basins, from tropical to arctic waters (Clapham 1992; Dawbin 1959; Secchi et al 2011). According to several studies, *M. novaeangliae* is one of the over-hunted species, which causes a significant decrease of its populations in the southern hemisphere, including the South Atlantic and South America (IWC 1998; Chittleborough 1965; Findlay et al 2011; Secchi et al 2011).

The waters of the Alor Strait and its surroundings are located in the Province of East Nusa Tenggara, which is one of the areas of the Indonesian Exclusive Economic Zone (ZEEI), next to the west coast. Timor Leste and Australia are on the trajectory of the Indonesian Throughflow, considered as the confluence of two current masses from the Pacific Ocean and Indian Ocean (Putra et al 2016).

The waters of the Alor Strait are unique, with dynamic oceanographic variables. Significant changes occur in the sea surface, due to the vertical temperature variability and salinity during the southeast monsoon season. The dynamics of the waters in the surface layer is influenced by the monsoon wind patterns, determining the upwelling in the Savu Sea waters. The process of stirring up the water masses (upwelling) affects the living conditions of the phytoplankton, the hydrology and the nutrient enrichment in the waters (Sediadi 2004; Packard et al 2015). Among the most significant impacts of the upwelling are an increased fertility (abundance of plankton as natural food) and an increased sea water temperature (warm), providing comfort to a group of cetacean species, including the *M. novaeangliae* and the bottle-nose dolphin (*Tursiops truncatus*), which are migrating to the waters of the Alor Strait (Mujiyanto et al 2017).

The Alor Strait is part of the Pantar Strait Conservation Area and its surrounding sea which has been arranged in the Ministerial Regulation No. 5/KEPMEN-KP/2015, by the Ministry of Marine Affairs and Fisheries Republic of Indonesia. The appearance of the *M. novaeangliae* in the Alor Strait waters was initially informed by some fishermen in the location. The fishermen reported the information to the researchers, so further monitoring and research on this species could be conducted. Therefore, the objective of this study was to design and implement a monitoring research experiment on *M. novaeangliae* in the Alor Strait waters, using the field observation of the bio-oceanography.

16

Material and Method

Description of the study site. The research was conducted in the Pantar Strait, Alor, East Nusa Tenggara from March to December 2020. The field survey method, modified from Dharmadi et al (2017) and Mujiyanto et al (2017), was the zig-zag track method using a boat of 10 GT with two observation decks, involving two different groups of on-deck observers. Water samples were taken at every cetaceans appearance event, in order to determine the plankton distribution. Field photography documentation of cetaceans appearance was also performed for further identification. Pictures were taken using Canon camera and drone. The ship was traveling at a speed of 7 to 8 knots on each survey trajectory. The map of the research area can be seen in Figure 1.

23

Regarding the processing of Alor Strait oceanographic data, the Aqua MODIS (Moderate Resolution Imaging Spectroradiometer) satellite image data, Copernicus Marine Service Data, ARGO (Array for Real-Time Geostrophic Oceanography) Float and GEBCO data (General Bathymetric Chart of the Oceans) were used. The data were processed using the ArcGIS 10.8, applying the kriging interpolation method (Wirasatriya et al 2020; Hartoko 2010; Hartoko et al 2019). Data processed in the study are the sea surface temperature, chlorophyll-a, current speed, wind speed and depth. Data processing was carried out at the Marine Geomatics Research Laboratory, Diponegoro University, Semarang.




Figure 1. Map of research locations for *Megaptera novaeangliae* in the Alor Strait (MMAF Decree 2015).

Results and Discussion

Appearance and distribution of the *M. novaeangliae* in the Alor Strait. The study discovered three points of the *M. novaeangliae* appearance in the Alor Strait, they can be seen in Figure 2.




Figure 2. Appearance of *Megaptera novaeangliae* at Alor Strait.

Figure 2 shows the of appearance of *M. novaeangliae* at each observation point. The appearance of the *M. novaeangliae* in Alor waters occurred in the afternoon from 05.00 to 05.45 pm on December 1st 2020, which was precisely during the rainy season in Indonesia. *M. novaeangliae* appearance event at each point was recorded in the pictures presented in Figures 3, 4 and 5.



Figure 3. (A, B, C & D) The appearance of a *Megaptera novaeangliae* with jumping behavior at the Point 1 (coordinates 124.063298° longitude and 8.123086° latitude) (original).




Figure 4. (A, B, C & D) The appearance of a *Megaptera novaeangliae* with jumping behavior at the Point 2 (coordinates: 124.019435° longitude and 8.152566° latitude) (original).




Figure 5. (A, B, C & D) Appearance of the *Megaptera novaeangliae* with Salto's demeanor and threw his body into the body of water at the Point 2 (coordinates: 124.019435° longitude and 8.152566° latitude) (original).

The appearance of the *M. novaeangliae* at three points in the Alor Strait waters had the same behavior: it was jumping over the sea surface, slamming its tail into the water, flipping and hitting its body in the water, which can be seen in Figures 4, 5, and 6. The *M. novaeangliae* was jumping over the sea surface using its elongated pectoral fins during maneuver and swam back into the water, then came back to the water surface for about five minutes, jumped back for about five seconds and sang (made a sound at the sea level). The map of the zone of the *M. novaeangliae* appearance is shown in Figure 6.




Figure 6. Map of the appearance of the *Megaptera novaeangliae* in the Alor Strait.

According to several studies, *M. novaeangliae* leaps out of the water (jumping through the seawater), hits the surface of the water with its long fins, then hits the water surface with its tail (lobtail) and freezes in a vertical position with its head above the water (spyhop); it breaks through by swimming from the depth to the surface of the water at an oblique angle, then jumps into the air at a variable angle (of slope upto 70 degrees), over the sea surface (Clapham 2009; Galvin 2006; Fish & Battle 1995; Parsons et al 2008; McSweeney et al 1989; Zerbini et al 2004; Calambokidis & Barlow 2004; Kügler et al 2020; Mohebbi-Kalkhoran et al 2019). Then, it rotates on its long axis, landing on its back with its belly rising up and eventually singing. Singing is an important point of the *M. novaeangliae* social behavior, considering that the sound can be related to the intrasexual competition and/or intersexual selection.

The tubercles of *M. novaeangliae* in front of its body are used as an enhanced lifting device to control the water flow over its fins and maintain the lift at high angles of attack. The morphology of the fin exhibits high maneuverability related to its unique feeding behavior. Its eyeballs are similar to other cetaceans and show adaptation to diving and migration, contributing to the perception of temperature, pressure and light differences (Woodward et al 2006; Hampe et al 2015; Rodrigues et al 2014).

The group of the *M. novaeangliae* found in the Alor Strait during the rainy season in December 2020 was assumed to have migrated from the Atlantic Ocean and Australian seawaters. This was due to the appearance of the *M. novaeangliae* in the Alor Strait waters with a distance of ± 700 km from Australian waters, migrating for food and mating. According to Palsbøll et al (1997) and to the IWC (1998), *M. novaeangliae* could cover a distance of 8,000 km from the breeding grounds, in the tropical zones, from July to October in the southern hemisphere and from December to March in the northern hemisphere. In order to gain enough strength and body mass for giving birth and intensively breastfeeding their babies for several months, the pregnant *M. novaeangliae* female could swim thousands kilometers to the nutrient-rich arctic or mid-temperature waters, where they could find their food. Usually, the pregnant *M. novaeangliae* female arrives earlier than other whales which are not pregnant or in their adolescence period (IWC.int/humpback-whale 2021).

According to Craig et al (2014), the pregnant *M. novaeangliae* female seeks for recessed waters (bays) to give birth so that its newborn can be protected. Based on this research, considering the condition of the Alor Strait waters which are protected and close to coastal areas, it is assumed that one of the reasons of the *M. novaeangliae* appearance in the Alor Strait waters in the rainy season of December 2020, was the migration for reproduction or giving birth. Furthermore, considering its group behavior (throwing its body in the air and diving by poking its tail in the water opening its mouth at the water surface and releasing large bubbles in the waters) it can be assumed that the *M. novaeangliae* migrated for foraging and mating in the Alor Strait waters. This research found and identified zooplankton from three classes of different phyla, at three points where *M. novaeangliae* specimens were observed. The class of Crustacea of the phylum Arthropod was represented with the highest density of zooplankton. A total of 1,957 individuals L^{-1} were found at the three points, as it can be seen in Table 1 and Figure 7.

Table 1

Density of zooplankton at the Point 3 on the appearance of *Megaptera novaeangliae* at Alor Strait

Zooplankton	Class	ST1	ST2	ST3	Total
		Ind L^{-1}			
Protozoa	Sarcodina	130	127	145	402
Annelida (=annulata)	Polychaeta	95	30	72	197
Arthropoda	Crustacea	677	647	633	1957
Mollusca	Gastropoda	112	145	105	362
Echinodermata	Ophiuroidea	6	10	23	39




Figure 7. (A, B, C & D) Diversity of crustacea dominant at the Point 3 of the *Megaptera novaeangliae* appearance (original).

According to Nybakken (1992), the zooplankton species which are the most important as fish food are from the class of the crustaceans, belonging to the arthropod phylum. Crustaceans are animals that have cells consisting of chitin or calcium, which are difficult to digest. Crustaceans can be divided into 2 groups: Entomostraca or low-level crustaceans and Malacostracea or high-level crustaceans.

According to Werth et al (2019), Burkard et al (2015) and Chen (2012), *M. novaeangliae* has cranial elevation, i.e. the expulsion of filtered water begins with a small splash in the anterior of the mouth, followed by a continuous outflow in the middle or posterior area of the mouth apart from the splash in the mouth which is free of turbulence during swallowing.

The submersion of the *M. novaeangliae* head creates a vortex in the undersea and water surface, large enough for the purpose of gathering prey, such as groups of small shrimp and small fish to eat. Due to the abundance of zooplankton of the class Crustacea at the three points of the *M. novaeangliae* specimens appearance, it can be assumed that one of the reasons of their appearance in the Alor waters is migrating for food. This is because the type of plankton from the crustacean class is part of the cetacean diet, including the *M. novaeangliae*.

Oceanographic variables supporting the appearance of the *M. novaeangliae* in the Alor Strait. Oceanographic variables such as sea depth temperature, chlorophyll-a, current velocity and the waters depth are important indicators in supporting the presence of cetaceans in a waterbody (Ballance et al 2006; Tynan et al 2005; Cañadas et al 2002; Hamazaki 2002; Praca et al 2009).

Alor Strait is still influenced by the water mass of the Indian Ocean and the Pacific Ocean, which makes its waters quite sensitive to temperature changes, especially at the points where the *M. novaeangliae* appears in Alor Waters. Research results obtained through ARGO (Array for Real-Time Geostrophic Oceanography) data analysis showed that the monthly average of the sea water temperature from 2006 to 2020, at a depth of 0 to 200 m, ranged from 18.0 to 28.0°C, which can be seen in Figure 8. The monthly average of the sea surface temperature data processed through Aqua MODIS (Moderate Resolution Imaging Spectroradiometer) data ranged from 25.0 to 31.0°C, as in Figure 9.




Figure 8. Yearly average of seawater temperature at a depth of 0 to 200 m, at Point 3 of the *Megaptera novaeangliae* sighting in Alor Strait.




Figure 9. Monthly (1-12) average of sea surface temperature 2020.

From the ARGO and Aqua MODIS data, the research determined that the average depth and sea surface temperature in the Alor Strait waters, including its surrounding seas at the tropics, are still in the normal category for the emergence of cetaceans, especially the *M. novaeangliae*. This is supported by the research of Houser et al (2004), which explains that at the tropics the average temperature range at the time of the appearance of cetaceans is between 20 to 31°C. Forcada et al (1996) documented the preferred temperature for cetaceans as ranging between 22.3 and 26.3°C (average 24.2°C). Consistently, the same sea surface and depth temperature values for the presence of cetaceans in water bodies were determined by other studies (Laran & Gannier 2008; Gregor & Trites 2001; Hamazaki 2002; Doniol-Valcroze et al 2011), as ranging from 22.4 to 26.7°C (average 23.08°C). Putra et al (2016) found the sea surface temperature of the appearance of cetaceans in the Savu sea waters in the range of 30 to 31°C. The average distribution of chlorophyll-a and the average monthly sea surface flow velocity, based on the results obtained during 2020 from the Aqua MODIS data analysis in the Alor Strait seawater, ranged from 0.2 to 0.8 mm m^{-3} and from 0.2 to 0.6 m s^{-1} , respectively, as in Figure 10 and Figure 11.




Figure 10. Monthly average of seawater chlorophyll-a 2020 at Alor Strait.




Figure 11. Monthly (1 to 12) Alor strait's current speed average in 2020.

The research found that the average chlorophyll-a and the mean sea surface flow velocities for the emergence of cetaceans in Alor waters were ranging from 0.1 to 0.6 mm m^{-3} and from 0.2 to 0.6 m s^{-1} , respectively. To this extent, it can be concluded that the average range of chlorophyll-a and the average flow velocity in the Alor Strait are still categorized as normal for the emergence of cetaceans in the Alor Strait.

The appearance of the *M. novaeangliae* in the Alor Strait was located at approximately 7 to 12 km from the coast, with a depth of more than 50 m. According to certain studies (Putra et al 2016; Salim 2011), cetaceans, especially whales, were found in the Savu Sea waters at a depth of less than 100 m. The depth modeling of the waters where the *M. novaeangliae* was found in the Alor Strait used the GEBCO data, as in Figure 12.




Figure 12. Depth at three observation points at Alor Strait.

Conclusions. The results of the research showed that around 15 *M. novaeangliae* crossed the Alor waters in the afternoon of December 1st 2020, from 05.00 to 05.45 pm with a mating behavior. The coordinates of the appearance were in: 8.123086° latitude and 124.063298° longitude. The average physical factors of the waters included: the sea surface temperature ranging from 25.0 to 31.0°C, the chlorophyll content ranging from 0.2 to 0.8 mm m⁻³, the waters depth >50 m, the surface water currents 0.2 to 0.6 m s⁻¹. The dominant type of zooplankton found consisted of crustaceans from the Arthropoda phylum, with a density of 1,957 individuals L⁻¹. This findings will support the *M. novaeangliae* species protection in Alor waters- the Eastern of Indian Ocean, Indonesia.

Acknowledgments. The authors would like to thank the Local Government of East Nusa Tenggara Province and the Office of World Wildlife Fund for Nature (WWF) at Solor-Alor. Sincere thanks to the LPDP Indonesian Government for the scholarship attributed to Jahved Ferianto Maro, in the framework of the Doctoral Program of Coastal Management, Faculty of Fisheries and Marine Sciences and the Institute of Research and Community Services, and also to the Diponegoro University, Semarang, Indonesia for the funding support through the contract no. 233-39/UN7.6.1/PP/2020. Special thanks to the National Aeronautics and Space Administration (NASA) for making available the MODIS AquaTerra satellite data, <https://argo.ucsd.edu/>, Copernicus Marine Service, the SeaDas software application and MIKE 21, to the ESRI for the permit to use the ArcGis 10.8 software for processing the satellite data and to all other parties for their help and contributions.

Conflict of interest. The authors declare no conflict of interest.

References

- Ballance L. T., Pitman R. L., Fiedler P. C., 2006 Oceanographic influences on seabirds and cetaceans of the eastern tropical Pacific: A review. *Progress in Oceanography* 69:360-390.
- Bejder L., Vidensen S., Hermensen L., Simon M., Hanf D., Medsen P. T., 2019 Low energy expenditure and resting behaviour of humpback whale mother-calf pairs highlights conservation importance of sheltered breeding areas. *Scientific Report* 9 (711):1-11.

- Burkard M., Whitworth D., Schirmer K., Nash S. B., 2015 Establishment of the first humpback whale fibroblast cell lines and their application in chemical risk assessment. *Aquatic Toxicology* 167:240-247.
- Calambokidis J., Barlow J., 2004 Abundance of blue and humpback whales in the eastern north pacific estimated by capture-recapture and line-transect methods. *Marine Mammal Science* 20(1):63-85.
- Cañadas A., Sagarminaga R., García-Tiscar S., 2002 Cetacean distribution related with depth and slope in the Mediterranean waters off southern Spain. *Deep Sea Research Part I: Oceanographic Research Papers* 49(11):2053-2073.
- Chen J. H., 2012 The effect of humpback whale-like leading edge protuberances on hydrofoil performance. 15th International Symposium on Flow Visualization 1-76.
- Chittleborough R. G., 1965 Dynamics of two populations of the humpback whale, *Megaptera novaeangliae* (Borowsk). *Australian Journal of Marine and Freshwater Research* 16:33-128.
- Clapham P. J., 2009 Humpback whale: *Megaptera novaeangliae*. *Encyclopedia of marine mammals* (Second Edition). Academic Press 582-585.
- Clapham P. J., 1992 The attainment of sexual maturity in humpback whales. *Canadian Journal of Zoology* 70:1470-1472.
- Craig A. S., Herman L. M., Pack A. A., Waterman J. O., 2014 Habitat segregation by female humpback whales in Hawaiian waters: avoidance of males? *Behavior* 151(5):613-631.
- Dawbin W. H., 1959 Evidence on growth-rates obtained from two marked humpback whales. *Nature* 183:1749-1750.
- Dharmadi, Faizah R., Wiadnyana N. N., 2017 [Appearance frequency, behavior, and distribution of marine mammals in the Savu Sea, East Nusa Tenggara]. *BAWAL Widya Capture Fisheries Research* 3(3):209-216. [In Indonesian].
- Doniol-Valcroze T., Lesage V., Giard J., Michaud R., 2011 Optimal foraging theory predicts diving and feeding strategies of the largest marine predator. *Behavioral Ecology* 22(4):880-888
- Dréo R., Bouffaut L., Leroy E., Barruol G., Samaran F., 2019 Baleen whale distribution and seasonal occurrence revealed by an ocean bottom seismometer network in the western Indian Ocean. *Deep Sea Research Part II: Topical Studies in Oceanography* Volume 161:132-144
- Findlay K., Meyer M., Elwen S., Kotze D., Johnson R., Truter P., Uamusse C., Siteo S., Wilke C., Kerwath S., Swanson S., Staverees L., Westhuizen J. V. D., 2011 Distribution and abundance of humpback whales, *Megaptera novaeangliae*, off the coast of Mozambique. *Journal of Cetacean Research and Management* 3:163-174.
- Fish F. E., Battle J. M., 1995 Hydrodynamic design of the humpback whale flipper. *Journal of Morphology* 255(1):51-60.
- Forcada J., Aguilar A., Hammond P., Pastor X., Aguilar R., 1996 Distribution and abundance of fin whales (*Balaenoptera physalus*) in the western Mediterranean sea during the summer. *Journal of Zoology* 238(1):23-34.
- Galvin C., 2006 Surface-piercing activities of the humpback whale, *Megaptera*, related to parasites and mechanics. *American Geophysical Union Eos* 87(52).
- Gregg E. J., Trites A. W., 2001 Predictions of critical habitat for five whale species in the waters of coastal British Columbia. *Canadian Journal of Fisheries and Aquatic Sciences* 58:1265-1285.
- Hamazaki T., 2002 Spatiotemporal prediction models of cetacean habitats in the mid-western North Atlantic Ocean (from Cape Hatteras, North Carolina, U.S.A. to Nova Scotia, Canada). *Marine Mammal Science* 18(4):920-939.
- Hampe O., Franke H., Hipsley C. A., Kardjilov, Müller J., 2015 Prenatal cranial ossification of the humpback whale (*Megaptera novaeangliae*). *Journal of Morphology* 276(5):564-582.
- Hartoko A., 2010 Spatial distribution of *Thunnus.sp*, vertical and horizontal sub-surface multilayer temperature profiles of in-situ agro float data in Indian Ocean. *Journal of Coastal Development* 14(1):61-74.

- Hartoko A., Febrianto A., Pamungkas A., Fachruddin I., 2019 The myth and legend of Sadai and Gaspar strait Bangka Belitung (Banca-Billiton) and oceanographic conditions. *International Journal of GEOMATE* 17(62):212-218.
- Houser D. S., Finneran J., Carder D., Bonn W. V., Smith C., Hoh C., Mattrey R., Ridgway S., 2004 Structural and functional imaging of bottlenose dolphin (*Tursiops truncatus*) cranial anatomy. *Journal of Experimental Biology* 207(21):3657-3665.
- Kügler A., Lammers M., Zang E., Kaplan M., Mooney T., 2020 Fluctuations in Hawaii's humpback whale *Megaptera novaeangliae* population inferred from male song chorusing off Maui. *Endangered Species Research* 43:421-434.
- Laran S., Gannier A., 2008 Spatial and temporal prediction of fin whale distribution in the northwestern Mediterranean Sea. *ICES Journal of Marine Science* 65(7):1260-1269.
- McSweeney D. J., Chu K. C., Dolphin W. F., Guinee L. N., 1989 North Pacific humpback whale songs: a comparison of Southeast Alaskan feeding ground songs with Hawaiian wintering ground songs. *Marine Mammal Science* 5(2):139-184.
- Mohebbi-Kalkhoran H., Zhu C., Schinault M., Ratilal P., 2019 Classifying humpback whale calls to song and non-song vocalizations using bag of words descriptor on acoustic data. 18th IEEE International Conference on Machine Learning and Applications (ICMLA) 865-870.
- Mujiyanto M., Riswanto R., Nastiti A. S., 2017 [Effectiveness of cetacean protection sub zone in marine protected areas TNP Sawu Sea, East Nusa Tenggara]. *Coastal and Ocean Journal* 1(2):1-12. [In Indonesian].
- Nybakken J. W., 1992 [Marine biology: An ecological approach (text book)]. Gramedia Pustaka Utama, Jakarta, pp. 104-106. [In Indonesian].
- Packard T., Osma N., Fernández-Urruzola I., Codispoti L. A., Christensen J. P., Gómez M., 2015 Peruvian upwelling plankton respiration: calculations of carbon flux, nutrient retention efficiency, and heterotrophic energy production. *Biogeosciences* 12: 2641-2654.
- Palsbøll P. J., Allen J., Bérubé M., Clapham P. J., Feddersen T. P., Hammond P. S., Hudson R. R., Jørgensen H., Katona S., Larsen A. H., Larsen F., Lien J., Mattila D. K., Sigurjónsson J., Sears R., Smith T., Sponer R., Stevick P., Øien N., 1997 Genetic tagging of humpback whales. *Nature* 388:767-769
- Parsons E. C. M., Wright A. J., Gore M. A., 2008 The nature of humpback whale (*Megaptera novaeangliae*) song. *Journal of Marine Animals and Their Ecology* 1(1):21-30.
- Praca E., Gannier A., Das K., Laran S., 2009 Modelling the habitat suitability of cetaceans: Example of the sperm whale in the northwestern Mediterranean Sea. *Deep Sea Research Part I: Oceanographic Research Papers* 56(4):648-657.
- Putra M. I. H., Lewis S. A., Kurniasih E. M., Prabuning D., Faiqoh E., 2016 Plankton Biomass Models Based on GIS and Remote Sensing Technique for Predicting Marine Megafauna Hotspots in the Solor Waters. *IOP Conference Series: Earth and Environmental Science* 47:1-19.
- Rodrigues F. M., Silva F. M. O., Trompieri-Silveira A. C., Vergara-Parente J. E., Miglino M. A., Guimarães J. P., 2014 Morphology of the eyeball from the Humpback whale (*Megaptera novaeangliae*). *Microscopy Research and Technique* 77(5):348-55.
- Rosas C. L., Gil M. N., Uhart M. M., 2012 Trace metal concentrations in Southern Right whale (*Eubalaena australis*) at Península Valdés, Argentina. *Marine Pollution Bulletin* 64(6):1255-1260.
- Salim D., 2011 [Conservation of marine mammals (*Cetaceans*) in the Savu Sea, East Nusa Tenggara]. *Marine Journal* 4(1):24-41. [In Indonesian].
- Secchi E. R., Rosa L. D., Kinan P. G., Nicolette R. F., Rufino A. M. N., Azevedo A. F., 2011 Encounter rates and abundance of humpback whales (*Megaptera novaeangliae*) in Gerlache and Bransfield Straits, Antarctic Peninsula. *Journal of Cetacean Research and Management (Special Issue)* 3:107-111.
- Sediadi A., 2004 [The effect of upwelling on the abundance and distribution of phytoplankton in Banda Sea and surrounding waters]. *MAKARA of Science Series* 8(2):43-51. [In Indonesian].

- Tynan C. T., Ainley D. G., Barth J. A., Cowles T. J., Pierce S. D., Spear L. B., 2005 Cetacean distributions relative to ocean processes in the northern California current system. *Deep Sea Research Part II: Topical Studies in Oceanography* 52(1):145-167.
- Werth A. J., Kosma M. M., Chenoweth E. M., Straley J. M., 2019 New views of humpback whale flow dynamics and oral morphology during prey engulfment. *Marine Mammal Science* 35(4).
- Wirasatriya A., Setiawan J. D., Sugianto D. N., Rosyadi I. A., Haryadi H., Winarso G., Setiawan R. Y., Susanto R. D., 2020 Ekman dynamics variability along the southern coast of Java revealed by satellite data. *International Journal of Remote Sensing* 41(21):8475-8496.
- Woodward B. L., Winn J. P., Fish F. E., 2006 Morphological specializations of baleen whales associated with hydrodynamic performance and ecological niche. *Journal of Morphology* 267(11):1284-1294.
- Zerbini A., Andriolo A., da Mata J., Simões P., Siciliano S., Pizzorno J., Waite J., DeMaster D., VanBlaricom G., 2004 Winter distribution and abundance of humpback whales (*Megaptera novaeangliae*) of Northeastern Brazil. *Journal of Cetacean Research and Management* 6(1):101-107.
- *** International Whaling Commission, IWC, 1998 Report of the scientific committee. Report of International Whale Commission 48(53):118.
- *** International Whaling Commission, IWC.int/humpback-whale
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2015 [Republic of Indonesia. Ministerial Regulation No. 5/KEPMEN-KP/2015]. [In Indonesian].

Received: 11 April 2021. Accepted: 30 May 2021. Published online: 11 June 2021.

Authors:

Jahved Ferianto Maro, Diponegoro University, Faculty of Fisheries and Marine Science, Coastal Resources Management Doctoral Programe, Prof. Sudarto Street, 50275 Semarang, Central Java, Indonesia, e-mail: yantogisuntrib@gmail.com

Agus Hartoko, Diponegoro University, Faculty of Fisheries and Marine Science, Coastal Resources Management, Prof. Sudarto Street, 50275 Semarang, Central Java, Indonesia, e-mail: agushartoko.undip@gmail.com

Sutrisno Anggoro, Diponegoro University, Faculty of Fisheries and Marine Science, Coastal Resources Management, Prof. Sudarto Street, 50275 Semarang, Central Java, Indonesia, e-mail: sutrisno.anggoro@yahoo.co.id

Max Rudolf Muskananfolo, Diponegoro University, Faculty of Fisheries and Marine Science, Coastal Resources Management, Prof. Sudarto Street, 50275 Semarang, Central Java, Indonesia, e-mail: maxmuskananfolo@yahoo.com

Lumban Nauli Lumban Toruan, Nusa Cendana University, Faculty of Fisheries and Marine Science, Coastal Resources Management, Jl. Adi Sucipto, Penfui, Kelapa Lima, 85001 Kupang City, East Nusa Tenggara, Indonesia, e-mail: naulitoruan@gmail.com

Micael Dakahamapu, Environmental Practitioners, Kalabahi City street, Air Kenari Kalabahi, 85819 District Alor, East Nusa Tenggara, Indonesia, e-mail: mikeinspire@yahoo.com

Erick Nugraha, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street, 12520 Pasarminggu, South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Maro J. F., Hartoko A., Anggoro S., Muskananfolo M. R., Toruan L. N. L., Dakahamapu M., Nugraha E., 2021 The appearance of the humpback whale (*Megaptera novaeangliae*) during the rainy season in the Alor Waters, Indonesia. *AACL Bioflux* 14(3):1596-1608.

The appearance of the humpback whale (*Megaptera novaeangliae*) during the rainy season in the Alor Waters, Indonesia.

ORIGINALITY REPORT

18%

SIMILARITY INDEX

16%

INTERNET SOURCES

9%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1

bioflux.com.ro

Internet Source

5%

2

Submitted to Universitas Jenderal Soedirman

Student Paper

2%

3

www.science.gov

Internet Source

1%

4

Submitted to Universitas Diponegoro

Student Paper

1%

5

scholarworks.alaska.edu

Internet Source

1%

6

journal.iwc.int

Internet Source

1%

7

www.readkong.com

Internet Source

<1%

8

Submitted to West Los Angeles College

Student Paper

<1%

9

B M Rehatta, M M Kamal, M Boer, A Fahrudin, Zairion, J S R Ninef. "Growth, mortality, recruitment pattern, and exploitation rate of shared stock flying fish (Exocoetidae) at border area of Indonesia and Timor Leste in Ombai Strait", IOP Conference Series: Earth and Environmental Science, 2021

Publication

<1 %

10

Mochamad Iqbal Herwata Putra, Putu Liza Kusuma Mustika. "Maximum entropy model: Estimating the relative suitability of cetacean habitat in the northern Savu Sea, Indonesia", Marine Mammal Science, 2020

Publication

<1 %

11

iwccentre.org

Internet Source

<1 %

12

Submitted to Brooksby Melton College

Student Paper

<1 %

13

Dian Novianto, Eko Susilo. "ROLE OF SUB SURFACE TEMPERATURE, SALINITY AND CHLOROPHYLL TO ALBACORE TUNA ABUNDANCE IN INDIAN OCEAN", Indonesian Fisheries Research Journal, 2016

Publication

<1 %

14

core.ac.uk

Internet Source

<1 %

pureadmin.uhi.ac.uk

15

Internet Source

<1 %

16

cronfa.swan.ac.uk

Internet Source

<1 %

17

jonuns.com

Internet Source

<1 %

18

Bambang Sukresno, Agus Hartoko, Budi Sulisty, Subiyanto. "Empirical Cumulative Distribution Function (ECDF) Analysis of Thunnus.sp Using ARGO Float Sub-surface Multilayer Temperature Data in Indian Ocean South of Java", Procedia Environmental Sciences, 2015

Publication

<1 %

19

O Sagnol, C Richter, LH Field, F Reitsma. "Spatio-temporal distribution of sperm whales () off Kaikoura, New Zealand, in relation to bathymetric features ", New Zealand Journal of Zoology, 2014

Publication

<1 %

20

Yayuk Astuti, Dhina Anggraeni, Adi Darmawan. "Photocatalytic Performance of Bismuth Oxide Prepared by Citric Acid-Fueled Solution Combustion on Decolorisation of Organic Dye Molecules", IOP Conference Series: Materials Science and Engineering, 2020

Publication

<1 %

21	ejurnal.ung.ac.id Internet Source	<1 %
22	research-repository.st-andrews.ac.uk Internet Source	<1 %
23	www.globalscientificjournal.com Internet Source	<1 %
24	www.int-res.com Internet Source	<1 %
25	ebin.pub Internet Source	<1 %
26	epubs.scu.edu.au Internet Source	<1 %
27	hdl.handle.net Internet Source	<1 %
28	navfac.navy.mil Internet Source	<1 %
29	repository.up.ac.za Internet Source	<1 %
30	A K Dewi, J L Gaol, V P Siregar, A S Atmadipoera. "Bigeye tuna fishing ground analysis using oceanographic features in Eastern Indian Ocean off Southern Java", IOP Conference Series: Earth and Environmental Science, 2020 Publication	<1 %

31

Sudaryono. "Evaluation of potential of lupin meal as an alternative to fish meal in juvenile *Penaeus monodon* diets", *Aquaculture Nutrition*, 12/1999

Publication

<1 %

32

uludag.edu.tr

Internet Source

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

The appearance of the humpback whale (Megaptera novaeangliae) during the rainy season in the Alor Waters, Indonesia.

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11

PAGE 12

PAGE 13