

Alkaloids piperine in dichloromethane fraction of red galangal rizhome (*Alpinia purpurata*)

by Bambang Cahyono

Submission date: 13-Nov-2019 09:19AM (UTC+0700)

Submission ID: 1212670274

File name: Suzery_2019_IOP_Conf._Ser._Mater._Sci._Eng._509_012076.pdf (917.55K)

Word count: 1899

Character count: 10276

PAPER • OPEN ACCESS

Alkaloids piperine in dichloromethane fraction of red galangal rizhome (*Alpinia purpurata*)

To cite this article: Meiny Suzery *et al* 2019 *IOP Conf. Ser.: Mater. Sci. Eng.* **509** 012076

View the [article online](#) for updates and enhancements.



IOP | ebooks™

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

Alkaloids piperine in dichloromethane fraction of red galangal rizhome (*Alpinia purpurata*)

Meiny Suzery^{1*}, Resti Yuyun Septembe Ria¹, Bambang Cahyono¹

¹ Department of Chemistry, Faculty of Science and Mathematics, Diponegoro University, Semarang Indonesia 50267

* Corresponding author: meiny.suzery@live.undip.ac.id

Abstract. Red galangal rizhome (*Alpinia purpurata*) was extracted with dichloromethane using maceration methods. The crude extract was fractionated using vacuum column chromatography with a mixture of ethyl acetate:n-hexane (2:3) to give two fractions (F1 and F2). Fraction 2 was fed into the silica gel of column chromatography tailed by recrystallization to produce crystalline (17 mg). The purity was tested by using melting point apparatus and thin-layer chromatography method for any solvent. Their structures were established by spectroscopic methods (UV-Vis, FTIR, ¹H-NMR and ¹³C-NMR). Result of analysis gave melting point of crystals around 130°C. The UV-Vis spectrometer obtained maximum wavelength 344 nm. FTIR analysis showed C=O carbonyl, C-O ether, C-N, and C=C aromatic. Analysis by ¹H-NMR spectrometer showed that bioactive compounds have a 19 protons. Analysis by ¹³C-NMR spectrometer showed that bioactive compounds have a 17 carbon atoms. The results of identification showed that bioactive compounds in dichloromethane fraction of red galangal rhizome was a compound piperine alkaloids types with a chemical formula of C₁₇H₁₉O₃N.

Keywords: *Alpinia purpurata*, red galangal rizhome, Piperine, alkaloids

1. Introduction

Red galangal rhizome (*Alpinia purpurata* K. Schum) is one of herbs that broadly been used as raw materials for bio-pharmacy products or material sources of the drugs [1]. *Alpinia purpurata* K. Schum (family Zingiberaceae) is typical plant for tropical and subtropic regions of Asia.[2].

Phytochemical analysis for Rhizome *Alpinia purpurata* showed positive results for alkaloids (14.9%), phenols (9.5%), flavonoids (0.85%) and tannins (13.8%) [3]. Investigation of the flavonoids compound contained by kaempferol-3-owned glucuronide, kaempferol-3-ulicronide and rutin of ethanol extract has been done using HPLC [4]. Ethanolic extracts of *Alpinia purpurata* showed highest inhibitory activity against *Enterobacter aerogens* [5]. Other chemical contents contained within hexane and dichloromethane are fatty alcohol, kumatakerin, sitosteryl-3-O-6-palmitoyl-β-D-glucoside and β-galactoside sitosteryl [6]. Based on literature above, despite the limitations of phytochemical studies of red galangal rhizome (*Alpinia purpurata*) species that grows in Indonesia, in this research we started further exploration about isolation and bioactive compounds identification from red galangal dichloromethane fraction.

2. Materials and Methods

The equipment used in this research were vials bottle, capillary pipettes, thin layer chromatography, column chromatography, vacuum rotary evaporator (IKA basic RF 10), blender, mixer, spatula, analytical balance (Ohaus), Fisher Jones melting point apparatus, UV-Vis Spectroscopy (Spectroquant

® Pharo 300), FTIR (Government employees), and NMR (Agilent 500 MHz with console system DD2 operated on the frequency of 500 MHz (^1H) and 125 MHz (^{13}C). The materials used in this research were methanol, acetone, ethyl acetate, dichloromethane, n-hexane, chloroform, ether, silica gel G60 F₂₅₄ from *e. Merck*. Red galangal's rhizomes (*Alpinia purpurata*) collected from Semarang, Central Java, Indonesia. Rhizomes specimen was stored in Wet Laboratory of Tropical Disease Diponegoro University.

2.1. Isolation and Constituents Identification

Dried rhizomes of *Alpinia purpurata* (2000 g) were extracted with maceration methods using dichloromethane solvent followed by reduced pressure evaporation to give 75,91 g crude product. The crude product was fractionated with column vacuum chromatography using a dichloromethane to give two fractions (F1 and F2). We used F2 for further treatment because the amount of crude fraction obtained was higher than crude fraction of F1. We partitioned F2 using gravity column chromatography with mixture of ethyl acetate: n-hexane (2:3) as eluent, followed by recrystallization to produce crystals approximately 17 mg.

2.2. Characterization of Bioactive Compounds

Crystals obtained from isolation of F2 (17 mg) characterized using UV-Vis spectrometer, FTIR, ^1H -NMR, and ^{13}C -NMR.

3. Results and Discussion

3.1. Isolation and Identification

The extraction process was done using maceration method using methanol solvent, producing thick reddish brown extract for about 75.91 grams, obtained from 2 kg dry powder of red galangal rhizome with a yield of 4.2172% (w/w). Condensed extracts were then fractionated using vacuum column chromatography with dichloromethane as eluent, resulting two fractions: (F1=0.12 g and F2=1.83 g). F2 fraction was then separated using gravity column chromatography (GCC) method using ethyl acetate: n-hexane (2:3) as eluent to acquired 0.2136 grams of solid. Recrystallization was carried out with ether to produce 17 mg of yellow blanch crystals. Identification of crystal purity was done by measurement of melting point and acquired melting point of 130°C.

3.2. Characterization of Bioactive Compounds

Characterization result of pure compound using UV Spectrometer shown in Fig. 1.




Figure 1. UV-Vis spectrum of pure compound.

UV-Vis analysis resulting maximum wavelength ($\lambda_{\text{methanol}}=344$ nm). The analysis result using Spectrophotometer UV-Vis reinforced past studies from Adosraku *et al.* [7] and Deepthi *et al.* [8] which showed that piperine compounds have maximum wavelength at 343 nm using methanol solvent. From this result, we assume that the pure compound is piperine but further analysis need to be done.

Further analysis using IR spectroscopy shown in Fig. 2.




Figure 2. Infrared spectrum of pure compound.

The absorption patterns of FTIR of pure compound showed in table 1.

Table 1. Absorption patterns of pure compound using FTIR spectrometry.

Wavenumber (cm ⁻¹)	Vibration types
3435.29	O-H stretch
2937.27	Asymmetrical aliphatic C-H stretch
2863.05	Symmetrical aliphatic C-H stretch
1631.66	Amide C=O stretch
1584.59	Aromatic C=C stretch
1443.91	Aliphatic C-H bending
1365.28	C-N bending
1027.30	Ether C-O stretch
929.10; 845.26; 704.65	para, meta, orto substitution on aromatic ring

FTIR result showed that pure compound contains functional groups: C-N, amide C=O, aromatic C=C, ethers C-O, and aliphatic C-H. These data reinforced structure of piperine compound. Our assumption was confirmed from previous study from Deepthi *et al.* [8] which identify the piperine alkaloid from *Piper longium* that shows the functional group absorption of C=O at 1635 cm⁻¹ wavenumbers and aromatic C=C at 1580 cm⁻¹. As well as research from Sirat and Liamen [9] which has been identified piperine from *Alpinia purpurata* which shows absorption of amide C=O at 1640 cm⁻¹ as well as C=C absorption at 1595 cm⁻¹ and 1500 cm⁻¹. Based on data obtained, it was believed that pure crystalline contains alkaloid compound known as piperine whose structure is presented in Fig. 3.




Figure 3. Piperine structure.

NMR characterization of pure compound (piperine) was done in order to confirmed the data obtained beforehand. NMR characterization of pure compound shown in Fig. 4.




Figure 4. ^{13}C -NMR spectrum.

The spectrum of ^{13}C -NMR from compound isolation informed existence of 4 carbon atoms as C-quartener appearing at δ 165.551 ppm (C-14); δ 148.241 ppm (C-5); δ 148.319 ppm (C-4), and δ 131.150 ppm (C-7). There are seven =CH- carbon atoms appearing at δ 142.620 ppm (C-12), δ 138.350 ppm (C-10), δ 125.488 ppm (C-8), δ 122.648 ppm (C-11), δ 120.190 ppm (C-13), δ 108.627 ppm (C-9), and δ 105.794 ppm (C-6). There are also six -CH₂- carbon atoms at δ 101.410 ppm (C-2), δ 47.051 ppm (C-17), δ 43.384 ppm (C-21), δ 26.880 ppm (C-18), δ 25.774 ppm (C-20), and δ 24.816 ppm (C-19).

Characterization by H NMR produce spectra shown in Fig. 5.




Figure 5. $^1\text{H-NMR}$ spectrum.

Fig. 5 shows the $^1\text{H-NMR}$ spectrum, which give several peaks, at δ 1,5625-1,6677 ppm (6H, m, overlapping peaks from H-19, H-20 and H-18); δ 3,5222-3,6330 ppm (4H, d, H-21 and H-17); δ 5.9724 ppm (2H, s, H-2); δ 6,4198-6,4491 ppm (1H, d, H-13); δ 6.6968 -6,7829 ppm (3H, d, overlapping peaks from H-11, H-10 and H-9); δ 6.8773-6.8964 ppm (1H, dd, H-8); δ 6,9762- 6,9792 ppm (1H, d, H-6); δ 7,3733- 7,4228 ppm (1H, ddd, H-12). The chemical shifts of $^1\text{H-NMR}$ and $^{13}\text{C-NMR}$ are in accordance with the position of the proton and carbon atoms from piperine.

4. Conclusion

Bioactive compound isolated from dichloromethane fraction of red galangal is a type of alkaloids known as piperine which has 130°C melting point, 344 nm maximum wavelength and chemical formula $\text{C}_{17}\text{H}_{19}\text{O}_3\text{N}$.

References

- [1] Wardana HD, Barwa N, Kongsjahju A, Iqbal M, Khalid M and Taryadi R 2002 *Budi Daya secara Organik Tanaman Obat Rimpang* (Jakarta: Penebar Swadaya)
- [2] Kress W J, Prince L M and Williams K J 2002 The phylogeny and a new classification of the gingers (Zingiberaceae): evidence from molecular data *Am. J. Bot.* **89** 10 1682-96
- [3] Subramanian V and Suja S 2011 Phytochemical screening of *Alpinia purpurata* (Vieill.) *Res. J. Pharm. Biol. Chem. Sci.* **2** 3 866
- [4] Victório C, Kuster R and Lage C 2009 Detection of flavonoids in *Alpinia purpurata* (Vieill.) K. Schum. leaves using high-performance liquid chromatography *Revista Brasileira de Plantas Mediciniais* **11** 2 147-53
- [5] Kochuthressia K, Britto S J, Jaseentha M, Raj L J M and Senthilkumar S 2010 Antimicrobial efficacy of extracts from *Alpinia purpurata* (Vieill.) K. Schum. against human pathogenic bacteria and fungi *Agriculture and Biology Journal of North America* **1** 6 1249-52
- [6] Villaflores O B, Macabeo A P G, Gehle D, Krohn K, Franzblau S G and Aguinaldo A M 2010 Phytoconstituents from *Alpinia purpurata* and their in vitro inhibitory activity against *Mycobacterium tuberculosis* *Pharmacogn. Mag.* **6** 24 339
- [7] Adosraku R K, Kyekyeku J O and Attah I Y 2013 Characterization and HPLC quantification of

- piperine isolated from Piper guineense (fam. Piperaceae) *Int. J. Pharm. Pharm. Sci.* **5** 1 252-6
- [8] Deepthi R, Junise V, Shubin S and Rajesh S 2012 Isolation, identification and antimycobacterial evaluation of piperine from Piper longum *Pharm. Lett.* **4** 3 863-8
- [9] Sirat H M and Liamen M R 1995 Chemical constituents of Alpinia purpurata *Pertanika J. Sci. Technol.* **3** 1 67-71

Alkaloids piperine in dichloromethane fraction of red galangal rizhome (*Alpinia purpurata*)

ORIGINALITY REPORT

2%

SIMILARITY INDEX

1%

INTERNET SOURCES

2%

PUBLICATIONS

0%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Brent J. Yoder, Shugeng Cao, Andrew Norris, James S. Miller et al. " Antiproliferative Prenylated Stilbenes and Flavonoids from from the Madagascar Rainforest ", Journal of Natural Products, 2007
Publication <1%
- 2** J.Thomas Mckinnon, Eleanor Meyer, Jack B Howard. "Infrared analysis of flame-generated PAH samples", Combustion and Flame, 1996
Publication <1%
- 3** link.springer.com
Internet Source <1%
- 4** stuartxchange.org
Internet Source <1%
- 5** Thuc-Huy Duong, Xuan-Phong Ha, Warinthorn Chavasiri, Mehdi A. Beniddir et al. " Sanctis A-C: Three Racemic Procyanidin Analogues from The Lichen ", European Journal of Organic Chemistry, 2018 <1%

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On