

5. Expression of Stanniocalcin-1 as a Predictor of Recurrence in Breast Cancer Patients

by Muflihatul Muniroh

Submission date: 07-May-2023 06:34PM (UTC+0700)

Submission ID: 2086411163

File name: cin-1_as_a_Predictor_of_Recurrence_in_Breast_Cancer_Patients.pdf (1,009.42K)

Word count: 3416

Character count: 16681



Expression of Stanniocalcin-1 as a Predictor of Recurrence in Breast Cancer Patients

Yan Wisnu Prajoko^{1*}, Rizka Hastari², Muflihatal Muniroh³, Dik Puspasari⁴

¹Department of Surgical Oncology, Faculty of Medicine, Diponegoro University, Kariadi Hospital, Semarang, Indonesia; ²Department of Biomedical Science, Faculty of Medicine, Diponegoro University, Semarang, Indonesia; ³Department of Physiology, Faculty of Medicine, Diponegoro University, Semarang, Indonesia; ⁴Department of Anatomical Pathology, Faculty of Medicine, Diponegoro University, Kariadi Hospital, Semarang, Indonesia

Abstract

Edited by: Ksenija Bojković-Kostovska
Citation: Prajoko YW, Hastari R, Muniroh M, Puspasari D. Expression of Stanniocalcin-1 as a Predictor of Recurrence in Breast Cancer Patients. Open Access Maced J Med Sci. 2022 Feb 23; 10(B): 346-350. <https://doi.org/10.3889/oamjms.2022.8124>
Key words: Stanniocalcin-1; Recurrence; Breast cancer
Correspondence: Yan Wisnu Prajoko, Department of Surgical Oncology, Faculty of Medicine, Diponegoro University, Kariadi Hospital, Semarang, Indonesia. E-mail: yanprajoko@gmail.com
Received: 30-Nov-2021
Revised: 11-Feb-2022
Accepted: 14-Feb-2022
Copyright: © 2022 Yan Wisnu Prajoko, Rizka Hastari, Muflihatal Muniroh, Dik Puspasari
Funding: This research did not receive any financial support
Competing Interests: The authors have declared that no competing interests exist
Open Access: This is an open-access article distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 International License (CC BY-NC 4.0)

BACKGROUND: About 40% of breast cancer patients will experience a recurrence and the life expectancy will decrease from 95% to 40%. Stanniocalcin-1 (STC1) is a biomarker that is thought to have potential as a risk factor for recurrence.

AIM: The objective of the study is to determine the role of STC1 expression as a risk factor for recurrence in breast cancer patients.

METHODS: Cross-sectional observational study on breast cancer patients, assessed the clinicopathological character and seen whether the patient's condition had a recurrence. Immunohistochemical staining with STC1 antibody was performed on the patient's paraffin block. Reading of immunohistochemical preparations using the Allred score method by looking at five fields of view. Data analysis regarding the relationship between STC1 and recurrence was carried out using the Chi-square test or Fisher's Exact test, then Logistic regression analysis was performed to determine the significant relationship between risk factors for recurrence.

RESULTS: Expression of STC1 was significantly different in patients who had recurrence and who did not experience recurrence ($P = 0.000$). There was a significant relationship between STC1 expression with age, tumor size, lymph node status, type of cancer, estrogen receptor status, and stage at diagnosis ($P < 0.005$).

CONCLUSION: It can be concluded that STC1 has a strong relationship with recurrence, so it has the potential to be used as a predictor of recurrence in breast cancer.

Introduction

Breast cancer is one of the most common types of cancer in women in the world. In 2018 there were around 2.89 million women diagnosed with breast cancer with a mortality rate of 627,000 people [1]. In Indonesia, in 2018 there were 59,256 women diagnosed with breast cancer, with the mortality rate reaches 12.75% [1], [2]. Breast cancer treatment is a multidisciplinary treatment that includes systemic therapy, local therapy, and supportive therapy. However, all of these efforts still have a risk of recurrence in the future, especially in patients with advanced stages [3], [4]. Approximately 40% of patients who have undergone a series of breast cancer therapy will experience a recurrence.

Recurrence of breast cancer will reduce the patient's life expectancy by up to 40% whereas patients who do not experience recurrence have a life expectancy of 95% [5], [6]. The risk of recurrence in breast cancer should be known early because in addition to affecting life expectancy, recurrence will also affect the patient's quality of life and increase

hospital costs. Several factors that influence the recurrence of breast cancer include young age, tumor size, biomarker status of Human Epidermal Growth Factor Receptor 2 (HER-2), Estrogen Receptor (ER), negative Progesterone Receptor (PR), presence of lymph node involvement, high grade in anatomical pathology [6], [7], [8], [9].

One of the biomarkers that are thought to be a predictor of recurrence in breast cancer is Stanniocalcin-1 (STC1) [10], [11]. STC1 which is a protein excreted by several organs including the thyroid, ovaries, liver, prostate, and kidneys. STC1 is strongly expressed in breast cancer tissue and is not expressed in normal breasts [12]. STC1 has been investigated to play a role in cancer signaling in the PI3K/Akt pathway which will affect the invasiveness of cancer cells [13]. Several studies have also shown that therapy with STC1 antibody can inhibit the aggressiveness of cancer cells in cases of triple-negative breast cancer (TNBC) *in vitro*.

Based on this, it is suspected that STC1 has the potential as a predictor of cancer. However, until now there has been no further research on this matter.

Therefore, research is needed on the expression of STC1 as a risk factor for recurrence in breast cancer patients and to determine the relationship between STC1 expression on the incidence of recurrence in breast cancer patients.

Methods

This research is an observational study with a cross-sectional study model. The research sample was paraffin block from breast cancer patients who underwent surgery at Kariadi Hospital from January 2016 to December 2016. Anatomical pathology and immunohistochemistry examinations were carried out at the Anatomical Pathology Department of Faculty of Medicine Diponegoro University/Kariadi Hospital Semarang. Demographic data, staging, and biomarker status (Estrogen Receptor, Progesterone Receptor, and Human Epidermoid Receptor-2) were taken from the patient's medical records. Further confirmation of the patient's current condition through medical records and telephone.

The population of this study was invasive breast cancer patients in women with Stage I, II, and III who came and were treated at Kariadi Hospital Semarang Indonesia.

After the data is collected, then tabulation and coding are carried out. Statistical analysis was performed using SPSS version 25 software (SPSS, IBM). Analysis of the data regarding the relationship of STC1 with recurrence using the Chi-square test or Fisher's Exact test, then Logistic regression analysis was carried out to determine the relationship between significant risk factors for recurrence.

Results

This study involved fifty-two breast cancer patients who met the requirements for completeness of medical records and paraffin blocks were still available in the Anatomical Pathology Department of the Kariadi Hospital (Figure 1), fifty-two breast cancer patients were divided into two groups, namely the group with recurrence of thirty-two patients and the group without recurrence as many as twenty patients. Most of the patients who experienced recurrence were patients aged more than 46-years-old, the most character in tumor size was T3, lymph node status was N2 and N3, mostly stage III, most subtype was triple-negative, the most histopathological grade is intermediate and most STC1 scoring was >6.7 (Table 1). There was a significant relationship between ER, PR, and STC1




Figure 1: Flow of getting research sample

expression on recurrence and there was no significant relationship between HER-2 expression and recurrence in breast cancer patients (Table 2). There was a significant relationship between STC1 expression and cancer subtype (TNBC) and the patient's lymph node status (N3) (Table 3).

Table 1 shows that several risk factors for recurrence in breast cancer include age, tumor size, lymph node involvement, stage, tumor type (Luminal, HER-2, and TNBC), and STC1 expression. The results of 6 statistical tests with Chi-Square found several risk factors that were significantly associated with recurrence, i.e., age more than 45-years-old, tumor size > T2, with lymph node involvement, TNBC subtype, and strong STC1 expression.

Several significant risk factors in the Chi-Square test will then be tested with logistic regression, previously several risk factors were categorized into 2 categories such as stage, tumor size, nodule, and age. From the results of the logistic regression test, it was found that Crude OR was greater than Adjusted OR for STC1 expression > 6.7, stage > 2, age > 45 and TNBC cancer type, while tumor size < T2, KGB N0 status, and ER status (+) has a smaller Crude OR than the Adjusted OR. This shows that recurrence is influenced by STC1 expression > 6.7, stage > 2, age > 45 and type of TNBC cancer, while tumor size < 2, N0 lymph node status and ER (+) status are protective factors against recurrence (Table 4).

In IHC examination, STC1 is expressed on the cell membrane and cytoplasm which will appear brownish in color, as in Figure 2a according to Allred

Table 1: Patient characteristics and recurrence

Characteristics	Recurrence			No recurrences	p	Current condition		p
	Locoregional	Distant	Mixed			Dead	Alive	
Age								
<45	1 (5.9)	3 (17.6)	1 (5.9)	12 (70.6)	0.003*	5 (29.4)	12 (70.6)	0.014*
46-59	3 (10)	18 (60)	3 (10)	6 (20)		22 (73.3)	8 (26.7)	
≥60	1 (20)	2 (40)	0 (0)	2 (40)		3 (60)	2 (40)	
Tumor Size								
T1	0 (0)	0 (0)	0 (0)	1 (100)	0.004*	0 (0)	1 (100%)	0.000*
T2	2 (12.5)	2 (12.5)	0 (0)	12 (75)		3 (18.8)	13 (81.3)	
T3	1 (5)	11 (55)	1 (5)	7 (35)		12 (60)	8 (40)	
T4	2 (13.3)	10 (10)	3 (20)	0 (100)		15 (100)	0 (0)	
Node Status								
N0	1 (3.6)	7 (25)	2 (7.1)	18 (64.3)	0.001*	10 (35.7)	18 (64.3)	0.006*
N1	2 (25)	9 (69.2)	0 (0)	2 (15.4)		10 (76.9)	3 (23.1)	
N2	2 (25)	5 (62.5)	1 (12.5)	0 (0)		7 (87.5)	1 (12.5)	
N3	0 (0)	2 (66.7)	1 (33.3)	0 (0)		3 (100)	0 (0)	
Stage								
I	0 (0)	0 (0)	0 (0)	1 (100)	0.001*	0 (0)	1 (100)	0.000*
II	1 (4.5)	4 (18.2)	1 (4.5)	16 (72.7)		6 (27.3)	16 (72.7)	
III	4 (13.8)	19 (65.5)	3 (10.3)	3 (10.3)		24 (82.8)	5 (17.2)	
Subtype								
Luminal	10 (28.6)	4 (11.4)	2 (5.7)	19 (54.3)	0.015*	14 (40)	21 (60)	0.001*
HER-2	0 (0)	4 (80)	0 (0)	1 (20)		4 (80)	1 (20)	
TNBC	1 (8.3)	9 (75)	2 (16.7)	0 (0)		12 (100)	0 (0)	
Histopathological grade								
Grade II	4 (9.5)	18 (42.9)	3 (7.1)	17 (40.5)	0.938*	23 (54.8)	19 (45.2)	0.305**
Grade III	1 (10)	5 (50)	1 (10)	3 (30)		7 (70)	3 (30)	
STC-1								
Allred score>6.7	3 (14.8)	19 (70.4)	1 (3.7)	4 (14.8)	0.001**	23 (85.2)	4 (14.8)	0.001**
Allred score≤6.7	1 (4)	4 (16)	3 (12)	18 (72)		7 (28)	18 (22)	

HER-2: Human epidermal growth factor receptor 2, TNBC: Triple-negative breast cancer, STC-1: Stanniocalcin-1, *P value <0.05

Table 2: Relationship between STC1, ER, PR, and HER-2 expression with recurrence

Immunology profile	Recurrence n (%)	No recurrence n (%)	p
ER			
ER (+)	16 (44.4)	20 (55.6)	0.001**
ER (-)	16 (100)	0 (0)	
PR			
PR (+)	16 (44.4)	20 (55.6)	0.001**
PR (-)	16 (100)	0 (0)	
HER 2			
HER 2 (+)	5 (71.4)	2 (28.6)	0.402**
HER 2 (-)	26 (57.8)	19 (42.2)	
STC-1			
Allred score>6.7	23 (85.2)	4 (14.8)	0.001**
Allred score≤6.7	7 (28)	18 (72)	

*Statistical test using Spearman's correlation. Significant value $P < 0.05$. **Statistical test using Fisher's Exact Test. Significant value if $P < 0.05$. STC-1: Stanniocalcin-1, ER: Estrogen receptor, PR: Progesterone receptor, HER-2: Human epidermal growth factor receptor 2.

Score > 6.7 while in Figure 2b and c with Allred Score ≤ 6.7.

Table 3: Relationship of STC-1 expression and patient characteristics

Characteristics	STC-1		p
	Allred score>6.7 n (%)	Allred score≤6.7 n (%)	
Age			
<45	5 (29.4)	12 (70.6)	0.055*
46-59	18 (60)	12 (40)	
≥60	4 (80)	1 (20)	
Tumor Size			
T1	0 (0)	1 (100)	0.284*
T2	6 (37.5)	10 (62.5)	
T3	11 (55)	9 (45)	
T4	10 (66.7)	5 (33.3)	
Node Status			
N0	10 (35.7)	18 (64.3)	0.030*
N1	9 (69.2)	4 (30.8)	
N2	7 (87.5)	1 (12.5)	
N3	1 (33.3)	2 (66.7)	
Stage			
I	0 (0)	1 (100)	0.069*
II	8 (36.4)	14 (63.6)	
III	19 (65.5)	10 (34.5)	
Subtypes			
Luminal	14 (40)	21 (60)	0.047*
HER-2	4 (80)	1 (20)	
TNBC	9 (75)	3 (25)	
Histopathologic grades			
Grade II	21 (50)	21 (50)	0.729
Grade III	6 (60)	11 (40)	

*Statistics test using Chi-square. The value is significant $P < 0.05$. **Statistical test using Fisher's Exact Test. Significant value if $P < 0.05$. STC-1: Stanniocalcin-1, HER-2: Human epidermal growth factor receptor 2, TNBC: Triple-negative breast cancer.

Discussion

STC1 is a protein that is an oncogene if the amount is excessive in the human body. Under normal conditions, STC1 will be expressed in several tissues such as the thyroid, ovary, liver, prostate, and kidney [14]. Expression of STC1 is associated with recurrence in breast cancer because of the nature of STC1 that can affect cancer cell growth, angiogenesis, and metastasis [11], [13], [15].

Table 4: Relationship of recurrence with breast cancer characteristics and expression of STC-1

Risk factors	Recurrence		Crude OR	Adjusted OR	95% CI
	Yes	No			
STC1 score>6.7	23	4	14.79	31.03	2.115–455.511
Tumor size<T2	25	9	7.22	0.49	0.19–12.559
Stage	24	5	13.6	164.7	1.426–19037.7
ER Status (+)	14	2	8.75	0.93	0.003–3.017
Nodal Status N0	20	4	9.00	0.42	0.030–6.072
Age>45	25	11	0.2	0.40	0.032–4.797
TNBC	11	1	15	294.1	1.840–47003.5

STC-1: Stanniocalcin-1, ER: Estrogen receptor, TNBC: Triple-negative breast cancer.

In this study, the majority of breast cancer patients were luminal types (67.3%) this is in accordance with the data that most breast cancers in Indonesia are luminal types, which is around 41.3% both luminal A and luminal B.20 In this study as many as 30.8% of patients with luminal type A had a recurrence in the first 2 years after the patient finished treatment, this exceeded the estimated recurrence in patients with luminal type, which was about 20%. In this study, it was found that the type of TNBC cancer was the type that experienced the most recurrences compared to the others. In this study it was found that the type of TNBC cancer will express STC1 strongly, this is in accordance with research conducted by Brantley which states STC1 as a predictor of recurrence in TNBC subtype, as previously known that TNBC subtype is a cancer with worst prognosis amongst all types of breast cancer [10], [16], [17], [18].




Figure 2: Microscopic expression of Stanniocalcin-1

In a previous study, it was found that patients with STC1 (+) will mostly experience locoregional recurrence at 3 years after completion of treatment, but in this study, most of them experienced distant recurrence, this is possible because the patients in the previous study were mostly diagnosed in stage I and II while the patients in this study were mostly diagnosed at stage II and III.

Conclusion

Recurrence in breast cancer can be detected with the expression of STC1, besides that patient characteristics are also closely related to recurrence, in this study patients with a strong expression of STC1, stage > 2, age > 45 years and the type of TNBC are patients who are susceptible to recurrence. This study can be used as the basis for further research using STC1 as a therapeutic target to prevent breast cancer recurrence.

Acknowledgment

Thank you to RSUP Dr Kariadi Semarang and the PA Section of RSUP Dr. Kariadi who has helped the smooth running of this research.

References

- Sung H, Ferlay J, Siegel RL, et al. Global Cancer Statistics 2020: GLOBOCAN Estimates of Incidence and Mortality Worldwide for 36 Cancers in 185 Countries. *CA Cancer J Clin*. 2021;71(3). doi:10.3322/caac.21660
PMid: 33538338
- Penyakit Kanker S. InfoDATIN 4 Februari-Hari Kanker Sedunia.; 2015.
- Ng CH, Bhoo Pathy N, Taib NA, et al. Comparison of breast cancer in Indonesia and Malaysia - A clinico-pathological study between dharmas cancer centre Jakarta and university Malaya medical centre, Kuala Lumpur. *Asian Pacific J Cancer Prev*. 2011;12(11):2943-6.
- Colleoni M, Sun Z, Price KN, et al. Annual hazard rates of recurrence for breast cancer during 24 years of follow-up: Results from the international breast cancer study group trials I to V. *J Clin Oncol*. 2016;34(9). doi:10.1200/JCO.2015.62.3504
PMid: 26786933
- Montalvo DN, González NM, Montalvo MTV, Jiménez AA, Echiburú-Chau C, Calaf GM. Patterns of recurrence and survival in breast cancer. *Oncol Rep*. 2008;20(3).
- Joensuu K, Leidenius M, Kero M, Andersson LC, Horwitz KB, Heikkilä P. ER, PR, HER2, Ki-67 and CK5 in early and late relapsing breast cancer-reduced CK5 expression in metastases. *Breast Cancer Basic Clin Res*. 2013;7(1). doi:10.4137/BCBCR.S10701
PMid: 23514931
- Song F, Zhang J, Li S, Wu J, Jin T, Qin J, et al. ER-positive breast cancer patients with more than three positive nodes or grade 3 tumors are at high risk of late recurrence after 5-year adjuvant endocrine therapy. *Onco Targets Ther*. 2017;10:4859-67. http://doi.org/10.2147/OTT.S142698
PMid:29042797
- Kabel AM. Tumor Marker of Breast Cancer: New Prospective Journal of Oncological Sciences; 2017. p. 3.
- van der Leij F, Elkhuizen PH, Bartelink H, van de Vijver MJ. Predictive factors for local recurrence in breast cancer. *Semin Radiat Oncol*. 2012;22(2):100-7. http://doi.org/10.1016/j.semradonc.2011.12.001
PMid:22385917
- Brantley KD, Kjærsgaard A, Cronin-Fenton D, Yacoub R, Nielsen AS, Lauridsen KL, et al. Stanniocalcin expression as a predictor of late breast cancer recurrence. *Cancer Epidemiol Biomarkers Prev*. 2018;27(6):653-9. http://doi.org/10.1158/1055-9965.EPI-17-0905
PMid:29593009
- Hou Y, Wang XZ, Hang YK, Liu JB, Wang MJ. High expression of STC1 is associated with poor prognosis of breast cancer patients. *Int J Clin Exp Pathol*. 2016;9:11826-31.
- Wascher RA, Huynh KT, Giuliano AE, Hansen NM, Singer FR, Elashoff D, et al. Stanniocalcin-1: A novel molecular blood and bone marrow marker for human breast cancer. *Clin Cancer Res*. 2003;9(4):1427-35.
- Jeon M, Han J, Nam SJ, Lee JE, Kim S. STC-1 expression is upregulated through an Akt/NF- κ B-dependent pathway in triple-negative breast cancer cells. *Oncol Rep*. 2016;36(3):1717-22.
- Fang Z, Tian Z, Luo K, Song H, Yi J. Clinical significance of stanniocalcin expression in tissue and serum of gastric cancer patients. *Chin J Cancer Res*. 2014;26(5):602-10. http://doi.

- [org/10.3978/j.issn.1000-9604.2014.10.08](http://doi.org/10.3978/j.issn.1000-9604.2014.10.08)
PMid:25400427
15. Chang AC, Doherty J, Huschtscha LI, Redvers R, Restall C, Reddel RR, *et al.* STC1 expression is associated with tumor growth and metastasis in breast cancer. *Clin Exp Metastasis*. 2015;32(1):15-27. <http://doi.org/10.1007/s10585-014-9687-9>
PMid:25391215
16. Wickberg Å, Magnuson A, Holmberg L, Adami HO, Liljegren G. Influence of the subtype on local recurrence risk of breast cancer with or without radiation therapy. *Breast*. 2018;42:54-60. <http://doi.org/10.1016/j.breast.2018.08.097>
PMid:30176553
17. Senkus E, Kyriakides S, Ohno S, *et al.* Primary breast cancer: ESMO Clinical Practice Guidelines for diagnosis, treatment and follow-up. *Ann Oncol*. 2015;26. doi:10.1093/annonc/mdv298
PMid: 26314782
18. Rosa Mendoza ES, Moreno E, Caguioa PB. Predictors of early distant metastasis in women with breast cancer. *J Cancer Res Clin Oncol*. 2013;139(4). doi:10.1007/s00432-012-1367-z
PMid: 23283528

5. Expression of Stanniocalcin-1 as a Predictor of Recurrence in Breast Cancer Patients

ORIGINALITY REPORT

23%
SIMILARITY INDEX

19%
INTERNET SOURCES

21%
PUBLICATIONS

9%
STUDENT PAPERS

PRIMARY SOURCES

1 Submitted to Badan PPSDM Kesehatan
Kementerian Kesehatan **2%**
Student Paper

2 www.researchsquare.com **1%**
Internet Source

3 ir.lib.hiroshima-u.ac.jp **1%**
Internet Source

4 Nilam Smaradhania, Septiman Rahman,
Salman Ardi Syamsu, Prihantono Prihantono.
"Urokinase type plasminogen activator
receptor (uPAR) and human epidermal growth
factor receptor 2 (HER2) expression in
metastasis of breast cancer", Breast Disease,
2021 **1%**
Publication

5 clonerresources.com **1%**
Internet Source

6 www.science.gov **1%**
Internet Source

7	9dok.org Internet Source	1 %
8	Dhai Rashid, Raghda Bardan. "Anti-mullerian Hormone Serum Level as a Predictor of Responsiveness to Clomiphene Citrate Therapy in Iraqi Women with PCOS", Open Access Macedonian Journal of Medical Sciences, 2022 Publication	1 %
9	Anastasya Siregar, Dian Novita Chandra, Ikhwan Rinaldi. "Correlation of Patient Generated-subjective Global Assessment with Serum C-reactive Protein Level in Stage I-IV Head-and-neck Cancer", Open Access Macedonian Journal of Medical Sciences, 2022 Publication	1 %
10	Siqin Wang, Jin Hu, Yanting Zhang, Jian Shen, Fang Dong, Ximeng Zhang, Chong Lu, Dan Shang. "Presentation and survival by hormonal receptor status in metaplastic breast cancer: A propensity score-matched analysis", The Breast, 2021 Publication	1 %
11	www.communityhealthjournal.org Internet Source	1 %
12	Submitted to Universitas Diponegoro Student Paper	1 %

13 Elridho Sampepajung, William Hamdani, Daniel Sampepajung, Prihantono Prihantono. "Overexpression of NF-kB as a predictor of neoadjuvant chemotherapy response in breast cancer", Breast Disease, 2021
Publication 1 %

14 synapse.koreamed.org
Internet Source 1 %

15 Submitted to CSU, San Jose State University
Student Paper 1 %

16 Littman, E.. "Role of laparoscopic treatment of endometriosis in patients with failed in vitro fertilization cycles", Fertility and Sterility, 200512
Publication 1 %

17 Mohamed Soliman, Ahmed Magdi, Moataz Fatthy, Rania El-Sherif. "The Validity of Carotid Doppler Peak Velocity and Inferior Vena Cava Collapsibility Index in Identifying the Fluid Responders in Mechanically Ventilated Septic Shock Patients", Open Access Macedonian Journal of Medical Sciences, 2022
Publication 1 %

18 aging-us.com
Internet Source 1 %

19 Submitted to Udayana University
Student Paper 1 %

20	www.labroots.com Internet Source	1 %
21	www.ncbi.nlm.nih.gov Internet Source	<1 %
22	biblio.ugent.be Internet Source	<1 %
23	ir.ymlib.yonsei.ac.kr Internet Source	<1 %
24	pdfs.semanticscholar.org Internet Source	<1 %
25	C. Kent Osborne, Patrick Neven, Luc Y. Dirix, John R. Mackey et al. "Gefitinib or Placebo in Combination with Tamoxifen in Patients with Hormone Receptor-Positive Metastatic Breast Cancer: A Randomized Phase II Study", <i>Clinical Cancer Research</i> , 2011 Publication	<1 %
26	Hanif Afkari, Firdian Makrufardi, Basuki Hidayat, Hendra Budiawan, Achmad Hussein Sundawa Kartamihardja. "Correlation between ER, PR, HER-2, and Ki-67 with the risk of bone metastases detected by bone scintigraphy in breast cancer patients: A cross sectional study", <i>Annals of Medicine and Surgery</i> , 2021 Publication	<1 %

27 Jung Hoon Cho, Ji Min Park, Hyung Seok Park, Seho Park, Seung Il Kim, Byeong-Woo Park. "Oncologic safety of breast-conserving surgery compared to mastectomy in patients receiving neoadjuvant chemotherapy for locally advanced breast cancer", Journal of Surgical Oncology, 2013
Publication

28 doc-pak.undip.ac.id
Internet Source

29 jeccr.biomedcentral.com
Internet Source

30 Mahsa Ahadi, Motahareh Heibatollahi, Sara Zahedifard. "Comparison of ER, PR, Ki67 and HER-2/neu Reactivity Pattern with Patients' Age, Histologic Grade, Tumor Size and Lymph Node Status in Invasive Ductal Breast Cancer", International Journal of Cancer Management, 2020
Publication

31 1library.co
Internet Source

32 Dawei Chen, Haiyong Wang, Xinyu Song, Fang Shi, Li Kong, Jinming Yu. "Prognostic score model to determine receipt of radiotherapy after modified radical mastectomy", Oncotarget, 2017

33 Wei-de Zhong, Zhao-dong Han, Hui-chan He, Xue-cheng Bi et al. "CD147, MMP-1, MMP-2 and MMP-9 Protein Expression as Significant Prognostic Factors in Human Prostate Cancer", *Oncology*, 2008

Publication

<1 %

34 e-journal.unair.ac.id

Internet Source

<1 %

35 repository.poltekkes-kdi.ac.id

Internet Source

<1 %

36 "Abstracts—APASL 2013", *Hepatology International*, 2013

Publication

<1 %

37 "Original Research Abstract", *European Heart Journal Supplements*, 2019

Publication

<1 %

38 Olmez, A.. "Risk of recurrence after drug withdrawal in childhood epilepsy", *Seizure: European Journal of Epilepsy*, 200905

Publication

<1 %

39 Samarth Kansara, Vijay Pandey, Peter E. Lobie, Gautam Sethi, Manoj Garg, Amit Kumar Pandey. "Mechanistic Involvement of Long Non-Coding RNAs in Oncotherapeutics Resistance in Triple-Negative Breast Cancer", *Cells*, 2020

<1 %

Publication

Exclude quotes Off

Exclude matches Off

Exclude bibliography On