

Indoleamine 2, 3-dioxygenase (IDO) levels in leprosy patients with positive serology

by M. Zen Rahfiludin

Submission date: 04-May-2021 11:43AM (UTC+0700)

Submission ID: 1577500641

File name: A11_-_Indoleamin_2,_3-dioxygenase_IDO.pdf (136.03K)

Word count: 2057

Character count: 11411

Indoleamine 2, 3-dioxygenase (IDO) levels in leprosy patients with positive serology

Ago Harlim, M. Zen Rahfiludin*

Faculty of Medicine, The Christian University of Indonesia, Jl. Mayjen Sutoyo, Cawang . Jakarta Timur (50275), Indonesia

* Faculty of Public Health, Diponegoro University, Indonesia
Jl. Prof. Soedarto, Sh, Tembalang, Semarang (50239). Indonesia

Abstract

Background Seropositive leprosy individuals have a tendency to experience a change into leprosy lepromatous type. Previous studies have demonstrated that the activity of indoleamine 2, 3-dioxygenase (IDO) in patients with lepromatous type leprosy is higher than those with tuberculoid type leprosy. However, no study has been available showing how IDO level can be correlated to seropositive subjects.

Methods Our observational cross-sectional study was conducted in Brebes, Central Java, Indonesia. Leprosy seropositivity was determined by measuring the level of anti-phenolic glycolipid-1 IgM and IDO level using ELISA.

Results Result of The IDO level in seropositive leprosy subjects was higher compared to those who were seronegative ($p = 0.048$).

Conclusion In the early stage of leprosy infection, a high level of IDO level can already be found; therefore, it can affect immune response and ultimately affecting further course of the disease .

Key words

Seropositive leprosy, indoleamine 2, 3-dioxygenase, anti-phenolic glycolipid-1 IgM.

Introduction

Leprosy still remains a major public health problem in Indonesia since the incidence of leprosy is relatively high.¹ It is possibly due to contact with seropositive leprosy patients that have been turning into manifest leprosy.

Measuring the level of anti-phenolic glycolipid-1 immunoglobulin M (PGL-1 IgM) is one of the methods to detect seropositive leprosy subjects.^{2,3} The level of such cellular immune response will determine leprosy spectrum. The

cellular immunity in leprosy tuberculoid type is still good, on the contrary, there is decreased cellular immunity in leprosy lepromatous type.⁴ Individuals with seropositive leprosy contact (IgM anti-PGL-1 level more than 600 unit/ml), have a tendency to experience a change into leprosy lepromatous type.

Previous studies have demonstrated that leprosy patients with lepromatous type have a higher activity of indoleamine 2, 3-dioxygenase (IDO) compared to those with tuberculoid type. The role of IDO levels in leprosy has been studied and an increased number of macrophages/dendritic cells (DC-lineage IDO+ cells were found in lepromatous (LL) compared to tuberculoid (BT) and reversal reaction (RR) patients. LL lesions were characterized by

Address for correspondence

M. Zen Rahfiludin
Faculty of Public Health,
Universitas of Diponegoro, Jl. Prof Soedarto Sh,
Tembalang, Semarang, Indonesia
Email: rafiludin@yahoo.com

massive macrophage infiltration containing a variable number of Virchow cells, full of bacilli accompanied by few lymphocytes, separated from the flat epidermis by a clear zone. BT lesions were characterized by an organized dermal granuloma composed of phagocytic cells with evident epithelioid differentiation surrounded by lymphocytes of giant cells. RR lesions showed activated macrophages with an epithelioid appearance, organized or not as granuloma.⁵

However, no study has been conducted to evaluate how IDO level can be correlated to seropositive leprosy subjects (IgM anti PGL – 1 level more than 600 unit/ml) and have not showed a clinical manifestation on the skin. Our study aimed to identify the difference of IDO level between seropositive and seronegative leprosy subjects.

Methods

Our study was an observational study using a cross-sectional method. In August 2015, the study was conducted in Brebes, Central Java, Indonesia. Ethical clearance was approved by the Ethics Committee of Medical and Public Health Research, Faculty of Public Health, Diponegoro University.

The eligible subjects participated in our study were those who fulfilled the inclusion criteria: individuals who were living together with a leprosy patient, age between 15-55 years, showed no clinical symptom of leprosy, were not taking immunosuppressants during the last three months prior to the study and those who were willing to participate in our study. They also did not suffer from tuberculosis and did not take any anti-tuberculosis medications. The subjects were divided into 2 groups, i.e. those with seropositive leprosy (anti PGL-1 IgM >

600 unit/mL) 6 and those who were seronegative.

Five milliliters of venous blood was drawn from each subject. The whole blood was centrifuged to separate the serum, which was then stored at –70°C. Seropositive leprosy was determined by measuring the levels of anti-PGL-1 IgM antibodies using enzyme-linked immunosorbent assay (ELISA) (Polyclonal rabbit anti-human IgM/HRP (Dako ®) and analyzing IDO levels using ELISA commercial kits (Cusabio®, Catalog No: CSB-E09966h). All tests were conducted at the Institute Tropical Disease Laboratory, Airlangga University, Surabaya.

The education level of subjects was categorized into 2 groups, i.e. basic and moderate education. Subjects in basic education group were those who had completed their elementary and junior high school; while moderate education group involved those with senior high school educational background

Results

Table 1 shows that the gender, level of education and age had no effect on the seropositivity (anti-PGL-1 IgM antibodies levels) in leprosy patients; but the levels of IDO were significantly different between seropositive leprosy with seronegative subjects ($p = 0.048$).

In sex variable, there were 20.7% male and 79.3% female with seropositive leprosy and 33.3% male and 66.7% female with seronegative leprosy. There is no significant difference between seropositive leprosy subjects and seronegative leprosy subjects in sex variable ($p = 0.275$).

In level of education variable, seropositive leprosy subjects in primary education is 82.8% and 17.2% is in higher education while in

Table 1 Subject characteristics and indoleamine 2, 3-dioxygenase (IDO) levels.

Variables	Seropositive leprosy (n=29)	Seronegative leprosy (n =30)	P
Sex			
Male (%)	20.7	33.3	0.275 ^a
Female (%)	79.3	66.7	
Level of education			
Primary education (%)	82.8	80.0	0.786 ^a
Higher education (%)	17.2	20.0	
Age (years), (mean + SD)	33 ± 10	35 ± 11	0.399 ^b
IDO (pg/mL), (median + SD)	13.80 ± 9.13	9.10 ± 4.60	0.048 ^c

a=chi-square test, b= independent t-test, c=Mann-Whitney test

seronegative leprosy subjects there were 80% in primary education and 20% in higher education ($p = 0.786$)

The mean age of seropositive leprosy subjects was 33 ± 10 years and the mean age of seronegative leprosy subjects was 35 ± 11 years. ($p = 0.399$)

The mean IDO level in seropositive leprosy subject was 13.80 ± 9.13 pg/mL while that in seronegative leprosy subject was 9.10 ± 4.60 pg/mL. There was significant difference between seropositive leprosy and seronegative leprosy subjects in terms of IDO level ($p=0.048$).

Discussion

The first body mechanical defense against *Mycobacterium leprae* is by activating the antigen presenting cells (APCs) of immune system, i.e. the dendritic cells, monocytes and macrophages. IDO is predominantly expressed in APCs. Interferon gamma (IFN- γ) and dependent and/or independent signal pathways can activate and mediate IDO.⁷

Higher level of IDO found in seropositive leprosy subjects of our study supported the results of previous studies which indicate a high IDO protein expression in lepromatous skin lesions and high IDO activity in the sera of lepromatous leprosy patients.⁵ People who live

together under one and the same roof with leprosy patients with seropositive anti-PGL-I antibody have higher risk for developing leprosy (particularly the lepromatous type) compared to those who have contact with seronegative patients. The anti-PGL-I antibody level in people who live together under one and the same roof with leprosy indicates increased risk of having leprosy in the next 4 years. Another study in Philippines showed an increased risk of developing leprosy to 24-fold is caused by seropositive contacts.^{8,9}

In addition due to *M. leprae*, the increased IDO activity in the sera of lepromatous leprosy patients also depends on IFN- γ . There is also a statement that IDO activity was at least partially dependent upon IFN- γ from a study in dengue patients.¹⁰ IDO can be induced in response to various stimuli. The most potent inducer is IFN- γ , although other inflammatory stimuli, including type-I interferons (IFN- α/β) and bacterial lipopolysaccharide (LPS), and some viruses and intracellular pathogens, can induce IDO expression.¹⁵ Support for the "two signal" requirement for IDO-expression has also been obtained in human samples.¹⁴ For example, human monocytes expressed IDO in lepromatous leprosy due to the synergistic actions of IFN- γ and the bacterium (likely through pattern recognition receptor signaling).⁵ *M. leprae* and IFN- γ is a cause of an increased of IDO expression in monocytes leprosy patients lepromatous type.⁵ Leprosy patients with clinical

manifestation of leprosy or those with seronegative leprosy have a less IFN- γ level than seropositive leprosy.¹¹

A higher IDO level in seropositive group found in our study supports the results of some previous studies, which showed that IDO induction after bacterial infection could be beneficial to the host. Amino acid tryptophan will break into stable metabolite called kynurenine when the IDO level increases.¹² Moreover, tryptophan depletion and starvation will occur since bacteria need tryptophan.^{13,14} The growth of bacteria is dependent on the greater amount of tryptophan rather than T cells.¹³

20
IDO could be considered to be a double-edged sword in the immune response against pathogens. Immune cells, including antigen-presenting cells like DCs, are able to express IDO in response to various stimuli.¹⁰

Acknowledgement

Our grateful thank to Prof. Indropo Agustini, MD, DV and Leprosy Study Group of Tropical Disease Centre for using the laboratory for IgM anti-PGL-1 and IDO assay test at University of Airlangga, Surabaya. Indonesia and also to the technician of the Institute Tropical Disease (ITD) Laboratory, Iswayudi SKM; Dinar Adriaty, S.Si, M. Kes; Ratna Wahyuni S.Si, M. Kes who examined IgM anti-PGL-1 and IDO.

References

- 12
1. World Health Organization (WHO). Global leprosy update, 2013; reducing diseases burden. *Wkly Epidemiol Res*. 2014;89:389-400.
2. Gusni I, Adriaty D, Wahyuni R, Izumi S. Correlation between the sero-positivity level and new case detection rate (NCDR) in some leprosy endemic areas of Indonesia. *Jpn J Lepr*. 2009;78:140.

- 13
3. Izumi S. Subclinical infection by *Mycobacterium* *prae*. *Int J Lepr*. 1999;67 (4) (Suppl):S67-71.
4. Scollard DM, Adams LB, Gillis TP, Krahenbuhl JL, Truman RW, Williams DL. The continuing challenges of leprosy. *Clin Microbiol Rev*. 2006;19:338-81.
5. de Souza Sales J, Lara FA, Amadeu TP, de Oliveira Fulco T, da Costa Nery JA, Sampaio EP *et al*. The role of indoleamine 2, 3-dioxygenase in lepromatous leprosy immunosuppression. *Clin Exp Immunol*. 2011;165:251-63. 3
6. Rahfiludin MZ, Pangestuti DR. Phyllanthus niruri extract could improve immunoglobulin-M anti phenolic glycolipid-1 level in seropositive contact of Hansen's disease patients. *J Health Res*. 2012;26:55-8.
7. Murakami Y, Hoshi M, Imamura Y, Arima Y, Yamamoto Y, Saito K. Remarkable role of indoleamine 2,3-dioxygenase and tryptophan metabolites in infectious diseases: potential role in macrophage-mediated inflammatory diseases. *Mediators Inflamm*. 2013;2013.
8. Poet FJ, Meima A, Oskam L, Richardus JH. Risk factors for the development of clinical leprosy among contacts, and their relevance for targeted interventions. *Lepr Rev*. 2004;75:310-32.
9. Douglas JT, Cellona RV, Fajardo TT Jr, Abalos RM, Balagon MV, Klaster PR. Prospective study of serological conversion as a risk factor for development of leprosy among household contacts. *Clin Diag Lab Immunol*. 2004;11:897-900.
10. Becerra A, Warke RV, Xhaja K, Evans B, Evans Martin K *et al*. Increased activity of indoleamine 2, 3-dioxygenase in serum from acutely infected dengue patients linked to gamma interferon antiviral function. *J Gen Virol*. 2009;90:810-7.
11. Rahfiludin MZ, Kartasurya MI, Purwaningsih E. The different levels of interferon gamma capacity production on several stages of leprosy. *Med J Indones*. 2007;16:224-7.
12. Harden JL, Egilmez NK. Indoleamine 2, 3-dioxygenase and dendritic cell tolerogenicity. *Immunol Invest*. 2012;41:738-64.
13. Muller AJ, Heselger K, Schmidt SP, Specker K, Mackenzie CR, Däubener W. The missing link between indoleamine 2, 3-dioxygenase mediated antibacterial and immunoregulatory effects. *J Mol Med*. 2009;13:1125-35.
14. Chau F, Geffers R, Thalmann J, Haller H, Wagner AD. Restriction of *Chlamydia pneumoniae* replication in human dendritic cell by activation of indoleamine 2, 3-dioxygenase. *Microbes Infect*. 2009;11:1002-10.
15. King NJ, Thomas SR. Molecules in focus: indoleamine 2,3-dioxygenase. *Int J Biochem Cell Biol*. 2007;39:2167-72.

Indoleamine 2, 3-dioxygenase (IDO) levels in leprosy patients with positive serology

ORIGINALITY REPORT

17%

SIMILARITY INDEX

13%

INTERNET SOURCES

10%

PUBLICATIONS

6%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to University of Warwick Student Paper	1%
2	www.nejm.org Internet Source	1%
3	www.tci-thaijo.org Internet Source	1%
4	journals.sagepub.com Internet Source	1%
5	Ioannis D. Bassukas, Georgios Gaitanis, Max Hundeiker. "Leprosy and the natural selection for psoriasis", Medical Hypotheses, 2012 Publication	1%
6	M. S. Duthie, W. Goto, G. C. Ireton, S. T. Reece et al. "Use of Protein Antigens for Early Serological Diagnosis of Leprosy", Clinical and Vaccine Immunology, 2007 Publication	1%
7	sinta3.ristekdikti.go.id Internet Source	1%

8	www.wjgnet.com Internet Source	1 %
9	www.neliti.com Internet Source	1 %
10	Amanda Carreira Devides, Patricia Sammarco Rosa, Andréa de Faria Fernandes Belone, Neusa Maria Broch Coelho et al. "Can anti-PGL-1 and anti-NDO-LID-1 antibody titers be used to predict the risk of reactions in leprosy patients?", <i>Diagnostic Microbiology and Infectious Disease</i> , 2018 Publication	1 %
11	livedna.net Internet Source	1 %
12	www.dovepress.com Internet Source	1 %
13	Russell E. Bartt. "Leprosy (Hansen's disease)", <i>Current Treatment Options in Neurology</i> , 2004 Publication	1 %
14	www.microbiologyresearch.org Internet Source	1 %
15	inabj.org Internet Source	1 %
16	pdfs.semanticscholar.org Internet Source	1 %

17	ashpublications.org Internet Source	1 %
18	jeccr.biomedcentral.com Internet Source	1 %
19	unair.ac.id Internet Source	<1 %
20	Tim T. Lögters, Maurice D. Laryea, Marcus Jäger, Michael Schädel-Höpfner et al. "Kynurenine inhibits chondrocyte proliferation and is increased in synovial fluid of patients with septic arthritis", Journal of Orthopaedic Research, 2010 Publication	<1 %
21	www.rug.nl Internet Source	<1 %
22	Eleftheriadis, Theodoros, Georgios Pissas, Georgia Antoniadis, Vassilios Liakopoulos, and Ioannis Stefanidis. "Indoleamine 2,3-dioxygenase depletes tryptophan, activates general control nonderepressible 2 kinase and downregulates key enzymes involved in fatty acid synthesis in primary human CD4+ T-cells", Immunology, 2015. Publication	<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On

Indoleamine 2, 3-dioxygenase (IDO) levels in leprosy patients with positive serology

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4
