

glaucoma

by Maharani Maharani

Submission date: 05-May-2023 04:55PM (UTC+0700)

Submission ID: 2084955642

File name: C-1_2022_BMJ_Glaukoma.pdf (253.26K)

Word count: 2343

Character count: 12805

Aqueous Humour Malondialdehyde Level as Oxidative Stress Marker In Types Of Glaucoma



Maharani¹, Puspita Kusuma Dewi¹, Riski Prihatningtias¹,
Arief Wildan¹, Trilaksana Nugroho¹,
Edward Kurnia Setiawan Limijadi^{2*}, Fifin L Rahmi¹

¹Department of Ophthalmology, Medical Science Faculty, Universitas Diponegoro, Indonesia.

²Department of Clinical Pathology, Medical Science Faculty, Universitas Diponegoro, Indonesia.

*Corresponding to:

Edward Kurnia Setiawan Limijadi;
Department of Clinical Pathology,
Medical Science Faculty, Universitas
Diponegoro, Indonesia;
edwardksti@fk.undip.ac.id

Received: 2021-07-31

Accepted: 2022-02-25

Published: 2022-03-14

ABSTRACT

Background: Angle-closure glaucoma is Asia's most common type of glaucoma, leading to worse damage due to its oxidative stress. Another type is open-angle glaucoma in the second line. The oxidative stress level can be known by malondialdehyde level examination. The study about MDA levels in aqueous humour in primary angle-closure and open angle glaucoma has not been done before. Therefore, the objective of this study was to determine the difference of malondialdehyde levels in aqueous humour between primary open-angle glaucoma and primary angle-closure glaucoma.

Methods: This study was an analytical observational study. Nine subjects were primary open-angle glaucoma patients, and ten subjects were primary angle-closure glaucoma patients. The aqueous humour was aspirated from the anterior chamber during trabeculectomy and measured the MDA levels using a spectrophotometer. Data were analyzed by the Mann-Whitney test.

Results: There was no significant difference ($p=0.095$) of MDA levels in aqueous humour between primary open-angle glaucoma ($62.913 \pm 17.259 \mu\text{mol/L}$) and primary angle-closure glaucoma ($53.485 \pm 9.670 \mu\text{mol/L}$).

Conclusion: The oxidative stress level in primary open-angle glaucoma and primary angle-closure glaucoma is the same, indicating that the inflammation status in both conditions could be similar.

Keywords: Malondialdehyde (MDA), primary open-angle glaucoma, primary angle-closure glaucoma

Cite This Article: Maharani, Dewi, P.K., Prihatningtias, R., Wildan, A., Nugroho, T., Limijadi, E.K.S., Rahmi, F.L. 2022. Aqueous Humour Malondialdehyde Level as Oxidative Stress Marker In Types Of Glaucoma. *Bali Medical Journal* 11(1): 103-105. DOI: 10.15562/bmj.v11i1.2599

INTRODUCTION

Glaucoma is the second largest cause of blindness after Cataracts.¹ The blindness due to glaucoma in Indonesia reached 0.16% of the total population.² The glaucoma prevalence in Indonesia, according to the Basic Health Research (Riset Kesehatan Dasar or Riskesdas) in 2007, was around 4.6%. In Central Java, the glaucoma prevalence in 2007 was 2,3%.³

Definition of glaucoma is a group of progressive optic neuropathies characterized by degeneration of retinal ganglion cells and resulting optic nerve head.⁴ Degeneration of these nerves results in cupping, a characteristic appearance of the optic disc and visual loss.⁵ Glaucomas can be classified into open-angle glaucoma (POAG) and angle-closure glaucoma (PACG).⁴ Primary open-angle glaucoma (POAG) is a chronic,

progressive optic neuropathy in adults characterized by acquired atrophy of the optic nerve and loss of retinal ganglion cells and their axons.⁶ Primary Angle Closure Glaucoma (PACG) is responsible for half of glaucoma-related blindness. PACG is characterized by appositional approximation or contact between the iris and trabecular meshwork.⁷ Pathogenesis of glaucoma is not fully understood, but some factors are associated with this condition, such as high intraocular pressure (IOP), increased oxidative stress, aging, glutamate neurotoxicity, and susceptibility genes such as optineurin and myocilin.⁸⁻¹⁰

Oxidative stress is an important risk factor in the pathogenesis of glaucoma. Increasing ROS (Reactive Oxygen Species) production and imbalance between oxidative and antioxidant capacity have been postulated as the crucial factors in early retinal injury, together with the

reduced ocular perfusion pressure in the blood vessels (the vascular theory of glaucoma).¹¹ Studies conducted on aqueous humour and the blood of glaucoma patients reported an increase in the level of oxidative stress of malondialdehyde (MDA) and decreased total capacity of antioxidants.¹² Analysis of some studies showed an increase in oxidative damage to the nucleus and mitochondria of DNA in patients with glaucoma compared to normal people.¹³⁻¹⁴ The increase of oxidative stress in patients with glaucoma is the cause of apoptosis reaction in TM.¹⁵⁻¹⁶ The study by Ghanem *et al.* (2010), which assessed the correlation between oxidative stress and glaucoma severity, found higher MDA levels in glaucoma patients compared to the control group and a positive correlation between high MDA levels and the damage to the visual field.¹⁷ Based on the study of Faschinger *et*

al. (2006), there was a positive correlation between the MDA serum levels and MDA levels in aqueous humour of patients with angle-closure glaucoma. The study also found no significant difference between MDA levels in aqueous humour compared with levels in serum and plasma.¹⁸

Both Primary open-angle glaucoma (POAG) and primary angle-closure glaucoma (PACG) are glaucoma types that need to be studied about oxidative stress, such as MDA. The fact is that MDA has been increasing in glaucoma condition, but the level in POAG and PACG has not been understood. Therefore, this study tried to explain the MDA level in POAG and PACG for early study in Indonesia.

METHODS

This was an analytical observational study with a cross-sectional design. We conducted the study in the Ophthalmology Department of Dr. Kariadi Hospital and the GAKI laboratory of the Faculty of Medicine of Diponegoro University from November 2016 to July 2017.

The subject of this study was patients with POAG and PACG who underwent trabeculectomy. Patients with diseases that cause increased MDA levels such as diabetes mellitus, infection or inflammation and cancer, also patients who consumed antioxidants were excluded in this study.

Nine patients with POAG and 10 patients with PACG met with exclusion criteria. Aqueous humour was taken by aspiration during trabeculectomy surgery. It needed at least 0.025 ml using a 1cc disposable syringe. Samples were stored in tubes at a -20° C temperature for the examination. The samples were examined for MDA level in GAKI laboratory using the Quantikine Immunoassay Kit and read the results with a spectrophotometer at the excitation wavelength of 515 nm and an emission wavelength of 553 nm. Computer programs analyzed data. We performed the normality test and Mann-Whitney test. $P < 0.05$ was significant.

This study obtained ethical approval from the Research Ethics Commission of the Faculty of Medicine of Diponegoro University. All the subjects in this study have declared to join and signed the informed consent form.

RESULTS

Table 1. shows the subject of the characteristics in this study. Subjects with both primary open-angle glaucoma (POAG) types consist of 7 male (77.8%) and 2 (22.2%) females; meanwhile, subjects with primary angle-closure glaucoma (PACG) types consist of 2 male (20%) and 8 (80%) female. The median age subject with POAG is 57.78 ± 9.271 , and PACG is 58.8 ± 9.852 . The next analysis performed the difference test for MDA level in aqueous humour between POAG and PACG. Table 2 showed no significant difference in MDA levels in aqueous humour between POAG and PACG. MDA levels of the subject with POAG are higher than subjects with PACG.

DISCUSSION

Malondialdehyde (MDA) is one of the free radicals used as a marker for glaucoma.¹⁹ Nucci et al., (2013) showed an increase in the level of malondialdehyde and decreased total capacity of antioxidants in the blood and aqueous humor of patients with primary open-angle glaucoma.¹² Yildirim et al. (2005) showed an increase in MDA plasma levels in patients with glaucoma compared to the control group, and there was no significant difference in MDA levels in the blood and aqueous humour.²⁰

The study by Ghanem et al. (2010) assessed the correlation between the oxidative stress marker and the severity

of glaucoma in the primary open-angle glaucoma showed MDA levels in aqueous humor patients with aqueous humor glaucoma were higher compared to the control group. The study also showed a positive correlation between MDA level and the visual field damage.¹⁷

Glaucoma is a multifactorial disease and intraocular pressure is one of the most suspected risk factors causing damage to the retinal ganglion cells. Increased intraocular pressure and the visual field abnormalities occurring in glaucoma are associated with oxidative damage to DNA in the trabecular meshwork.¹⁴ Both POAG and PACG have suffered damage in the retinal ganglion cells, visual field abnormalities, and damage in the trabecular meshwork due to increased intraocular pressure. This almost identical pathophysiology and damage process cause the non-significant difference in MDA levels in the aqueous humour.

Primary angle-closure glaucoma consists of several stages, and most patients who come to the hospital are patients suffering from acute stage episodes and chronic stage patients. In this study, we selected the chronic stage patients who didn't show signs of inflammation to avoid bias from high MDA levels due to the inflammatory process in the acute stage. Therefore, we found clinical features and damages almost the same as those found in the primary open-angle glaucoma.

This study was preliminary that needs to explore more deeply with a larger

Table 1. The characteristics of subjects in the study

Characteristics	POAG	PACG	P
Gender			
Male N (%)	7 (77.8)	2 (20.0)	
Female N (%)	2 (22.2)	8 (80.0)	
Age (Median \pm SE)	57.78 ± 9.271	58.8 ± 9.852	0.455

Table 2. MDA level in POAG and PACG

Variable	Glaucoma groups (Median \pm SE)		P
	POAG	PACG	
MDA level ($\mu\text{mol/L}$)	62.913 ± 17.259	53.485 ± 9.670	0.095*

* = not significant

number of patients, both POAG and PACH. Furthermore, it is difficult to analyze the MDA level at various POAG and PACG. This study also did not determine the correlation between MDA level and variable of clinical visual damage. These limitations need to be evaluated in the next study.

CONCLUSIONS

The oxidative stress level in primary open-angle glaucoma and primary angle-closure glaucoma is the same, indicating that the inflammation status in both status could be similar.

CONFLICT OF INTEREST

No conflict of interest

ACKNOWLEDGEMENTS

The researchers would like to thank nurses at Dr. Kariadi Semarang Hospital, who had assisted in this research in sampling and laboratory technicians at GAKY laboratory who had examined the sample. The Medical Science Faculty funding granted this research in 2017.

CONFLICT OF INTEREST

All authors declared that there is no conflict of interest regarding this article

ETHICS APPROVAL

Research Ethics Commission has ethically approved this study of the Faculty of Medicine of Diponegoro University with ethical clearance number 224/EC/FK/RSDK 2020. All the subjects in this study have declared to join and signed the informed consent form

FUNDING

This article was self-funded

AUTHOR CONTRIBUTION

All authors contributed equally in the research and writing process of this article

REFERENCES

1. Bulletin of the World Health Organization : The International Journal of Public Health 2004 ; 82(11) : 844-851.
2. Ilyas S. 2007. Glaucoma. Ed 3. Jakarta: Sagung Seto, pp:2-3.
3. Ministry of Health Republic Indonesia. 2013. Basic Health Research. Jakarta: Data and Information Center.
4. Weinreb RN, Aung T, Medeiros FA, The Pathophysiology and Treatment of Glaucoma: A Review. JAMA. 2014 May 14; 311(18): 1901–1911.
5. Weinreb RN, Khaw PT. Primary open-angle glaucoma. Lancet. 2004; 363(9422):1711–1720.
6. American Academy of Ophthalmology. 2015. Primary Open-Angle Glaucoma. America: Elsevier Inc. pp 41-111.
7. Wright C, Tawfik MA, Waisbourd M and Katz LJ, Primary angle-closure glaucoma: an update. Acta Ophthalmol. 2016; 94: 217–225
8. M. Seki and S. A. Lipton, "Targeting excitotoxic/free radical signaling pathways for therapeutic intervention in glaucoma," Progress in Brain Research, vol. 173, pp. 495–510, 2008.
9. S. F. Janssen, T. G. M. F. Gorgels, W. D. Ramdas et al., "The vast complexity of primary open angle glaucoma: disease genes, risks, molecular mechanisms and pathobiology," Progress in Retinal and Eye Research, vol. 37, pp. 31–67, 2013.
10. N. Osborne and S. Del Olmo-Aguado, "Maintenance of retinal ganglion cell mitochondrial functions as a neuroprotective strategy in glaucoma," Current Opinion in Pharmacology, vol. 13, no. 1, pp. 16–22, 2013.
11. Kimura A, Namekata K, Guo X, Noro T, Harada C, and Harada T. Targeting Oxidative Stress for Treatment of Glaucoma and Optic Neuritis. Hindawi Oxidative Medicine and Cellular Longevity. 2017;1-8
12. Nucci C, Di Piero D, Varesi C, Ciuffoletti E, Russo R, Gentile R, et al. Increased malondialdehyde level and reduced total antioxidant capacity in aqueous humour and blood samples from patients with glaucoma. Mol Vis, 2013;7;19:1841-6.
13. Izzoti A, Sacca SC, Cartiglia C, De Flora S. Oxidative deoxyribonucleic acid damage in the eyes of glaucoma patients. Am J Med, 2003;114(8):638-46.
14. Sacca SC, Pascotto A, Camicione P, Capris P, Izzotti A. Oxidative DNA damage in the human trabecular meshwork: clinical correlation in patients with primary open-angle glaucoma. Arc Ophthalmol, 2005;123(4):458-63.
15. Baleriola J, Julian GE, Jose MM, Arturo FC, Enrique JR, Raquel FD. Apoptosis in the trabecular meshwork of glaucomatous patients. Mol Vis, 2008;14:1513-6.
16. Nurwasis, Suhendro G, Sudiana IK. The Role of SOD, Catalase, HSP-27, HSP- 70, and TNF- α Expression in Apoptosis of Retinal Ganglion Cells After Intra Ocular Pressure Increase on Rattus Norvegicus. IJPHRD, 2019;10(7):1174-8.
17. Ghanem AA, Arafa LF, El-Baz A. Oxidative stress markers in patients with primary open-angle glaucoma. Curr Eye Res. 2010;35(4):295-301.
18. Faschinger C, Schmut O, Wachswender C, Mossbock G. Glaucoma and oxidative stress determination of malondialdehyde a product of lipid peroxidation. Ophthalmologie. 2006;103(11):953-9.
19. Rokicki W , Zaleska-Fiolka J , Pojda-Wilczek D , Hampel A , Majewski W , Ogultekin S, Mrukwa-Kominek E. Differences in serum oxidative status between glaucomatous and nonglaucomatous cataract patients. BMC Ophthalmology. 2017;17(13):1-7.
20. Yildirim O, Ates NA, Ercan B, Muslu N, Unlu A, Tamer L, et al. Role of oxidative stress enzyme in open angle glaucoma. Laboratory study. Eye (Lond). 2005;19(5):580-3.



This work is licensed under a Creative Commons Attribution

glaucoma

ORIGINALITY REPORT

25%

SIMILARITY INDEX

18%

INTERNET SOURCES

15%

PUBLICATIONS

13%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Nahdlatul Ulama Surabaya Student Paper	3%
2	roj.igb.ru Internet Source	2%
3	www.journalofoptometry.org Internet Source	2%
4	Syaiful Anwar, Purwanto Adhipireno, Ria Triwardhani, Edward Kurnia Setiawan Limijadi. "The Correlation between Serum Amyloid A, Mean Platelet Volume, and Creatine Kinase Myocardial B in Acute Coronary Syndrome", Open Access Macedonian Journal of Medical Sciences, 2021 Publication	2%
5	ijbs-udayana.org Internet Source	2%
6	www.dynamed.com Internet Source	1%

7	Wenjuan Zhuang, Shaolin Wang, Juan Hao, Manyun Xu, Hao Chi, Shunyu Piao, Jianqing Ma, Xiaolong Zhang, Shaoping Ha. "Genotype-ocular biometry correlation analysis of eight primary angle closure glaucoma susceptibility loci in a cohort from Northern China", PLOS ONE, 2018 Publication	1 %
8	www.ijfans.org Internet Source	1 %
9	worldwidescience.org Internet Source	1 %
10	Submitted to Baskent University Student Paper	1 %
11	jama.jamanetwork.com Internet Source	1 %
12	medworm.com Internet Source	1 %
13	www.ijo.cn Internet Source	1 %
14	Mansi Vishal, Anchal Sharma, Lalit Kaurani, Giovanna Alfano et al. "Genetic association and stress mediated down-regulation in trabecular meshwork implicates MPP7 as a novel candidate gene in primary open angle glaucoma", BMC Medical Genomics, 2016	1 %

15 Ireneusz Majsterek, Katarzyna Malinowska, Malgorzata Stanczyk, Michal Kowalski et al. "Evaluation of oxidative stress markers in pathogenesis of primary open-angle glaucoma", Experimental and Molecular Pathology, 2011

Publication

16 Submitted to New Jersey Institute of Technology

Student Paper

17 www.jci.org

Internet Source

18 Submitted to Queen's University of Belfast

Student Paper

19 bioclima.ro

Internet Source

20 eprints.ucm.es

Internet Source

21 ijbms.mums.ac.ir

Internet Source

22 link.springer.com

Internet Source

23 www.ofthalmoloji.org

Internet Source

24

G Gazzard. "Intraocular pressure and visual field loss in primary angle closure and primary open angle glaucomas", British Journal of Ophthalmology, 2003

Publication

<1 %

25

Chaobin Lin, Fujin Wu, Tongmei Zheng, Xiuchun Wang, Yiwei Chen, Xiaomin Wu. "Kaempferol attenuates retinal ganglion cell death by suppressing NLRP1/NLRP3 inflammasomes and caspase-8 via JNK and NF-κB pathways in acute glaucoma", Eye, 2018

Publication

<1 %

26

Yoshida, Kaori, Ikuyo Ohguro, and Hiroshi Ohguro. "Black Currant Anthocyanins Normalized Abnormal Levels of Serum Concentrations of Endothelin-1 in Patients with Glaucoma", Journal of Ocular Pharmacology and Therapeutics, 2012.

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

glaucoma

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3
