

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artike) : Effectiveness Cream and Ointment of Spirulina platensis extract against amount of Fibroblas and Wound Area: Study on white rats whose skin is incised

Jumlah Penulis : 3 orang

Status Pengusul : Bimby Irenesia, Renni Yuniarti, Endang Mahati

Identitas Jurnal Ilmiah :

Indonesian Journal of Environmental Management and Sustainability (IJEMS)

a. Nama Jurnal : Sustainability (IJEMS)

b. Nomor ISSN : p-ISSN: 2598-6260 e-ISSN: 2598-6279

c. Volume/ nomor /Hal : Vol.4 no.2, p: 39-42

d. Edisi : Juni th 2020

e. Penerbit : Pascasarjana FMIPA Universitas Sriwijaya

f. Jumlah halaman : 4

g.DOI artikel (Jika ada) : <https://doi.org/10.26554/ijems.2020.4.2.39-42>

h. Alamat web Jurnal : <http://ijoems.com/index.php/ijems/article/view/116>

i. Terindeks di : SINTA 3

j. Online Turnitine : <https://doc-pak.undip.ac.id/1925/1/Turnitin-Effectiveness-Cream.pdf>

Kategori Publikasi Jurnal Ilmi (beri ✓ pada kategori yang tepat)

☐ Jurnal Ilmiah Internasional / internasional bereputasi *

☒ Jurnal Ilmiah Nasional Terakreditasi

☐ Jurnal Ilmiah Nasional/ Nasional terindeks di DOAJ CABI COPERNICUS** Tidak Terakreditasi

Hasil Penilaian Peer Review :

NO	KOMPONEN YG DINILAI	Nilai Maksimal Jurnal Ilmiah			Nilai Akhir Yang Diperoleh
		Internasional/internasional bereputasi	Nasional Terakreditasi	Nasional ***	
		<input type="text"/>	20	<input type="text"/>	
a	Kelengkapan unsur isi artikel (10%)		2		2
b	Ruang lingkup dan kedalaman pembahasan (30%)		6		5
c	Kecukupan dan kemutahiran data/informasi dan metodologi (30%)		6		5
d	Kelengkapan unsur dan berkualitas terbitan/ jurnal (30%)		6		5,5
Nilai Total = (100%)			20		17,5
Nilai Pengusul =				(17,5x 40%/2) = 3,5	3,5

Catatan Penilaian artikel oleh Reviewer

a. Kelengkapan unsur isi artikel (10%) : unsur isi artikel sudah lengkap mulai abstrak-pendahuluan-metode-hasil-pembahasan & referensi

b. Ruang lingkup dan kedalaman pembahasan (30%) :Ruang lingkup keilmuan sudah sesuai dg bidang ilmu pengusul. Pembahasan cukup dalam dan telah membandingkan dengan referensi2 yang sesuai

c. Kecukupan dan kemutahiran data/informasi dan metodologi (30%) : Data cukup lengkap untuk penelitian eksperimental dengan hewan coba, metode dijelaskan secara lengkap.

d. Kelengkapan unsur dan berkualitas terbitan/ jurnal (30%) :Jurnal nasional berbahasa Inggris terindex Sinta 3

Semarang,
Reviewer 1

Prof. Dr. dr.Tri Nur Kristina ,DMM,M.Kes
NIP 194902091979012001
Unit kerja: Fakultas Kedokteran Undip
Bidang ilmu : Ilmu Kedokteran
Jabatan/pangkat: Guru Besar

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel)	: Effectiveness Cream and Ointment of Spirulina platensis extract against amount of Fibroblas and Wound Area: Study on white rats whose skin is incised
Jumlah Penulis	: 3 orang
Status Pengusul	: Bimby Irenesia, Renni Yuniarti, Endang Mahati
Identitas Jurnal Ilmiah	: a. Nama Jurnal : Indonesian Journal of Environmental Management and Sustainability (IJEMS) b. Nomor ISSN : p-ISSN: 2598-6260 e-ISSN: 2598-6279 c. Volume/ nomor /Hal : Vol.4 no.2, p: 39-42 d. Edisi : Juni th 2020 e. Penerbit : Pascasarjana FMIPA Universitas Sriwijaya f. Jumlah halaman : 4 g.DOI artikel (Jika ada) : https://doi.org/10.26554/ijems.2020.4.2.39-42 h. Alamat web Jurnal : http://ijoems.com/index.php/ijems/article/view/116 i. Terindeks di : SINTA 3 j. Online Turnitine : https://doc-pak.undip.ac.id/1925/1/Turnitin-Effectiveness-Cream.pdf

Kategori Publikasi Jurnal Ilmial	: <input type="checkbox"/> Jurnal Ilmiah Internasional / internasional bereputasi *
(beri ✓ pada kategori yang tepat)	: <input checked="" type="checkbox"/> Jurnal Ilmiah Nasional Terakreditasi
	: <input type="checkbox"/> Jurnal Ilmiah Nasional/ Nasional terindeks di DOAJ CABI COPERNICUS** Tidak Terakreditasi

Hasil Penilaian Peer Review :

NO	KOMPONEN YG DINILAI	Nilai Maksimal Jurnal Ilmiah			Nilai Akhir Yang Diperoleh
		Internasional/internasional bereputasi	Nasional Terakreditasi	Nasional ***	
		<input type="text"/>	<input type="text" value="20"/>		
a	Kelengkapan unsur isi artikel (10%)		2		2
b	Ruang lingkup dan kedalaman pembahasan (30%):		6		5,7
c	Kecukupan dan kemutahiran data/informasi dan metodologi (30%)		6		5,8
d	Kelengkapan unsur dan berkualitas terbitan/ jurnal (30%)		6		5,0
Nilai Total = (100%)			20		18,5
Nilai Pengusul =				(..... x 40%/2 =	3,7

Catatan Penilaian artikel oleh Reviewer	
a.	Kelengkapan unsur isi artikel : Abstrak disajikan lengkap, jelas dan informatif. Kesimpulan, pembahasan, hasil dan pembahasan, serta saran dan rekomendasi. Bahas dan metode. Hasil dan pembahasan, serta saran dan rekomendasi. Bahas dan metode. Hasil dan pembahasan, serta saran dan rekomendasi.
b.	Ruang lingkup dan kedalaman pembahasan: Ruang lingkup sesuai dengan bidang ilmu pengusul dengan pembahasan yang luas dan dalam dan dengan dibuktikan dengan penelitian-penelitian terdahulu.
c.	Kecukupan dan kemutahiran data/informasi dan metodologi: Penelitian eksperimental laboratorium pada hewan coba, data dan hasil dari uji primerik dan histopatologis di perkuat, dan gambar perlatuan.
d.	Kelengkapan unsur dan berkualitas terbitan/ jurnal : Jurnal Nasional Terakreditasi S3

Semarang, 25 Agustus 2020
Reviewer 2

Prof. Dr. drg. Oedijani, M.S.
NIP 194902091979012001
Unit kerja: Fakultas Kedokteran Undip
Bidang ilmu : Ilmu Kedokteran
Jabatan/pangkat: Guru Besar



Home (<http://ijoems.com/index.php/ijems/index>) > Editorial Team

Editorial Team

Editor-In-Chief

Dr. rer. nat. Risfidian Mohadi, Sriwijaya University, Indonesia (Scopus ID = 55991505100) (<https://www.scopus.com/authid/detail.uri?authorId=55991505100>)

Managing Editors

Dr. Laila Hanum, Sriwijaya University, Indonesia (Scopus ID = 57194601647) (<https://www.scopus.com/authid/detail.uri?authorId=57194601647>)

Dr. Tarmizi Taher, Sriwijaya University, Indonesia (Scopus ID = 56104271500) (<https://www.scopus.com/authid/detail.uri?authorId=56104271500>)

Advisory International Editorial Boards

Dr. Ng Law Yong, Universiti Tunku Abdul Rahman, Malaysia (Scopus ID = 36680687200) (<https://www.scopus.com/authid/detail.uri?authorId=36680687200>)

Prof. Aldes Lesbani, Universitas Sriwijaya, Indonesia (Scopus ID = 15056199800) (<https://www.scopus.com/authid/detail.uri?authorId=15056199800>)

Dr. Roy Andreas, Universitas Jenderal Soedirman, Indonesia (Scopus ID = 56009301800) (<https://www.scopus.com/authid/detail.uri?authorId=56009301800>)

Dr. Akil Ahmad, University of KwaZulu- Natal, Durban-4041, South Africa (Scopus ID = 36022073300) (<https://www.scopus.com/authid/detail.uri?authorId=36022073300>)

Dr. Muneer Mohammed Awadh Ba-Abbad, Hadharamout University of Science and Technology, Yemen (Scopus ID = 39761137100) (<https://www.scopus.com/authid/detail.uri?authorId=39761137100>)

Dr. Koo Chai Hoon, Universiti Tunku Abdul Rahman, Department of Civil Engineering, Kajang, Malaysia (Scopus ID = 36721525200) (<https://www.scopus.com/authid/detail.uri?authorId=36721525200>)

Dr. M. Lutfi Firdaus, Universitas Bengkulu, Indonesia (Scopus ID = 56426642700) (<https://www.scopus.com/authid/detail.uri?authorId=56426642700>)

Prof. Hilda Zulkifli, Sriwijaya University, Indonesia (Scopus ID = 55234932600) (<https://www.scopus.com/authid/detail.uri?authorId=55234932600>)

Dr. Muhammad Said, Sriwijaya University, Indonesia (Scopus ID = 57189487421) (<https://www.scopus.com/authid/detail.uri?authorId=57189487421>)

Technical Editors

Neza Rahayu Palapa, Sriwijaya University, Indonesia (Scopus ID = 57204586233) (<https://www.scopus.com/authid/detail.uri?authorId=57204586233>)

JOURNAL MENU

Aim and Scope (<http://ijoems.com/index.php/ijems/ans>)

Publication Ethics (<http://ijoems.com/index.php/ijems/pe>)

Abstracting Indexing (<http://ijoems.com/index.php/ijems/indexing>)

Manuscript Template (<http://sciencetechindonesia.com/public/site/files/Article%20Template.docx>)

Copyright Notice (<http://ijoems.com/index.php/ijems/copyright>)

Home (<http://ijoems.com/index.php/ijems/index>) > Archives (<http://ijoems.com/index.php/ijems/issue/archive>)
> Vol. 4 No. 2 (2020): June



(<http://ijoems.com/index.php/ijems/issue/view/11>)

DOI: <https://doi.org/10.26554/ijems> (<https://doi.org/10.26554/ijems>)

Published: 2020-06-30

Articles

[Spirulina platensis Extract Reduces Serum TNF- \$\alpha\$, Neutrophils, and Increases Macrophage Count in Skin Incisional Mice Model](http://ijoems.com/index.php/ijems/article/view/115) (<http://ijoems.com/index.php/ijems/article/view/115>)

Riski Dwi Utami, Tri Nur Kristina, Renni Yuniati

DOI <https://doi.org/10.26554/ijems.2020.4.2.34-38> (<https://doi.org/10.26554/ijems.2020.4.2.34-38>)
34-38

Groups	Mean \pm SD	Min-Max	p Values
X1	15.50 \pm 2.429	12-19	p < 0.001
X2	19.17 \pm 2.317	16-22	
C1	6.33 \pm 1.862	4-9	
C2	11.17 \pm 1.329	9-13	

(<http://ijoems.com/index.php/ijems/article/view/115>)

 Abstract 36 times

PDF (<http://ijoems.com/index.php/ijems/article/view/115/72>)

Effectiveness Cream and Ointment of *Spirulina platensis* extract against amount of Fibroblas and Wound Area: Study on white rats whose skin is incised (<http://ijoems.com/index.php/ijems/article/view/116>)

Bimby Irenesia, Renni Yuniarti, Endang Mahati

DOI <https://doi.org/10.26554/ijems.2020.4.2.39-42> (<https://doi.org/10.26554/ijems.2020.4.2.39-42>)
39-42

Post-injury day	Group	Mean	SD	p-value
Day 14	Negative control	3.83	2.229	P<0.001
	Positive control	10.33	1.506	
	S.platensis extract 0.1% cream	15.00	2.757	
	S.platensis extract 0.1% ointment	17.50	1.378	

(<http://ijoems.com/index.php/ijems/article/view/116>)

👁 Abstract 33 times

PDF (<http://ijoems.com/index.php/ijems/article/view/116/68>)

Determinants Analysis of Public Perception of Waste Bank As An Alternative of Settlement Waste Management (<http://ijoems.com/index.php/ijems/article/view/107>)

Tri Hikmah Abdi Ningrum, Hilda Zulkifli, Rahmi Susanti

DOI <https://doi.org/10.26554/ijems.2020.4.2.43-47> (<https://doi.org/10.26554/ijems.2020.4.2.43-47>)

43-47

Education Level	Public Perception Good		Public Perception Bad		Total n	% n	p-value
	n	%	n	%			
High	43	54.4	36	45.6	79	100	0.423
Low	4	36.4	7	63.6	11	100	

(<http://ijoems.com/index.php/ijems/article/view/107>)

👁 Abstract 22 times


PDF (<http://ijoems.com/index.php/ijems/article/view/107/69>)

Alternative scenarios to utilise excess biogas in Palm Oil Mill (<http://ijoems.com/index.php/ijems/article/view/114>)

Novelita Wahyu Mondamina, Deni Rachmat, Mochammad Waris Tegar Laksono

DOI <https://doi.org/10.26554/ijems.2020.4.2.48-54> (<https://doi.org/10.26554/ijems.2020.4.2.48-54>)

48-54



(<http://ijoems.com/index.php/ijems/article/view/114>)

👁 Abstract 22 times

PDF (<http://ijoems.com/index.php/ijems/article/view/114/70>)

Effectiveness of *Spirulina platensis* Extract on Wound Area and TNF- α Levels on Blood: Experimental Studies In Wistar Rats Made Artificially by *Vulnus Scissum* and Infected by *Staphylococcus aureus* (<http://ijoems.com/index.php/ijems/article/view/117>)

Olvaria Misfa, Rennu Yuniarti, Yan Wisnu Prajoko

DOI <https://doi.org/10.26554/ijems.2020.4.2.55-58> (<https://doi.org/10.26554/ijems.2020.4.2.55-58>)

55-58

Effectiveness Cream and Ointment of Spirulina platensis extract against amount of Fibroblas and Wound Area: Study on white rats whose skin is incised

by Renni Yuniati

Submission date: 07-Aug-2020 06:58PM (UTC+0700)

Submission ID: 1366933233

File name: Artikel_Effectiveness_Cream.pdf (282.82K)

Word count: 2814

Character count: 14949

Effectiveness Cream and Ointment of *Spirulina platensis* extract against amount of Fibroblasts and Wound Area: Study on white rats whose skin is incised

Bimby Irenesia¹, Renni Yuniati^{2*}, Endang Mahati²

¹Department of Biomedicine, School of medicine, Medical Faculty of Diponegoro University, Semarang, Central Java, Indonesia

²Department of Dermatology-Venerology, Medical Faculty of Diponegoro University, Semarang, Central Java, Indonesia

²Department of Pharmacology, Medical Faculty of Diponegoro University, Semarang, Central Java, Indonesia

*Corresponding author e-mail: renniyuniati@yahoo.com

Abstract

Spirulina platensis has been well-known as an abundant protein and phycocyanin. *Spirulina platensis* has the effect of anti-inflammatory, so it is widely used to apply cosmetics and biomedicine in the healing process of wounds. To prove that the administration of an extract of *Spirulina platensis* can increase the number of fibroblasts and accelerate the wound healing process in Wistar rats with an incision on their skin. This study used a randomized post-test only control group design. The skins of thirty (30) male Wistar rats were incised. This treatment was divided into 4 groups, namely the group that received the 0,1% cream (X1) of *S. platensis* extract, and the group that was administered with the 0,1% ointment (X2) of *S. platensis* extract; the negative control group was only treated by a saline solution (C1), and the positive control group was with the administration of Gentamycin 0,1% ointment (C2). The total numbers of fibroblasts were examined on day 14 and calculated using the method of tissue biopsy and hematoxylin-eosin (HE) staining. The wound was checked after 14-day Data analysis was performed with a one-way ANOVA test and was continued by Post Hoc Test LSD. The total numbers of fibroblasts on day 14 from the group of X1, X2, C1, and C2 were 15,00; 17,50; 3,83, 10,33 with $p < 0.001$. The average size of the wounds on day 14 from the groups of X1, X2, C1, and C2 was 551,78; 435,76; 1795,95; 1193,57 with $p < 0.001$, respectively. Post Hoc Test results on the total number of fibroblasts and extensive wound area showed significant differences between them. Administration of the 0.1% ointment of *Spirulina platensis* extract believed to potentially increasing the number of fibroblasts and accelerate the process of wound healing.

Keywords

Spirulina platensis, fibroblasts, extensive wound area

Received: 3 June 2020, Accepted: 18 June 2020

<https://doi.org/10.26554/ijems.2020.4.2.39-42>

1. INTRODUCTION

A wound is a form of tissue damage on the skin caused by contact with a heat source (such as chemicals, water, heat, fire, radiation, and electricity), including the preventive actions or changes in physiological conditions (Fithriani et al., 2015). A wound causes disturbance in the function and anatomical structure of the body. Normal wound undergoes a series of events that involves a process of wound healing. (Syarina et al., 2015)

Wound healing is a dynamic process that involves complex interactions between cellular, molecular, biochemical, and physiological activity that produces the regeneration and the replacement of injured connective tissues at the wound site. Connective tissue cells, which are very important in the remodeling and healing of the damaged tissue, are fibroblasts. Wound healing can be helped in the form of

antiseptics, antibiotics, and wound care in general. While the treatments can be executed by giving drugs that are commercial or using alternative specialized materials that are founded in nature. (Syarina et al., 2015; Yu, 2017)

Natural products considered healthier and safer for the body. The World Health Organization estimated that 80% of the world population utilizes herbal medicine to maintain health. One of the commodities that have not yet been widely explored in Indonesia in terms of the development of herbal medicine is a microalga (Fithriani et al., 2015).

Algae has been a growing interest for the community as a drug and food, as it produces products that are harmless and nutritious to the health supplement market. One of the biodiversity examples of microalgae in Indonesia, which has various potency, is *Spirulina platensis*. (Fithriani et al., 2015; Syarina et al., 2015) *Spirulina platensis* is a blue-green

6
alga that has a spiral-shaped filament. (Fithriani et al., 2015; Syarina et al., 2015; Somchit et al., 2014; Yu, 2017). *Spirulina platensis* contains protein, chlorophyll, carotenes, and phycobilin (phycocyanin, allophycocyanin, and phycoerythrin). (Somchit et al., 2014; Konicková et al., 2014; Ridlo et al., 2016; Jung et al., 2016; Gur et al., 2013)

Provision of packaged ointment, based on wound healing, is much quicker than the cream one – this is due to the fast absorption ability of the lotion. However, there has been no research data on the effectiveness of *Spirulina platensis* administered in the form of an ointment. Therefore, it is scientifically necessary to research the efficacy of the *Spirulina platensis* extract of lotion on the number of fibroblasts and extensive wound area on Wistar white rats.

2. EXPERIMENTAL SECTION

2.1 Extracts of *Spirulina platensis*

The *S. platensis* used in this study were obtained as *S. platensis* powder with US FDA registration number of 15594742028 and CERES number of 50OGA1200043 (9241). The powdered *S. platensis* microalgae was macerated in 95% ethanol solution with 1:10 concentration (one part of *S. platensis* powder macerated in 10 regions of 95% ethanol solution). The maceration process was done for five days in a glass container. The glass container was stirred every day to make sure the uniformity of the maceration process. After five days, the solution was filtered through Whatman, and was evaporated using a rotary evaporator machine at the ethanol boiling point temperature until a thick extract is obtained. This extract of *S. platensis* was used as the material in the subsequent tests, further explained below.

2.2 Experimental Animals

30 male Wistar rats aged 2-3 months were acclimatized for 7 days. After 7 days, the experimental animals were randomly divided into 4 groups and were performed an incision on the skin of their backs. The first treatment group was administered with the 0.1% cream of *Spirulina platensis* extracts. The second treatment group was treated with the 0.1% ointment of *Spirulina platensis* extracts for 14 days. Third, the negative control group was given only 0.9% saline solution. The fourth, the positive control group, was treated with Gentamycin 0.1% ointment.

2.3 Incision Procedure

All rats were anesthetized with 0.1 cc ketamine per individual, then the backs of the rats were shaved with an area of 3 cm x 2.5 cm. Furthermore, an incision was made in the regions cut using a scalpel for 2 cm-length and 0.25 cm-depth by stretching skins with the index finger, while thumbs of the left-hand acted as a relaxant and a suppressor. The scalpel was held using its handle with the right hand to form an angle of 300-400 to the skin. The incision was executed by pulling the scalpel towards the caudal. After that, the wound was closed adequately with gauze.

2.4 The numbers of fibroblasts

Observation of the histology was taken on day 14, which performed on the skin samples. Then, the calculation on the number of fibroblasts was observed in the histopathological examination with the skin preparations that have been stained with the HE staining. This calculation was by using a binocular microscope (Olympus CX-21, Japan) with an objective magnification of 40x. Imaging was performed using the video photo with three times of repetitions.

2.5 Extensive Wound Area

Extensive injuries areas were observed on day 14 preliminary, starting from the extracts' administration on the first day of treatment after the rats were incised. The size of the wounds was assessed by photographing them and then measuring them using Imageraster 3.

2.6 Statistical Analysis

The test used in this study was a One-Way ANOVA test to determine the difference in the number of fibroblasts and extensive wound area against all treatment groups. The magnitude of the difference in the name of fibroblasts and broad wound area on each group was necessary to perform further analysis using Post Hoc Test LSD. The significance value in this study was if the analysis variable has a value of $p < 0.05$. All statistical analysis was analyzed using the computer program SPSS 25.00.

3. RESULTS AND DISCUSSION

All used samples had similar average weights among the groups up to day 14. All treatment rats lived until the end of the study, then were terminated at the end.

Table 1. The numbers of fibroblasts

Post-injury day	Group	Mean	SD	p-value
Day 14	Negative control	3.83	2.229	P<0.001
	Positive control	10.33	1.506	
	<i>S. platensis</i> extract 0.1% cream	15.00	2.757	
	<i>S. platensis</i> extract 0.1% ointment	17.50	1.378	

The analysis showed significant difference in the number of fibroblasts among the treatment groups on the 14th-day (Table 1), which had been discovered between the negative control group and the groups of 0.1% cream and 0.1% ointment of *Spirulina platensis* extracts. Also, the number of fibroblasts was also significantly different between the groups of 0.1% cream and 0.1 % ointment of *Spirulina platensis* extracts and the positive control group.

The analysis exhibited a significant difference in wound size amongst the treatment groups on the 14th-day (Table

Table 2. 14th-day of extensive wound area

Group	Mean	SD	p-value
Negative control	1795,95	119,78	P < 0.001
Positive control	1193,57	375,69	
S.platensis extract 0.1% cream	5551,78	135,97	
S.platensis extract 0.1% ointment	435,76	92,23	

2), which had been disclosed between the negative control group and the groups of 0.1% cream and 0.1% ointment of *Spirulina platensis* extracts. In addition, the size of the cuts was also significantly different between the group of 0.1% cream and 0.1% ointment of *Spirulina platensis* extracts and the positive control group.

Fibroblasts have a very prominent role in the wound healing process, especially in the proliferative phase. They undergo proliferation process and collagen synthesis. The collagen fibers that are formed may lead to the existence of restraint reassembling the edges of the wound (Purnama and Ratnawulan, 2014; Palumpun et al., 2017; Sihotang and Yulianti, 2018).

In this study, it was revealed that the administration of the extract of *Spirulina platensis* might potentially increase the number of fibroblasts in mice that performed incision on their skins. The highest level of fibroblasts accordingly belonged to the group with the treatment of 0.1% ointment of *Spirulina platensis* extracts on day 14. The analysis showed a significant difference in the number of fibroblasts among the treatment groups on the 14th-day which had been discovered between the negative control group and the groups receiving 0.1% cream and 0.1% ointment of *Spirulina platensis* extracts. In addition, the number of fibroblasts was also significantly different between the group of 0.1% cream and 0.1% ointment of *Spirulina platensis* extracts and the positive control group. This result was in accordance with the studies done by Ibtisam Ahmed Baheue-Eldin et al, where *Spirulina platensis* significantly affects, in a positive way, the process of wound healing via formation of granulation tissues (the number of polymorphonuclear cells, lymphocytes, macrophages, and fibroblast) and enhances neovascularization in the area of the wounds.

Based on this research, the hypothesis was provenly admitted because the number of fibroblasts on the administration of the ointment extract of *Spirulina platensis* was much higher compared to the administration of *Spirulina platensis* extract cream. Wound healing is a dynamic process that involves complex interactions between cellular, molecular, biochemical, and physiological activity that produces the regeneration and replacement of injured connective tissues in the location of the wound (Gur et al., 2013; Pur-

nama and Ratnawulan, 2014). There are 3 phases of the wound healing process: the inflammatory phase, proliferation phase, and maturation (remodeling) phase. (Purnama and Ratnawulan, 2014; Palumpun et al., 2017; Sihotang and Yulianti, 2018) Based on the research conducted by Syarina et al. (2015) stated that the water extract of *Spirulina* plants demonstrated the highest wound healing activity and may be considered a potential source of therapeutic agents to chronic wounds and related complications.

This study suggested that the administration of *Spirulina platensis* extract may promote the wound closure of mice that performed incision on their skins. The wound healing rate of mice in the group with 0.1% ointment of *Spirulina platensis* extracts, on day 14, had a faster healing activity compared to the group with 0.1% cream of *Spirulina platensis* extracts and the groups of positive control (KP) and negative control (KN). The analysis also exhibited a significant difference in terms of the size of the wounds amongst the treatment groups on the 14th-day which had been disclosed between the negative control group and the groups of 0.1% cream and 0.1% ointment of *Spirulina platensis* extracts. In addition, the size of the cuts was correspondingly significantly different between the groups of 0.1% cream and 0.1% ointment of *Spirulina platensis* extracts and the positive control group. This indicated that the extract of *Spirulina platensis* has a significant effect of anti-inflammatory to the wound healing acceleration. Similar reports by Seda Gunes et al. observed that the cell viability, the activity of wound healing and genotoxicity results reveals that the skin cream that was mixed with the ingredients of *Spirulina platensis* extracts was believed to be potential value in terms of cosmetic and biomedical application.

Based on this finding, the hypothesis was reportedly proved that the wound area on the administration of the ointment extract of *Spirulina platensis* was smaller compared to the administration of the cream extracts of *Spirulina platensis*.

4. CONCLUSIONS

The number of fibroblasts in the group receiving 0.1% ointment of *Spirulina platensis* extracts on day 14 was much more abundant than the group of 0.1% cream of *Spirulina platensis* extracts. However, no significant difference was revealed between both treatment groups. Extensive wound area in the group with 0.1% ointment of *Spirulina platensis* extracts was smaller in comparison with the group of 0.1% cream of *Spirulina platensis* extracts, and no significant difference was proven between both groups of treatment.

REFERENCES

- Fithriani, D., S. Amini, S. Melanie, and R. Susilowati (2015). Uji Fitokimia, Kandungan Total Fenol Dan Aktivitas Antioksidan Mikroalga *Spirulina* Sp., *Chlorella* Sp., dan

- Nannochloropsis Sp. *Jurnal Pascapanen dan Bioteknologi Kelautan dan Perikanan*, **10**(2); 101–109
- Gur, C. S., D. K. Erdogan, I. Onbasilar, P. Atilla, N. Cakar, and I. D. Gurhan (2013). In vitro and in vivo investigations of the wound healing effect of crude Spirulina extract and C-phycoerythrin. *Journal of Medicinal Plants Research*, **7**(8); 425–433
- Jung, S.-M., S. K. Min, H. C. Lee, Y. S. Kwon, M. H. Jung, and H. S. Shin (2016). Spirulina-PCL nanofiber wound dressing to improve cutaneous wound healing by enhancing antioxidative mechanism. *Journal of Nanomaterials*, **2016**
- Konícková, R., K. Vanková, J. Vaníková, K. Vánová, L. Muchová, I. Subhanová, M. Zadinová, J. Zelenka, A. Dvůrák, M. Kolár, et al. (2014). Anti-cancer effects of blue-green alga Spirulina platensis, a natural source of bilirubin-like tetrapyrrolic compounds. *Annals of Hepatology*, **13**(2); 273–283
- Palumpun, E. F., A. A. Wiraguna, and W. Pangkahila (2017). Pemberian ekstrak daun sirih (Piper betle) secara topikal meningkatkan ketebalan epidermis, jumlah fibroblas, dan jumlah kolagen dalam proses penyembuhan luka pada tikus jantan galur Wistar (Rattus norvegicus). *eBiomedik*, **5**(1)
- Purnama, H. and S. Ratnawulan (2014). Review Sistematis: Proses Penyembuhan dan Perawatan Luka. *Jurnal Fakultas Farmasi. Bandung. Universitas Padjadjaran*
- Ridlo, A., S. Sedjati, and E. Supriyanti (2016). Aktivitas Anti Oksidan Fikosianin Dari Spirulina Sp. Menggunakan Metode Transfer Elektron Dengan DPPH (1, 1-difenil-2-pikrilhidrazil). *Jurnal Kelautan Tropis*, **18**(2)
- Sihotang, H. M. and H. Yulianti (2018). Faktor-Faktor Yang Mempengaruhi Proses Penyembuhan Luka Post Sectio Caesarea. *Care: Jurnal Ilmiah Ilmu Kesehatan*, **6**(2); 175–183
- Somchit, M. N., N. A. Mohamed, Z. Ahmad, Z. A. Zakaria, L. Shamsuddin, M. S. Omar-Fauzee, and A. A. Kadir (2014). Anti-inflammatory and anti-pyretic properties of Spirulina platensis and Spirulina lonar: a comparative study. *Pak J Pharm Sci*, **27**(5); 1277–80
- Syarina, P. N. A., G. Karthivashan, F. Abas, P. Arulselvan, and S. Fakurazi (2015). Wound healing potential of Spirulina platensis extracts on human dermal fibroblast cells. *EXCLI journal*, **14**; 385
- Yu, J. (2017). Application of an Ultrafine Shearing Method for the Extraction of C-Phycocyanin from Spirulina platensis. *Molecules*, **22**(11); 2023

Effectiveness Cream and Ointment of Spirulina platensis extract against amount of Fibroblas and Wound Area: Study on white rats whose skin is incised

ORIGINALITY REPORT

6%

SIMILARITY INDEX

1%

INTERNET SOURCES

4%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to University Tun Hussein Onn Malaysia

Student Paper

1%

2

I Purnamasari, M Yerizam, A Hasan, R Junaidi. "Oxygen Adsorption Kinetics Study used Pressure Swing Adsorber (PSA) for Nitrogen Production", Journal of Physics: Conference Series, 2019

Publication

1%

3

Sariya Mapoung, Punnida Arjsri, Pilaiporn Thippraphan, Warathit Semmarath et al. "Photochemoprotective effects of Spirulina platensis extract against UVB irradiated human skin fibroblasts", South African Journal of Botany, 2020

Publication

1%

4

Submitted to Curtin University of Technology

Student Paper

1%

5

"The 22nd Conference of the Asian Pacific Association for the Study of the Liver",
Hepatology International, 2012

Publication

1 %

6

Asma Fraj, Fadhel Jaâfar, Meritxell Marti, Luisa Coderch, Neji Ladhari. "A comparative study of oregano (*Origanum vulgare* L.) essential oil-based polycaprolactone nanocapsules/microspheres: Preparation, physicochemical characterization, and storage stability", Industrial Crops and Products, 2019

Publication

<1 %

7

Qinghua Wu, Lian Liu, Anca Miron, Blanka Klímová, Dan Wan, Kamil Kuča. "The antioxidant, immunomodulatory, and anti-inflammatory activities of *Spirulina*: an overview", Archives of Toxicology, 2016

Publication

<1 %

8

www.ijoems.com

Internet Source

<1 %

9

Roberto Aquilani, Alfredo Costa, Roberto Maestri, Matteo Cotta Ramusino et al. "Mini Nutritional Assessment May Identify a Dual Pattern of Perturbed Plasma Amino Acids in Patients with Alzheimer's Disease: A Window to Metabolic and Physical Rehabilitation?", Nutrients, 2020

<1 %

10

sipil.ft.unsri.ac.id

Internet Source

<1 %

11

Submitted to Universitas Diponegoro

Student Paper

<1 %

12

Minuljo, T. Tedjo, H. Nugroho, and T.G.D. Pemayun. "PO365 LIMB INFECTION IN PATIENT WITH UNCONTROLLED TYPE 2 DIABETES: CASE REPORT", Diabetes Research and Clinical Practice, 2014.

Publication

<1 %

13

Fouad Al-Bayaty, Mahmood Ameen Abdulla. "A Comparison of Wound Healing Rate Following Treatment with Aftamed and Chlorine Dioxide Gels in Streptozotocin-Induced Diabetic Rats", Evidence-Based Complementary and Alternative Medicine, 2012

Publication

<1 %

14

E. Dharmana. "HLA-DRB1*12 is associated with protection against complicated typhoid fever, independent of tumour necrosis factor alpha", European Journal of Immunogenetics, 8/2002

Publication

<1 %

Exclude bibliography On

Effectiveness Cream and Ointment of Spirulina platensis extract against amount of Fibroblas and Wound Area: Study on white rats whose skin is incised

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4