

KORESPONDENSI JURNAL

Judul Artikel : Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java

Nama Jurnal : KEMAS: Jurnal Kesehatan Masyarakat



Penulis : 1. Oktavia Beni Kujariningrum
2. **Sri Winarni* (Penulis Korespondensi)**
3. Atik Mawarni
4. Najib Najib

No	Kegiatan	Tanggal	Keterangan	Halaman
1	Submission File	11 Oktober 2021	OJS submission	2-4
2	Revisi 1	12 April 2022	OJS dan File Revisi	5-11
3	Pengiriman hasil perbaikan 1	17 April 2022	OJS	12
4	Revisi 2	31 Agustus 2022	OJS dan File Revisi	13-19
5	Pengiriman hasil revisi 2	31 Agustus 2022	OJS	20
6	Revisi 3	05 September 2022	OJS dan File Revisi	21-27
7	Pengiriman hasil revisi 2	27 September 2022	OJS	28
8	Accept Submission	07 November 2022	OJS	29
9	Tagihan biaya publikasi	09 November 2022	E-mail	30-31
10	Permintaan Surat Bebas Plagiat & Kesiediaan Membayar	10 November 2022	E-mail	32
11	Pengiriman bukti transfer biaya publikasi	14 November 2022	E-mail	33
12	Pengiriman Surat Bebas Plagiat & Kesiediaan Membayar	15 November 2022	E-mail	34
13	Pengiriman LoA	21 November 2022	E-mail dan LoA	35-36
14	Artikel Terbit	Januari 2023	Website KEMAS: Jurnal Kesehatan Masyarakat https://journal.unnes.ac.id/nju/index.php/kemas/article/view/32644	37

#32644 Summary

SUMMARY REVIEW EDITING

Submission

Authors	Oktavia Beni Kujariningrum, Sri Winarni, Atik Mawarni, Najib Najib
Title	Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java
Original file	32644-83644-2-SM.DOCX 2021-10-11
Supp. files	32644-83646-1-SP.PDF 2021-10-11 32644-83649-1-SP.PDF 2021-10-11
Submitter	Oktavia Beni Kujariningrum 
Date submitted	October 11, 2021 - 11:13 AM
Section	Articles
Editor	Efa Nugroho, S.K.M., M.Kes 
Abstract Views	585

Status

Status	Published Vol 18, No 3 (2023)
Initiated	2023-01-05
Last modified	2023-03-06

Submission Metadata

Authors

Name	Oktavia Beni Kujariningrum 
ORCID iD	https://orcid.org/0000-0003-0730-0420
Affiliation	Biostatistics and Population, Faculty of Public Health, Diponegoro University
Country	Indonesia
Competing interests	CI POLICY —
Bio Statement	—
Name	Sri Winarni 
ORCID iD	https://orcid.org/0000-0002-9436-1581
Affiliation	Biostatistics and Population, Faculty of Public Health, Diponegoro University
Country	Indonesia
Competing interests	CI POLICY —
Bio Statement	—
Principal contact for editorial correspondence.	
Name	Atik Mawarni 
ORCID iD	https://orcid.org/0000-0001-8272-0009
Affiliation	Biostatistics and Population, Faculty of Public Health, Diponegoro University
Country	Indonesia
Competing interests	CI POLICY —
Bio Statement	—
Name	Najib Najib 
Affiliation	National Innovation Research Agency (BRIN)
Country	—
Competing interests	CI POLICY —
Bio Statement	—

Title and Abstract

Title	Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java
Abstract	Abstract. In Central Java, the prevalence of LBW (Low Birth Weight) has increased from 4.3 (2018) to 4.7 (2019) and be the biggest cause of neonatal mortality (46.4%) and infant mortality (40.5%). The aim of this research is to analyze the relationship between quality of ANC (Antenatal Care), iron supplementation, pregnancy complications, and maternal smoking status with LBW in Central Java. This research was an analytical study used secondary data from the 2017 IDHS (Indonesian Demographic and Health Survey). The sampling design used purposive sampling. Population study was 1205 babies born in Central Java. The sample comprised 952 babies. Independent variables were the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status, with the incidence of LBW as dependent variable. Data analysis was performed by chi-square continuity correction and logistic regression. Pregnancy complications have been associated with the incidences of LBW in Central Java (p -value = 0.0001). Iron supplementation (OR = 2.474) and pregnancy complications (OR = 4.869) had an effect on the incidence of LBW in Central Java. Iron supplementation and pregnancy complications influenced the incidence of LBW in Central Java.

Indexing

Keywords	LBW; Pregnancy Complications; Iron Supplementation
----------	--

ABOUT THE JOURNAL

- Focus and Scope
- Manuscript Submission
- Guide for Authors
- Editorial Board
- Reviewer Team
- Abstracting/Indexing
- Ethics Statement
- Policy of Screening for Plagiarism
- Contact
- 2,143,151
- View Visitor Stats

USER

You are logged in as...
oktaviabeni66
 » My Journals
 » My Profile
 » Log Out

JOURNAL CONTENT

Search


Search Scope

Browse

- » By Issue
- » By Author
- » By Title
- » Other Journals

COLABORATION WITH


Ikatan Ahli Kesehatan Masyarakat Indonesia
 IAKMI (The Indonesian Public Health Association) is an independent professional organization for the benefit of public health, based on Pancasila based on the 1945 Constitution. Mu
 Agreement No:
 402/UN.37.1.6/IKM/2012


Jejaring Nasional Pendidikan Kesehatan (JNPK)
 JNPK is an organization that gathers experts and observers in the field of health education, which was established on September 1, 2014. The founder of this organization is the University of Teacher Training Education Institut (LPTK) which organizes public health education, namely Universitas Negeri

Supporting Agencies

Agencies United States Agency for International Development; Diponegoro University; The Health Research Ethics Committee

OpenAIRE Specific Metadata

ProjectID —

References

- References
- Ardelia, K.I.A., Hardianto, G., Nuswantoro, D., 2019. Passive smoker during pregnancy is a risk factor of low birth weight. *Maj. Obstet. Ginekol.* 27, 12. <https://doi.org/10.20473/mog.v1i12019.12-16>
- Bappenas, UNICEF, 2017. Laporan Baseline SDG tentang Anak-Anak di Indonesia. Jakarta.
- BKKBN, BPS, Kemenkes, International, I., 2017. Indonesia Demographic and Health Survey 2017. DHS Program, Jakarta, Indonesia.
- Chhabra, S., Chopra, S., 2016. Mid Pregnancy Fetal Growth Restriction and Maternal Anaemia a Prospective Study. *J. Nutr. Disord. Ther.* 06. <https://doi.org/10.4172/2161-0509.1000187>
- Darwis, A., Abdullah, A., Maidar, Adamy, A., Nurjannah, 2020. Hubungan Komponen Pelayanan Antenatal Care (10T) dengan Kejadian Bayi Berat Lahir Rendah di Indonesia (Analisa Data Sekunder SDKI 2017). *JUKEMA* 6, 13–19.
- Dinas Kesehatan Provinsi Jawa Tengah, 2019. Profil Kesehatan Provinsi Jawa Tengah Tahun 2019, Dinkes Jateng.
- Dirjen Kesmas Kemenkes, 2018. Laporan Kinerja Ditjen Kesehatan Masyarakat Tahun 2017.
- Fang, F., Luo, Z.C., Dejemli, A., Delvin, E., Zhang, J., 2015. Maternal smoking and metabolic health biomarkers in newborns. *PLoS One.* <https://doi.org/10.1371/journal.pone.0143660>
- Huang, Shih-hui, Weng, K., Huang, Shih-ming, Liou, H., 2017. The effects of maternal smoking exposure during pregnancy on postnatal outcomes : A cross sectional study. *J. Chinese Med. Assoc.* 1–7. <https://doi.org/10.1016/j.jcma.2017.01.007>
- Isnaini, N., 2015. Analisis Faktor Yang Mempengaruhi Kejadian Bayi Berat Lahir Rendah. *J. Dunia Kesmas* 4, 196–202.
- Karjatin, A., 2016. Keperawatan Maternitas. Pusdik SDM Kesehatan, Jakarta.
- Kataoka, M.C., Paula, A., Carvalheira, P., Ferrari, A.P., Antonieta, M., Leite, D.B., Maria, C., Lima, G. De, 2018. Smoking during pregnancy and harm reduction in birth weight : a cross-sectional study. *BMC Pregnancy Childbirth* 1–10.
- Kementerian Kesehatan RI Badan Penelitian dan Pengembangan, 2018. Hasil Utama Riset Kesehatan Dasar 2018, Kementrian Kesehatan Republik Indonesia. Jakarta, Indonesia.
- Ko, T.J., Tsai, L.Y., Chu, L.C., Yeh, S.J., Leung, C., Chen, C.Y., Chou, H.C., Tsao, P.N., Chen, P.C., Hsieh, W.S., 2014. Parental smoking during pregnancy and its association with low birth weight, small for gestational age, and preterm birth offspring: A birth cohort study. *Pediatr. Neonatol.* 55, 20–27. <https://doi.org/10.1016/j.pedneo.2013.05.005>
- Kujarinigrum, O.B., Winarni, S., Najib, N., 2021. Multiple pregnancy : the biggest risk factor of low birth weight in Central Java , Indonesia (2017 IDHS secondary data study). *Ann. Trop. Med. Public Heal.* 24, 24–184. <https://doi.org/10.36295/ASRO.2021.24184>
- Kusumawati, D.D., Septiyaningsih, R., Kania, 2016. Faktor-Faktor Ibu yang Mempengaruhi Kejadian Bayi Berat Lahir Rendah (BBLR). *J. Kesehat. Al-Irsyad* 9, 8–16.
- Long, H., Yi, J.M., Hu, P.L., Li, Z. Bin, Qiu, W.Y., Wang, F., Zhu, S., 2012. Benefits of Iron supplementation for low birth weight infants: A systematic review. *BMC Pediatr.* 12. <https://doi.org/10.1186/1471-2431-12-99>
- Meiriza, W., Aladin, A., Edison, E., 2018. The Correlation of Maternal Factors and The Quality of Antenatal Care Services With Low Birth Weight Babies In Health Facilities Level I. *J. Midwifery* 3, 103. <https://doi.org/10.25077/jom.1.1.103-114.2018>
- Miranti, Mutiarasari, D., Arsin, A.A., Hadju, V., Mallongi, A., Nur, R., Amri, I., Haruni, H., Wahyuni, R.D., Rahma, Faris, A., 2020. Determinants of the incidence of stunting in the working area of Kinovaro Sigi Health Center. *Enfermería Clínica* 30, 246–252. <https://doi.org/10.1016/j.enfcli.2019.10.077>
- Neiger, R., 2017. Long-Term Effects of Pregnancy Complications on Maternal Health: A Review. *J. Clin. Med.* 6, 76. <https://doi.org/10.3390/jcm6080076>
- OECD, WHO, 2019. Health at a glance: Asia/Pacific 2018: Measuring progress towards universal health coverage. OECD Publishing, Paris.
- Owa, K., 2019. Chronic Energy Deficiency , Anemia as a Risk Factor for Low Birth Weight Babies in East Nusa Tenggara Kekurangan Energi Kronis , Anemia sebagai Faktor Risiko Bayi Berat Lah. *J. Kesehat. Prim.* 4, 13–22.
- Owa, K., Putra, I.W.G.A.E., Windiani, I.G.A.T., 2017. Risk factors for low birth weight infants in East Nusa Tenggara. *Public Heal. Prev. Med. Arch.* 5, 39. <https://doi.org/10.15562/phpma.v5i1.40>
- Paul, P., Zaveri, A., Chouhan, P., 2019. Assessing the Impact of Antenatal Care Utilization on Low Birthweight in India: Analysis of the 2015–2016 National Family Health Survey. *Child. Youth Serv. Rev.* 106, 104459. <https://doi.org/10.1016/j.childyouth.2019.104459>
- Phowira, J., Elvina, F.T., Wiguna, I.I., Wahyudi, F.R.H.B., Medise, B.E., 2020. The Association between Tobacco Exposure during Pregnancy and Newborns' Birth Weight in DKI Jakarta Community Members. *medRxiv* 1–15. <https://doi.org/10.1101/2020.10.29.20222059>
- Purwanto, A.D., Wahyuni, C.U., 2016. Hubungan Antara Umur Kehamilan, Kehamilan Ganda, Hipertensi dan Anemia dengan Kejadian Bayi Berat Lahir Rendah (BBLR). *J. Berk. Epidemiol.* 4, 349–359. <https://doi.org/10.20473/jbe.v4i3>
- Restu, S., Dasuki, D., Nurdianti Z, R.D.S., 2014. The influence of iron supplementation in pregnant women to the occurrence of low birth weight (LBW) babies in Palu, Central Sulawesi. *J. thee Med. Sci. (Berkala Ilmu Kedokteran)* 46, 41–51. <https://doi.org/10.19106/jmedscie004601201406>
- Sarah, S., Irianto, I., 2018. Pengaruh Tingkat Kepatuhan Minum Tablet Fe Terhadap Kejadian Anemia Pada Ibu Hamil Trimester III di Puskesmas Pejeruk Tahun 2017. *Yars. Med. J.* 26, 075. <https://doi.org/10.33476/jky.v26i2.392>
- Setiati A R Rahayu S 2017 Faktor Yang Mempengaruhi Kejadian RRI R (Berat Badan Lahir Rendah) Di Ruang Perawatan



LINK

Universitas Negeri Semarang
Pengembang Jurnal
Faculty of Sport Science



KEYWORDS

Adolescent Attitude COVID-19
Covid-19 Diarrhea HIV/AIDS
Hospital Hypertension Indonesia
Jurnal Kesehatan Masyarakat
Knowledge Management
Motivation Nutrition Stress Stunting
Toddler Tuberculosis Waste morta
stunting

Shahar, R.H., Rahmawati, D., 2017. Faktor yang mempengaruhi kejadian asfiksia pada bayi baru lahir. *Intensif Neonatus RSUD DR Moewardi Di Surakarta*. J. Keperawatan Glob. 2, 9–20. <https://doi.org/10.37341/jkg.v2i1.27>

Sharami, S.H., Darkhaneh, R.F., Zahiri, Z., Milani, F., Asgharnia, M., Shakiba, M., Didar, Z., 2013. The relationship between vaginal bleeding in the first and second trimester of pregnancy and preterm labor. *Int. J. Reprod. Biomed.* 11, 385–390.

Siramaneeerat, I., Agusbyana, F., Meebunmak, Y., 2018. Maternal Risk Factors Associated with Low Birth Weight in 376–383. <https://doi.org/10.2174/1874944501811010376>

Sulistiyani, S., Darmawati, A.T., Setiani, O., 2019. Correlation between local-specific traditions of women agricultural workers and the Incidence of low birth weight in South Metro Sub-District, Metro City, Lampung Province, Indonesia. *Ann. Trop. Med. Public Heal.* 22, 1–5. <https://doi.org/10.36295/ASRO.2019.221118>

Tyastuti, S., Wahyuningsih, H.P., 2016. *Asuhan Kebidanan Kehamilan, Pertama*, ed. Pusdik SDM Kesehatan, Badan Pengembangan dan Pemberdayaan Sumber Daya Kesehatan, Kementerian Kesehatan RI. BPPSDMK, Jakarta, Indonesia. <https://doi.org/10.1017/CBO9781107415324.004>

U.S. Department of Health and Human Services, 2014. *The Health Consequences of Smoking: 50 Years of Progress A Report of the Surgeon General*, A Report of the Surgeon General.



Upadhyay, R.P., Naik, G., Choudhary, T.S., Chowdhury, R., Taneja, S., Bhandari, N., Martinez, J.C., Bahl, R., Bhan, M.K., 2019. Cognitive and Motor Outcomes in Children Born Low Birth Weight: A Systematic Review and Meta-analysis of Studies from South Asia. *BMC Pediatr.* 19, 1–15.

WHO, UNICEF, 2019. *Low Birthweight Estimates: Level and Trends 2000-2015*, UNICEF and WHO.

#32644 Review

SUMMARY **REVIEW** EDITING

Submission



Authors Oktavia Beni Kujariningrum, Sri Winarni, Atik Mawarni, Najib Najib 
 Title Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java
 Section Articles
 Editor Efa Nugroho, S.K.M, M.Kes 

Peer Review

Round 1

Review Version 32644-83650-1-RV.DOCX 2021-10-11
 Initiated —
 Last modified —
 Uploaded file None

Editor Decision

Decision Accept Submission 2022-11-07
 Notify Editor  Editor/Author Email Record  2022-08-21
 Editor Version 32644-91924-1-ED.DOCX 2022-04-12
 32644-91924-2-ED.DOCX 2022-08-31
 32644-91924-3-ED.DOCX 2022-09-05
 Author Version 32644-91956-1-ED.DOCX 2022-04-17 DELETE
 32644-91956-2-ED.DOCX 2022-08-31 DELETE
 32644-91956-3-ED.DOCX 2022-09-27 DELETE
 Upload Author Version No file chosen

ISSN: 2355-3596


ABOUT THE JOURNAL

Focus and Scope
 Manuscript Submission
 Guide for Authors
 Editorial Board
 Reviewer Team
 Abstracting/Indexing
 Ethics Statement
 Policy of Screening for Plagiarism
 Contact
 2,143,152
[View Visitor Stats](#)

USER

You are logged in as...
oktaviabeni66
 » [My Journals](#)
 » [My Profile](#)
 » [Log Out](#)

JOURNAL CONTENT

Search
 Search Scope
 All 

Browse

» [By Issue](#)
 » [By Author](#)
 » [By Title](#)
 » [Other Journals](#)

COLABORATION WITH



Ikatan Ahli Kesehatan Masyarakat Indonesia

IAKMI (The Indonesian Public Health Association) is an independent professional organization for the betterment of public health, based on Pancasila based on the 1945 Constitution. Mutual Agreement No:
 402/UN.37.1.6/IKM/2012



Jejaring Nasional Pendidikan Kesehatan (JNPK)

JNPK is an organization that gathers experts and observers in the field of health education, which was established on September 1, 2014. The founder of this organization is the University of Teacher Training Education Institut Teknologi Sepuluh Nopember (ITS) which organizes public health education, namely Universitas Negeri

Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java

Oktavia Beni Kujariningrum^{1, a)}, Sri Winarni^{1, b)}, Atik Mawarni^{1, c)}

¹*Biostatistics and Population, Faculty of Public Health, Diponegoro University, Building C, 1st Floor, Main Campus of UNDIP 50275, Indonesia*

^{a)}Corresponding author: oktaviabeni66@gmail.com

^{b)}winarniwiwin1975@gmail.com

^{c)}atikm246@gmail.com

Abstract. In Central Java, the prevalence of LBW (Low Birth Weight) has increased from 4.3 (2018) to 4.7 (2019) and be the biggest cause of neonatal mortality (46.4%) and infant mortality (40.5%). The aim of this research is to analyze the relationship between quality of ANC (Antenatal Care), iron supplementation, pregnancy complications, and maternal smoking status with LBW in Central Java. This research was an analytical study used secondary data from the 2017 IDHS (Indonesian Demographic and Health Survey). The sampling design used purposive sampling. Population study was 1205 babies born in Central Java. The sample comprised 952 babies. Independent variables were the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status, with the incidence of LBW as dependent variable. Data analysis was performed by chi-square continuity correction and logistic regression. Pregnancy complications have been associated with the incidences of LBW in Central Java (p -value = 0.0001). Iron supplementation (OR = 2.474) and pregnancy complications (OR = 4.869) had an effect on the incidence of LBW in Central Java. Iron supplementation and pregnancy complications influenced the incidence of LBW in Central Java.

Keyword: LBW, Pregnancy Complications, Iron Supplementation

INTRODUCTION

In 2015, 14.6% the incidences of LBW were found in the world and the highest prevalence occurring in Asia (17.3%) (WHO and UNICEF, 2019). The incidence of LBW in developing countries in the Asia Pacific region experienced an increase of ≤ 2 babies per 100 live births in 2014 compared to 2000 and an increase of 0.8% occurred in Indonesia (OECD and WHO, 2019). The results of the 2018 Riskesdas (Basic Health Research) showed that 6.2% of babies born with LBW in Indonesia and 6.1% were found in Central Java (Kementerian Kesehatan RI Badan Penelitian dan Pengembangan, 2018). The Central Java Health Profile in 2019 showed an increase in the incidence of LBW from 4.3 (2018) to 4.7 (2019) (Dinas Kesehatan Provinsi Jawa Tengah, 2019). A total of 5.9% of LBW events were found in single (Kujariningrum et al., 2021).

More than 37% of toddler were stunted in 2013 (Bappenas and UNICEF, 2017). That condition had relationship with a history of LBW (p -value = 0.037; OR = 5.294) (Miranti et al., 2020). A study in South Asia showed that 17% of children with a history of LBW had a total IQ of less than 85 (Upadhyay et al., 2019). In Central Java, LBW is the biggest cause of neonatal mortality (46.4%) and infant mortality (40.5%) in 2019 (Dinas Kesehatan Provinsi Jawa Tengah, 2019).

Visits and completeness of ANC services affect the incidence of LBW (Darwis et al., 2020; Paul et al., 2019). The incidence of LBW was also related to maternal anemia status (OR = 4.03) (Purwanto and Wahyuni, 2016). As much as 50% of the 73.3% pregnant women at Pejerkuk Health Center had adherence to consuming low Fe tablets (Sarah and Irianto, 2018). The incidence of LBW in Cilacap Regional Hospital was associated with pregnancy complications (OR = 3.393) (Kusumawati et al., 2016). The entry of nicotine into the body's mechanism of pregnant

Comment [EN1]: Sorry for the inconvenience. Our journal system has just completed an overhaul and some of the article metadata is missing and disrupts the review process.

In general, this article is good and deserves to be published. There are only minor revision in the introduction, results and discussion sections.

women has an impact on fetal growth and development (U.S. Department of Health and Human Services, 2014). The incidence of LBW in Taiwan was associated with maternal smoking status (OR = 3.46)(Ko et al., 2014).

Based on the facts described above, this study aims to analyze the relationship between the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status with the incidence of LBW in Central Java.

Comment [EN2]: Before explaining the objective, add the novelty of the research. whether this is new research, continuation, or comparing with research that has been running before.

METHOD

This study used a cross-sectional design. Research data from the 2017 IDHS. The study population consisted of 1205 babies born to WUS (Women of Childbearing Age) in 2012-2017 in Central Java. The research sample was taken used a purposive sampling based on inclusion and exclusion criteria for the total population so that a sample of 952 babies was obtained. The inclusion criteria included babies who were weighed in birth, babies who were last born by respondents in the range of 2012-2017, single births, and received ANC while in the womb. Infants with mothers who did not know the number of ANC visits, history of blood draws, history of consultations, and history of receiving Fe tablets were excluded from the study sample list as exclusion criteria.

The research variables consisted of the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status as independent variables with the incidence of LBW as the dependent variable. This study used the weighting of the sample according to the ethics of using the 2017 IDHS raw data and referring to a tutorial published by the you tube account The DHS Program on August 25, 2015 entitled " Part IV: Demonstration of How to Weight DHS Data in SPSS & SAS ". The bivariate analysis used was a statistical analysis of chi-square continuity correction. The independent variables were included in the multivariate logistic regression method enter are the independent variables with *p-value* ≤ 0.25. The analysis was carried out using a significance level of 0.05 and a confidence level of 95%. The likelihood of LBW occurrences can be seen from the calculation of the predicted value using the following formula:

$$f(Z) = \frac{1}{1 + e^{-(b_0 + b_1x_1 + b_2x_2 + \dots + b_ix_i)}}$$

Formula description:

$f(Z)$ = probability

b_0 = constant

b_1, b_2, \dots, b_i = partial regression coefficient

x_1, x_2, \dots, x_i = independent variable

e = exponent function with constant value 2.72

This study has passed the ethical clearance number: 12 / EA / KEPK-FKM / 2020 issued by the Health Research Ethics Committee, Faculty of Public Health, Diponegoro University on January 26, 2021.

RESULTS

Comment [EN3]: results and discussion are combined into one section

TABLE 1. Correlation of Quality of Antenatal Care, Iron Supplementation, Pregnancy Complications, and Mother's Smoking Status with LBW incidence (N=952)

Independent Variable	LBW				Total		<i>p-value</i>
	Not		Yes		f	%	
	n	%	n	%			
1. Quality of <i>Antenatal Care</i>							0.460
Good	278	94.2	17	5.8	295	100	
Poor	624	95.1	33	4.9	657	100	
2. Iron Supplementation							0.059
Yes	854	95.1	45	4.9	899	100	
Not	48	90.6	5	9.4	53	100	
3. Pregnancy Complications							0.0001
Not	749	96.8	26	3.2	775	100	
Yes	153	86.5	24	13.5	177	100	
4. Mother's Smoking Status							0.712
Not	893	94.8	50	5.2	943	100	
Yes	9	100	0	0	9	100	

Most (94.8%) babies born in Central Java in 2012-2017 had a NBW (Normal Birth Weight). As many as 68.9 % of babies were born to mothers who received poor quality of ANC. Most (94.3%) babies were born to mothers who received iron supplementation during pregnancy. As many as 81.3% of babies were born to mother who did not experience complications during pregnancy. Most (99.1%) babies were born to mother who had never smoked.

Based on chi-square continuity correction, pregnancy complications were related to the incidences of LBW in Central Java (p -value = 0.0001). There were no relationship between the quality of ANC, iron supplementation, and maternal smoking status with the incidence of LBW in Central Java (Table 1).

Multivariate analysis used logistic regression with enter method gave result that iron supplementation and pregnancy complications affect the incidence of LBW in Central Java. Mothers who did not receive iron supplementation during their pregnancy had 2.474 times higher risk of giving birth to LBW babies than mothers who received iron supplementation (OR = 2.474). Mothers who experienced pregnancy complications had a 4.869 times higher risk of giving birth to LBW babies than mothers who did not experience complications (OR = 4.869) (Table 2).

TABLE 2. Results of Multivariate Analysis The Effect of Iron Supplementation and Pregnancy Complications on the Incidence of LBW in Central Java 2012-2017 (N=952)

Variables		B	SE	Wald	Sig	Exp (B)
Model 1	Iron Supplementation	0.906	0.366	6.125	0.013	2.474
	Pregnancy Complications	1.583	0.218	52.670	0.0001	4.869
	Constant	-3.484	0.155	506.665	0.0001	0.031

Based on the calculation above, $f(Z) = 0.27$ can be conclude that mothers who didn't receive iron supplementation and had pregnancy complications history have a chance 27% to give the incidence of LBW.

DISCUSSION

The quality of ANC is a risk factor for LBW incidence (Owa et al., 2017). Most of the mothers access ANC services with low quality and this condition is related to the incidence of LBW in Indonesia (Darwis et al., 2020). In contrast to the conditions found in Central Java. Chi-square result showed that there was no relationship between the quality of ANC and the incidences of LBW in Central Java (p -value = 0.488). This is in line with Meiriza (2018), which concluded that there was no relationship between the quality of ANC at level I health facilities and the incidence of LBW in Padang City (Meiriza et al., 2018). This study found as many as 68.9 % of infants born to mothers who received ANC with poor quality. The incidence of LBW was more found in the group of infants with mothers who received ANC with good quality (5.8%) compared to poor quality (4.9). This showed that pregnant women who receive good quality antenatal care can also deliver babies with LBW. This condition is possible because of the limited variables found in the secondary data of the 2017 IDHS and the information needed to assess the quality of ANC is not enough just by questionnaires result, but requires in-depth interviews. Owa (2019) conducted in-depth interviews and found that pregnant women who received less ANC quality had a 3.5 times higher risk for having a baby with LBW (OR = 3.5) (Owa, 2019).

Iron supplementation is an effort made to respond the high rates of iron deficiency anemia in pregnant women (Tyastuti and Wahyuningsih, 2016). Chi-square result showed that there was no relationship between iron supplementation and the incidences of LBW (p -value = 0.076). Different from the multivariate analysis result which showed the effect of iron supplementation on the incidence of LBW (p -value = 0.013). Mothers who didn't receive iron supplementation during their pregnancy had a 2.474 times higher risk for having a baby with LBW than mothers who received iron supplementation (OR = 2.474). In line with Restu et al (2014) which found that iron supplementation had an effects on the incidences of LBW (OR = 3.82) (Restu et al., 2014). This shows that mothers who get iron supplementation can avoid anemia. Iron deficiency anemia causes an increase in serum norepinephrine concentration which results in maternal and fetal stress, stimulates the synthesis of CRH (Corticotrophin-Releasing Hormone) which will increase fetal cortisol productions thus impacting IUGR (Intrauterine Growth Restriction) and resulted in LBW (Chhabra and Chopra, 2016). Iron supplementation can reduce the prevalence of iron deficiency anemia (Long et al., 2012). Each tablet contains FeSO_4 mg (iron 30 mg) which is useful as a reserve of iron, red blood cell synthesis and muscle blood synthesis during pregnancy (Tyastuti and Wahyuningsih, 2016).

Pregnancy complications are collection of symptoms of health problems during pregnancy that can affect the health conditions of the baby and mother (BKKBN et al., 2017; Neiger, 2017). Some symptoms of pregnancy complications such as prolonged nausea and vomiting, hypotension and hypertension are associated with a lack of nutritional intake to the fetus (Dirjen Kesmas Kemenkes, 2018; Karjatin, 2016). The results of this study found that the incidences of LBW was more in the group of infants with mothers who had complications during their pregnancy (13.5%) than in the group of infants with mothers who had no complications during their pregnancy (3.2%). Chi-square result showed that there was a significant relationship between pregnancy complications and the incidence of LBW (p -value = 0.0001). Mothers who experienced pregnancy complications had a 4,869 times higher risk for having a baby with LBW than mothers who did not experience pregnancy complications (OR = 4.869). Siramaneerat (2018) mentions the same finding that pregnancy complications are associated with the incidences of LBW (p -value = 0.0001; OR = 1.731) (Siramaneerat et al., 2018). The most common complication found in this study was bleeding (5.5%). Antepartum hemorrhage associated with the incidences of LBW (Setiati and Rahayu, 2017). Antepartum hemorrhage can increase the likelihood of preterm birth which is a risk of LBW (Sharami et al., 2013). Pregnant women who experience antepartum bleeding have a 2.23 times higher risk of giving birth to LBW babies than mothers who don't experience antepartum hemorrhage (Isnaini, 2015). There was a difference between birth weight in the group of mothers with and without antepartum hemorrhage. Antepartum hemorrhage causes blood flow that distributes oxygen and nutrients to the placenta from the mother to the fetus to be disturbed. Impaired delivery of oxygen and nutrients will cause fetal anemia, shock to fetal death. Fetuses that survive until birth will experience various disorders, including LBW (Setiati and Rahayu, 2017).

Receptors in the placental blood vessels that mix with nicotine cause a decrease in blood flow in the placenta and fetal vasoconstriction which results in impaired delivery of oxygen and nutrients to the fetus so that the fetus experiences malnutrition which results in impaired fetal growth (U.S. Department of Health and Human Services, 2014). The entry of nicotine in the body's mechanism is proven to have an impact on the low production of the hormone Insulin-Like Growth Factor-1 in pregnant women who smoke will affect fetal growth and development (Fang et al., 2015). Pregnant women that smoke will give birth to babies with birth weight 320-435 grams lower than pregnant women who do not smoke (Kataoka et al., 2018). Outcome of this study shows that the incidences of LBW is more common in the group of infants whose mothers didn't smoke (5.2%), while NBW more common in the group of infants with mothers who smoked (100%). Chi-square result showed that there was no relationship between the mother's smoking status and the incidences of LBW (p -value = 0.706). In line with Phowira et al (2020) which stated that the mother's smoking status was not related to the incidence of LBW in DKI Jakarta (p -value = 0.448) (Phowira et al., 2020). There was no relationship between the frequency of smoking per day and the incidences of LBW in Lampung Province (Sulistiyani et al., 2019). It can be said that pregnant women who do not smoke can give birth to babies with LBW. This mechanism shows that the impact of nicotine entering the body cannot be described only by the status of a pregnant woman as an active smoker or not, as information is available in the 2017 IDHS data. LBW was found in mothers with high levels of nicotine > 143 µg/g keratinize. This condition is not only found in pregnant women who smoke actively. Pregnant women who don't smoke have an average nicotine level of 153.2 ± 96.0 µg/g keratinize (>143 µg/g keratinize) as a result of exposure to cigarette smoke from the environment (passive smoking) (Huang et al., 2017). There was a relationship between passive smoking mothers and the incidences of LBW (OR = 3.04) (Ardelia et al., 2019).

CONCLUSION

Most babies are born at a normal weight. There was a relationship between pregnancy complications and the incidences of LBW in Central Java. The quality of antenatal care, iron supplementation, and maternal smoking status were not related with the incidences of LBW. Pregnant woman who don't get iron supplementation (OR = 2.474) and complications of pregnancy (OR = 4.869) had chances of 0.84 for having a baby with LBW. The pregnant women who experience signs of complications immediately take action and visit health facilities for further assistance. Pregnant women are also advised to take Fe tablets regularly at least 90 tablets during their pregnancy.

ACKNOWLEDGMENTS

Thank you to the United States Agency for International Development for providing access to secondary data for the 2017 IDHS through the dhsprogram.com website, Diponegoro University for facilitating us in accessing various

e-journal portals, and the Health Research Ethics Committee, Faculty of Public Health, Diponegoro University. has issued a permit for the ethical feasibility of this research.

REFERENCES



- Ardelia, K.I.A., Hardianto, G., Nuswantoro, D., 2019. Passive smoker during pregnancy is a risk factor of low birth weight. *Maj. Obstet. Ginekol.* 27, 12. <https://doi.org/10.20473/mog.v1i12019.12-16>
- Bappenas, UNICEF, 2017. Laporan Baseline SDG tentang Anak-Anak di Indonesia. Jakarta.
- BKKBN, BPS, Kemenkes, International, I., 2017. Indonesia Demographic and Health Survey 2017. DHS Program, Jakarta, Indonesia.
- Chhabra, S., Chopra, S., 2016. Mid Pregnancy Fetal Growth Restriction and Maternal Anaemia a Prospective Study. *J. Nutr. Disord. Ther.* 06. <https://doi.org/10.4172/2161-0509.1000187>
- Darwis, A., Abdullah, A., Maidar, Adamy, A., Nurjannah, 2020. Hubungan Komponen Pelayanan Antenatal Care (10T) dengan Kejadian Bayi Berat Lahir Rendah di Indonesia (Analisa Data Sekunder SDKI 2017). *JUKEMA* 6, 13–19.
- Dinas Kesehatan Provinsi Jawa Tengah, 2019. Profil Kesehatan Provinsi Jawa Tengah Tahun 2019, Dinkes Jateng.
- Dirjen Kesmas Kemenkes, 2018. Laporan Kinerja Ditjen Kesehatan Masyarakat Tahun 2017.
- Fang, F., Luo, Z.C., Dejemli, A., Delvin, E., Zhang, J., 2015. Maternal smoking and metabolic health biomarkers in newborns. *PLoS One*. <https://doi.org/10.1371/journal.pone.0143660>
- Huang, Shih-hui, Weng, K., Huang, Shih-ming, Liou, H., 2017. The effects of maternal smoking exposure during pregnancy on postnatal outcomes : A cross sectional study. *J. Chinese Med. Assoc.* 1–7. <https://doi.org/10.1016/j.jcma.2017.01.007>
- Isnaini, N., 2015. Analisis Faktor Yang Mempengaruhi Kejadian Bayi Berat Lahir Rendah. *J. Dunia Kesmas* 4, 196–202.
- Karjatin, A., 2016. Keperawatan Maternitas. Pusdik SDM Kesehatan, Jakarta.
- Kataoka, M.C., Paula, A., Carvalheira, P., Ferrari, A.P., Antonieta, M., Leite, D.B., Maria, C., Lima, G. De, 2018. Smoking during pregnancy and harm reduction in birth weight : a cross-sectional study. *BMC Pregnancy Childbirth* 1–10.
- Kementerian Kesehatan RI Badan Penelitian dan Pengembangan, 2018. Hasil Utama Riset Kesehatan Dasar 2018, Kementrian Kesehatan Republik Indonesia. Jakarta, Indonesia.
- Ko, T.J., Tsai, L.Y., Chu, L.C., Yeh, S.J., Leung, C., Chen, C.Y., Chou, H.C., Tsao, P.N., Chen, P.C., Hsieh, W.S., 2014. Parental smoking during pregnancy and its association with low birth weight, small for gestational age, and preterm birth offspring: A birth cohort study. *Pediatr. Neonatol.* 55, 20–27. <https://doi.org/10.1016/j.pedneo.2013.05.005>
- Kujariningrum, O.B., Winarni, S., Najib, N., 2021. Multiple pregnancy : the biggest risk factor of low birth weight in Central Java , Indonesia (2017 IDHS secondary data study). *Ann. Trop. Med. Public Heal.* 24, 24–184. <https://doi.org/10.36295/ASRO.2021.24184>
- Kusumawati, D.D., Septiyaningsih, R., Kania, 2016. Faktor-Faktor Ibu yang Mempengaruhi Kejadian Bayi Berat Lahir Rendah (BBLR). *J. Kesehat. Al-Irsyad* 9, 8–16.
- Long, H., Yi, J.M., Hu, P.L., Li, Z. Bin, Qiu, W.Y., Wang, F., Zhu, S., 2012. Benefits of Iron supplementation for low birth weight infants: A systematic review. *BMC Pediatr.* 12. <https://doi.org/10.1186/1471-2431-12-99>
- Meiriza, W., Aladin, A., Edison, E., 2018. The Correlation of Maternal Factors and The Quality of Antenatal Care Services With Low Birth Weight Babies In Health Facilities Level I. *J. Midwifery* 3, 103. <https://doi.org/10.25077/jom.1.1.103-114.2018>
- Miranti, Mutiarasari, D., Arsin, A.A., Hadju, V., Mallongi, A., Nur, R., Amri, I., Haruni, H., Wahyuni, R.D., Rahma, Faris, A., 2020. Determinants of the incidence of stunting in the working area of Kinovaro Sigi Health Center. *Enfermeria Clínica* 30, 246–252. <https://doi.org/10.1016/j.enfcli.2019.10.077>
- Neiger, R., 2017. Long-Term Effects of Pregnancy Complications on Maternal Health: A Review. *J. Clin. Med.* 6, 76. <https://doi.org/10.3390/jcm6080076>
- OECD, WHO, 2019. Health at a glance: Asia/Pacific 2018: Measuring progress towards universal health coverage. OECD Publishing, Paris.
- Owa, K., 2019. Chronic Energy Deficiency , Anemia as a Risk Factor for Low Birth Weight Babies in East Nusa Tenggara Kekurangan Energi Kronis , Anemia sebagai Faktor Risiko Bayi Berat Lah. *J. Kesehat. Prim.* 4, 13–22.

- Owa, K., Putra, I.W.G.A.E., Windiani, I.G.A.T., 2017. Risk factors for low birth weight infants in East Nusa Tenggara. *Public Heal. Prev. Med. Arch.* 5, 39. <https://doi.org/10.15562/phpma.v5i1.40>
- Paul, P., Zaveri, A., Chouhan, P., 2019. Assessing the Impact of Antenatal Care Utilization on Low Birthweight in India: Analysis of the 2015–2016 National Family Health Survey. *Child. Youth Serv. Rev.* 106, 104459. <https://doi.org/10.1016/j.childyouth.2019.104459>
- Phowira, J., Elvina, F.T., Wiguna, I.I., Wahyudi, F.R.H.B., Medise, B.E., 2020. The Association between Tobacco Exposure during Pregnancy and Newborns' Birth Weight in DKI Jakarta Community Members. *medRxiv* 1–15. <https://doi.org/10.1101/2020.10.29.20222059>
- Purwanto, A.D., Wahyuni, C.U., 2016. Hubungan Antara Umur Kehamilan, Kehamilan Ganda, Hipertensi dan Anemia dengan Kejadian Bayi Berat Lahir Rendah (BBLR). *J. Berk. Epidemiol.* 4, 349–359. <https://doi.org/10.20473/jbe.v4i3>
- Restu, S., Dasuki, D., Nurdianti Z, R.D.S., 2014. The influence of iron supplementation in pregnant women to the occurrence of low birth weight (LBW) babies in Palu, Central Sulawesi. *J. thee Med. Sci. (Berkala Ilmu Kedokteran)* 46, 41–51. <https://doi.org/10.19106/jmedscie004601201406>
- Sarah, S., Irianto, I., 2018. Pengaruh Tingkat Kepatuhan Minum Tablet Fe Terhadap Kejadian Anemia Pada Ibu Hamil Trimester III di Puskesmas Pejeruk Tahun 2017. *Yars. Med. J.* 26, 075. <https://doi.org/10.33476/jky.v26i2.392>
- Setiati, A.R., Rahayu, S., 2017. Faktor Yang Mempengaruhi Kejadian BBLR (Berat Badan Lahir Rendah) Di Ruang Perawatan Intensif Neonatus RSUD DR Moewardi Di Surakarta. *J. Keperawatan Glob.* 2, 9–20. <https://doi.org/10.37341/jkg.v2i1.27>
- Sharami, S.H., Darkhaneh, R.F., Zahiri, Z., Milani, F., Asgharnia, M., Shakiba, M., Didar, Z., 2013. The relationship between vaginal bleeding in the first and second trimester of pregnancy and preterm labor. *Int. J. Reprod. Biomed.* 11, 385–390.
- Siramaneerat, I., Agushyana, F., Meebunmak, Y., 2018. Maternal Risk Factors Associated with Low Birth Weight in 376–383. <https://doi.org/10.2174/1874944501811010376>
- Sulistiyani, S., Darmawati, A.T., Setiani, O., 2019. Correlation between local-specific traditions of women agricultural workers and the Incidence of low birth weight in South Metro Sub-District, Metro City, Lampung Province, Indonesia. *Ann. Trop. Med. Public Heal.* 22, 1–5. <https://doi.org/10.36295/ASRO.2019.221118>
- Tyastuti, S., Wahyuningsih, H.P., 2016. Asuhan Kebidanan Kehamilan, Pertama. ed, Pusdik SDM Kesehatan, Badan Pengembangan dan Pemberdayaan Sumber Daya Kesehatan, Kementerian Kesehatan RI. BPPSDMK, Jakarta, Indonesia. <https://doi.org/10.1017/CBO9781107415324.004>
- U.S. Department of Health and Human Services, 2014. The Health Consequences of Smoking: 50 Years of Progress A Report of the Surgeon General, A Report of the Surgeon General.
- Upadhyay, R.P., Naik, G., Choudhary, T.S., Chowdhury, R., Taneja, S., Bhandari, N., Martines, J.C., Bahl, R., Bhan, M.K., 2019. Cognitive and Motor Outcomes in Children Born Low Birth Weight : A Systematic Review and Meta-analysis of Studies from South Asia. *BMC Pediatr.* 19, 1–15.
- WHO, UNICEF, 2019. Low Birthweight Estimates: Level and Trends 2000-2015, UNICEF and WHO.

#32644 Review

[SUMMARY](#)
[REVIEW](#)
[EDITING](#)

Submission



Authors Oktavia Beni Kujariningrum, Sri Winarni, Atik Mawarni, Najib Najib 
Title Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java
Section Articles
Editor Efa Nugroho, S.K.M, M.Kes 

Peer Review

Round 1

Review Version 32644-83650-1-RV.DOCX 2021-10-11
Initiated —
Last modified —
Uploaded file None

Editor Decision

Decision Accept Submission 2022-11-07
Notify Editor  Editor/Author Email Record  2022-08-21
Editor Version 32644-91924-1-ED.DOCX 2022-04-12
 32644-91924-2-ED.DOCX 2022-08-31
 32644-91924-3-ED.DOCX 2022-09-05
Author Version 32644-91956-1-ED.DOCX 2022-04-17 [DELETE](#)
 32644-91956-2-ED.DOCX 2022-08-31 [DELETE](#)
 32644-91956-3-ED.DOCX 2022-09-27 [DELETE](#)
Upload Author Version No file chosen

ISSN: 2355-3596

ABOUT THE JOURNAL

[Focus and Scope](#)
[Manuscript Submission](#)
[Guide for Authors](#)
[Editorial Board](#)
[Reviewer Team](#)
[Abstracting/Indexing](#)
[Ethics Statement](#)
[Policy of Screening for Plagiarism](#)
[Contact](#)
 2,143,152
[View Visitor Stats](#)

USER

You are logged in as...
oktaviabeni66
[» My Journals](#)
[» My Profile](#)
[» Log Out](#)

JOURNAL CONTENT

Search

Search Scope

Browse

[» By Issue](#)
[» By Author](#)
[» By Title](#)
[» Other Journals](#)

COLABORATION WITH



Ikatan Ahli Kesehatan Masyarakat Indonesia
 IAKMI (The Indonesian Public Health Association) is an independent professional organization for the better of public health, based on Pancasila based on the 1945 Constitution. Mu

Agreement No:
 402/UN.37.1.6/IKM/2012





Jejaring Nasional Pendidikan Kesehatan (JNPK)
 JNPK is an organization that gathers experts and observers in the field of health education, which was established on September 1, 2014. The founder of this organization is the University of Teacher Training Education Institut (LPTK) which organizes public health education, namely Universitas Negeri

#32644 Review

[SUMMARY](#)
[REVIEW](#)
[EDITING](#)

Submission



Authors Oktavia Beni Kujariningrum, Sri Winarni, Atik Mawarni, Najib Najib 
Title Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java
Section Articles
Editor Efa Nugroho, S.K.M, M.Kes 

Peer Review

Round 1

Review Version 32644-83650-1-RV.DOCX 2021-10-11
Initiated —
Last modified —
Uploaded file None

Editor Decision

Decision Accept Submission 2022-11-07
Notify Editor  Editor/Author Email Record  2022-08-21
Editor Version

32644-91924-1-ED.DOCX	2022-04-12	
32644-91924-2-ED.DOCX	2022-08-31	
32644-91924-3-ED.DOCX	2022-09-05	

Author Version

32644-91956-1-ED.DOCX	2022-04-17	DELETE
32644-91956-2-ED.DOCX	2022-08-31	DELETE
32644-91956-3-ED.DOCX	2022-09-27	DELETE

Upload Author Version

No file chosen

ISSN: 2355-3596

ABOUT THE JOURNAL

[Focus and Scope](#)
[Manuscript Submission](#)
[Guide for Authors](#)
[Editorial Board](#)
[Reviewer Team](#)
[Abstracting/Indexing](#)
[Ethics Statement](#)
[Policy of Screening for Plagiarism](#)
[Contact](#)
 2,143,152
[View Visitor Stats](#)

USER

You are logged in as...
oktaviabeni66
[» My Journals](#)
[» My Profile](#)
[» Log Out](#)

JOURNAL CONTENT

Search

Search Scope

All

Browse

[» By Issue](#)
[» By Author](#)
[» By Title](#)
[» Other Journals](#)

COLABORATION WITH



Ikatan Ahli Kesehatan Masyarakat Indonesia

IAKMI (The Indonesian Public Health Association) is an independent professional organization for the betterment of public health, based on Pancasila based on the 1945 Constitution. Mutual Agreement No:
 402/UN.37.1.6/IKM/2012



Jejaring Nasional Pendidikan Kesehatan (JNPK)

JNPK is an organization that gathers experts and observers in the field of health education, which was established on September 1, 2014. The founder of this organization is the University of Teacher Training Education Institut Teknologi Sepuluh Nopember (ITS) which organizes public health education, namely Universitas Negeri

Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java

Oktavia Beni Kujariningrum^{1, b)}, Sri Winarni^{1, a)*}, Atik Mawarni^{1, c)}, Najib Najib^{2, d)}

¹*Biostatistics and Population, Faculty of Public Health, Diponegoro University, Building C, 1st Floor, Main Campus of UNDIP 50275, Indonesia*

²*National Innovation Research Agency (BRIN)*

^{a)}Corresponding author: winarni@live.undip.ac.id*

^{b)}oktaviabeni66@gmail.com

^{c)}atikm246@gmail.com

^{d)}jibpenkb@gmail.com

Abstract.In Central Java, the prevalence of LBW (Low Birth Weight) has increased from 4.3 (2018) to 4.7 (2019) and be the biggest cause of neonatal mortality (46.4%) and infant mortality (40.5%).The aim of this research isto analyze the relationship between quality of ANC (Antenatal Care), iron supplementation, pregnancy complications, and maternal smoking status with LBW in Central Java.This research was an analytical study used secondary data from the 2017 IDHS (Indonesian Demographic and Health Survey). The sampling design used purposive sampling.Population study was 1205 babies born in Central Java. The sample comprised 952 babies. Independent variables were the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status, with the incidence of LBW as dependent variable.Data analysis was performed by chi-square continuity correction and logistic regression.Pregnancy complications have been associated with the incidences of LBW in Central Java (p -value = 0.0001). Iron supplementation (OR = 2.474) and pregnancy complications (OR = 4.869) had an effect on the incidence of LBW in Central Java.Iron supplementation and pregnancy complications influenced the incidence of LBW in Central Java.

Keyword: LBW, Pregnancy Complications, Iron Supplementation

INTRODUCTION

In 2015, 14.6% the incidences of LBW were found in the world and the highest prevalence occurring in Asia (17.3%)(WHO and UNICEF, 2019).The incidence of LBW in developing countries in the Asia Pacific region experienced an increase of ≤ 2 babies per 100 live births in 2014 compared to 2000 and an increase of 0.8% occurred in Indonesia(OECD and WHO, 2019). The results of the 2018 Riskesdas (Basic Health Research) showed that 6.2% of babies born with LBW in Indonesia and 6.1% were found in Central Java(Kementerian Kesehatan RI Badan Penelitian dan Pengembangan, 2018). The Central Java Health Profile in 2019 showed an increase in the incidence of LBW from 4.3 (2018) to 4.7 (2019)(Dinas Kesehatan Provinsi Jawa Tengah, 2019). A total of 5.9% of LBW events were found in single (Kujariningrum et al., 2021).

More than 37% of toddler were stunted in 2013(Bappenas and UNICEF, 2017).That condition had relationship with a history of LBW (p -value = 0.037; OR = 5.294)(Miranti et al., 2020).A study in South Asia showed that 17% of children with a history of LBW had a total IQ of less than 85(Upadhyay et al., 2019). In Central Java, LBW is the biggest cause of neonatal mortality (46.4%) and infant mortality (40.5%) in 2019 (Dinas Kesehatan Provinsi Jawa Tengah, 2019).

Visits and completeness of ANC services affect the incidence of LBW(Darwis et al., 2020; Paul et al., 2019) The incidence of LBW was also related to maternal anemia status (OR = 4.03) (Purwanto and Wahyuni, 2016). As much as 50% of the 73.3% pregnant women at Pejerk Health Center had adherence to consuming low Fe tablets(Sarah

and Irianto, 2018). The incidence of LBW in Cilacap Regional Hospital was associated with pregnancy complications (OR = 3.393) (Kusumawati et al., 2016). The entry of nicotine into the body's mechanism of pregnant women has an impact on fetal growth and development (U.S. Department of Health and Human Services, 2014). The incidence of LBW in Taiwan was associated with maternal smoking status (OR = 3.46) (Ko et al., 2014).

The incidence of LBW was related to many factors. There were differences in the causes of LBW in each region. A study to determine the risk factor of LBW in an area is urgently needed. There were been a lot of previously research about LBW, but for research with the scope of research in Central Java used the 2017 IDHS secondary data analysis with sample weighting has never been done. Based on the facts described above, this study aims to analyze the relationship between the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status with the incidence of LBW in Central Java.

METHOD

This study used a cross-sectional design. Research data from the 2017 IDHS. The study population consisted of 1205 babies born to WUS (Women of Childbearing Age) in 2012-2017 in Central Java. The research sample was taken used a purposive sampling based on inclusion and exclusion criteria for the total population so that a sample of 952 babies was obtained. The inclusion criteria included babies who were weighed in birth, babies who were last born by respondents in the range of 2012-2017, single births, and received ANC while in the womb. Infants with mothers who did not know the number of ANC visits, history of blood draws, history of consultations, and history of receiving Fe tablets were excluded from the study sample list as exclusion criteria.

The research variables consisted of the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status as independent variables with the incidence of LBW as the dependent variable. This study used the weighting of the sample according to the ethics of using the 2017 IDHS raw data and referring to a tutorial published by the you tube account The DHS Program on August 25, 2015 entitled " Part IV: Demonstration of How to Weight DHS Data in SPSS & SAS ". The bivariate analysis used was a statistical analysis of chi-square continuity correction. The independent variables were included in the multivariate logistic regression method enter are the independent variables with $p\text{-value} \leq 0.25$. The analysis was carried out using a significance level of 0.05 and a confidence level of 95%. The likelihood of LBW occurrences can be seen from the calculation of the predicted value using the following formula:

$$f(Z) = \frac{1}{1 + e^{-(b_0 + b_1x_1 + b_2x_2 + \dots + b_ix_i)}}$$

Formula description:

$f(Z)$	= probability
b_0	= constant
b_1, b_2, \dots, b_i	= partial regression coefficient
x_1, x_2, \dots, x_i	= independent variable
e	= exponent function with constant value 2.72

This study has passed the ethical clearance number: 12 / EA / KEPK-FKM / 2020 issued by the Health Research Ethics Committee, Faculty of Public Health, Diponegoro University on January 26, 2021.

RESULTS AND DISCUSSION

Most (94.8%) babies born in Central Java in 2012-2017 had a NBW (Normal Birth Weight). As many as 68.9 % of babies were born to mothers who received poor quality of ANC. Most (94.3%) babies were born to mothers who received iron supplementation during pregnancy. As many as 81.3% of babies were born to mother who did not experience complications during pregnancy. Most (99.1%) babies were born to mother who had never smoked.

Based on chi-square continuity correction, pregnancy complications were related to the incidences of LBW in Central Java ($p\text{-value} = 0.0001$). There were no relationships between the quality of ANC, iron supplementation, and maternal smoking status with the incidence of LBW in Central Java (Table 1).

TABLE 1. Correlation of Quality of Antenatal Care, Iron Supplementation, Pregnancy Complications, and Mother's Smoking Status with LBW incidence (N=952)

Independent Variable	LBW				f	Total %	p-value
	Not		Yes				
	n	%	n	%			
1. Quality of <i>Antenatal Care</i>							0.460
Good	278	94.2	17	5.8	295	100	
Poor	624	95.1	33	4.9	657	100	
2. Iron Supplementation							0.059
Yes	854	95.1	45	4.9	899	100	
Not	48	90.6	5	9.4	53	100	
3. Pregnancy Complications							0.0001
Not	749	96.8	26	3.2	775	100	
Yes	153	86.5	24	13.5	177	100	
4. Mother's Smoking Status							0.712
Not	893	94.8	50	5.2	943	100	
Yes	9	100	0	0	9	100	

Multivariate analysis used logistic regression with enter method gave result that iron supplementation and pregnancy complications affect the incidence of LBW in Central Java. Mothers who did not receive iron supplementation during their pregnancy had 2.474 times higher risk of giving birth to LBW babies than mothers who received iron supplementation (OR = 2.474). Mothers who experienced pregnancy complications had a 4.869 times higher risk of giving birth to LBW babies than mothers who did not experience complications (OR = 4.869) (Table 2). Based on the calculation above, $f(Z) = 0.27$ can be conclude that mothers who didn't receive iron supplementation and had pregnancy complications history have a chance 27% to give the incidence of LBW.

TABLE 2. Results of Multivariate Analysis The Effect of Iron Supplementation and Pregnancy Complications on the Incidence of LBW in Central Java 2012-2017 (N=952)

Variables		B	SE	Wald	Sig	Exp (B)
Model 1	Iron Supplementation	0.906	0.366	6.125	0.013	2.474
	Pregnancy Complications	1.583	0.218	52.670	0.0001	4.869
	Constant	-3.484	0.155	506.665	0.0001	0.031

The quality of ANC is a risk factor for LBW incidence (Owa et al., 2017). Most of the mothers access ANC services with low quality and this condition is related to the incidence of LBW in Indonesia (Darwis et al., 2020). In contrast to the conditions found in Central Java. Chi-square result showed that there was no relationship between the quality of ANC and the incidences of LBW in Central Java (p -value = 0.488). This is in line with Meiriza (2018), which concluded that there was no relationship between the quality of ANC at level I health facilities and the incidence of LBW in Padang City (Meiriza et al., 2018). This study found as many as 68.9 % of infants born to mothers who received ANC with poor quality. The incidence of LBW was more found in the group of infants with mothers who received ANC with good quality (5.8%) compared to poor quality (4.9). This showed that pregnant women who receive good quality antenatal care can also deliver babies with LBW. This condition is possible because of the limited variables found in the secondary data of the 2017 IDHS and the information needed to assess the quality of ANC is not enough just by questionnaires result, but requires in-depth interviews. Owa (2019) conducted in-depth interviews and found that pregnant women who received less ANC quality had a 3.5 times higher risk for having a baby with LBW (OR = 3.5) (Owa, 2019).

Iron supplementation is an effort made to respond the high rates of iron deficiency anemia in pregnant women (Tyastuti and Wahyuningsih, 2016). Chi-square result showed that there was no relationship between iron supplementation and the incidences of LBW (p -value = 0.076). Different from the multivariate analysis result which showed the effect of iron supplementation on the incidence of LBW (p -value = 0.013). Mothers who didn't receive iron supplementation during their pregnancy had a 2.474 times higher risk for having a baby with LBW than

mothers who received iron supplementation (OR = 2.474). In line with Restu et al (2014) which found that iron supplementation had an effects on the incidences of LBW (OR = 3.82)(Restu et al., 2014). This shows that mothers who get iron supplementation can avoid anemia. Iron deficiency anemia causes an increase in serum norepinephrine concentration which results in maternal and fetal stress, stimulates the synthesis of CRH (Corticotrophin-Releasing Hormone) which will increase fetal cortisol productions thus impacting IUGR (Intrauterine Growth Restriction) and resulted in LBW (Chhabra and Chopra, 2016). Iron supplementation can reduce the prevalence of iron deficiency anemia(Long et al., 2012).Each tablet contains FeSO₄ mg (iron 30 mg) which is useful as a reserve of iron, red blood cell synthesis and muscle blood synthesis during pregnancy(Tyastuti and Wahyuningsih, 2016).

Pregnancy complications are collection of symptoms of health problems during pregnancy that can affect the health conditions of the baby and mother (BKKBN et al., 2017; Neiger, 2017). Some symptoms of pregnancy complications such as prolonged nausea and vomiting, hypotension and hypertension are associated with a lack of nutritional intake to the fetus (Dirjen Kesmas Kemenkes, 2018; Karjatin, 2016). The results of this study found that the incidences of LBW was more in the group of infants with mothers who had complications during their pregnancy (13.5%) than in the group of infants with mothers who had no complications during their pregnancy (3.2%). Chi-square result showed that there was a significant relationship between pregnancy complications and the incidence of LBW (p -value = 0.0001). Mothers who experienced pregnancy complications had a 4,869 times higher risk for having a baby with LBW than mothers who did not experience pregnancy complications (OR = 4.869). Siramaneerat (2018) mentions the same finding that pregnancy complications are associated with the incidences of LBW (p -value = 0.0001; OR = 1.731) (Siramaneerat et al., 2018). The most common complication found in this study was bleeding (5.5%). Antepartum hemorrhage associated with the incidences of LBW (Setiati and Rahayu, 2017). Antepartum hemorrhage can increase the likelihood of preterm birth which is a risk of LBW (Sharami et al., 2013). Pregnant women who experience antepartum bleeding have a 2.23 times higher risk of giving birth to LBW babies than mothers who don't experience antepartum hemorrhage (Isnaini, 2015). There was a difference between birth weight in the group of mothers with and without antepartum hemorrhage. Antepartum hemorrhage causes blood flow that distributes oxygen and nutrients to the placenta from the mother to the fetus to be disturbed. Impaired delivery of oxygen and nutrients will cause fetal anemia, shock to fetal death. Fetuses that survive until birth will experience various disorders, including LBW(Setiati and Rahayu, 2017).

Receptors in the placental blood vessels that mix with nicotine cause a decrease in blood flow in the placenta and fetal vasoconstriction which results in impaired delivery of oxygen and nutrients to the fetus so that the fetus experiences malnutrition which results in impaired fetal growth (U.S. Department of Health and Human Services, 2014). The entry of nicotine in the body's mechanism is proven to have an impact on the low production of the hormone Insulin-Like Growth Factor-1 in pregnant women who smoke will affect fetal growth and development (Fang et al., 2015). Pregnant women that smoke will give birth to babies with birth weight 320-435 grams lower than pregnant women who do not smoke(Kataoka et al., 2018). Outcome of this study shows that the incidences of LBW is more common in the group of infants whose mothers didn't smoke (5.2%), while NBW more common in the group of infants with mothers who smoked (100%). Chi-square result showed that there was no relationship between the mother's smoking status and the incidences of LBW (p -value = 0.706). In line with Phowira et al (2020) which stated that the mother's smoking status was not related to the incidence of LBW in DKI Jakarta (p -value = 0.448) (Phowira et al., 2020).There was no relationship between the frequency of smoking per day and the incidences of LBW in Lampung Province (Sulistiyani et al., 2019). It can be said that pregnant women who do not smoke can give birth to babies with LBW. This mechanism shows that the impact of nicotine entering the body cannot be described only by the status of a pregnant woman as an active smoker or not, as information is available in the 2017 IDHS data. LBW was found in mothers with high levels of nicotine > 143 µg/g keratinize. This condition is not only found in pregnant women who smoke actively. Pregnant women who don't smoke have an average nicotine level of 153.2 ± 96.0 µg/g keratinize (>143 µg/g keratinize) as a result of exposure to cigarette smoke from the environment (passive smoking)(Huang et al., 2017).There was a relationship between passive smoking mothers and the incidences of LBW (OR = 3.04) (Ardelia et al., 2019).

CONCLUSION

Most babies are born at a normal weight. There was a relationship between pregnancy complications and the incidences of LBW in Central Java. The quality of antenatal care, iron supplementation, and maternal smoking status were not related with the incidences of LBW. Pregnant woman who don't get iron supplementation (OR = 2.474) and complications of pregnancy (OR = 4.869) had chances of 0.84 for having a baby with LBW.The pregnant

women who experience signs of complications immediately take action and visit health facilities for further assistance. Pregnant women are also advised to take Fe tablets regularly at least 90 tablets during their pregnancy.

ACKNOWLEDGMENTS

Thank you to the United States Agency for International Development for providing access to secondary data for the 2017 IDHS through the dhsprogram.com website, Diponegoro University for facilitating us in accessing various e-journal portals, and the Health Research Ethics Committee, Faculty of Public Health, Diponegoro University. has issued a permit for the ethical feasibility of this research.

REFERENCES


- Ardelia, K.I.A., Hardianto, G., Nuswantoro, D., 2019. Passive smoker during pregnancy is a risk factor of low birth weight. *Maj. Obstet. Ginekol.* 27, 12. <https://doi.org/10.20473/mog.v1i12019.12-16>
- Bappenas, UNICEF, 2017. *Laporan Baseline SDG tentang Anak-Anak di Indonesia*. Jakarta.
- BKKBN, BPS, Kemenkes, International, I., 2017. *Indonesia Demographic and Health Survey 2017. DHS Program*, Jakarta, Indonesia.
- Chhabra, S., Chopra, S., 2016. Mid Pregnancy Fetal Growth Restriction and Maternal Anaemia a Prospective Study. *J. Nutr. Disord. Ther.* 06. <https://doi.org/10.4172/2161-0509.1000187>
- Darwis, A., Abdullah, A., Maidar, Adamy, A., Nurjannah, 2020. Hubungan Komponen Pelayanan Antenatal Care (10T) dengan Kejadian Bayi Berat Lahir Rendah di Indonesia (Analisa Data Sekunder SDKI 2017). *JUKEMA* 6, 13–19.
- Dinas Kesehatan Provinsi Jawa Tengah, 2019. *Profil Kesehatan Provinsi Jawa Tengah Tahun 2019*, Dinkes Jateng.
- Dirjen Kesmas Kemenkes, 2018. *Laporan Kinerja Ditjen Kesehatan Masyarakat Tahun 2017*.
- Fang, F., Luo, Z.C., Dejemli, A., Delvin, E., Zhang, J., 2015. Maternal smoking and metabolic health biomarkers in newborns. *PLoS One*. <https://doi.org/10.1371/journal.pone.0143660>
- Huang, Shih-hui, Weng, K., Huang, Shih-ming, Liou, H., 2017. The effects of maternal smoking exposure during pregnancy on postnatal outcomes : A cross sectional study. *J. Chinese Med. Assoc.* 1–7. <https://doi.org/10.1016/j.jcma.2017.01.007>
- Isnaini, N., 2015. Analisis Faktor Yang Mempengaruhi Kejadian Bayi Berat Lahir Rendah. *J. Dunia Kesmas* 4, 196–202.
- Karjatin, A., 2016. *Keperawatan Maternitas*. Pusdik SDM Kesehatan, Jakarta.
- Kataoka, M.C., Paula, A., Carvalheira, P., Ferrari, A.P., Antonieta, M., Leite, D.B., Maria, C., Lima, G. De, 2018. Smoking during pregnancy and harm reduction in birth weight : a cross-sectional study. *BMC Pregnancy Childbirth* 1–10.
- Kementerian Kesehatan RI Badan Penelitian dan Pengembangan, 2018. *Hasil Utama Riset Kesehatan Dasar 2018*, Kementerian Kesehatan Republik Indonesia. Jakarta, Indonesia.
- Ko, T.J., Tsai, L.Y., Chu, L.C., Yeh, S.J., Leung, C., Chen, C.Y., Chou, H.C., Tsao, P.N., Chen, P.C., Hsieh, W.S., 2014. Parental smoking during pregnancy and its association with low birth weight, small for gestational age, and preterm birth offspring: A birth cohort study. *Pediatr. Neonatol.* 55, 20–27. <https://doi.org/10.1016/j.pedneo.2013.05.005>
- Kujariningrum, O.B., Winarni, S., Najib, N., 2021. Multiple pregnancy : the biggest risk factor of low birth weight in Central Java , Indonesia (2017 IDHS secondary data study). *Ann. Trop. Med. Public Heal.* 24, 24–184. <https://doi.org/10.36295/ASRO.2021.24184>
- Kusumawati, D.D., Septiyaningsih, R., Kania, 2016. Faktor-Faktor Ibu yang Mempengaruhi Kejadian Bayi Berat Lahir Rendah (BBLR). *J. Kesehat. Al-Irsyad* 9, 8–16.
- Long, H., Yi, J.M., Hu, P.L., Li, Z. Bin, Qiu, W.Y., Wang, F., Zhu, S., 2012. Benefits of Iron supplementation for low birth weight infants: A systematic review. *BMC Pediatr.* 12. <https://doi.org/10.1186/1471-2431-12-99>
- Meiriza, W., Aladin, A., Edison, E., 2018. The Correlation of Maternal Factors and The Quality of Antenatal Care Services With Low Birth Weight Babies In Health Facilities Level I. *J. Midwifery* 3, 103. <https://doi.org/10.25077/jom.1.1.103-114.2018>
- Miranti, Mutiarasari, D., Arsin, A.A., Hadju, V., Mallongi, A., Nur, R., Amri, I., Haruni, H., Wahyuni, R.D., Rahma, Faris, A., 2020. Determinants of the incidence of stunting in the working area of Kinovaro Sigi Health Center. *Enfermeria Clínica* 30, 246–252. <https://doi.org/10.1016/j.enfcli.2019.10.077>

- Neiger, R., 2017. Long-Term Effects of Pregnancy Complications on Maternal Health: A Review. *J. Clin. Med.* 6, 76. <https://doi.org/10.3390/jcm6080076>
- OECD, WHO, 2019. Health at a glance: Asia/Pacific 2018: Measuring progress towards universal health coverage. OECD Publishing, Paris.
- Owa, K., 2019. Chronic Energy Deficiency , Anemia as a Risk Factor for Low Birth Weight Babies in East Nusa Tenggara Kekurangan Energi Kronis , Anemia sebagai Faktor Risiko Bayi Berat Lah. *J. Kesehat. Prim.* 4, 13–22.
- Owa, K., Putra, I.W.G.A.E., Windiani, I.G.A.T., 2017. Risk factors for low birth weight infants in East Nusa Tenggara. *Public Heal. Prev. Med. Arch.* 5, 39. <https://doi.org/10.15562/phpma.v5i1.40>
- Paul, P., Zaveri, A., Chouhan, P., 2019. Assessing the Impact of Antenatal Care Utilization on Low Birthweight in India: Analysis of the 2015–2016 National Family Health Survey. *Child. Youth Serv. Rev.* 106, 104459. <https://doi.org/10.1016/j.childyouth.2019.104459>
- Phowira, J., Elvina, F.T., Wiguna, I.I., Wahyudi, F.R.H.B., Medise, B.E., 2020. The Association between Tobacco Exposure during Pregnancy and Newborns' Birth Weight in DKI Jakarta Community Members. *medRxiv* 1–15. <https://doi.org/10.1101/2020.10.29.20222059>
- Purwanto, A.D., Wahyuni, C.U., 2016. Hubungan Antara Umur Kehamilan, Kehamilan Ganda, Hipertensi dan Anemia dengan Kejadian Bayi Berat Lahir Rendah (BBLR). *J. Berk. Epidemiol.* 4, 349–359. <https://doi.org/10.20473/jbe.v4i3>.
- Restu, S., Dasuki, D., Nurdianti Z, R.D.S., 2014. The influence of iron supplementation in pregnant women to the occurrence of low birth weight (LBW) babies in Palu, Central Sulawesi. *J. thee Med. Sci. (Berkala Ilmu Kedokteran)* 46, 41–51. <https://doi.org/10.19106/jmedscie004601201406>
- Sarah, S., Irianto, I., 2018. Pengaruh Tingkat Kepatuhan Minum Tablet Fe Terhadap Kejadian Anemia Pada Ibu Hamil Trimester III di Puskesmas Pejeruk Tahun 2017. *Yars. Med. J.* 26, 075. <https://doi.org/10.33476/jky.v26i2.392>
- Setiati, A.R., Rahayu, S., 2017. Faktor Yang Mempengaruhi Kejadian BBLR (Berat Badan Lahir Rendah) Di Ruang Perawatan Intensif Neonatus RSUD DR Moewardi Di Surakarta. *J. Keperawatan Glob.* 2, 9–20. <https://doi.org/10.37341/jkg.v2i1.27>
- Sharami, S.H., Darkhaneh, R.F., Zahiri, Z., Milani, F., Asgharnia, M., Shakiba, M., Didar, Z., 2013. The relationship between vaginal bleeding in the first and second trimester of pregnancy and preterm labor. *Int. J. Reprod. Biomed.* 11, 385–390.
- Siramaneerat, I., Agushyana, F., Meebunmak, Y., 2018. Maternal Risk Factors Associated with Low Birth Weight in 376–383. <https://doi.org/10.2174/1874944501811010376>
- Sulistiyani, S., Darmawati, A.T., Setiani, O., 2019. Correlation between local-specific traditions of women agricultural workers and the Incidence of low birth weight in South Metro Sub-District, Metro City, Lampung Province, Indonesia. *Ann. Trop. Med. Public Heal.* 22, 1–5. <https://doi.org/10.36295/ASRO.2019.221118>
- Tyastuti, S., Wahyuningsih, H.P., 2016. Asuhan Kebidanan Kehamilan, Pertama. ed, Pusdik SDM Kesehatan, Badan Pengembangan dan Pemberdayaan Sumber Daya Kesehatan, Kementerian Kesehatan RI. BPPSDMK, Jakarta, Indonesia. <https://doi.org/10.1017/CBO9781107415324.004>
- U.S. Department of Health and Human Services, 2014. The Health Consequences of Smoking: 50 Years of Progress A Report of the Surgeon General, A Report of the Surgeon General.
- Upadhyay, R.P., Naik, G., Choudhary, T.S., Chowdhury, R., Taneja, S., Bhandari, N., Martines, J.C., Bahl, R., Bhan, M.K., 2019. Cognitive and Motor Outcomes in Children Born Low Birth Weight : A Systematic Review and Meta-analysis of Studies from South Asia. *BMC Pediatr.* 19, 1–15.
- WHO, UNICEF, 2019. Low Birthweight Estimates: Level and Trends 2000-2015, UNICEF and WHO.

#32644 Review


[SUMMARY](#) [REVIEW](#) [EDITING](#)

Submission

Authors Oktavia Beni Kujariningrum, Sri Winarni, Atik Mawarni, Najib Najib 

Title Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java

Section Articles

Editor Efa Nugroho, S.K.M, M.Kes 

Peer Review

Round 1

Review Version 32644-83650-1-RV.DOCX 2021-10-11



Initiated —

Last modified —

Uploaded file None

Editor Decision

Decision Accept Submission 2022-11-07

Notify Editor  Editor/Author Email Record  2022-08-21

Editor Version 32644-91924-1-ED.DOCX 2022-04-12
32644-91924-2-ED.DOCX 2022-08-31
32644-91924-3-ED.DOCX 2022-09-05

Author Version 32644-91956-1-ED.DOCX 2022-04-17 [DELETE](#)
32644-91956-2-ED.DOCX 2022-08-31 [DELETE](#)
32644-91956-3-ED.DOCX 2022-09-27 [DELETE](#)

Upload Author Version No file chosen

ISSN: 2355-3596

ABOUT THE JOURNAL

[Focus and Scope](#)

[Manuscript Submission](#)

[Guide for Authors](#)

[Editorial Board](#)

[Reviewer Team](#)

[Abstracting/Indexing](#)

[Ethics Statement](#)

[Policy of Screening for Plagiarism](#)

[Contact](#)
2,143,152
[View Visitor Stats](#)

USER

You are logged in as...

oktaviabeni66

» [My Journals](#)

» [My Profile](#)

» [Log Out](#)

JOURNAL CONTENT

Search

Search Scope

[Browse](#)

» [By Issue](#)

» [By Author](#)

» [By Title](#)

» [Other Journals](#)

COLABORATION WITH



Ikatan Ahli Kesehatan Masyarakat Indonesia

IAKMI (The Indonesian Public Health Association) is an independent professional organization for the betterment of public health, based on Pancasila based on the 1945 Constitution. Mutual Agreement No: 402/UN.37.1.6/IKM/2012




Jejaring Nasional Pendidikan Kesehatan (JNPK)

JNPK is an organization that gathers experts and observers in the field of health education, which was established on September 1, 2014. The founder of this organization is the University of Teacher Training Education Institut Teknologi Sepuluh Nopember (ITS) which organizes public health education, namely Universitas Negeri

#32644 Review


[SUMMARY](#) [REVIEW](#) [EDITING](#)

Submission

Authors Oktavia Beni Kujariningrum, Sri Winarni, Atik Mawarni, Najib Najib 

Title Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java

Section Articles

Editor Efa Nugroho, S.K.M, M.Kes 

Peer Review

Round 1

Review Version 32644-83650-1-RV.DOCX 2021-10-11



Initiated —

Last modified —

Uploaded file None

Editor Decision

Decision Accept Submission 2022-11-07

Notify Editor  Editor/Author Email Record  2022-08-21

Editor Version 32644-91924-1-ED.DOCX 2022-04-12
32644-91924-2-ED.DOCX 2022-08-31
32644-91924-3-ED.DOCX 2022-09-05

Author Version 32644-91956-1-ED.DOCX 2022-04-17 [DELETE](#)
32644-91956-2-ED.DOCX 2022-08-31 [DELETE](#)
32644-91956-3-ED.DOCX 2022-09-27 [DELETE](#)

Upload Author Version No file chosen

ISSN: 2355-3596

ABOUT THE JOURNAL

[Focus and Scope](#)

[Manuscript Submission](#)

[Guide for Authors](#)

[Editorial Board](#)

[Reviewer Team](#)

[Abstracting/Indexing](#)

[Ethics Statement](#)

[Policy of Screening for Plagiarism](#)

[Contact](#)
2,143,152
[View Visitor Stats](#)

USER

You are logged in as...

oktaviabeni66

» [My Journals](#)

» [My Profile](#)

» [Log Out](#)

JOURNAL CONTENT

Search

Search Scope

Browse

» [By Issue](#)

» [By Author](#)

» [By Title](#)

» [Other Journals](#)

COLABORATION WITH



Ikatan Ahli Kesehatan Masyarakat Indonesia

IAKMI (The Indonesian Public Health Association) is an independent professional organization for the betterment of public health, based on Pancasila based on the 1945 Constitution. Mutual Agreement No: 402/UN.37.1.6/IKM/2012



Jejaring Nasional Pendidikan Kesehatan (JNPK)

JNPK is an organization that gathers experts and observers in the field of health education, which was established on September 1, 2014. The founder of this organization is the University of Teacher Training Education Institut Teknologi Sepuluh Nopember (ITS) which organizes public health education, namely Universitas Negeri

Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java

Comment [WHC1]: Judultanpasingkata
n

Oktavia Beni Kujariningrum^{1, b)}, Sri Winarni^{1, a)*}, Atik Mawarni^{1, c)}, Najib Najib^{2, d)}

¹*Biostatistics and Population, Faculty of Public Health, Diponegoro University, Building C, 1st Floor, Main Campus of UNDIP 50275, Indonesia*

²*National Innovation Research Agency (BRIN)*

^{a)}Corresponding author: winarni@live.undip.ac.id*

^{b)}oktaviabeni66@gmail.com

^{c)}atikm246@gmail.com

^{d)}jibpenkb@gmail.com

Abstract. In Central Java, the prevalence of LBW (Low Birth Weight) has increased from 4.3 (2018) to 4.7 (2019) and be the biggest cause of neonatal mortality (46.4%) and infant mortality (40.5%). The aim of this research is to analyze the relationship between quality of ANC (Antenatal Care), iron supplementation, pregnancy complications, and maternal smoking status with LBW in Central Java. This research was an analytical study used secondary data from the 2017 IDHS (Indonesian Demographic and Health Survey). The sampling design used purposive sampling. Population study was 1205 babies born in Central Java. The sample comprised 952 babies. Independent variables were the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status, with the incidence of LBW as dependent variable. Data analysis was performed by chi-square continuity correction and logistic regression. Pregnancy complications have been associated with the incidences of LBW in Central Java (p -value = 0.0001). Iron supplementation (OR = 2.474) and pregnancy complications (OR = 4.869) had an effect on the incidence of LBW in Central Java. Iron supplementation and pregnancy complications influenced the incidence of LBW in Central Java.

Keyword: LBW, Pregnancy Complications, Iron Supplementation

INTRODUCTION

In 2015, 14.6% the incidences of LBW were found in the world and the highest prevalence occurring in Asia (17.3%) (WHO and UNICEF, 2019). The incidence of LBW in developing countries in the Asia Pacific region experienced an increase of ≤ 2 babies per 100 live births in 2014 compared to 2000 and an increase of 0.8% occurred in Indonesia (OECD and WHO, 2019). The results of the 2018 Riskesdas (Basic Health Research) showed that 6.2% of babies born with LBW in Indonesia and 6.1% were found in Central Java (Kementerian Kesehatan RI Badan Penelitian dan Pengembangan, 2018). The Central Java Health Profile in 2019 showed an increase in the incidence of LBW from 4.3 (2018) to 4.7 (2019) (Dinas Kesehatan Provinsi Jawa Tengah, 2019). A total of 5.9% of LBW events were found in single (Kujariningrum et al., 2021).

More than 37% of toddler were stunted in 2013 (Bappenas and UNICEF, 2017). That condition had relationship with a history of LBW (p -value = 0.037; OR = 5.294) (Miranti et al., 2020). A study in South Asia showed that 17% of children with a history of LBW had a total IQ of less than 85 (Upadhyay et al., 2019). In Central Java, LBW is the biggest cause of neonatal mortality (46.4%) and infant mortality (40.5%) in 2019 (Dinas Kesehatan Provinsi Jawa Tengah, 2019).

Visits and completeness of ANC services affect the incidence of LBW (Paul et al., 2019). The incidence of LBW was also related to maternal anemia status (OR = 1.23) (Figueiredo et al., 2018). As much as 50% of the 73.3% pregnant women at Pejerkut Health Center had adherence to consuming low Fe tablets (Sarah and Irianto, 2018). The

incidence of LBW was associated with pregnancy complications (Bener et al., 2012). The entry of nicotine into the body's mechanism of pregnant women has an impact on fetal growth and development (Nemoto et al., 2021). The incidence of LBW in Taiwan was associated with maternal smoking status (OR = 3.46) (Ko et al., 2014).

The incidence of LBW was related to many factors. There were differences in the causes of LBW in each region. A study to determine the risk factor of LBW in an area is urgently needed. There were a lot of previously research about LBW, but for research with the scope of research in Central Java used the 2017 IDHS secondary data analysis with sample weighting has never been done. Based on the facts described above, this study aims to analyze the relationship between the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status with the incidence of LBW in Central Java.

METHOD

This study used a cross-sectional design. Research data from the 2017 IDHS. The study population consisted of 1205 babies born to WUS (Women of Childbearing Age) in 2012-2017 in Central Java. The research sample was taken used a purposive sampling based on inclusion and exclusion criteria for the total population so that a sample of 952 babies was obtained. The inclusion criteria included babies who were weighed in birth, babies who were last born by respondents in the range of 2012-2017, single births, and received ANC while in the womb. Infants with mothers who did not know the number of ANC visits, history of blood draws, history of consultations, and history of receiving Fe tablets were excluded from the study sample list as exclusion criteria.

The research variables consisted of the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status as independent variables with the incidence of LBW as the dependent variable. This study used the weighting of the sample according to the ethics of using the 2017 IDHS raw data and referring to a tutorial published by the YouTube account The DHS Program on August 25, 2015 entitled "Part IV: Demonstration of How to Weight DHS Data in SPSS & SAS". The bivariate analysis used was a statistical analysis of chi-square continuity correction. The independent variables were included in the multivariate logistic regression method enter are the independent variables with $p\text{-value} \leq 0.25$. The analysis was carried out using a significance level of 0.05 and a confidence level of 95%. The likelihood of LBW occurrences can be seen from the calculation of the predicted value using the following formula:

$$f(Z) = \frac{1}{1 + e^{-(b_0 + b_1x_1 + b_2x_2 + \dots + b_ix_i)}}$$

Formula description:

$f(Z)$ = probability

b_0 = constant

b_1, b_2, \dots, b_i = partial regression coefficient

x_1, x_2, \dots, x_i = independent variable

e = exponent function with constant value 2.72

This study has passed the ethical clearance number: 12 / EA / KEPK-FKM / 2020 issued by the Health Research Ethics Committee, Faculty of Public Health, Diponegoro University on January 26, 2021.

RESULTS AND DISCUSSION

Most (94.8%) babies born in Central Java in 2012-2017 had a NBW (Normal Birth Weight). As many as 68.9 % of babies were born to mothers who received poor quality of ANC. Most (94.3%) babies were born to mothers who received iron supplementation during pregnancy. As many as 81.3% of babies were born to mother who did not experience complications during pregnancy. Most (99.1%) babies were born to mother who had never smoked.

Based on chi-square continuity correction, pregnancy complications were related to the incidences of LBW in Central Java ($p\text{-value} = 0.0001$). There were no relationships between the quality of ANC, iron supplementation, and maternal smoking status with the incidence of LBW in Central Java (Table 1).

Comment [WHC2]: Pembahasan di perdalam

TABLE 1. Correlation of Quality of Antenatal Care, Iron Supplementation, Pregnancy Complications, and Mother's Smoking Status with LBW incidence (N=952)

Independent Variable	LBW				Total		p-value
	Not		Yes		f	%	
	n	%	n	%			
1. Quality of <i>Antenatal Care</i>							0.460
Good	278	94.2	17	5.8	295	100	
Poor	624	95.1	33	4.9	657	100	
2. Iron Supplementation							0.059
Yes	854	95.1	45	4.9	899	100	
Not	48	90.6	5	9.4	53	100	
3. Pregnancy Complications							0.0001
Not	749	96.8	26	3.2	775	100	
Yes	153	86.5	24	13.5	177	100	
4. Mother's Smoking Status							0.712
Not	893	94.8	50	5.2	943	100	
Yes	9	100	0	0	9	100	

Multivariate analysis used logistic regression with enter method gave result that iron supplementation and pregnancy complications affect the incidence of LBW in Central Java. Mothers who did not receive iron supplementation during their pregnancy had 2.474 times higher risk of giving birth to LBW babies than mothers who received iron supplementation (OR = 2.474). Mothers who experienced pregnancy complications had a 4.869 times higher risk of giving birth to LBW babies than mothers who did not experience complications (OR = 4.869) (Table 2). Based on the calculation above, $f(Z) = 0.27$ can be conclude that mothers who didn't receive iron supplementation and had pregnancy complications history have a chance 27% to give the incidence of LBW.

TABLE 2. Results of Multivariate Analysis The Effect of Iron Supplementation and Pregnancy Complications on the Incidence of LBW in Central Java 2012-2017 (N=952)

Variables	B	SE	Wald	Sig	Exp (B)
Model 1					
Iron Supplementation	0.906	0.366	6.125	0.013	2.474
Pregnancy Complications	1.583	0.218	52.670	0.0001	4.869
Constant	-3.484	0.155	506.665	0.0001	0.031

The quality of ANC is a risk factor for LBW incidence (Owa et al., 2017). Most (93.27%) of the mothers access ANC services with low quality and this condition is related to the incidence of LBW in Indonesia (Darwis et al., 2020). In contrast to the conditions found in Central Java. Chi-square result showed that there was no relationship between the quality of ANC and the incidences of LBW in Central Java (p -value = 0.488). This is in line with Meiriza (2018), which concluded that there was no relationship between the quality of ANC at level I health facilities and the incidence of LBW in Padang City (Meiriza et al., 2018). This study found as many as 68.9 % of infants born to mothers who received ANC with poor quality. The incidence of LBW was more found in the group of infants with mothers who received ANC with good quality (5.8%) compared to poor quality (4.9). This showed that pregnant women who receive good quality antenatal care can also deliver babies with LBW. This condition is possible because of the limited variables found in the secondary data of the 2017 IDHS and the information needed to assess the quality of ANC is not enough just by questionnaires result, but requires in-depth interviews. Owa (2019) conducted in-depth interviews and found that pregnant women who received less ANC quality had a 3.5 times higher risk for having a baby with LBW (OR = 3.5) (Owa, 2019).

Iron supplementation is an effort made to respond the high rates of iron deficiency anemia in pregnant women (Seu et al., 2019). Chi-square result showed that there was no relationship between iron supplementation and the incidences of LBW (p -value = 0.076). Different from the multivariate analysis result which showed the effect of iron supplementation on the incidence of LBW (p -value = 0.013). Mothers who didn't receive iron supplementation during their pregnancy had a 2.474 times higher risk for having a baby with LBW than mothers who received iron supplementation (OR = 2.474). In line with Restu et al (2014) which found that iron supplementation had an effects

on the incidences of LBW (OR = 3.82)(Restu et al., 2014). This shows that mothers who get iron supplementation can avoid anemia. Iron deficiency anemia causes an increase in serum norepinephrine concentration which results in maternal and fetal stress, stimulates the synthesis of CRH (Corticotrophin-Releasing Hormone) which will increase fetal cortisol productions thus impacting IUGR (Intrauterine Growth Restriction) and resulted in LBW (Chhabra and Chopra, 2016). Iron supplementation can reduce the prevalence of iron deficiency anemia(Long et al., 2012).Each tablet contains FeSO_4 mg (iron 30 mg) which is useful as a reserve of iron, red blood cell synthesis and muscle blood synthesis during pregnancy(Seu et al., 2019).

Pregnancy complications are collection of symptoms of health problems during pregnancy that can affect the health conditions of the baby and mother (BKKBN et al., 2017; Neiger, 2017).Some symptoms of pregnancy complications such as prolonged nausea and vomiting, hypotension and hypertension are associated with a lack of nutritional intake to the fetus and LBW (Marshall et al., 2022). The results of this study found that the incidences of LBW was more in the group of infants with mothers who had complications during their pregnancy (13.5%) than in the group of infants with mothers who had no complications during their pregnancy (3.2%). Chi-square result showed that there was a significant relationship between pregnancy complications and the incidence of LBW (p -value = 0.0001). Mothers who experienced pregnancy complications had a 4,869 times higher risk for having a baby with LBW than mothers who did not experience pregnancy complications (OR = 4.869). Siramaneerat (2018) mentions the same finding that pregnancy complications are associated with the incidences of LBW (p -value = 0.0001; OR = 1.731) (Siramaneerat et al., 2018). The most common complication found in this study was bleeding (5.5%). Antepartum hemorrhage associated with the incidences of LBW(Bener et al., 2012). Antepartum hemorrhage can increase the likelihood of preterm birth which is a risk of LBW (Sharami et al., 2013). Pregnant women who experience antepartum hemorrhage have a 1.6 times higher risk of giving birth to LBW babies than mothers who don't experience antepartum hemorrhage(Bener et al., 2012). There was a difference between birth weight in the group of mothers with and without antepartum hemorrhage. Antepartum hemorrhage causes blood flow that distributes oxygen and nutrients to the placenta from the mother to the fetus to be disturbed. Impaired delivery of oxygen and nutrients will cause fetal anemia, shock to fetal death. Fetuses that survive until birth will experience various disorders, including LBW(Kuribayashi et al., 2021).

Receptors in the placental blood vessels that mix with nicotine cause a decrease in blood flow in the placenta and fetal vasoconstriction which results in impaired delivery of oxygen and nutrients to the fetus so that the fetus experiences malnutrition which results in impaired fetal growth(Nemoto et al., 2021). The entry of nicotine in the body's mechanism is proven to have an impact on the low production of the hormone Insulin-Like Growth Factor-1 in pregnant women who smoke will affect fetal growth and development (Fang et al., 2015). Pregnant women that smoke will give birth to babies with birth weight 320-435 grams lower than pregnant women who do not smoke(Kataoka et al., 2018). Outcome of this study shows that the incidences of LBW is more common in the group of infants whose mothers didn't smoke (5.2%), while NBW more common in the group of infants with mothers who smoked (100%). Chi-square result showed that there was no relationship between the mother's smoking status and the incidences of LBW (p -value = 0.706). In line with Phowira et al (2020) which stated that the mother's smoking status was not related to the incidence of LBW in DKI Jakarta (p -value = 0.448) (Phowira et al., 2020).There was no relationship between the frequency of smoking per day and the incidences of LBW in Lampung Province (Sulistiyan et al., 2019). It can be said that pregnant women who do not smoke can give birth to babies with LBW. This mechanism shows that the impact of nicotine entering the body cannot be described only by the status of a pregnant woman as an active smoker or not, as information is available in the 2017 IDHS data. LBW was found in mothers with high levels of nicotine > 143 $\mu\text{g/g}$ keratinize. This condition is not only found in pregnant women who smoke actively. Pregnant women who don't smoke have an average nicotine level of 153.2 ± 96.0 $\mu\text{g/g}$ keratinize (>143 $\mu\text{g/g}$ keratinize) as a result of exposure to cigarette smoke from the environment (passive smoking)(Huang et al., 2017).There was a relationship between passive smoking mothers and the incidences of LBW (OR = 3.04) (Ardelia et al., 2019).

CONCLUSION

Most babies are born at a normal weight. There was a relationship between pregnancy complications and the incidences of LBW in Central Java. The quality of antenatal care, iron supplementation, and maternal smoking status were not related with the incidences of LBW. Pregnant woman who don't get iron supplementation (OR = 2.474) and complications of pregnancy (OR = 4.869) had chances of 0.84 for having a baby with LBW.The pregnant

women who experience signs of complications immediately take action and visit health facilities for further assistance. Pregnant women are also advised to take Fe tablets regularly at least 90 tablets during their pregnancy.

ACKNOWLEDGMENTS

Thank you to the United States Agency for International Development for providing access to secondary data for the 2017 IDHS through the dhsprogram.com website, Diponegoro University for facilitating us in accessing various e-journal portals, and the Health Research Ethics Committee, Faculty of Public Health, Diponegoro University. has issued a permit for the ethical feasibility of this research.

REFERENCES


- Ardelia, K.I.A., Hardianto, G., Nuswantoro, D., 2019. Passive smoker during pregnancy is a risk factor of low birth weight. *Maj. Obstet. Ginekol.* 27, 12. <https://doi.org/10.20473/mog.v1i12019.12-16>
- Bappenas, UNICEF, 2017. *Laporan Baseline SDG tentang Anak-Anak di Indonesia*. Jakarta.
- Bener, A., Salameh, K.M.K., Yousafzai, M.T., Saleh, N.M., 2012. Pattern of Maternal Complications and Low Birth Weight: Associated Risk Factors among Highly Endogamous Women. *ISRN Obstet. Gynecol.* 2012, 1–7. <https://doi.org/10.5402/2012/540495>
- BKKBN, BPS, Kemenkes, International, I., 2017. *Indonesia Demographic and Health Survey 2017*. DHS Program, Jakarta, Indonesia.
- Chhabra, S., Chopra, S., 2016. Mid Pregnancy Fetal Growth Restriction and Maternal Anaemia a Prospective Study. *J. Nutr. Disord. Ther.* 6. <https://doi.org/10.4172/2161-0509.1000187>
- Darwis, A., Abdullah, A., Maidar, Adamy, A., Septiani, R., Nurjannah, 2020. The Relationship Between Service Quality Antenatal Care and Low Birth Weight in Indonesia: IDHS in 2017. *Adv. Heal. Sci. Res.* 22, 337–340. <https://doi.org/10.2991/ahsr.k.200215.064>
- Dinas Kesehatan Provinsi Jawa Tengah, 2019. *Profil Kesehatan Provinsi Jawa Tengah Tahun 2019*, Dinkes Jateng.
- Fang, F., Luo, Z.C., Dejemli, A., Delvin, E., Zhang, J., 2015. Maternal smoking and metabolic health biomarkers in newborns. *PLoS One*. <https://doi.org/10.1371/journal.pone.0143660>
- Figueiredo, A.C.M.G., Gomes-Filho, I.S., Silva, R.B., Pereira, P.P.S., Da Mata, F.A.F., Lyrio, A.O., Souza, E.S., Cruz, S.S., Pereira, M.G., 2018. Maternal anemia and low birth weight: A systematic review and meta-analysis. *Nutrients* 10, 1–17. <https://doi.org/10.3390/nu10050601>
- Huang, S., Weng, K., Huang, S., Liou, H., 2017. The effects of maternal smoking exposure during pregnancy on postnatal outcomes : A cross sectional study. *J. Chinese Med. Assoc.* 1–7. <https://doi.org/10.1016/j.jcma.2017.01.007>
- Kataoka, M.C., Paula, A., Carvalheira, P., Ferrari, A.P., Antonieta, M., Leite, D.B., Maria, C., Lima, G. De, 2018. Smoking during pregnancy and harm reduction in birth weight : a cross-sectional study. *BMC Pregnancy Childbirth* 1–10.
- Kementerian Kesehatan RI Badan Penelitian dan Pengembangan, 2018. *Hasil Utama Riset Kesehatan Dasar 2018*, Kementrian Kesehatan Republik Indonesia. Jakarta, Indonesia.
- Ko, T.J., Tsai, L.Y., Chu, L.C., Yeh, S.J., Leung, C., Chen, C.Y., Chou, H.C., Tsao, P.N., Chen, P.C., Hsieh, W.S., 2014. Parental smoking during pregnancy and its association with low birth weight, small for gestational age, and preterm birth offspring: A birth cohort study. *Pediatr. Neonatol.* 55, 20–27. <https://doi.org/10.1016/j.pedneo.2013.05.005>
- Kujariningrum, O.B., Winarni, S., Najib, N., 2021. Multiple pregnancy : the biggest risk factor of low birth weight in Central Java , Indonesia (2017 IDHS secondary data study). *Ann. Trop. Med. Public Heal.* 24, 24–184. <https://doi.org/10.36295/ASRO.2021.24184>
- Kuribayashi, M., Tsuda, H., Ito, Y., Tezuka, A., Ando, T., Tamakoshi, K., Mizuno, K., 2021. Evaluation of the risk factors for antepartum hemorrhage in cases of placenta previa: a retrospective cohort study. *J. Int. Med. Res.* 49, 1–9. <https://doi.org/10.1177/03000605211054706>
- Long, H., Yi, J.M., Hu, P.L., Li, Z. Bin, Qiu, W.Y., Wang, F., Zhu, S., 2012. Benefits of Iron supplementation for low birth weight infants: A systematic review. *BMC Pediatr.* 12. <https://doi.org/10.1186/1471-2431-12-99>
- Marshall, N.E., Abrams, B., Barbour, L.A., Catalano, P., Christian, P., Friedman, J.E., Hay, W.W., Hernandez, T.L., Krebs, N.F., Oken, E., Purnell, J.Q., Roberts, J.M., Soltani, H., Wallace, J., Thornburg, K.L., 2022. The importance of nutrition in pregnancy and lactation: lifelong consequences. *Am. J. Obstet. Gynecol.* 226, 607–

632. <https://doi.org/10.1016/j.ajog.2021.12.035>
- Meiriza, W., Aladin, A., Edison, E., 2018. The Correlation of Maternal Factors and The Quality of Antenatal Care Services With Low Birth Weight Babies In Health Facilities Level I. *J. Midwifery* 3, 103. <https://doi.org/10.25077/jom.1.1.103-114.2018>
- Miranti, Mutiarasari, D., Arsin, A.A., Hadju, V., Mallongi, A., Nur, R., Amri, I., Haruni, H., Wahyuni, R.D., Rahma, Faris, A., 2020. Determinants of the incidence of stunting in the working area of Kinovaro Sigi Health Center. *Enfermeria Clínica* 30, 246–252. <https://doi.org/10.1016/j.enfcli.2019.10.077>
- Neiger, R., 2017. Long-Term Effects of Pregnancy Complications on Maternal Health: A Review. *J. Clin. Med.* 6, 76. <https://doi.org/10.3390/jcm6080076>
- Nemoto, T., Ando, H., Nagao, M., Kakinuma, Y., Sugihara, H., 2021. Prenatal Nicotine Exposure Induces Low Birthweight and Hyperinsulinemia in Male Rats. *Front. Endocrinol. (Lausanne)*. 12, 1–11. <https://doi.org/10.3389/fendo.2021.694336>
- OECD, WHO, 2019. Health at a glance: Asia/Pacific 2018: Measuring progress towards universal health coverage. OECD Publishing, Paris.
- Owa, K., 2019. Chronic Energy Deficiency , Anemia as a Risk Factor for Low Birth Weight Babies in East Nusa Tenggara. *J. Kesehat. Prim.* 4, 13–22.
- Owa, K., Putra, I.W.G.A.E., Windiani, I.G.A.T., 2017. Risk factors for low birth weight infants in East Nusa Tenggara. *Public Heal. Prev. Med. Arch.* 5, 39. <https://doi.org/10.15562/phpma.v5i1.40>
- Paul, P., Zaveri, A., Chouhan, P., 2019. Assessing the Impact of Antenatal Care Utilization on Low Birthweight in India: Analysis of the 2015–2016 National Family Health Survey. *Child. Youth Serv. Rev.* 106, 104459. <https://doi.org/10.1016/j.childyouth.2019.104459>
- Phowira, J., Elvina, F.T., Wiguna, I.I., Wahyudi, F.R.H.B., Medise, B.E., 2020. The Association between Tobacco Exposure during Pregnancy and Newborns' Birth Weight in DKI Jakarta Community Members. *medRxiv* 1–15. <https://doi.org/10.1101/2020.10.29.20222059>
- Restu, S., Dasuki, D., Nurdianti Z, R.D.S., 2014. The influence of iron supplementation in pregnant women to the occurrence of low birth weight (LBW) babies in Palu, Central Sulawesi. *J. thee Med. Sci. (Berkala Ilmu Kedokteran)* 46, 41–51. <https://doi.org/10.19106/jmedscie004601201406>
- Sarah, S., Irianto, I., 2018. Pengaruh Tingkat Kepatuhan Minum Tablet Fe Terhadap Kejadian Anemia Pada Ibu Hamil Trimester III di Puskesmas Pejeruk Tahun 2017. *Yars. Med. J.* 26, 75. <https://doi.org/10.33476/jky.v26i2.392>
- Seu, M.M.V., Mose, J.C., Panigoro, R., Sahiratmadja, E., 2019. Anemia Prevalence after Iron Supplementation among Pregnant Women in Midwives Practice of Primary Health Care Facilities in Eastern Indonesia. *Hindawi* 2019, 1–8. <https://doi.org/10.1155/2019/1413906>
- Sharami, S.H., Darkhaneh, R.F., Zahiri, Z., Milani, F., Asgharnia, M., Shakiba, M., Didar, Z., 2013. The relationship between vaginal bleeding in the first and second trimester of pregnancy and preterm labor. *Int. J. Reprod. Biomed.* 11, 385–390.
- Siramaneerat, I., Agushyana, F., Meebunmak, Y., 2018. Maternal Risk Factors Associated with Low Birth Weight in 376–383. <https://doi.org/10.2174/1874944501811010376>
- Sulistiyani, S., Darmawati, A.T., Setiani, O., 2019. Correlation between local-specific traditions of women agricultural workers and the Incidence of low birth weight in South Metro Sub-District, Metro City, Lampung Province, Indonesia. *Ann. Trop. Med. Public Heal.* 22, 1–5. <https://doi.org/10.36295/ASRO.2019.221118>
- Upadhyay, R.P., Naik, G., Choudhary, T.S., Chowdhury, R., Taneja, S., Bhandari, N., Martines, J.C., Bahl, R., Bhan, M.K., 2019. Cognitive and Motor Outcomes in Children Born Low Birth Weight : A Systematic Review and Meta-analysis of Studies from South Asia. *BMC Pediatr.* 19, 1–15.
- WHO, UNICEF, 2019. Low Birthweight Estimates: Level and Trends 2000-2015, UNICEF and WHO.

#32644 Review


[SUMMARY](#) [REVIEW](#) [EDITING](#)

Submission

Authors Oktavia Beni Kujariningrum, Sri Winarni, Atik Mawarni, Najib Najib 

Title Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java

Section Articles

Editor Efa Nugroho, S.K.M, M.Kes 

Peer Review

Round 1

Review Version 32644-83650-1-RV.DOCX 2021-10-11



Initiated —

Last modified —

Uploaded file None

Editor Decision

Decision Accept Submission 2022-11-07

Notify Editor  Editor/Author Email Record  2022-08-21

Editor Version 32644-91924-1-ED.DOCX 2022-04-12
32644-91924-2-ED.DOCX 2022-08-31
32644-91924-3-ED.DOCX 2022-09-05

Author Version 32644-91956-1-ED.DOCX 2022-04-17 [DELETE](#)
32644-91956-2-ED.DOCX 2022-08-31 [DELETE](#)
32644-91956-3-ED.DOCX 2022-09-27 [DELETE](#)

Upload Author Version No file chosen

ISSN: 2355-3596

ABOUT THE JOURNAL

[Focus and Scope](#)

[Manuscript Submission](#)

[Guide for Authors](#)

[Editorial Board](#)

[Reviewer Team](#)

[Abstracting/Indexing](#)

[Ethics Statement](#)

[Policy of Screening for Plagiarism](#)

[Contact](#)
2,143,152
[View Visitor Stats](#)

USER

You are logged in as...

oktaviabeni66

» [My Journals](#)

» [My Profile](#)

» [Log Out](#)

JOURNAL CONTENT

Search

Search Scope

[Browse](#)

» [By Issue](#)

» [By Author](#)

» [By Title](#)

» [Other Journals](#)

COLABORATION WITH



Ikatan Ahli Kesehatan Masyarakat Indonesia

IAKMI (The Indonesian Public Health Association) is an independent professional organization for the betterment of public health, based on Pancasila based on the 1945 Constitution. Mutual Agreement No: 402/UN.37.1.6/IKM/2012




Jejaring Nasional Pendidikan Kesehatan (JNPK)

JNPK is an organization that gathers experts and observers in the field of health education, which was established on September 1, 2014. The founder of this organization is the University of Teacher Training Education Institut Teknologi Sepuluh Nopember (ITS) which organizes public health education, namely Universitas Negeri

#32644 Review


[SUMMARY](#) [REVIEW](#) [EDITING](#)

Submission

Authors Oktavia Beni Kujariningrum, Sri Winarni, Atik Mawarni, Najib Najib 

Title Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java

Section Articles

Editor Efa Nugroho, S.K.M, M.Kes 

Peer Review

Round 1

Review Version 32644-83650-1-RV.DOCX 2021-10-11



Initiated —

Last modified —

Uploaded file None

Editor Decision

Decision Accept Submission 2022-11-07

Notify Editor  Editor/Author Email Record  2022-08-21

Editor Version 32644-91924-1-ED.DOCX 2022-04-12
32644-91924-2-ED.DOCX 2022-08-31
32644-91924-3-ED.DOCX 2022-09-05

Author Version 32644-91956-1-ED.DOCX 2022-04-17 [DELETE](#)
32644-91956-2-ED.DOCX 2022-08-31 [DELETE](#)
32644-91956-3-ED.DOCX 2022-09-27 [DELETE](#)

Upload Author Version No file chosen

ISSN: 2355-3596

ABOUT THE JOURNAL

[Focus and Scope](#)

[Manuscript Submission](#)

[Guide for Authors](#)

[Editorial Board](#)

[Reviewer Team](#)

[Abstracting/Indexing](#)

[Ethics Statement](#)

[Policy of Screening for Plagiarism](#)

[Contact](#)
2,143,152
[View Visitor Stats](#)

USER

You are logged in as...
oktaviabeni66

- » [My Journals](#)
- » [My Profile](#)
- » [Log Out](#)

JOURNAL CONTENT

Search

Search Scope

Browse

- » [By Issue](#)
- » [By Author](#)
- » [By Title](#)
- » [Other Journals](#)

COLABORATION WITH



Ikatan Ahli Kesehatan Masyarakat Indonesia

IAKMI (The Indonesian Public Health Association) is an independent professional organization for the betterment of public health, based on Pancasila based on the 1945 Constitution. Mutual Agreement No: 402/UN.37.1.6/IKM/2012



Jejaring Nasional Pendidikan Kesehatan (JNPK)

JNPK is an organization that gathers experts and observers in the field of health education, which was established on September 1, 2014. The founder of this organization is the University of Teacher Training Education Institut Teknologi Sepuluh Nopember (ITS) which organizes public health education, namely Universitas Negeri

Administrasi Publikasi 18.3 Januari 2023

Jurnal Kemas <kemas@mail.unnes.ac.id>

Wed, Nov 9, 2022 at 10:50 PM

To: LUBERTA.RD@gmail.com, Diffah Hanim <diffah@staff.uns.ac.id>, donoIND323@gmail.com, intan mutiaraputri <intan.mutiaraputri@unisayogya.ac.id>, sri.handayani@dsn.dinus.ac.id, titisristi98@gmail.com, andarini.ea@gmail.com, noersaudah15@gmail.com, ns.indah@yahoo.com, humeyra.dewi@gmail.com, cehazetabidin@gmail.com, yudisadiaz@gmail.com, henrisetiawan1989@gmail.com, ri.syahniar@umj.ac.id, Farsida Farsida Farsida <farsida@umj.ac.id>, nadyalutfi3095@gmail.com, Raestafebrianti@gmail.com, mulyani1898@gmail.com, rahmini@umj.ac.id, Mirna Widiyanti <ninawidhy@gmail.com>, tyoadining@gmail.com, eviiriani83@gmail.com, tongatmania@gmail.com, vandrastea@ymail.com, winarni@live.undip.ac.id, oktaviabeni66@gmail.com, atikm246@gmail.com, Najib Hn <jibpenkb@gmail.com>

Yth. Penulis di Jurnal KEMAS

Dengan hormat,

Kami beritahukan bahwa artikel Bapak/ Ibu dengan:

1. No. 32899 Judul: Level of Knowledge and Self Efficacy Improve Breast Self-Examination (BSE) Behaviours
2. No. 24782 Judul: The Association Between Family Support and Physical Activity with Cognitive Function among Older Adults Aged 60-79 Years Old
3. No. 36701 Judul: Sri Handayani, Ririn Nurmandhani, Edi Jaya Kusuma, Sadono Wiwoho
4. No. 38234 Judul: Peran Pengetahuan dan Persepsi terhadap Penggunaan Kontrasepsi Modern Pria di Indonesia
5. No. 36740 Judul: Communication, Information, and Education on Self Awareness about Marriage Age Maturation among Adolescent Girls
6. No. 35343 Judul: Relationship between nutritional status and living conditions with the risk of tuberculosis in children
7. No. 34942 Judul: The Utilization of Voluntary Counseling and Testing by Women of Reproductive Age in West Papua
8. No. 32644 Judul: Analysis of Maternal Predisposing Factors with The Incidence of Low Birth Weight in Central Java

akan segera kami proses untuk penerbitan di Jurnal KEMAS (volume 18.3 Januari 2023).

Jika dalam 7 hari kerja belum merespon email ini, maka redaksi berhak memundurkan penerbitan artikel ke edisi berikutnya

Sesuai dengan syarat administratif, bagi artikel yang akan dimuat:

1. Penulis dikenakan biaya publish dan translet/proofread sebesar Rp. 2.500.000,00 dengan ketentuan sebagai berikut:
 - a. Biaya tersebut dikirim ke Jurnal KEMAS ke no rek. BNI 0328641658 a.n Widya Hary Cahyati untuk mendapatkan surat LAYAK MUAT.
 - b. Bukti transfer harap dikirim ke email : kemas@mail.unnes.ac.id
 - c. Apabila penulis menghendaki versi cetak, maka bisa melakukan konfirmasi ke Redaksi KEMAS dan mengganti biaya cetak Rp 200.000/ eksemplar.

Kenaikan biaya pada edisi ini dikarenakan mulai edisi ini (vo. 18, no. 3) **jurnal KEMAS sudah terindeks scopus.**

2. Penulis menyertakan Surat Bebas Plagiat dan Surat Kesediaan Membayar Biaya Submit, dan dikirim melalui email kemas@mail.unnes.ac.id (Form terlampir, mohon bubuhkan materai 10.000 pada kedua form tersebut).

Demikian pemberitahuan kami, atas perhatian dan kerjasamanya kami ucapkan terimakasih.

Ketua Dewan Redaksi

Prof. Dr. dr. Oktia Woro K.H., M.Kes.

KEMAS Journal

F5 Building, 2nd Floor, Public Health Department, Sport Science Faculty, Semarang State University, Semarang,
Central Java, Indonesia, 50229

<http://journal.unnes.ac.id/nju/index.php/kemas>

DISCLAIMER

This electronic mail and/ or any files transmitted with it may contain confidential or copyright information of Universitas Negeri Semarang and/ or its Subsidiaries. If you are not an intended recipient, you must not keep, forward, copy, use, or rely on this electronic mail, and any such action is unauthorized and prohibited. If you have received this electronic mail in error, please reply to this electronic mail to notify the sender of its incorrect delivery, and then delete both it and your reply. Finally, you should check this electronic mail and any attachments for the presence of viruses. Universitas Negeri Semarang accepts no liability for any damages caused by any viruses transmitted by this electronic mail.

Surat Bebas Plagiat & Kesediaan Membayar

Jurnal Kemas <kemas@mail.unnes.ac.id>

Thu, Nov 10, 2022 at 3:40 PM

To: LUBERTA.RD@gmail.com, Diffah Hanim <diffah@staff.uns.ac.id>, donoIND323@gmail.com, intan mutiaraputri <intan.mutiaraputri@unisayogya.ac.id>, sri.handayani@dsn.dinus.ac.id, titisristi98@gmail.com, andarini.ea@gmail.com, noersaudah15@gmail.com, ns.indah@yahoo.com, humeyra.dewi@gmail.com, cehazetabidin@gmail.com, yudisadiaz@gmail.com, henrisetiawan1989@gmail.com, Rike Syahniar <ri.syahniar@umj.ac.id>, Farsida Farsida Farsida <farsida@umj.ac.id>, nadyalutfi3095@gmail.com, Raestafebrianti@gmail.com, mulyani1898@gmail.com, rahmini@umj.ac.id, Mirna Widiyanti <ninawidhy@gmail.com>, tyoadining@gmail.com, eviiriani83@gmail.com, tongatmania@gmail.com, vandrastea@gmail.com, winarni@live.undip.ac.id, oktaviabeni66@gmail.com, atikm246@gmail.com, Najib Hn <jibpenkb@gmail.com>

Berkut kami kirimkan form Surat bebas plagiat & kesediaan membayar.
Mohon setelah diisi bisa dikirimkan kembali via email.
Terima kasih

KEMAS Journal

F5 Building, 2nd Floor, Public Health Department, Sport Science Faculty, Semarang State University, Semarang, Central Java, Indonesia, 50229

<http://journal.unnes.ac.id/nju/index.php/kemas>

DISCLAIMER

This electronic mail and/ or any files transmitted with it may contain confidential or copyright information of Universitas Negeri Semarang and/ or its Subsidiaries. If you are not an intended recipient, you must not keep, forward, copy, use, or rely on this electronic mail, and any such action is unauthorized and prohibited. If you have received this electronic mail in error, please reply to this electronic mail to notify the sender of its incorrect delivery, and then delete both it and your reply. Finally, you should check this electronic mail and any attachments for the presence of viruses. Universitas Negeri Semarang accepts no liability for any damages caused by any viruses transmitted by this electronic mail.



SURAT PERNYATAAN BEBAS PLAGIAT dan KESEDIAAN PEMBAYARAN BIAYA SUBMIT-2020.doc
33K



Sri Winarni <winarni@live.undip.ac.id> Mon, Nov 14, 2022 at 4:43 PM
To: kemas@mail.unnes.ac.id

Selamat sore

Berikut saya lampirkan bukti transfer biaya publikasi artikel No. 32644 dengan judul "Analysis of Maternal Predisposing Factors with The Incidence of Low Birth Weight in Central Java".

Terima kasih





Sri Winarni <winarni@live.undip.ac.id>

Surat Bebas Plagiat & Kesediaan Membayar_No. 32644

Sri Winarni <winarni@live.undip.ac.id>
To: kemas@mail.unnes.ac.id

Tue, Nov 15, 2022 at 9:27 AM

Selamat pagi

Berikut saya lampirkan surat bebas plagiasi dan kesediaan membayar untuk artikel No. 32644 dengan judul "Analysis of Maternal Predisposing Factors with The Incidence of Low Birth Weight in Central Java".

Terima kasih



Surat Bebas Plagiasi dan Kesediaan Membayar.pdf
2689K



Sri Winarni <winarni@live.undip.ac.id>

Surat Bebas Plagiat & Kesediaan Membayar_No. 32644

Jurnal Kemas <kemas@mail.unnes.ac.id>
To: Sri Winarni <winarni@live.undip.ac.id>

Mon, Nov 21, 2022 at 4:18 PM

Berikut kami kirimkan LoA nya.
Terima kasih

KEMAS Journal

F5 Building, 2nd Floor, Public Health Department, Sport Science Faculty, Semarang State University, Semarang,
Central Java, Indonesia, 50229
<http://journal.unnes.ac.id/nju/index.php/kemas>

[Quoted text hidden]

DISCLAIMER

This electronic mail and/ or any files transmitted with it may contain confidential or copyright information of Universitas Negeri Semarang and/ or its Subsidiaries. If you are not an intended recipient, you must not keep, forward, copy, use, or rely on this electronic mail, and any such action is unauthorized and prohibited. If you have received this electronic mail in error, please reply to this electronic mail to notify the sender of its incorrect delivery, and then delete both it and your reply. Finally, you should check this electronic mail and any attachments for the presence of viruses. Universitas Negeri Semarang accepts no liability for any damages caused by any viruses transmitted by this electronic mail.



LoA Januari 2023_Sri Winarni.pdf
27K



REDAKSI JURNAL KEMAS
JURUSAN ILMU KESEHATAN MASYARAKAT
FAKULTAS ILMU KEOLAHRAHAGAN
UNIVERSITAS NEGERI SEMARANG
Kampus Sekaran Gunungpati Semarang 50229
Telp (024) 8508107 Fax (024) 8508007
Laman: <http://journal.unnes.ac.id/nju/index.php/kemas>,
[e-mail: kemas@mail.unnes.ac.id](mailto:kemas@mail.unnes.ac.id)

Nomor : 6/KEMAS/XI/2022
Lampiran : -
Perihal : **Artikel Layak Muat**


Yth:

Bapak/ Ibu Oktavia Beni Kujariningrum, Sri Winarni, Atik Mawarni, Najib

Berdasarkan artikel Sejawat yang diajukan ke Redaksi Jurnal KEMAS dengan judul:

Analysis of Maternal Predisposing Factors with The Incidence of Low Birth Weight in Central Java

Bersama ini kami sampaikan bahwa hasil penilaian dari dewan redaksi, artikel Sejawat layak untuk dimuat di Jurnal KEMAS Volume 18 No.3 (Januari 2023).
Atas perhatian dan kerjasama yang baik diucapkan terima kasih.

Semarang, 21 November 2022
Ketua Dewan Redaksi,

Prof. Dr. dr. Oktia Woro K.H., M.Kes.
NIP. 19591001 198703 2 001

Analysis of Maternal Predisposing Factors with The Incidence of LBW in Central Java

Oktavia Beni Kujariningrum⁽¹⁾, Sri Winarni⁽²⁾, Atik Mawarni⁽³⁾, Najib Najib⁽⁴⁾,

DOI: <https://doi.org/10.15294/kemas.v18i3.32644>

(1) Biostatistics and Population, Faculty of Public Health, Diponegoro University
(2) Biostatistics and Population, Faculty of Public Health, Diponegoro University
(3) Biostatistics and Population, Faculty of Public Health, Diponegoro University
(4) National Innovation Research Agency (BRIN)

Abstract

Abstract. In Central Java, the prevalence of LBW (Low Birth Weight) has increased from 4.3 (2018) to 4.7 (2019) and be the biggest cause of neonatal mortality (46.4%) and infant mortality (40.5%). The aim of this research is to analyze the relationship between quality of ANC (Antenatal Care), iron supplementation, pregnancy complications, and maternal smoking status with LBW in Central Java. This research was an analytical study used secondary data from the 2017 IDHS (Indonesian Demographic and Health Survey). The sampling design used purposive sampling. Population study was 1205 babies born in Central Java. The sample comprised 952 babies. Independent variables were the quality of ANC, iron supplementation, pregnancy complications, and maternal smoking status, with the incidence of LBW as dependent variable. Data analysis was performed by chi-square continuity correction and logistic regression. Pregnancy complications have been associated with the incidences of LBW in Central Java (p -value = 0.0001). Iron supplementation (OR = 2.474) and pregnancy complications (OR = 4.869) had an effect on the incidence of LBW in Central Java. Iron supplementation and pregnancy complications influenced the incidence of LBW in Central Java.

Keywords

LBW; Pregnancy Complications; Iron Supplementation



?	Total citations
?	Recent citations
n/a	Field Citation Ratio
n/a	Relative Citation Ratio

Full Text:

PDF

References

- Ardelia, K.I.A., Hardianto, G., Nuswantoro, D., 2019. Passive smoker during pregnancy is a risk factor of low birth weight. *Maj. Obstet. Ginekol.* 27, 12. <https://doi.org/10.20473/mog.v1i12019.12-16>
- Bappenas, UNICEF, 2017. Laporan Baseline SDG tentang Anak-Anak di Indonesia. Jakarta.
- BKKBN, BPS, Kemenkes, International, I., 2017. Indonesia Demographic and Health Survey 2017. DHS Program, Jakarta, Indonesia.
- Chhabra, S., Chopra, S., 2016. Mid Pregnancy Fetal Growth Restriction and Maternal Anaemia a Prospective Study. *J. Nutr. Disord. Ther.* 06. <https://doi.org/10.4172/2161-0509.1000187>
- Darwis, A., Abdullah, A., Maidar, Adamy, A., Nurjannah, 2020. Hubungan Komponen Pelayanan Antenatal Care (10T) dengan Kejadian Bayi Berat Lahir Rendah di Indonesia (Analisa Data Sekunder SDKI 2017). *JUKEMA* 6, 13–19.
- Dinas Kesehatan Provinsi Jawa Tengah, 2019. Profil Kesehatan Provinsi Jawa Tengah Tahun 2019, Dinkes Jateng.
- Dirjen Kesmas Kemenkes, 2018. Laporan Kinerja Ditjen Kesehatan Masyarakat Tahun 2017.
- Fang, F., Luo, Z.C., Dejemli, A., Delvin, E., Zhang, J., 2015. Maternal smoking and metabolic health biomarkers in newborns. *PLoS One*. <https://doi.org/10.1371/journal.pone.0143660>
- Huang, Shih-hui, Weng, K., Huang, Shih-ming, Liou, H., 2017. The effects of maternal smoking exposure during pregnancy on postnatal outcomes : A cross sectional study. *J. Chinese Med. Assoc.* 1–7. <https://doi.org/10.1016/j.jcma.2017.01.007>
- Isnaini, N., 2015. Analisis Faktor Yang Mempengaruhi Kejadian Bayi Berat Lahir Rendah. *J. Dunia Kesmas* 4, 196–202.
- Karjatin, A., 2016. Keperawatan Maternitas. Pusdik SDM Kesehatan, Jakarta.
- Kataoka, M.C., Paula, A., Carvalheira, P., Ferrari, A.P., Antonieta, M., Leite, D.B., Maria, C., Lima, G. De, 2018. Smoking during pregnancy and harm reduction in birth weight : a cross-sectional study. *BMC Pregnancy Childbirth* 1–10.
- Kementerian Kesehatan RI Badan Penelitian dan Pengembangan, 2018. Hasil Utama Riset Kesehatan Dasar 2018, Kementrian Kesehatan Republik Indonesia. Jakarta, Indonesia.
- Ko, T.J., Tsai, L.Y., Chu, L.C., Yeh, S.J., Leung, C., Chen, C.Y., Chou, H.C., Tsao, P.N., Chen, P.C., Hsieh, W.S., 2014. Parental smoking during pregnancy and its association with low birth weight, small for gestational age, and preterm birth offspring: A birth cohort study. *Pediatr. Neonatol.* 55, 20–27. <https://doi.org/10.1016/j.pedneo.2013.05.005>
- Kujariningrum, O.B., Winarni, S., Najib, N., 2021. Multiple pregnancy : the biggest risk factor of low birth weight in Central Java , Indonesia (2017 IDHS secondary data study). *Ann. Trop. Med. Public Heal.* 24, 24–184. <https://doi.org/10.36295/ASRO.2021.24184>
- Kusumawati, D.D., Septiyaningsih, R., Kania, 2016. Faktor-Faktor Ibu yang Mempengaruhi Kejadian Bayi Berat Lahir Rendah (BBLR). *J. Kesehat. Al-Irsyad* 9, 8–16.

ARTICLE TOOLS

- Print this article
- Indexing metadata
- How to cite item
- Supplementary files
- Finding References
- Email this article
- Email the author

ABOUT THE JOURNAL

- Focus and Scope
- Manuscript Submission
- Guide for Authors
- Editorial Board
- Reviewer Team
- Abstracting/Indexing
- Ethics Statement
- Policy of Screening for Plagiarism
- Contact
- 2,143,144
- View Visitor Stats

USER

- You are logged in as...
- 1waviq_1**
- » My Journals
 - » My Profile
 - » Log Out

JOURNAL CONTENT

Search

Search Scope

All

Search

Browse

- » By Issue
- » By Author
- » By Title
- » Other Journals

COLABORATION WITH



Ikatan Ahli Kesehatan Masyarakat Indonesia

IAKMI (The Indonesian Public Health Association) is an independent professional organization for the benefit of public health, based on Pancasila based on the 1945 Constitution. Mu

Agreement No:
402/UN.37.1.6/IKM/2012

