

Abdominal Diameter Profiles have Relationship with Insulin Resistance in Obese Female Adolescents

by Fahmy Arif Tsani

Submission date: 01-Sep-2020 02:40PM (UTC+0700)

Submission ID: 1377479914

File name: tionship_with_Insulin_Resistance_in_Obese_Female_Adolescents.pdf (265.27K)

Word count: 4430

Character count: 24340

Abdominal Diameter Profiles have Relationship with Insulin Resistance in Obese Female Adolescents

Fillah Fithra Diény^{1,2*}, Ria Fitri Setyaningsih¹, Deny Yudi Fitranti^{1,2}, ⁸Idananda Fikri Jauharany¹,
Yehezkiel Dwi Putra¹, A Fahmy Arif Tsani^{1,2**}

¹Department of Nutrition Science, ¹⁴Faculty of Medicine, Universitas Diponegoro, Semarang, Central Java, INDONESIA

²Center of Nutrition Research, Faculty of Medicine, Universitas Diponegoro, Semarang, Central Java, INDONESIA

*Corresponding Author: fillahdieny@gmail.com

**Corresponding Author: fahmi_tsani@yahoo.com

Citation: Diény FF, Setyaningsih RF, Fitranti DY, Jauharany FF, Putra YD, Tsani AFA. Abdominal Diameter Profiles have Relationship with Insulin Resistance in Obese Female Adolescents. *Electron J Gen Med.* 2020;17(5):em219. <https://doi.org/10.29333/ejgm/7882>

ARTICLE INFO

Received: 8 Nov. 2019

Accepted: 12 Feb. 2020

ABSTRACT

Purpose: This study aimed to determine the relationship of abdominal diameter consist of Sagittal Abdominal Diameter (SAD), Ratio SAD/Height and Abdominal Diameter Index (ADI) with insulin resistance in obese female adolescents.

Method: This study was conducted in June-August 2019 at Universitas Diponegoro, Central Java. this study used cross sectional design with a total of 120 female students aged 17-21 years and selected using purposive sampling method. Study variables included SAD, SAD/Height, ADI and insulin resistance level. SAD was measured by abdominal caliper, SAD/Height was ratio of SAD divided by height. ADI was ratio of SAD divided by thigh circumference. The value of insulin resistance was obtained from the calculation of **homeostasis model assessment insulin resistance (HOMA-IR)**. Data normality was analyzed by Kolmogorov-smirnov test, all data were not normal distribution so the bivariate test analyzed by Rank spearman test.

Results: As many as 42.5% subjects had SAD which included at risk and 83.3% subjects had suffering insulin resistance. There was significant relationship between SAD, SAD/Height, ADI with insulin resistance (SAD (p=0.16), SAD/Height (p<0.001) and ADI (p=0.003). In addition, SAD, SAD/Height, ADI were significant relationship with **fasting blood glucose** (SAD (p=0.048), SAD/Height (p=0.022) and ADI (p=0.003).

Conclusion: There were significant relationship between SAD, SAD/Height, ADI with **insulin resistance and fasting blood glucose in obese adolescent female**.

Keywords: Sagittal Abdominal Diameter (SAD), SAD/Height, Abdominal Diameter Index (ADI), insulin resistance

INTRODUCTION

Obesity is one factor of that cause various diseases such as cardiovascular disease, hypertension, cancer, and diabetes mellitus. Most countries in Asia have increasing of overweight and obesity prevalence in this last decade. Increased of obesity prevalence does not only occurred in adult group but also occurred in adolescent group. Indonesia is the one of the countries that have high prevalence of abdominal obesity in the Southeast Asian region (1). The prevalence of visceral/abdominal obesity at age ≥ 15 years based on Basic Health Research 2013 as many as 26.6% and increase to 31% in 2018 (2,3). A study of adolescent in Semarang showed that 46 of 566 adolescents suffering abdominal obesity (4). The prevalence of abdominal obesity in female higher than male, the result data of Basic Health Research 2013 showed that female group who suffering abdominal obesity is 56.3% while the male group is 43.7% (5).

Abdominal obesity can reflect excess of visceral fat tissue or subcutaneous or both. Visceral fat contributes to the pathogenesis of diabetes mellitus, glucose intolerance and

insulin resistance. Insulin resistance is a impaired condition of glucose absorption that induced by insulin (6). Fat tissue distribution in intraabdominal area caused increasing metabolic such as lipogenesis and lipolysis activity. Lipolysis produce *free fatty acids* (FFA), FFA from visceral fat brought to the portal circulation and liver. Excessive of FFA secretion from visceral fat through the liver cause toxic effect for example gluconeogenesis and insulin resistance (6,7). Study of 220 obese children and adolescents in Brazil shows that 33.2% subjects suffering insulin resistance (8). Study of 52 obese adolescents in Semarang also shows that 91.6% subjects suffering insulin resistance (9).

Excessive of visceral fat tissue in abdominal obesity can measure by *Computed Tomography*(CT) as gold standard, but measurement of Waist Circumference (WC), and abdominal diameter also can be used as alternative. Abdominal diameter measurement consist of three components, namely Sagittal Abdominal Diameter (SAD), Sagittal Abdominal Diameter/Height (SAD/Height), Abdominal Diameter Index (ADI) (10). SAD can describe the visceral fat size, moreover it can also predicting metabolic syndrome from visceral fat area. SAD is closely related with visceral fat mass than other

3




Figure 1. The flow diagram of subject inclusion and exclusion in this research

anthropometric measurements. SAD measured by supine position when subcutaneous fat move to side of waist. SAD measurement at this position reflect size of intraabdominal fat in antero-posterior like in CT image measurement. SAD strongly related with glucose intolerance, cardiovascular disease and good indicator for predicting insulin resistance (10-13).

SAD/Height is derivative of SAD measurement, which is use height as divisor of SAD. SAD/Height is anthropometric indicator that significantly related to visceral fat tissue (10,14). ADI is derivative of SAD measurement divided by thigh circumference. ADI is significantly related to several indicators of metabolic syndrome, and maybe has significant relationship with insulin resistance (15). A study in Brazil also showed that SAD/Height and ADI is good indicator for measurement visceral fat tissue in women (16). Study on abdominal diameter (SAD, SAD/Height and ADI) as insulin resistance indicator has not been much done especially in Indonesia. Therefore, based on this background the researchers interest to identifying the relationship of abdominal diameter (SAD, SAD/Height and ADI) with insulin resistance in obese adolescent female.

METHODS

This study was conducted in June-August 2019 with female student subject at Diponegoro University Semarang. This study was an observational analytic with cross sectional design. As described in Figure 1, of 1260 participants from this study were selected by screening process which obtained 1260 subjects, after that 215 subjects were included to inclusion criteria and selected as many as 120 subjects who involved in this study using purposive sampling method. The inclusion criteria were female student in Diponegoro University Semarang, aged 17-21 years, having waist circumference >80 cm, not pregnant, not consuming alcohol, not consuming drug that can related to glucose and insulin level, willing to fast minimum 8 hours before taking blood sample. The exclusion criteria was the subject resign during the study. This study had received permission from the Health Research Ethics Committee with Number 373/EC/KEPK/FK UNDIP/III/2019.

The independent variables in this study were abdominal diameter consisting of SAD, SAD/Height, and ADI. SAD measured by abdominal caliper which is manually measurement. Measurement were conducted with supine position, subject relax on the table. One of caliper's arm was placed under the subject (iliac crest) and the other caliper's arm was slowly moving down on subject abdomen with normal breathing (14). SAD categorized as normal if <19.3 cm and at

Table 1. Subject characteristic

Variable	Min	Max	Median±SD
Age (year)	17	21	19± 0.8
Weight (kg)	47.80	107.40	66.60±11.38
Height (cm)	141.20	171.40	157.40±5.44
Sagittal Abdominal Diameter(cm)	14.90	26.10	18.68±2.12
Sagittal Abdominal Diameter/Height (ratio)	0.09	0.17	0.12±0.014
Abdominal Diameter Index/ADI	0.03	0.56	0.32±0.41
Fasting blood glucose(mg/dL)	68.00	206.00	87.00±13.15
Insulin (µU/mL)	2.71	52.60	10.82±6.92
HOMA-IR	0.54	18.32	2.33±2.09

risk if ≥19.3 cm (17). SAD/Height was derivative of SAD, which is use height (in centimeter) as SAD divisor. ADI was derivative of SAD measurement divided by thigh circumference (in centimeter). The thigh circumference was measured on the right side of the body, at the midpoint between inguinal crease and proximal border on the patella (16).

The dependent variable in this study was insulin resistance. Insulin resistance is a disruption of the biological response to insulin which cause body insulin requirement increase and then resulting hyperinsulinemia to maintain normal glucose level. Insulin resistance level obtained from calculating Homeostasis Model Assessment Insulin Resistance (HOMA-IR), with formula (18):

$$HOMA\ IR = \frac{fasting\ insulin\ \left(\frac{mU}{L}\right) \times fasting\ glucose\ \left(\frac{mmol}{L}\right)}{22.5} \quad (1)$$

Cut off the normal value of HOMA-IR in adolescent was <1.65 (9). Blood sample of subject was taken by laboratory officer. Before the blood was taken, the subject fasting for 8-12 hours and then blood was taken 5 cc.

Univariate data analysis was used to describe the characteristic of each variable in this study. Data normality test analyzed by Kolmogorov-smirnov test. Bivariate data analyzed by Rank Spearman test for abnormal data distribution. The purpose of bivariate analysis was to determine relationship of each abdominal diameter variable with insulin resistance.

RESULT

Subject Characteristic

Table 1 shows subject characteristic of this study. Age range of subjects were 17-21 years old. The highest fasting blood glucose of subject was 206 mg/dL while the lowest fasting blood glucose was 68 mg/dL. The median insulin level

Table 2. Frequency distribution of sagittal abdominal diameter, fasting blood glucose, fasting insulin and HOMA-IR

Characteristic	n	%
Sagittal Abdominal Diameter		
No risk	69	57.5
At risk	51	42.5
Fasting Blood Glucose		
Normal	117	97.5
High	3	2.5
Fasting Insulin		
Normal	107	89.2
High	13	10.8
HOMA-IR		
Normal	20	16.7
Resistance	100	83.3

Table 3. Relationship of abdominal diameter with insulin resistance and fasting blood glucose

Variable	Fasting blood glucose		HOMA-IR	
	r	p	r	p
Sagittal abdominal diameter	0.181	0.048	0.297	0.001
SAD/Height	0.209	0.022	0.345	<0.001
Abdominal diameter index	0.270	0.003	0.273	0.003

of subject was 0.82 μU/mL. The highest HOMA-IR value of subject was 18.32 with median value 2.33. The median in SAD measurement was 18.68 cm. The highest SAD measurement of subject was 26.10 cm.

Frequency distribution of sagittal abdominal diameter, fasting blood glucose, fasting insulin and HOMA-IR were presented in Table 2. Almost half of the subjects (42.5%) had SAD that categorized at risk. Majority of subjects (89.2%) had normal fasting insulin level. Almost of subjects (97.5%) had normal fasting blood glucose, whereas conversely most of subjects (83.3%) had suffering insulin resistance.

Relationship of Abdominal Diameter with Insulin Resistance

Bivariate analysis in this study used Rank Spearman test. Relationship of abdominal diameter with insulin resistant and fasting blood glucose were presented in Table 3. In this study, we found that SAD, SAD/Height and ADI had significant relationship with insulin resistance and fasting blood glucose (p<0.05), with positive correlation.

DISCUSSION

The result of this study shows that SAD median was 18.68±2.12cm with minimum value 14.90 cm and maximum value 26.10 cm. This result similar to study that conducted in Brazil by Feliciano et al. showed that in overweight adolescent the SAD measurement reached 26.3 cm (19). SAD is indicator that can describe visceral fat size (20). SAD measured by supine position, visceral abdominal fat tend to lift the abdominal wall toward sagittal abdominal subcutaneous fat pressure the stomach and subcutaneous fat moves to the waist, so SAD can reflect the visceral abdominal fat volume (21). As many as 42.5% subjects had SAD which categorized at risk. SAD in women closely related with hyperlipidemia, insulin resistance and cardiovascular risk, moreover it can also be used as a predictor of cardiometabolic risk score (22). The study of

children and adolescent in Arab shows that SAD significantly related with the severity of metabolic syndrome components (23).

The result of this study shows that HOMA-IR median was 2.33±2.09cm with minimum value 0.54 cm and maximum value 18.32 cm. The high score of HOMA-IR similar to González-Zavala et al.'s study that found high score of HOMA-IR in overweight adolescent subjects that reached 18.32 (24). This study showed that the HOMA-IR most of subjects (83.3%) included in insulin resistance category, however, almost of subjects had normal fasting blood glucose and normal fasting insulin. This result similar to Yi Kung Hye et al.'s study that found high prevalence of insulin resistant in obese adolescent subject was 47.1% (25). The study of 38 obese adolescent in Semarang also shows similar result with this study. This condition can occur because hyperglycemia is the last part from insulin resistance condition, which is long enough condition of impaired glucose intolerance and or impaired fasting blood glucose. Carbohydrate metabolism in adolescence also still good, so the body can maintaining normal glucose level (26). The high incidence of insulin resistance who suffering by subjects increased the risk of metabolic syndrome, because insulin resistance is a risk factor for metabolic syndrome that first arise (9). The highest HOMA-IR value in the subject was 18.32. Adolescent group with metabolic syndrome significantly had higher of HOMA-IR. HOMA-IR is indicator that strongly correlated with total and basal insulin. The HOMA-IR mean progressively increase with following sex maturation increase and weight from normal to obese. The study in India shows that the increase of HOMA-IR significantly related with body mass index, moreover there was significant different between HOMA-IR in adolescent with normal nutritional status, overweight and obesity (27).

Bivariate analysis in this study showed that SAD, SAD/Height, and ADI significant related to insulin resistance with positive correlation. A study of 75 overweight and obese adolescent showed that SAD and insulin resistance had significant relationship, with correlation that not much different from the study (r=0.485). The study also showed that SAD described a better correlation with insulin resistance than other anthropometric measurements. SAD can predict insulin resistance because it measures abdominal visceral fat than subcutaneous fat. Abdominal visceral fat also had strongly related with cardio metabolic disease (4). The study of women subject in Brazil comparing SAD with body mass index (BMI), waist circumference (WC), waist to hip ratio (WHR) as resistance insulin indicator, the result show that SAD and BMI was strongest indicator to indicate the occurrence of insulin resistance. BMI was widely used to identify metabolic risk but it can not be used alone, because it can not distinguish muscle adipose tissue or body fat distribution, whereas SAD had reported as a good marker for visceral adipose tissue in various ethnic group (21).

In contrast with SAD, study about relationship between SAD/Height ratio and ADI with insulin resistance still limited, but study of 194 subjects in Brazil shows that SAD/Height and ADI had significant related and positive correlation with visceral adipose tissue (16). A study of 1347 subjects in Korea showed that high visceral fat mass and low adiponectin level were correlated with increased risk of insulin resistance and β cell dysfunction. Obesity is well known as risk factor for insulin resistance. Abdominal obesity can cause insulin resistance by stimulating metabolic product formation which derived from

fat, hormone and cytokines. Insulin resistance can cause endothelial dysfunction and insulin signaling pathway changes. Several previous studies show that visceral adipose tissue can inhibit adiponectin secretion. Adiponectin mediates the insulin sensitivity system and glucose homeostasis (29). Insulin resistance is adaptive response of Free Fatty Acid (FFA) increase, and related with visceral adipose tissue level. FFA increase can change the liver, muscle and other tissue metabolism toward lipid deposition and oxidation. Insulin secretion increase offset the glucose capacity decrease. Subcutaneous adipose tissue taking FFA and storing excess calories in visceral adipose tissue (30). Excess FFA secretion from visceral fat will be oxidized, stored (lipids droplets) or metabolized to toxic derivatives (DAG and ceramides). Toxic derivatives cause insulin resistance, impaired cell function (lipotoxicity) or cause apoptosis. Apoptosis in the pancreas cause decrease and impairment of β cell capacities to secrete insulin which increase the risk of developing type 2 diabetes mellitus. The consequence of insulin resistance increase is endogenous glucose production by liver and glucose utilization decrease by peripheral tissue. As result are glycemia and insulin pancreas secretion increase, beside that occur hepatic insulin clearance impaired, and causing hyperinsulinemia (31).

This study also showed that significant relationship between abdominal diameter (SAD, SAD/Height, and ADI) with fasting blood glucose level. Abdominal diameter measurement can reflect visceral fat. Visceral/abdominal fat stores are considered as implication of insulin resistance, cardiovascular and other metabolic condition which related to type 2 diabetes mellitus. Excessive visceral fat related with impaired of insulin sensitivity. Visceral fat related with impaired glucose regulation, because it is closely related to insulin resistance (32). Lypolysis of visceral fat can cause free circulated of FFA in obese individual. This cause FFA enter to portal vein which can accumulated in the liver and cause impaired of glucose tolerance and insulin homeostasis. Visceral fat tissue and liver fat tissue related with insulin resistance in diabetic patient. A study of 297 subject showed that subject with isolated impaired fasting glucose and isolated impaired glucose tolerance had excess abdominal fat (visceral fat, abdominal subcutaneous fat and liver fat) (33).

CONCLUSION

As many as 42,5% subjects had SAD were categorized at risk, while 83,3% subject had suffering resistance insulin. There were significant relations between abdominal diameter (SAD, SAD/Height, ADI) with insulin resistance and fasting blood glucose in obese adolescent female, with positive correlation. As it is important to monitor the growth and development of adolescents over time, it is advisable to standardize the use of one measure of body fat location. Considering that abdominal fat, more than total fat, has been associated with cardiometabolic risk, it is recommended the use of abdominal diameter measurement (SAD, SAD/Height, ADI) as a measure that reflects the adipose tissue in this region, in the assessment of the nutritional status of adolescents.

ACKNOWLEDGEMENTS

This research was funded by the "Hibah Penelitian Dasar Unggulan Perguruan Tinggi (PDUPT) 2019, The Ministry of Research, Technology and Higher Education, Indonesia."

REFERENCES

- Ramachandran A, Snehalatha C. Rising burden of obesity in Asia. *Journal of Obesity*, 2010;1-7. <https://doi.org/10.1155/2010/868573> PMID:20871654 PMCID:PMC2939400
- Ministry of Health Republic of Indonesia. Riset Kesehatan 2013. Jakarta: Health Research and Development Agency Ministry of Health Republic of Indonesia. 2013.
- Ministry of Health Republic of Indonesia. Riset Kesehatan Dasar 2018. Jakarta: Health Research and Development Agency Ministry of Health Republic of Indonesia. 2018.
- Hanifah NID, Dieny FF. Hubungan Total Asupan Serat, Serat Larut Air (Soluble), Dan Serat Tidak Larut Air (Insoluble) Dengan Kejadian Sindrom Metabolik Pada Remaja Obesitas. *Journal of Nutrition College*, 2016;5(3):148-55. <https://doi.org/10.14710/jnc.v5i3.16385>
- Sudikno S, Syarif H, Dwiriani CM, Riyadi H. Faktor Risiko Obesitas Sentral Pada Orang Dewasa Umur 25-65 Tahun Di Indonesia (Analisis Data Riset Kesehatan Dasar 2013). *The Journal of Nutrition and Food Research*, 2016;38(2). <https://doi.org/10.22435/pgm.v38i2.5540.111-120>
- Ye J. Mechanisms of Insulin Resistance in Obesity. *Frontiers of Medicine in China*, 2013;7(1):14-24. <https://doi.org/10.1007/s11684-013-0262-6> PMID:23471659 PMCID:PMC3936017
- Westphal SA. Obesity, Abdominal Obesity, and Insulin Resistance. *Clinical Cornerstone*, 2008;9(1):23-31. [https://doi.org/10.1016/S1098-3597\(08\)60025-3](https://doi.org/10.1016/S1098-3597(08)60025-3)
- dos Santos Romualdo MC, de Nobrega FJ, Escrivao MAMS. Insulin Resistance in Obese Children and Adolescents. *Jornal de Pediatria*, 2014;90(6):600-7. <https://doi.org/10.1016/j.jped.2014.03.005> PMID:25019650
- Nuraini IS, Sulchan M, Dieny FF. Resistensi Insulin pada Remaja Stunted Obesity Usia 15-18 Tahun di Kota Semarang. *Journal of Nutrition College*, 2017;6(2):164-71. <https://doi.org/10.14710/jnc.v6i2.16906>
- Carneiro Roriz AK, Santana Passos LC, Cunha De Oliveira C, Eickemberg M, De Almeida Moreira P, Barbosa Ramos L. Anthropometric Clinical Indicators in the Assessment of Visceral Obesity: an Update. *Nutricion Clinica Dietetica Hospitalaria*, 2016;36(2):168-79. <https://doi.org/10.12873/362carneiroriz>
- Anjana M, Sandeep S, Deepa R, Vimalaswaran KS, Farooq S, Mohan V. Visceral and Central Abdominal Fat and Anthropometry in Relation to Diabetes in Asian Indians. *Diabetes Care*, 2004;27(12):2948-53. <https://doi.org/10.2337/diacare.27.12.2948> PMID:15562212
- Firouzi SA, Tucker LA, LeCheminant JD, Bailey BW. Sagittal abdominal diameter, waist circumference, and BMI as predictors of multiple measures of glucose metabolism: An NHANES investigation of US adults. *Journal of Diabetes Research*, 2018:1-14. <https://doi.org/10.1155/2018/3604108> PMID:30018985 PMCID:PMC6029495

13. Gletsu-Miller N, Kahn HS, Gasevic D, Liang Z, Frediani JK, Torres WE, et al. Sagittal abdominal diameter and visceral adiposity: Correlates of beta-cell function and dysglycemia in severely obese women. *Obesity Surgery*, 2013;23(7):874-81. <https://doi.org/10.1007/s11695-013-0874-6> PMID:23408092 PMCID:PMC3750719
14. Khan HS, Rissanen H, Bullard KM, Knekt P. The Population Distribution of the Sagittal Abdominal Diameter (SAD) and SAD/height Ratio Among Finnish Adults. *Clinical Obesity*, 2014;4(6):333-41. <https://doi.org/10.1016/j.physbeh.2017.03.040> PMID:28363838 PMCID:PMC5510739
15. Ehrlich AC, Smith DA. Abdominal Diameter Index and 12-year Cardiovascular Disease Incidence in Male Bridge and Tunnel Workers. *International Journal of Obesity*, 2011;35(3):409-15. <https://doi.org/10.1038/ijo.2010.143> PMID:20714330
16. Roriz AKC, Passos L, Oliveira C, Eickemberg M, Moreira P, Ramos L. Discriminatory Power of Indicators Predictors of Visceral Adiposity Evaluated by Computed Tomography in Adults and Elderly Individual. *Nutricion Hospitalaria*, 2014;29(6):1401-7. <https://doi.org/10.3305/nh.2014.29.6.7185>
17. Sampaio LR, Simões EJ, Assis AMO, Ramos LR. Validity and reliability of the sagittal abdominal diameter as a predictor of visceral abdominal fat. *Arquivos Brasileiros de Endocrinologia e Metabologia*, 2007;51(6):980-6. <https://doi.org/10.1590/s0004-27302007000600013> PMID:17934666
18. Matthews DR, Hosker JP, Rudenski AS, Naylor BA, Treacher DF, Turner RC. Homeostasis model assessment: insulin resistance and beta-cell function from fasting plasma glucose and insulin concentrations in man. *Diabetologia*, 1985;28:412-9. <https://doi.org/10.1007/BF00280883> PMID:3899825
19. Pereira PF, Serrano HMS, Carvalho GQ, Ribeiro SMR, et al. Measurements of body fat distribution: assessment of collinearity with body mass, adiposity and height in female adolescents. *Revista Paulista de Pediatria*, 2015;33(1):63-71. [https://doi.org/10.1016/S2359-3482\(15\)30032-4](https://doi.org/10.1016/S2359-3482(15)30032-4)
20. Stokic E, Srdic B, Brtko V, Tomic-Nagalic D. Sagittal Abdominal Diameter as the Anthropometric Measure of Cardiovascular Risk. 2012:319-340. <https://doi.org/10.5772/31677>
21. Vasques ACJ, Cassani RSL, Forti ACe, Viela BS, Pareja JC, Tambascia MA, et al. Sagittal Abdominal Diameter as a Surrogate Marker of Insulin Resistance in an Admixed Population — Brazilian Metabolic Syndrome Study (BRAMS). *PLOS ONE*, 2015;2:1-13. <https://doi.org/10.1371/journal.pone.0125365> PMID:25951458 PMCID:PMC4423830
22. Riserus U, Faire Ude, Berglund L, Hellenius M-L. Sagittal Abdominal Diameter as a Screening Tool in Clinical Research: Cutoffs for Cardiometabolic Risk. *Journal of Obesity*, 2010:1-7. <https://doi.org/10.1155/2010/757939> PMID:20798888 PMCID:PMC2925288
23. Al-attas OS, Al-daghri NM, Alokail MS, Alkharfy KM, Draz H, Yakout S, et al. Association of body mass index, sagittal abdominal diameter and waist-hip ratio with cardiometabolic risk factors and adipocytokines in Arab children and adolescents. *BMC Pediatrics*, 2012;12(119):1-6. <https://doi.org/10.1186/1471-2431-12-119> PMID:22871266 PMCID:PMC3470957
24. González-Zavala MA, Velasco-Morales A, Terrazas-Flores JJ, de la Cruz-Galicia MG. Levels of insulin and HOMA-IR in adolescents in Saltillo, Coahuila, Mexico. *Medicina Universtaria*, 2015;17(67):80-7. <https://doi.org/10.1016/j.rmu.2015.02.004>
25. Yi KH, Hwang JS, Kim EY, Lee SH, Kim DH, Lim SJ. Prevalence of insulin resistance and cardiometabolic risk in Korean children and adolescents: A population-based study. *Diabetes Research and Clinical Practice*, 2014;103(1):106-13. <https://doi.org/10.1016/j.diabres.2013.10.021> PMID:24290751
26. Ramadhan NR, Wijayanti HS. Dietary Acid Load, Keseimbangan Asam Basa Tubuh dan Resistensi Insulin pada Remaja Obesitas. *Journal of Nutrition College*, 2017;6(1):69-75. <https://doi.org/10.14710/jnc.v6i1.16895>
27. Singh Y, Garg MK, Tandon N, Marwaha RK. A Study of Insulin Resistance by HOMA-IR and its Cut-off Value to Identify Metabolic Syndrome in Urban Indian Adolescents Yashpal. *Journal of Clinical Research in Pediatric Endocrinology*, 2013;5(4):245-51. <https://doi.org/10.4274/Jcrpe.1127> PMID:24379034 PMCID:PMC3890224
28. Krishnappa SK, Yashoda HT, Boraiah G, Vishwa S. Sagittal Abdominal Diameter to Measure Visceral Adipose Tissue in Overweight or Obese Adolescent Children and Its Role as A Marker of Insulin Resistance. *Journal of Clinical and Diagnostic Research*, 2015;9(11):9-12. <https://doi.org/10.7860/JCDR/2015/15971.6742> PMID:26673888 PMCID:PMC4668495
29. Moon HU, Ha KH, Han SJ, Kim HJ, Kim DJ. The Association of Adiponectin and Visceral Fat with Insulin Resistance and β -Cell Dysfunction. *Journal of Korean Medical Science*, 2019;34(1):1-12. <https://doi.org/10.3346/jkms.2019.34.e7> PMID:30618514 PMCID:PMC6318440
30. Donohoe CL, Doyle SL, Reynolds JV. Visceral Adiposity, Insulin Resistance and Cancer Risk. *Diabetology & Metabolic Syndrome*, 2011;12(3):1-13. <https://doi.org/10.1186/1758-5996-3-12> PMID:21696633 PMCID:PMC3145556
31. Castro AVB, Koika CM, Kim SP, Bergman RN. Obesity, Insulin Resistance and Comorbidity - Mechanisms of Association. *Arq Bras Endocrinol Metabol*, 2015;58(6):600-9. <https://doi.org/10.1159/000398850>
32. Kantartzis K, MacHann J, Schick F, Fritsche A, Häring HU, Stefan N. The Impact of Liver Fat VS Visceral Fat in Determining Categories of Prediabetes. *Diabetologia*, 2010;53(5):882-9. <https://doi.org/10.1007/s00125-010-1663-6> PMID:20099057
33. Borel AL, Nazare JA, Smith J, Aschner P, Barter P, Van Gaal L, et al. Visceral, subcutaneous abdominal adiposity and liver fat content distribution in normal glucose tolerance, impaired fasting glucose and/or impaired glucose tolerance. *International Journal of Obesity*, 2015;39(3):495-501. <https://doi.org/10.1038/ijo.2014.163> PMID:25179244

Abdominal Diameter Profiles have Relationship with Insulin Resistance in Obese Female Adolescents

ORIGINALITY REPORT

16%

SIMILARITY INDEX

12%

INTERNET SOURCES

14%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	www.elsevier.es Internet Source	2%
2	www.aem-sbem.com Internet Source	1%
3	Submitted to Eastern Nazarene College Student Paper	1%
4	www.acnut.com Internet Source	1%
5	jkms.org Internet Source	1%
6	journals.plos.org Internet Source	1%
7	Muhammad Roomi, Muhammad Mahmood, Yaser Khan. "Identifying Therapeutic Compounds Targeting RNA-Dependent-RNA-Polymerase of Sars-Cov-2", American Chemical Society (ACS), 2020 Publication	1%

8

Valentina A. Manullang, Ayu Rahadiyanti, Syafira N. Pratiwi, Diana N. Afifah. " Glycemic Index, Starch, and Protein Digestibility in Cookies ", Journal of Food Quality, 2020

Publication

1%

9

Borel, A-L, J-A Nazare, J Smith, P Aschner, P Barter, L Van Gaal, C E Tan, H-U Wittchen, Y Matsuzawa, T Kadowaki, R Ross, C Brulle-Wohlhueter, N Alméras, S M Haffner, B Balkau, and J-P Després. "Visceral, subcutaneous abdominal adiposity and liver fat content distribution in normal glucose tolerance, impaired fasting glucose and/or impaired glucose tolerance", International Journal of Obesity, 2014.

Publication

<1%

10

Hyun Uk Moon, Kyoung Hwa Ha, Seung Jin Han, Hae Jin Kim, Dae Jung Kim. "The Association of Adiponectin and Visceral Fat with Insulin Resistance and β -Cell Dysfunction", Journal of Korean Medical Science, 2019

Publication

<1%

11

Ana Carolina J. Vasques, Roberta S. L. Cassani, Adriana C. e Forti, Brunna S. Vilela et al. "Sagittal Abdominal Diameter as a Surrogate Marker of Insulin Resistance in an Admixed Population—Brazilian Metabolic Syndrome Study (BRAMS)", PLOS ONE, 2015

<1%

12 Vanesa España Romero, Jonatan R. Ruiz, Francisco B. Ortega, Enrique G. Artero et al. "Body fat measurement in elite sport climbers: Comparison of skinfold thickness equations with dual energy X-ray absorptiometry", Journal of Sports Sciences, 2009

Publication

13 garuda.ristekdikti.go.id <1 %
Internet Source

14 Submitted to Universitas Diponegoro <1 %
Student Paper

15 www.jcdr.net <1 %
Internet Source

16 www.ncbi.nlm.nih.gov <1 %
Internet Source

17 Thomas Reinehr, Gideon de Sousa, Werner Andler. "Longitudinal Analyses among Overweight, Insulin Resistance, and Cardiovascular Risk Factors in Children", Obesity Research, 2005

Publication

18 www.hindawi.com <1 %
Internet Source

19 www.science.gov <1 %
Internet Source

20 Gary L. Pittenger. "Small Fiber Neuropathy Is Associated with the Metabolic Syndrome", Metabolic Syndrome and Related Disorders, 06/2005

Publication

<1%

21 "Original Research", European Heart Journal Supplements, 2016

Publication

<1%

22 dmsjournal.biomedcentral.com

Internet Source

<1%

23 www.e-sciencecentral.org

Internet Source

<1%

24 www.psychiatryinvestigation.org

Internet Source

<1%

25 Barbara Perez Vogt, Daniela Ponce, Jacqueline Costa Teixeira Caramori. "Anthropometric Indicators Predict Metabolic Syndrome Diagnosis in Maintenance Hemodialysis Patients", Nutrition in Clinical Practice, 2015

Publication

<1%

26 A C Ehrlich. "Abdominal diameter index and 12-year cardiovascular disease incidence in male bridge and tunnel workers", International Journal of Obesity, 08/17/2010

Publication

<1%

27

Makbul Aman Mansyur, Syakib Bakri, Ilham Jaya Patellongi, Ilham Akbar Rahman. "The association between metabolic syndrome components, low-grade systemic inflammation and insulin resistance in non-diabetic Indonesian adolescent male", *Clinical Nutrition ESPEN*, 2020

Publication

<1%

28

Katta Subraya Prakash Rao, Peraje Vasu Dinesh. "ASSOCIATION OF BLOOD PRESSURE IN ADULTS WITH SELECTED ANTHROPOMETRIC PARAMETERS AND CRP: A HOSPITAL BASED CROSS SECTIONAL STUDY", *Journal of Evolution of Medical and Dental Sciences*, 2016

Publication

<1%

29

Sharon H. Jackson, Anna Bellatorre, Timothy McNeel, Anna María Nápoles, Kelvin Choi. "Longitudinal Associations between Obesity, Inflammation, and the Incidence of Type 2 Diabetes Mellitus among US Black and White Adults in the CARDIA Study", *Journal of Diabetes Research*, 2020

Publication

<1%

30

Edita Stoki, Biljana Srdi, Vladimir Brtko, Dragana Tomi-Nagli. "Chapter 9 Sagittal Abdominal Diameter as the Anthropometric

<1%

Measure of Cardiovascular Risk", IntechOpen, 2012

Publication

31

"Minutes of the 44th Genral Assembly of the European Association for the Study of Diabetes", Diabetologia, 2009

Publication

<1%

32

eprints.skums.ac.ir

Internet Source

<1%

33

Kahn, H. S., H. Rissanen, K. M. Bullard, and P. Knekt. "The population distribution of the sagittal abdominal diameter (SAD) and SAD/height ratio among Finnish adults : SAD & SAD/height among adult Finns", Clinical Obesity, 2014.

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

Abdominal Diameter Profiles have Relationship with Insulin Resistance in Obese Female Adolescents

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
