

A Cross-Sectional Assessment of Dietary Patterns and Their Relationship to Hypertension and Obesity in Indonesia

by Martha Irene Kartasurya

Submission date: 22-Sep-2022 03:33PM (UTC+0700)

Submission ID: 1906085227

File name: Their_Relationship_to_Hypertension_and_Obesity_in_Indonesia.pdf (538.23K)

Word count: 9129

Character count: 45181



A Cross-Sectional Assessment of Dietary Patterns and Their Relationship to Hypertension and Obesity in Indonesia

Oyedolapo A Anyanwu,¹ Sara C Foltz,¹ Fang Fang Zhang,¹ Kenneth Chui,² Virginia R Chomitz,² Martha I Kartasurya,³ and Elena N Naumova¹

¹Friedman School of Nutrition Science, Tufts University, Boston, MA, USA; ²Public Health and Community Medicine, School of Medicine, Tufts University, Boston, MA, USA; and ³Department of Public Health Nutrition, Diponegoro University, Semarang, Indonesia

ABSTRACT

Background: There is a marked increase in the intake of foods associated with higher risks for hypertension and obesity in Indonesia. However, studies assessing the relationship between dietary patterns and health outcomes are few.

Objective: The purpose of this study was to characterize dietary patterns and investigate their relationship with hypertension and obesity in Indonesia.

Method: Exploratory factor analysis was used to derive dietary patterns from a brief food scanner filled by 31,160 respondents aged 15 y and older in the Indonesian Family Life Survey wave 5 (IFLS 5). Age- and gender-specific quintiles of consumption were created for each pattern and the association between quintiles of each dietary pattern and the odds for hypertension and obesity were assessed using multivariate logistic regression analyses.

Results: Two dietary patterns were identified: a modern dietary pattern characterized by fast foods, soft drinks, sweet snacks, and salty snacks and a traditional pattern characterized by fish, vegetables, and fruits. Younger age and being male were significantly correlated with higher consumption of the modern pattern ($P < 0.0001$ and $P = 0.03$, respectively). Analyses showed no association between hypertension and the modern pattern. However, the traditional pattern revealed lower odds for hypertension among those in the highest quintile compared with the lowest quintile (OR: 0.84; 95% CI: 0.74, 0.95; P -trend < 0.05). Individuals in the highest quintile of each dietary pattern had higher odds of obesity compared with those in the lowest quintile (modern pattern—OR: 1.31; 95% CI: 1.15, 1.49; P -trend < 0.00 ; traditional pattern—OR: 1.25; 95% CI: 1.10, 1.42; P -trend < 0.01).

Conclusions: More studies using gold-standard measures of dietary intake are needed to better understand the relationship between the modern dietary pattern and hypertension in Indonesia. Also, both modern and traditional dietary patterns in Indonesia may be energy dense, leading to higher risk for obesity. *Curr Dev Nutr* 2022;6:nzac091.

Keywords: nutrition transitions, exploratory factor analysis, dietary patterns, hypertension, obesity, LMICs

© The Author(s) 2022. Published by Oxford University Press on behalf of the American Society for Nutrition. This is an Open Access article distributed under the terms of the Creative Commons Attribution-NonCommercial License (<https://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact journals.permissions@oup.com

Manuscript received November 8, 2021. Initial review completed April 11, 2022. Revision accepted April 22, 2022. Published online May 3, 2022.

This research was funded in part by the Tufts Institute of Environment, 2020 Environmental Research Fellowship, and the National Science Foundation (NSF) International Research Experiences for Students (IRES) grant: Collaborative Research: US-Indonesian Research Experience for Students on Sustainable Adaptation of Coastal Areas to Environmental Change (1826939 – PI Naumova EN).

Author disclosures: The authors report no conflicts of interest.

Supplemental Tables 1–6 are available from the “Supplementary data” link in the online posting of the article and from the same link in the online table of contents at <https://academic.oup.com/cdn/>.

Address correspondence to OAA (e-mail: oyedola@tufts.edu).

Abbreviations used: CVD, cardiovascular disease; EA, enumeration area; IFLS, Indonesian Family Life Survey; IRB, Institutional Review Board; LMIC, lower-to middle-income country; PA, physical activity; PROMIS, Patient-Reported Outcome Measurement Information System; RAND, Research and Development Corporation; SES, socioeconomic status.

Introduction

Over the past 4 decades, diet-related noncommunicable diseases have soared in lower- to middle-income countries (LMICs), and now co-exist with infectious diseases and malnutrition (1–3). The sharp increase in chronic disease conditions in LMICs has been linked to demographic, economic, and epidemiological transitions (1, 4). LMICs are also experiencing nutritional transitions—substantial shifts in dietary patterns

(4, 5)—that are having a major public health impact. Consequently, diet-related cardiovascular disease (CVD) risk factors are increasing in LMICs to a level that the health systems are not fully equipped for, resulting in higher mortality rates than in higher-income countries (6, 7).

Dietary patterns rather than isolated nutrients provide a more realistic assessment of people's dietary intake behavior, and are likely to have interactive and synergistic effects that will be large enough to allow the detection of a relationship between diet and health outcomes,

where one exists (8). In recognition of this, the Scientific Committee for the 2020–2025 Dietary Guidelines for Americans made dietary patterns a major focus of its report and recommendations (9). In studies from high-income countries like the United States, findings have consistently linked higher prevalence of CVD events to consumption of dietary patterns that are high in saturated fat, sodium, and added sugars (10–12). However, for LMICs, individual-level data to help elucidate the relationship between dietary patterns and CVD risk factors are still lacking (3, 13).

Specifically for Indonesia, the world's fourth most populous and largest island country (14), the mortality rate due to CVD risk factors like hypertension and obesity has been on the rise in tandem with nutrition transition (15). Importantly, Indonesia presents an exception to the problem of data gaps highlighted for many LMICs to some extent, with the existence of ongoing, national-level survey data that capture dietary intake and health indicators at the individual level. Yet, relatively few studies have looked at the associations between individual-level dietary intake and CVD risk factors for the Indonesian population. The few studies that have assessed the links between diet and health outcomes in Indonesia have focused only on selected food groups rather than dietary patterns (16, 17). The aim of this study was to characterize the dietary patterns from a brief food screener and to assess the relationship between the dietary patterns and the odds for hypertension and obesity among survey respondents.

6 Methods

Study design and data source

This was a cross-sectional study design that used the fifth wave of the Indonesia Family Life Survey (IFLS). The IFLS is an ongoing, national survey that assesses dietary intake and health indicators of Indonesian adults and children. We followed the procedures of the Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) cross-sectional reporting guidelines in this study (18).

The IFLS was stratified by province and urban/rural location, and households were randomly selected from these strata based on a sampling frame derived from the Indonesian Bureau of Statistics (14). A total of 321 enumeration areas (EAs) were randomly selected from 13 provinces, and EAs in urban and smaller provinces were oversampled to facilitate urban-rural and Javanese–non-Javanese comparisons (19). The ethical approval for IFLS 5 was provided in the United States by the Institutional Review Board (IRB) of the Research and Development Corporation (RAND) and in Indonesia by the IRB of the University of Gadjah Mada (20). Prior to the start of the survey, each respondent provided a signed written consent to participate in the study (20). The IFLS data are publicly available upon request and creation of a free online account with RAND (<https://www.rand.org/well-being/social-and-behavioral-policy/data/IFLS/IFLS/ifls5.html>).

The first wave (IFLS 1) was conducted in 1993 among individuals living in 7224 households. The fifth wave (IFLS 5), used for this study, was conducted in 2014 and administered to 16,204 households consisting of 50,148 individuals (Figure 1) (14). Only wave 5 included questions addressing the intake of fast foods, with fried and sweet snacks added, to capture the intake of some unhealthy food options (14). For this study, we limited our analyses to respondents aged 15 y and older

($n = 31,160$) who had sampling weights for all survey modules and health measurements and responded to the demographic and food consumption modules of the survey (Figure 1). Analyses accounted for sample attrition since IFLS 1 for the oversampling of urban areas and included survey weights that were matched to the Indonesian population in 2014 in the surveyed provinces for all age groups of respondents surveyed (14). To reduce reverse causality, respondents who reported currently taking medication for any chronic condition ($n = 2551$) were excluded from the regression analyses (Figure 1).

Dietary patterns

The food-consumption module of IFLS 5 asked about the consumption of 17 food items (Figure 2 and Supplemental Table 1). Most of the items included were iron- and vitamin A-rich foods, which are two micronutrients of public health importance for Indonesia (14). Respondents were asked if they consumed each food item in the past week and how many days in the week they consumed it (14).

We used factor analysis to aggregate the 17 food items in the food screener into dietary patterns with orthogonal rotation using the procedure “proc factor” steps in SAS version 9.4 (SAS Institute). The number of factors retained was determined by considering the scree plot, eigenvalues (> 1), magnitude of factor loadings (≥ 0.3), and the interpretability of the factors (8). The factor score of each dietary pattern for each respondent was derived by multiplying the mean frequency of consumption of each food item by their derived factor loadings (8).

53 Outcomes

The main outcome of interest was hypertension, and the secondary outcome of interest was obesity. For hypertension, 3 measures of each respondent's blood pressure were taken, following study protocol (14). The average of the 3 measurements was used in our analyses. Hypertension was defined as a combination of measured mean systolic blood pressure ≥ 140 mm Hg and/or mean diastolic blood pressure ≥ 90 mm Hg (21, 22) and categorized as a binary variable. To assess obesity status, each respondent's weight (kg) and height (cm) were taken based on the study protocol (14). Height was recalculated into meters by dividing the centimeter values by 100, and BMI values were calculated based on the formula: kg/m^2 . Obesity was then assessed as a binary variable of BMI values ≥ 27.50 , based on the Asian population cutoffs (23).

Covariates

Sociodemographic information collected during the survey included age, gender, urban-rural location, employment status, and educational level. Behavioral factors included the following: smoking status, physical activity (PA), and sleep quality. Respondents were asked if they ever smoked or were former smokers and if they are currently smoking or using tobacco products. Respondents were asked about the amount of time they spent engaged in different forms of physical activities and their intensity in the past 7 d. The PA module was based on the International Physical Activity Questionnaire (IPAQ). PA was coded to capture intensity by estimating metabolic equivalent (MET) values (24). We further categorized the PA scores into 3 levels: low, medium, and high (21). The sleep scale had 10 items drawn from the Patient-Reported Outcome Measurement Information System (PROMIS), which is a validated instrument used internationally in clinical settings with diverse populations (14, 25). Eight items on the scale asked respondents to describe




FIGURE 1 Study flowchart of the IFLS 5 (2014). IFLS 5, Indonesian Family Life Survey Wave 5.

their sleep patterns using different statements like, “My sleep was refreshing,” “I had difficulty falling asleep,” and “I had problems during the day because of poor sleep.” Responses were measured on a 5-point scale: 1 = not at all, 5 = very much. One item directly addressed sleep quality: “my quality of sleep was...” and was scored on a 5-point scale: 1 = very poor, 5 = very good. Finally, 1 item assessed ease of falling asleep: “I had trouble sleeping” and was scored on a 5-point scale: 1 = never, 5 = always. Following the methodology used by PROMIS, we reversed the score for negative statements such as “I had difficulty falling asleep” and then combined the scores of each statement to derive an overall sleep score of 50 (25). We then categorized the sleep score into 3 levels: <30 = poor, 30–39 = fair, and 40 and above = good.

Statistical analyses

We estimated means and SDs for continuous variables and percentages for categorical variables. We conducted ANOVA tests (for continuous variables) and chi-square tests (for categorical variables) to assess associations between each quintile of the derived dietary patterns and sociodemographic (age, gender, urban-rural location, employment status, and educational level) and behavioral factors (smoking status, PA, and sleep quality). Factor scores for each derived dietary pattern were

categorized into age- and gender-specific quintiles. Logistic regression models were used to estimate ORs for dietary patterns in association with hypertension and obesity, with simultaneous adjustments of both dietary patterns in the model. A few models were developed, and the final model retained was adjusted for age, gender, BMI, smoking, PA, sleep quality, employment status, and educational level. We included employment status and educational level in separate models in order to avoid model overfit, since both variables are proxies for socioeconomic status. We could not adjust for energy intake in any of the models because the food-consumption module did not provide details in terms of the composition or quantity of each food item. We evaluated the statistical significance of the different models by performing the log-likelihood ratio test. We assessed effect modification by age and gender and plotted the log odds of having hypertension by gender and by different iterations of age in years (5, 25, 50, 75, and 95 percentiles, which correspond to ages 16, 25, 34, 45, and 64 y, respectively). Based on the results, we then conducted subanalyses with stratification by gender and age group. A *P* value of 0.05 or lower was considered statistically significant for all analyses. The factor analysis was carried out using SAS statistical software, version 9.4 (SAS Institute, Inc.). All other statistical analyses were performed using the “svyset” procedure for Stata SE

Factor Loadings of the Derived Dietary Patterns




FIGURE 2 Factor loadings of the derived dietary patterns. Factor loadings $> .3$ were retained for each factor. The orange line corresponds to the factor loadings of each food group for the traditional dietary pattern; the blue line corresponds to the factor loadings of each food group for the modern dietary pattern. The numbers in the middle are the numerical values of the factor loadings. The further the dot in a line from the center of the circle, the larger the factor loading for a particular food group.

version 17 (StataCorp LP), to account for the complex survey procedure and attrition.

Results

Table 1 shows the sociodemographic and health characteristics of survey respondents. Overall, the weight-adjusted mean \pm SD age of study participants was 42.4 \pm 15.8 y, women constituted 53.7% of the respondents, and urban versus rural breakdown was approximately 50:50. The prevalence of hypertension was 29.4% in the study population. Among those who have hypertension, 6.3% were 15–19 y, 18.9% were 20–49 y, and 51.3% were 50 y and older. Average BMI (in kg/m^2) for the whole sample was 23.4 ± 4.57 . Based on the Asian population cutoffs, the prevalence of overweight and obesity among the study population was 30.9% and 17.4%, respectively.

Figure 2 and Supplemental Table 2 show the results of the exploratory factor analysis. Factor 1 loaded highly on instant noodle, soft drink, eggs, sweet snacks, fast food, meat, fried snacks, and dairy and was labeled as “modern.” Factor 2 loaded on banana, leafy vegetables, papaya, carrots, sweet potatoes, and fish and was labeled “traditional.” Table 2 shows the distribution of quintiles of consumption of both dietary patterns based on respondents’ sociodemographic and behavioral factors. Respondents who reported higher adherence to the modern pattern were younger, and more likely to be male, and have higher levels of education (secondary educational level or higher) and moderate-to-high-level PA. They were also more likely to report poor air sleep quality. Individuals who reported higher adherence to the traditional pattern were older and more likely to be female and

nonsmokers. Overall, more rural dwellers, and people with primary educational level or lower, reported low adherence to both types of diet.

Table 3 shows the association between dietary patterns and hypertension from multivariable logistic regression. Compared with individuals in the lowest quintiles, those in the highest quintiles of the traditional pattern had a 16% lower odds for hypertension (Q5 vs. Q1: OR = 0.84; 95% CI: 0.74, 0.95; P -trend < 0.05). Further adjustments of other health-related and psychosocial variables (e.g., self-rated health and mental health) did not change the results (Supplemental Table 3). When we substituted educational level for employment status in the model, the pattern of relationship between the traditional pattern and hypertension remained unchanged, but the trend was no longer significant (Supplemental Table 4).

Table 3 shows results of the unadjusted and adjusted logistic regression analyses to assess the relationship between the odds of obesity and levels of consumption of the derived dietary patterns. Both patterns were associated with a higher odds for obesity. Compared with individuals in the highest quintile, those in the highest quintile of the modern dietary pattern and traditional dietary pattern had a 31% (OR = 1.31; 95% CI: 1.15, 1.49; P -trend < 0.001) and 25% (OR = 1.25; 95% CI: 1.10, 1.42; P -trend < 0.01) higher odds for obesity, respectively.

Figure 3 shows the margins plot for the log odds of hypertension by gender. In the adjusted model, among males, the highest level of consumption of the modern pattern relative to the lowest is associated with higher log odds of hypertension while, for women, consuming at the highest level relative to the lowest is associated with lower log odds of hypertension (P -interaction = 0.01). We found no significant gender effect for the traditional pattern. Figure 4 shows the margins plot for the log odds of hypertension by age. Irrespective of the level of consumption of the modern pattern, the log odds for hypertension increase with age (P -interaction = 0.01). Similar to gender, age effects were not significant in assessing the role of high versus low adherence of the traditional pattern. In stratified analyses by gender, the highest level of consumption of the modern pattern relative to the lowest was associated with 17% reduced odds for hypertension (OR: 0.83; 95% CI: 0.69, 0.99; P -trend = 0.036) among women, while we found no significant association for men (Supplemental Table 5). When stratified by age group, older age was associated with lower odds for hypertension only among respondents 50 y and older with the highest level of adherence to the modern pattern relative to the lowest (OR: 0.76; 95% CI: 0.62, 0.94; P -trend = 0.015), while no significant effect was observed for other age groups (Supplemental Table 6).

Discussion

Diet-related health risk factors like hypertension and obesity have been on the rise in Indonesia, parallel to nutrition transitions. Using dietary data collected from national surveys, we characterized 2 dietary patterns—modern and traditional—and their sociodemographic and behavioral correlates. We also examined the association between these patterns and 2 major CVD risk factors among respondents 15 y and older in the IFLS wave 5 data (2014). Our analyses revealed a significant variation in food-consumption patterns by population subgroups. We also found some indications that the traditional pattern

TABLE 1 Characteristics of study respondents in the Indonesian Family Life Survey Wave 5, stratified by age group¹

Characteristics	Total (n = 31,160)	15–19 y (n = 3562)	20–49 y (n = 20,943)	≥50 y (n = 6655)
Gender, n (%)				
Female	16,619 (53.7)	1857 (45.6)	11,221 (54.0)	3541 (54.9)
Male	14,541 (46.3)	1705 (54.4)	9722 (46.0)	3114 (45.1)
Age, mean ± SD, y	42.6 ± 15.8	16.8 ± 1.8	35.1 ± 9.4	59.6 ± 6.2
Place of residence, n (%)				
Rural	12,797 (50.3)	1374 (48.3)	8473 (49.3)	2950 (52.3)
Urban	18,363 (49.7)	2188 (51.7)	12,470 (50.7)	3705 (47.7)
Marital status, n (%)				
Single	6159 (15.0)	3232 (91.6)	2856 (13.6)	71 (0.9)
Married	22,602 (74.0)	319 (8.2)	17,309 (82.0)	4974 (75.0)
Divorced	797 (2.9)	11 (0.2)	537 (2.8)	249 (3.7)
Widowed	1602 (8.1)	0 (0.0)	241 (1.6)	1361 (20.3)
Educational level, n (%)				
Primary or less	10,413 (44.6)	222 (6.7)	5640 (32.7)	4551 (71.9)
Secondary	16,019 (43.2)	3017 (85.5)	11,463 (51.5)	1539 (20.8)
Beyond secondary	4708 (12.2)	323 (7.8)	3832 (15.8)	553 (7.3)
Employment status, n (%)				
Employed	18,326 (59.9)	668 (19.1)	13,637 (64.9)	4021 (60.5)
Student	2576 (5.6)	2216 (61.6)	359 (1.6)	1 (0.0)
Housekeeping	8406 (27.6)	408 (11.0)	6228 (29.8)	1770 (27.5)
Retired	566 (2.6)	0 (0.0)	4 (0.0)	562 (7.3)
Unemployed ²	671 (2.4)	96 (2.9)	389 (2.0)	186 (2.8)
Other ³	613 (2.0)	174 (5.5)	324 (1.6)	115 (1.8)
BMI, mean ± SD, kg/m ²	23.4 ± 4.6	20.4 ± 4.6	23.88 ± 4.9	23.4 ± 3.5
Weight status, ⁴ n (%)				
Underweight	3838 (12.2)	1127 (32.8)	1894 (9.0)	817 (12.8)
Normal	12,612 (39.6)	1779 (49.9)	8340 (38.9)	2493 (38.4)
Overweight	9440 (30.9)	472 (12.6)	6841 (33.0)	2127 (31.4)
Obese	1185 (17.4)	176 (4.7)	3822 (19.1)	1185 (17.3)
Blood pressure, mean ± SD, mmHg				
Systolic	131.5 ± 22.2	118.2 ± 14.4	125.1 ± 18.3	144.6 ± 19.5
Diastolic	80.1 ± 12.0	72.7 ± 10.5	78.9 ± 12.1	83.6 ± 9.9
Hypertension, n (%)				
No	23,851 (70.6)	3347 (93.7)	17,330 (81.2)	3174 (48.7)
Yes	6995 (29.4)	201 (6.3)	3412 (18.8)	3382 (51.3)

¹Percentages are weighted to account for sampling design and attrition factor.²Unemployed category includes respondents that are not currently employed or disabled.³Other: unspecified employment category.⁴Weight status derived from BMI categories based on Asian population cutoffs.

might be protective of hypertension and that both the modern and traditional dietary patterns were associated with higher odds for obesity.

Studies looking into dietary patterns and their sociodemographic correlates in Indonesia are few but reveal converging trends. In a series of focus group discussions on food culture among Indonesian adults in West Sumatra, Lipoeto et al. found low consumption of Western-type food relative to traditional food (26). Our findings corroborate this: overall, the study population reported very low consumption of fast foods and soft drinks (Figure 2 and Supplemental Table 2). This, however, does not mean that Indonesia is not undergoing nutrition transition. According to Lipoeto et al. (15), there is an increase in the expenditure for oil and fat, as well as ready-to-eat foods, especially in the urban areas; urban dwellers and people of higher socioeconomic status (SES) are at higher risk (16) the adverse health effects of nutrition transition. Treloar et al. (27) also found that younger people and people of higher SES were more likely to buy “snack” food or eat out.

Overall, our analyses revealed that there is a marked difference in adherence to each type of dietary pattern by population subgroups in the study population. This has important public health implications, especially with respect to considering which subpopulations are most vulnerable to the adverse health impacts of nutrition transitions. It may also provide useful information to guide the design of targeted nutrition interventions to address disparities in dietary intake by population subgroups.

Evidence from high-income countries suggests that a dietary pattern characterized by higher intake of processed foods with high saturated fat, sodium, and sugar contents significantly increases the risk for CVDs (28–30). Hypertension is the leading risk factor for CVD in Indonesia (21, 31, 32), with current prevalence figures at about 48% among adults 40 y and older, and about 70% of cases undiagnosed (21).

We found no association between the modern dietary pattern and hypertension, while the traditional dietary pattern was inversely

TABLE 2 Characteristics by dietary patterns among respondents of the Indonesian Family Life Survey Wave 5 (2014)¹

Characteristics	Modern dietary pattern						Additional dietary pattern					
	Quintile 1 (n = 6231)	Quintile 2 (n = 6249)	Quintile 3 (n = 6226)	Quintile 4 (n = 6232)	Quintile 5 (n = 6222)	P ²	Quintile 1 (n = 6233)	Quintile 2 (n = 6225)	Quintile 3 (n = 6246)	Quintile 4 (n = 6233)	Quintile 5 (n = 6223)	P ²
Age, mean ± SD, y	43.9 ± 15.9	43.0 ± 15.8	42.1 ± 15.7	41.8 ± 15.6	41.1 ± 15.9	<0.0001	42.1 ± 16.1	41.9 ± 15.7	42.2 ± 15.7	42.6 ± 15.7	43.3 ± 15.6	<0.0001
Age group, n (%)												
15–19 y	710 (7.8)	712 (7.5)	714 (7.4)	713 (7.6)	713 (7.4)	0.69	710 (7.4)	714 (7.8)	716 (7.6)	713 (7.5)	709 (7.5)	0.91
20–49 y	4192 (57.5)	4202 (57.9)	4185 (56.8)	4188 (56.6)	4176 (56.2)		4188 (56.5)	4187 (57.3)	4195 (57.0)	4189 (56.7)	4184 (57.7)	
≥50 y	1329 (34.7)	1335 (34.6)	1327 (35.8)	1331 (35.8)	1333 (36.4)		1335 (36.1)	1324 (34.9)	1335 (35.4)	1331 (35.9)	1330 (34.9)	
Sex, n (%)												
Female	3325 (55.5)	3327 (54.1)	3323 (53.7)	3328 (53.1)	3316 (52.2)	0.03	3325 (52.8)	3327 (53.7)	3327 (54.5)	3319 (53.4)	3321 (54.3)	0.55
Male	2906 (44.5)	2922 (45.9)	2903 (46.3)	2904 (46.9)	2906 (47.8)		2908 (47.2)	2898 (46.3)	2919 (45.5)	2914 (46.6)	2902 (45.7)	
Place of residence, n (%)												
Rural	3402 (62.7)	2787 (53.6)	2426 (48.4)	2139 (44.1)	2043 (42.2)	<0.0001	2639 (53.3)	2718 (52.0)	2767 (52.8)	2541 (49.4)	2132 (43.6)	<0.0001
Urban	2829 (37.3)	3462 (46.4)	3800 (51.6)	4093 (55.9)	4179 (57.8)		3594 (46.7)	3507 (48.0)	3479 (47.2)	3692 (50.6)	4091 (56.4)	
Marital status, n (%)												
Single	1006 (12.4)	1113 (13.5)	1238 (15.1)	1293 (16.2)	1509 (1.0)	<0.0001	1355 (16.6)	1260 (15.6)	1222 (14.9)	1182 (14.5)	1140 (13.2)	<0.0001
Married	4695 (75.6)	4632 (74.9)	4531 (73.9)	4475 (73.2)	4269 (72.3)		4303 (70.2)	4502 (73.8)	4576 (74.5)	4587 (74.9)	4634 (76.9)	
Divorced	166 (3.1)	164 (3.1)	144 (2.6)	164 (2.9)	159 (2.7)		210 (3.9)	152 (2.7)	141 (2.8)	146 (2.4)	148 (2.7)	
Widowed	364 (8.9)	340 (8.5)	313 (8.4)	300 (7.8)	285 (7.0)		365 (9.3)	311 (7.9)	307 (7.8)	318 (8.2)	301 (7.2)	
Educational level, n (%)												
Primary school or less	2875 (57.5)	2281 (47.6)	1914 (42.0)	1775 (39.1)	1568 (36.4)	<0.0001	2548 (52.8)	2296 (48.3)	2069 (43.8)	1867 (40.8)	1633 (36.5)	<0.0001
Secondary school	2795 (35.6)	3191 (42.5)	3302 (44.6)	3284 (45.8)	3447 (47.9)		3140 (40.3)	3186 (42.1)	3286 (44.7)	3223 (44.3)	3184 (44.8)	
Beyond secondary school	560 (6.9)	774 (10.0)	1006 (13.4)	1167 (15.1)	1201 (15.7)		541 (6.9)	738 (9.6)	887 (11.5)	1141 (14.9)	1401 (18.7)	
Employment status, n (%)												
Employed	3587 (57.9)	3599 (59.1)	3611 (59.7)	3749 (61.1)	3780 (62.0)	<0.0001	3501 (57.7)	3610 (59.5)	3665 (59.1)	3754 (61.7)	3796 (61.8)	<0.001

(Continued)

TABLE 2 (Continued)

Characteristics	Modern dietary pattern					Traditional dietary pattern					P ²
	Quintile 1 (n = 6231)	Quintile 2 (n = 6249)	Quintile 3 (n = 6226)	Quintile 4 (n = 6232)	Quintile 5 (n = 6222)	Quintile 1 (n = 6233)	Quintile 2 (n = 6225)	Quintile 3 (n = 6246)	Quintile 4 (n = 6233)	Quintile 5 (n = 6223)	
Student	406 (4.6)	485 (5.2)	527 (5.5)	579 (6.3)	579 (6.3)	518 (5.5)	514 (5.7)	555 (6.1)	494 (5.3)	495 (5.2)	
Housekeeping	1812 (29.7)	1750 (27.7)	1719 (28.0)	1590 (26.6)	1535 (25.7)	1793 (29.0)	1709 (27.4)	1674 (27.9)	1655 (27.0)	1575 (26.3)	
Retired	131 (2.9)	127 (2.9)	111 (2.6)	100 (2.3)	97 (2.3)	98 (2.4)	106 (2.5)	111 (2.6)	108 (2.5)	143 (3.0)	
Unemployed	166 (2.7)	146 (2.8)	130 (2.2)	107 (2.0)	122 (2.0)	173 (3.1)	149 (2.4)	128 (2.4)	110 (1.8)	111 (2.0)	
Other (unspecified)	129 (2.2)	141 (2.3)	128 (1.9)	107 (1.8)	108 (1.6)	149 (2.4)	137 (2.2)	113 (2.9)	112 (1.8)	102 (1.7)	
Smoking status, n (%)											
Nonsmoker	3961 (64.3)	4005 (62.8)	4048 (64.0)	4046 (63.6)	3958 (61.7)	3832 (59.5)	3966 (63.2)	4009 (64.1)	4060 (63.9)	4151 (66.1)	<0.0001
Former smoker	271 (4.6)	273 (4.8)	290 (4.8)	304 (5.9)	278 (4.8)	245 (4.5)	247 (4.2)	272 (4.5)	288 (4.8)	364 (6.3)	
Current smoker	1999 (31.1)	1971 (32.4)	1888 (31.2)	1882 (31.2)	1986 (33.5)	2156 (36.0)	2012 (32.6)	1965 (31.4)	1885 (31.3)	1708 (27.6)	
Level of physical activity, ³ n (%)											
Low	2159 (33.0)	2083 (30.6)	2068 (30.4)	2095 (31.5)	1974 (29.0)	2329 (34.6)	2180 (32.6)	2063 (30.8)	1931 (28.3)	1876 (27.9)	<0.0001
Moderate	1912 (30.9)	2097 (33.7)	2112 (33.9)	2123 (33.5)	2140 (34.5)	1979 (31.3)	2016 (31.3)	2063 (33.2)	2171 (35.0)	2155 (34.7)	
High	2160 (36.1)	2069 (35.8)	2046 (35.7)	2014 (35.0)	2108 (36.5)	1925 (34.1)	2029 (35.1)	2120 (36.0)	2131 (36.7)	2192 (37.4)	
Sleep quality, n (%)											
Poor	631 (10.0)	632 (9.6)	714 (11.2)	701 (11.1)	894 (13.4)	864 (13.4)	730 (11.0)	672 (10.2)	642 (9.8)	664 (10.4)	<0.0001
Fair	1978 (29.6)	1912 (31.5)	2177 (34.3)	2302 (34.3)	2364 (35.4)	2193 (31.9)	2194 (32.4)	2257 (33.7)	2186 (32.8)	2111 (31.4)	
Good	3622 (60.5)	3497 (58.8)	3335 (57.2)	3229 (54.6)	2964 (51.3)	3176 (54.7)	3301 (56.6)	3317 (56.1)	3405 (57.3)	3448 (58.2)	

¹Percentages are weighted to account for sampling design and attrition factor.²P-values based on Pearson's chi-square tests for categorical variables and ANOVA for continuous variables.³Level of physical activity was categorized based on tertiles of total minutes per day of metabolic equivalent tasks (METs) of physical activity: low if < 540, moderate if 540–2887, and high if > 2888.

TABLE 3 Logistic regression analyses to assess the relationship between dietary patterns, hypertension, and obesity¹

Factor scores	Hypertension, n (%)		OR (95% CI) ²	Obesity, n (%)		OR (95% CI) ³
	Yes (n = 5868)	No (n = 22,488)		Yes (n = 4527)	No (n = 24,012)	
Modern dietary pattern ⁴						
Q1 (≤ -0.51)	1276 (22.1)	4430 (20.1)	Ref.	795 (17.7)	4949 (21.1)	Ref.
Q2 ($-0.50, 0.02$)	1221 (21.1)	4454 (19.8)	1.00 (0.89–1.13)	884 (19.1)	4829 (20.3)	1.11 (0.98–1.27)
Q3 (0.03, 0.56)	1169 (19.6)	4535 (20.3)	0.94 (0.83–1.06)	957 (20.7)	4784 (20.0)	1.19 (1.05–1.36)
Q4 (0.57, 1.26)	1144 (19.5)	4510 (20.0)	0.98 (0.86–1.10)	965 (21.3)	4718 (19.5)	1.25 (1.10–1.43)
Q5 (≥ 1.27)	1058 (17.8)	4559 (19.9)	0.90 (0.79–1.02)	926 (21.1)	4732 (19.0)	1.31 (1.15–1.49)
P-trend			0.41			< 0.001
Traditional dietary pattern ⁴						
Q1 (≤ -0.65)	1230 (22.4)	4552 (21.6)	Ref.	853 (19.2)	4975 (22.4)	Ref.
Q2 ($-0.64, 0.15$)	1198 (20.6)	4497 (20.3)	0.99 (0.87–1.11)	862 (19.8)	4871 (20.5)	1.08 (0.95–1.23)
Q3 (0.16, 0.42)	1161 (19.3)	4574 (20.2)	0.90 (0.80–1.01)	857 (18.9)	4917 (20.2)	1.03 (0.90–1.17)
Q4 (0.43–1.22)	1177 (19.7)	4483 (19.5)	0.93 (0.83–1.05)	951 (20.4)	4746 (19.3)	1.13 (0.99–1.28)
Q5 (≥ 1.23)	1102 (18.0)	4382 (18.5)	0.84 (0.74–0.95)	1004 (21.7)	4503 (17.7)	1.25 (1.10–1.42)
P-trend			0.047			< 0.01

¹Percentages were weighted to account for complex sampling design and attrition factor. Q, quintile; Ref., reference.

²ORs and 95% CIs were adjusted for age, gender, BMI, physical activity, sleep quality, smoking status, urban/rural location, and employment status.

³ORs and 95% CIs were additionally adjusted for marital status.

⁴The 2 dietary patterns were simultaneously included in the same model.

associated with the odds of having hypertension in this population (Table 3). The null findings for the association between the modern pattern and hypertension may be due to residual confounding by gender and age. For instance, a greater proportion of participants who have hypertension were women and older (≥ 50 y), and both subgroups had lower consumption of the modern dietary pattern (Table 2). Conversely, a greater proportion of the younger participants (15–49 y), who had a lower prevalence of hypertension, also had a higher intake of the modern pattern. As has been established in the literature, gender and older age are causal factors for hypertension and this is also

true for Indonesia (16, 21). Our analyses showed a significant association between higher adherence to the traditional pattern and hypertension based on a model that included employment status as an indicator for SES, while in another model that retained educational level as a proxy for SES, this significant association was attenuated. It may be that the significant association that we found could be an indication of the “healthy worker effect,” rather than a real association (33, 34). The available data did not provide enough granularity on the types of employment for us to further assess the true influence of employment status on the observed relationship. Thus, more studies with robust




FIGURE 3 Modern dietary pattern and hypertension by sex. Results were adjusted for age, BMI, physical activity, sleep quality, smoking status, urban/rural location, and employment status. HTN, hypertension; Q, quintile.




FIGURE 4 Modern dietary pattern and hypertension by age. Results were adjusted for gender, BMI, physical activity, sleep quality, smoking status, urban/rural location, and employment status. HTN, hypertension; Q, quintile.

measures of SES are needed to better elucidate the relationship between the levels of adherence to the traditional pattern and hypertension in Indonesia.

We further examined whether the association between dietary patterns and hypertension differs among population with different gender and age. In these subanalyses, the modern pattern was significantly associated with lower odds for hypertension among females and older adults (Supplemental Tables 5 and 6). This was an unexpected finding that warrants further investigation. One possible explanation could be that the modern pattern could be a proxy for higher protein intake, since it is characterized by meat, including beef, pork, and chicken, as well as eggs. Previous studies have shown the beneficial effect of protein on blood pressure, although evidence is mixed concerning whether it is animal-source or plant-based protein that confers this protection (35–37). Another explanation may be the menopausal status of the women. Previous studies have shown that premenopausal women have lower systolic blood pressure compared with menopausal women (38, 39). A greater proportion of survey respondents are less than 50 y and predominantly female (Table 1). More and better designed studies (experimental and/or cohort) and better dietary assessment instruments like 24-h recalls are needed to better elucidate the relationship between higher intake of proteins and hypertension by gender and age group.

Overweight and obesity rates in Indonesia have also been on the rise across all population subgroups, with changing diets and sedentary lifestyle likely the major drivers (7, 40). Prevalence rates for overweight are particularly high for women (26–31%) compared to men (16–21%) (7, 40). More importantly, change in diets means that even poor households are not immune from obesity due to higher intake of low-nutrient, energy-dense foods (6, 17). We found that both types of patterns are associated with higher odds for obesity. The higher odds for obesity observed for both types of dietary patterns can be better

understood in the light of other relevant literature from Indonesia. Lipoeto, Lin, and Angeles-Agdeppa (26) concluded that the nutrition transition being experienced in Indonesia and other countries of South East Asia was not all due to consuming more Western food, since the populations in the region have largely preserved their traditional eating patterns. Rather, they postulated that it might be better explained by the changing composition of the traditional diets (26). This provides a good context for our findings and is a crucial point to consider, especially in categorizing dietary patterns as healthy versus unhealthy. In our study, although the traditional pattern was characterized by higher intakes of fish, leafy green vegetables, and fruits (foods typically considered to be healthier options), the odds for obesity on this type of diet was only slightly lower than with the modern pattern (Table 3). This is an indication that how food is prepared is equally as important as the type of food being consumed. Preliminary results from an ongoing qualitative study on diet quality and health outcomes in Indonesia reveal that frying, using the same oil multiple times, and the use of coconut milk are the most common methods for cooking many Indonesian dishes, including fish and vegetables (O Anyanwu¹, E Naumova¹, V Chomitz², F Zhang¹, K Chui², M Kartasurya³, S Folt¹, 2022; ¹Tufts University Friedman School of Nutrition and Policy, ²School of Medicine, Tufts University, ³Diponegoro University, Semarang, Indonesia; unpublished data). Healthcare providers interviewed during the study concluded that these practices are key drivers for the high number of CVD-related cases they treat in their practices (O Anyanwu¹, E Naumova¹, V Chomitz², F Zhang¹, K Chui², M Kartasurya³, S Folt¹, 2022; ¹Tufts University Friedman School of Nutrition and Policy, ²School of Medicine, Tufts University, ³Diponegoro University, Semarang, Indonesia; unpublished data). More studies addressing the sociodemographic factors related to changes in the composition of traditional diets, cooking practices, and their effects on CVD risk factors are warranted for Indonesia.

Challenges and limitations

This study has several limitations. First, our study uses a cross-sectional design and thus it may not indicate a strong causal argument between the exposures and outcomes of interest. Second, it is only the fifth wave of the IFLS that captured the intake of potentially unhealthy foods, and it was the only wave used in this study, so we were not able to investigate any longitudinal changes in the associations. Third, the instrument for assessing food intake in the survey was a brief food screener that was designed to capture specific micronutrients of public health importance for Indonesia and provided no details in quantities of consumption or methods of preparation. Thus, we could not estimate and adjust for total energy intake in our analyses, leaving room for measurement errors and confounding by total energy intake. Based on these limitations, we recommend caution about making any causal inference on any of our findings.

Strengths of the study

The sparseness of information available from the brief food screener used in our study notwithstanding, it is still comparatively a valid instrument for assessing food consumption at the individual level (41, 42), especially for an LMIC like Indonesia, where nationally representative dietary data at the individual level are scarce. Other available options such as per-capita expenditure estimates do not capture differences in food-consumption patterns at the individual and population subgroup levels as the brief screener was able to do (3). Brief food screeners have been shown in the literature to have adequate reproducibility and validity when compared with a longer food-frequency questionnaire and biomarkers (41, 43). Additionally, brief food screeners have low cost, low respondent burden and are appropriate for studies testing a limited set of hypotheses (41, 42). Further, conducting factor analysis afforded us a more rigorous and valid approach to quantifying the food-consumption patterns of study participants. By aggregating the food items into patterns rather than individual food items, we were able to detect a significant association that another study using the same dataset missed (16). Also, the large sample size allowed us to have more precise and robust estimates of the relationship between diet and CVD risk factors of respondents than a less-powered study would have done.

Conclusions

Our findings provided some initial evidence for the associations between dietary intake patterns, hypertension, and obesity in Indonesia. Future studies using a longitudinal design are warranted to further evaluate the role of dietary intake patterns in noncommunicable diseases in Indonesians, who have been experiencing a transition in dietary intake patterns in the past few decades. Further, our findings may be informative for public health and nutrition researchers in Indonesia to help identify vulnerable subgroups at higher risk of the adverse health effects of nutrition transitions and to tailor nutrition interventions to meet the needs of the different subpopulations in Indonesia.

Acknowledgments

The authors' responsibilities were as follows—OAA, FFZ, ENN, and SCF: designed the research; OAA: conducted the research, wrote the manuscript, and had primary responsibility for final content; OAA,

FFZ, and KC: performed statistical analyses; FFZ, SCF, KC, ENN, VRC, and MIK: critically reviewed and revised the manuscript; and all authors: read and approved the final manuscript.

Data Availability

Data described in the manuscript, code book, and analytic code will be made available upon request pending application and approval.

References

- Amuna P, Zotor FB. Epidemiological and nutrition transition in developing countries: impact on human health and development. *The epidemiological and nutrition transition in developing countries: evolving trends and their impact in public health and human development. Proc Nutr Soc* 2008;67(1):82–90.
- Nishida C, Shetty P, Uauy R. Diet, nutrition and the prevention of chronic diseases: scientific background papers of the Joint WHO/FAO Expert Consultation (Geneva, 28 January–1 February 2002—introduction. *Public Health Nutr* 2004;7(1):99–100.
- Khatibzadeh S, Kashaf MS, Micha R, Fahimi S, Shi P, Elmadfa I, et al. A global database of food and nutrient consumption. *Bull World Health Organ* 2016;94(12):931.
- Abrahams Z, Mchiza Z, Steyn NP. Diet and mortality rates in sub-Saharan Africa: stages in the nutrition transition. *BMC Public Health* 2011;11(1):801.
- Popkin BM. The nutrition transition and obesity in the developing world. *J Nutr* 2001;131(3):871S–8S.
- Schmidhuber J, Shetty P. The nutrition transition to 2030. Why developing countries are likely to bear the major burden. *Acta Agric Scand C* 2005;2(3–4):150–66.
- Schröders J, Wall S, Hakimi M, Dewi FST, Weinehall L, Nichter M, et al. How is Indonesia coping with its epidemic of chronic noncommunicable diseases? A systematic review with meta-analysis. *PLoS One* 2017;12(6):1–31.
- Moeller SM, Reedy J, Millen AE, Dixon LB, Newby P, Tucker KL, et al. Dietary patterns: challenges and opportunities in dietary patterns research: an Experimental Biology workshop, April 1, 2006. *J Acad Nutr Diet* 2007;107(7):1233–9.
- Dietary Guidelines Advisory Committee. Scientific report of the 2020 Dietary Guidelines Advisory Committee: advisory report to the Secretary of Agriculture and the Secretary of Health and Human Services. Washington (DC): Agricultural Research Service; 2020.
- Shan Z, Li Y, Baden MY, Bhupathiraju SN, Wang DD, Sun Q, et al. Association between healthy eating patterns and risk of cardiovascular disease. *JAMA Intern Med* 2020;180(8):1090–100.
- Yu E, Malik VS, Hu FB. Cardiovascular disease prevention by diet modification: JACC Health Promotion Series. *J Am Coll Cardiol* 2018;72(8):914–26.
- Estruch R, Ros E, Salas-Salvadó J, Covas M-I, Corella D, Arós F, et al. Primary prevention of cardiovascular disease with a Mediterranean diet supplemented with extra-virgin olive oil or nuts. *N Engl J Med* 2018;378(25):e34.
- Micha R, Coates J, Leclercq C, Charrondiere UR, Mozaffarian D. Global dietary surveillance: data gaps and challenges. *Food Nutr Bull* 2018;39(2):175–205.
- Strauss J, Witoelar F, Sikoki B. The fifth wave of the Indonesian Family Life Survey (IFLS 5): overview and field report. Vol. 1. Santa Monica (CA): RAND; 2016. Available from: <https://www.rand.org/well-being/social-and-behavioral-policy/data/IFLS/IFLS5.html>
- Lipoeto N, Wattanapenpaiboon N, Malik A, Wahlqvist M. Nutrition transition in west Sumatra, Indonesia. *Asia Pac J Clin Nutr* 2004;13(3):312–16.

16. Peltzer K, Pengpid S. The prevalence and social determinants of hypertension among adults in Indonesia: a cross-sectional population-based national survey. *Int J Hypertens* 2018;2018:1–9.
17. Pengpid S, Peltzer K. The prevalence of underweight, overweight/obesity and their related lifestyle factors in Indonesia, 2014–2015. *AIMS Public Health* 2017;4(6):633.
18. Erik von Elm M, Altman DG, Egger M, Pocock SJ, Gøtzsche PC, Vandenbroucke JP. The Strengthening of Reporting of Observational Studies in Epidemiology (STROBE) statement: guidelines for reporting observational studies. *Ann Intern Med* 2007;147(8):573, e7.
19. RAND Corporation. The Indonesia Family Life Survey (IFLS): note regarding ethical clearance [Internet]. Santa Monica (CA): RAND; 2016. Available from: <https://www.rand.org/well-being/social-and-behavioral-policy/data/IFLS/IFLS.html>
20. Strauss J, Wittoel F, Sikoki B. Household Survey Questionnaire for the Indonesia Family Life Survey, wave 5. Santa Monica (CA): RAND; 2016.
21. Hussain MA, Al Mamun A, Reid C, Huxley RR. Prevalence, awareness, treatment and control of hypertension in Indonesian adults aged ≥ 40 years: findings from the Indonesia Family Life Survey (IFLS). *PLoS One* 2016;11(8):1–16.
22. Chobanian AV, Bakris GL, Black HR, Cushman WC, Green LA, Izzo JL, Jr, et al. The seventh report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA* 2003;289(19):2560–71.
23. Hsu WC, Araneta MRG, Kanaya AM, Chiang JL, Fujimoto W. BMI cut points to identify at-risk Asian Americans for type 2 diabetes screening. *Diabetes Care* 2015;38(1):150–8.
24. Scoring the International Physical Activity Questionnaire (IPAQ) [Internet]. 2018. Available from: https://ugc.futurelearn.com/uploads/files/bc/c5/bcc53b14-ec1e-4d90-88e3-1568682f32ae/IPAQ_PDF.pdf (accessed 15 April 2020).
25. Ader DN. Developing the Patient-Reported Outcomes Measurement Information System (PROMIS). *Med Care J* 2007;45(5):1–2.
26. Lipoeto NI, Lin KG, Angeles-Agdeppa I. Food consumption patterns and nutrition transition in South-East Asia. *Public Health Nutr* 2013;16(9):1637–43.
27. Treloar C, Porteous J, Hassan F, Kasniyah N, Lakshmanudu M, Sama M, et al. The cross cultural context of obesity: an INCLEN multicentre collaborative study. *Health Place* 1999;5(4):279–86.
28. Moreira PV, Hyseni L, Moubarac J-C, Martins APB, Baraldi LG, Capewell S, et al. Effects of reducing processed culinary ingredients and ultra-processed foods in the Brazilian diet: a cardiovascular modelling study. *Public Health Nutr* 2018;21(1):181–8.
29. Srour B, Fezeu LK, Kesse-Guyot E, Allès B, Méjean C, Andrianasolo RM, et al. Ultra-processed food intake and risk of cardiovascular disease: prospective cohort study (NutriNet-Santé). *BMJ* 2019;365:11451.
30. Hall KD, Ayuketah A, Brychta R, Cai H, Cassimatis T, Chen KY, et al. Ultra-processed diets cause excess calorie intake and weight gain: an inpatient randomized controlled trial of ad libitum food intake. *Cell Metab* 2019;30(1):67–77, e3.
31. Prihartono NA, Fitriyani F, Riyadina W. Cardiovascular disease risk factors among blue and white-collar workers in Indonesia. *Acta Med Indones* 2018;50(2):96–103.
32. Hussain MA, Huxley RR, Al Mamun A. Multimorbidity prevalence and pattern in Indonesian adults: an exploratory study using national survey data. *BMJ Open* 2015;5(12):e009810.
33. Shah D. Healthy worker effect phenomenon. *Indian J Occup Environ Med* 2009;13(2):77.
34. Chowdhury R, Shah D, Payal AR. Healthy worker effect phenomenon: revisited with emphasis on statistical methods—a review. *Indian J Occup Environ Med* 2017;21(1):2.
35. Altorf-van der Kuil W, Engberink MF, Brink EJ, van Baak MA, Bakker SJ, Navis G, et al. Dietary protein and blood pressure: a systematic review. *PLoS One* 2010;5(8):e12102.
36. Liu R, Dang S, Yan H, Wang D, Zhao Y, Li Q, et al. Association between dietary protein intake and the risk of hypertension: a cross-sectional study from rural western China. *Hypertens Res* 2013;36(11):972–9.
37. Sun D, Zhou T, Li X, Heianza Y, Liang Z, Bray GA, et al. Genetic susceptibility, dietary protein intake, and changes of blood pressure: the pounds lost trial. *Hypertension* 2019;74(6):1460–7.
38. Portaluppi F, Pansini F, Manfredini R, Mollica G. Relative influence of menopausal status, age, and body mass index on blood pressure. *Hypertension* 1997;29(4):976–9.
39. Staessen JA, Ginocchio G, Thijs L, Fagard R. Conventional and ambulatory blood pressure and menopause in a prospective population study. *J Hum Hypertens* 1997;11(8):507–14.
40. Römling C, Qaim M. Obesity trends and determinants in Indonesia. *Appetite* 2012;58(3):1005–13. doi: 10.1016/j.appet.2012.02.053.
41. Byers T, Marshall J, Fiedler R, Zielezny M, Graham S. Assessing nutrient intake with an abbreviated dietary interview. *Am J Epidemiol* 1985;122(1):41–50.
42. Spencer EH, Elon LK, Hertzberg VS, Stein AD, Frank E. Validation of a brief diet survey instrument among medical students. *J Am Diet Assoc* 2005;105(5):802–6.
43. Rifas-Shiman SL, Willett WC, Lobb R, Kotch J, Dart C, Gillman MW. PrimeScreen, a brief dietary screening tool: reproducibility and comparability with both a longer food frequency questionnaire and biomarkers. *Public Health Nutr* 2001;4(2):249–54.

A Cross-Sectional Assessment of Dietary Patterns and Their Relationship to Hypertension and Obesity in Indonesia

ORIGINALITY REPORT

15%
SIMILARITY INDEX

12%
INTERNET SOURCES

11%
PUBLICATIONS

4%
STUDENT PAPERS

PRIMARY SOURCES

1 academic.oup.com 3%
Internet Source

2 Sang-Ah Lee. "Dietary patterns and blood pressure among middle-aged and elderly Chinese men in Shanghai", British Journal Of Nutrition, 07/2010 1%
Publication

3 www.ncbi.nlm.nih.gov 1%
Internet Source

4 www.science.gov <1%
Internet Source

5 www.mdpi.com <1%
Internet Source

6 pure.uva.nl <1%
Internet Source

7 nesr.usda.gov <1%
Internet Source

8

Patel, Sumaiya, Adnan Custovic, Jaclyn A Smith, Angela Simpson, Gina Kerry, and Clare S Murray. "Cross-sectional association of dietary patterns with asthma and atopic sensitisation in childhood - in a cohort study", *Pediatric Allergy and Immunology*, 2014.

Publication

<1 %

9

repositorio.uam.es

Internet Source

<1 %

10

Hui-Peng Liew, Sheena Gardner. "The interrelationship between smoking and depression in Indonesia", *Health Policy and Technology*, 2016

Publication

<1 %

11

www.dietaryguidelines.gov

Internet Source

<1 %

12

bmcpublichealth.biomedcentral.com

Internet Source

<1 %

13

link.springer.com

Internet Source

<1 %

14

Laura J. Botzet, Julia M. Rohrer, Ruben C. Arslan. "Analysing Effects of Birth Order on Intelligence, Educational Attainment, Big Five, and Risk Aversion in an Indonesian Sample", *European Journal of Personality*, 2020

Publication

<1 %

15	Submitted to University of Glasgow Student Paper	<1 %
16	www.tandfonline.com Internet Source	<1 %
17	big5.39kf.com Internet Source	<1 %
18	nutritionj.biomedcentral.com Internet Source	<1 %
19	digitalcommons.fiu.edu Internet Source	<1 %
20	Submitted to Albany College of Pharmacy Student Paper	<1 %
21	Shreyasi Deb, Nancy A. Miller. "Relations among race/ethnicity, gender, and mental health status in primary care use.", Psychiatric Rehabilitation Journal, 2017 Publication	<1 %
22	etd.repository.ugm.ac.id Internet Source	<1 %
23	Cai-Xia Zhang. "Greater vegetable and fruit intake is associated with a lower risk of breast cancer among Chinese women", International Journal of Cancer, 07/01/2009 Publication	<1 %

24	Dong Wang, Yuna He, Yanping Li, Dechun Luan, Xiaoguang Yang, Fengying Zhai, Guansheng Ma. "Dietary patterns and hypertension among Chinese adults: a nationally representative cross-sectional study", BMC Public Health, 2011 Publication	<1 %
25	bmjopensem.bmj.com Internet Source	<1 %
26	repositori.upf.edu Internet Source	<1 %
27	Submitted to KYUNG HEE UNIVERSITY Student Paper	<1 %
28	mhealth.jmir.org Internet Source	<1 %
29	news.csudh.edu Internet Source	<1 %
30	nsft.sbm.ac.ir Internet Source	<1 %
31	www.39kf.com Internet Source	<1 %
32	"Track 5: Populations and population health", Obesity Reviews, 2016. Publication	<1 %

33	Christin Heidemann, Christa Scheidt-Nave, Almut Richter, Gert B. M. Mensink. "Dietary patterns are associated with cardiometabolic risk factors in a representative study population of German adults", British Journal of Nutrition, 2011	<1 %
Publication		

34	Farah Naja, Laila Itani, Nahla Hwalla, Abla M. Sibai, Samer A. Kharroubi. "Identification of dietary patterns associated with elevated blood pressure among Lebanese men: A comparison of principal component analysis with reduced rank regression and partial least square methods", PLOS ONE, 2019	<1 %
Publication		

35	nutrition.tufts.edu	<1 %
Internet Source		

36	www.aans.org	<1 %
Internet Source		

37	www.ijphrd.com	<1 %
Internet Source		

38	www.ncchta.org	<1 %
Internet Source		

39	www.scielo.br	<1 %
Internet Source		

40	Frances A. Tylavsky, Assiamira Ferrara, Diane J. Catellier, Emily Oken et al. "Understanding childhood obesity in the US: the NIH environmental influences on child health outcomes (ECHO) program", International Journal of Obesity, 2019 Publication	<1 %
41	Makiko Sekiyama, Hong Jiang, Budhi Gunawan, Linda Dewanti, Ryo Honda, Hana Shimizu-Furusawa, Oekan Abdoellah, Chiho Watanabe. "Double Burden of Malnutrition in Rural West Java: Household-Level Analysis for Father-Child and Mother-Child Pairs and the Association with Dietary Intake", Nutrients, 2015 Publication	<1 %
42	assets.researchsquare.com Internet Source	<1 %
43	bmchealthservres.biomedcentral.com Internet Source	<1 %
44	journals.plos.org Internet Source	<1 %
45	mdpi-res.com Internet Source	<1 %
46	mro.massey.ac.nz Internet Source	<1 %

47

www.karger.com

Internet Source

<1 %

48

Beard, V.A.. "Household Contributions to Community Development in Indonesia", World Development, 200704

Publication

<1 %

49

Frances Gunner, Michael Lindsay, Pieter Brown, Anneliese Shaw, Trish Davey, Susan Lanham-New, Bruce Griffin, Joanne Fallowfield. "Impact of the occupational environment of a submerged submarine on cardiometabolic health of Royal Navy submariners", Occupational and Environmental Medicine, 2020

Publication

<1 %

50

Mohammad Ziaul Islam Chowdhury, Meshbahur Rahman, Tanjila Akter, Tania Akhter et al. "Hypertension prevalence and its trend in Bangladesh: evidence from a systematic review and meta-analysis", Clinical Hypertension, 2020

Publication

<1 %

51

Paulo Rogério Melo Rodrigues, Rita Adriana Gomes de Souza, Mara Lima De Cnop, Luana Silva Monteiro et al. "Dietary quality varies according to data collection instrument: a comparison between a food frequency

<1 %

questionnaire and 24-hour recall", Cadernos de Saúde Pública, 2016

Publication

52

Rachel Dankner, Michael Shanik, Jesse Roth, Ayala Luski, Flora Lubin, Angela Chetrit. "Sex and ethnic-origin specific BMI cut points improve prediction of 40-year mortality: the Israel GOH study", Diabetes/Metabolism Research and Reviews, 2015

Publication

<1 %

53

dash.harvard.edu

Internet Source

<1 %

54

equityhealthj.biomedcentral.com

Internet Source

<1 %

55

ijbnpa.biomedcentral.com

Internet Source

<1 %

56

jamanetwork.com

Internet Source

<1 %

57

s-space.snu.ac.kr

Internet Source

<1 %

58

theijoem.com

Internet Source

<1 %

59

www.cambridge.org

Internet Source

<1 %

60

www.disparitiesresearch.org

Internet Source

<1 %

61

www.hindawi.com

Internet Source

<1 %

62

www.orthopeden.org

Internet Source

<1 %

63

Carolina Batis, Daniela Sotres-Alvarez, Penny Gordon-Larsen, Michelle A. Mendez, Linda Adair, Barry Popkin. "Longitudinal analysis of dietary patterns in Chinese adults from 1991 to 2009", British Journal of Nutrition, 2013

Publication

<1 %

64

Chan Nie, Tingting Yang, Ziyun Wang, Deji Suolang et al. "Dietary patterns and gallstone risks in Chinese adults: a cross-sectional analysis of China Multi-Ethnic Cohort Study", Journal of Epidemiology, 2022

Publication

<1 %

65

Gerard Ngueta. "Caffeine and caffeine metabolites in relation to hypertension in U.S. adults", European Journal of Clinical Nutrition, 2019

Publication

<1 %

66

Nicole D. Fields, Brian W. Whitcomb, Elizabeth R. Bertone-Johnson, Airín D. Martínez, Nicole A. VanKim. "Race-specific associations between psychological distress and obesity: the role of social cohesion", Ethnicity & Health, 2022

Publication

<1 %

67 "Nutrition and Health in a Developing World", Springer Science and Business Media LLC, 2017 <1 %
Publication

68 Dorit T Stein, Nikkil Sudharsanan, Shita Dewi, Jennifer Manne-Goehler, Firman Witoelar, Pascal Geldsetzer. "Change in clinical knowledge of diabetes among primary healthcare providers in Indonesia: repeated cross-sectional survey of 5105 primary healthcare facilities", BMJ Open Diabetes Research & Care, 2020 <1 %
Publication

69 Preventive Nutrition, 2015. <1 %
Publication

70 Tabea K. Kirschbaum, Michaela Theilmann, Nikkil Sudharsanan, Jennifer Manne - Goehler et al. "Targeting Hypertension Screening in Low - and Middle - Income Countries: A Cross - Sectional Analysis of 1.2 Million Adults in 56 Countries", Journal of the American Heart Association, 2021 <1 %
Publication

71 Zumin Shi, Vijay Ganji. "Dietary patterns and cardiovascular disease risk among Chinese adults: a prospective cohort study", European Journal of Clinical Nutrition, 2020 <1 %
Publication

Exclude quotes Off

Exclude bibliography On

Exclude matches Off

A Cross-Sectional Assessment of Dietary Patterns and Their Relationship to Hypertension and Obesity in Indonesia

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8

PAGE 9

PAGE 10

PAGE 11