

# Anthropometric-Parameters and Total-Cholesterol to HDL-Cholesterol Ratio are Better in Long-Distance Cyclists

*by* Martha Irene Kartasurya

---

**Submission date:** 21-Feb-2020 10:14AM (UTC+0700)

**Submission ID:** 1261230549

**File name:** ists\_Indonesia\_North\_Coast\_and\_Tour\_de\_Borobudur\_2017\_Study.pdf (196.03K)

**Word count:** 5704

**Character count:** 28279



## Anthropometric-Parameters and Total-Cholesterol to HDL-Cholesterol Ratio are Better in Long-Distance Cyclists

Mahatul Azam<sup>1,2,3\*</sup>, Sri Ratna Rahayu<sup>1</sup>, Arulita Ika Fibriana<sup>1</sup>, Hardhono Susanto<sup>1</sup>, Martha Irene Kartasurya<sup>4</sup>, Udin Bahrudin<sup>5</sup>

<sup>1</sup>Public Health Department, Sport Sciences Faculty, Universitas Negeri Semarang

<sup>2</sup>Doctoral Program in Medicine and Health Sciences, Faculty of Medicine, Universitas Diponegoro

<sup>3</sup>Department of Anatomy, Faculty of Medicine, Universitas Diponegoro

<sup>4</sup>Public Health Nutrition Department, Faculty of Public Health, Universitas Diponegoro

<sup>5</sup>Department of Cardiology and Vascular Medicine, Faculty of Medicine, Universitas Diponegoro

### Article Info

Article History:  
Submitted May 2018  
Accepted June 2018  
Published November 2018

Keywords:  
Total-Cholesterol,  
HDL-Cholesterol,  
Coronary-Artery-Diseases,  
Long-Distance, Cyclism

DOI  
<https://doi.org/10.15204/kemas.v14i2.14506>

### Abstract

Total-Cholesterol (TC) to HDL-Cholesterol (HDL-C) ratio in athletes is well known to be better than sedentary people. However, information about comparison of TC/HDL-C in different groups of cyclists based on cycling touring characteristics and anthropometry parameters is lacking. This study aimed to compare TC/HDL-C ratio between groups based on the type of tour in cyclists population. Eighty-eight participants were recruited in this cross-sectional study. Mean differences of parameters between groups was analyzed by One-Way Anova and independent t-test, whereas multivariate analyses was conducted by binary logistic-regression. P value <0.05 was considered statistically significant. TC/HDL-C ratio in all groups were less than 4.5. There is no differences of TC-levels (NC240K: 216.6±55.04, TdB140K: 208.1±27.13, TdB100K: 203.1±31.95; p=0.427). But there is significantly different level of HDL-C (NC240K: 68.9±19.09, TdB140K: 52.1±13.9, TdB100K: 53.6±12.45; p=0.0001) and TC/HDL-C ratio (NC240K: 3.3±1.12, TdB140K: 4.2±1.07, TdB100K: 4.0±1.06; p=0.007). Between TC/HDL ratio groups (≥4 or <4), there were differences of weight, BMI, waist-circumference, hip-circumference and type of tour (p<0.05). Finally, BMI and type of tour were the most influential factor. Long-distance cyclists have a synergistic effect of lipid profile and anthropometry measurements, and heavier cycling tour participant, that represent cycling training habits, tend to have lower TC/HDL ratio (<4).

### Introduction

Numerous studies show that levels of HDL and total cholesterol ratio is a major predictor for Coronary Artery Disease (CAD) (Arquer, 2006; Lemieux, 2011; Riwanto, 2013). Increasing level of serum cholesterol will increase risk of CAD (Ahn, 2016; Rosjidi, 2017). The Framingham study about cardiovascular

risk factors has shown that a decrease in plasma HDL cholesterol levels are associated with increased risk of CAD, with levels of HDL less than 40 mg/dl as a major risk factor of CAD, while HDL levels of more than 60 mg/dl protects against CAD occurrence (Penalva, 2003; Natarajan, Ray, 2010; Azam, 2017). High total cholesterol level is a risk factor of CAD.

\*Correspondence Address:  
Public Health Department, Sport Sciences Faculty, Universitas Negeri Semarang,  
Indonesia.  
Email: mahatulazam@mail.unnes.ac.id

Optimal total cholesterol level in the body is 200 mg/dL (Penana, 2003; Health & Human Service, 2005). Total Cholesterol to HDL Cholesterol TC/HDL ratio is a well-known parameter of CAD risk factor (Millan, 2009; Endarti, 2016). TC/HDL ratio cut-off point of 4.5 is determined as risk factor of CAD (Millan, 2009).

It is generally known that athletes have higher levels of HDL compared to non-athletes. The relationship between physical activity and health and HDL cholesterol has been studied extensively. Some studies have shown that marathon runner, jogging (Scheers, 2008), sky (Taralov, 2000; Kannan, 2014) and other sports athlete have higher HDL cholesterol. Study in Spain for amateur and professional cyclists also found a difference in average total cholesterol and HDL level, and also TC/HDL ratio profile. These parameters are also compared to high intensity workouts for long periods (Aguiló, 2003).

It is known that lipid profiles, especially HDL cholesterol, are strongly influenced by the levels of physical activities on athletes or non-athletes alike. However, little is known about the information of HDL profile and TC/HDL ratio for long distance cyclists based on distance that will describe their daily workout intensity and cycling training habits. The aims of this present study was to investigate the difference in TC/HDL levels ratio between cyclists based on distance and intensity of training.

#### Methods

The study was carried out at Indonesia cycling touring event of 2017 North Coast (NC) and Tour de Borobudur (TdB), which involved 1,211 participants. They consist of 31 NC participants, 207 participants of TdB140K and 973 participants of TdB100K. Study participation was offered through leaflets, social media, and the internet along with the publication of touring event. Inclusion criterias were as follow: male, between 20 to 75 years old. A minimum sample size of 78 participants was required, which was then divided into groups consisting of at least 26 participants into each group: NC240K, TdB140K, and TdB 100K. Exclusion criterias were the following: clinical conditions that

is harmful for participating in the events and a medical history of the previous acute coronary syndrome.

Eighty-eight volunteer and eligible participants of long-distance cycling (LDC) touring, i.e. NC and TdB 2017 (Indonesia), were recruited to this observational study. The study protocol was approved by the local ethical board (no. 607/EC/FK-RSDK/X/2017) and conformed to the concepts of the Declaration of Helsinki. This study was registered in [clinicaltrials.gov](http://clinicaltrials.gov) (ID number NCT03310450). The study was funded by the Directorate of Research and Public Service, Ministry of Research, Technology and Higher Education of the Republic of Indonesia, and delegates all the research procedures to the researchers.

All participants completed a questionnaire about subject characteristics, health conditions, medical history and the intensity of daily cycling activities. Participants have their anthropometric profile measured, i.e. weight, height, body mass index (BMI), waist circumference, hip circumference and waist to hip ratio. Body weight was measured using body weight scale SECA 813<sup>®</sup> and body height was measured using microtoise SECA Bodymeter 206<sup>®</sup>, whereas waist/hip circumferences were measured using ergonomic circumference measuring tape SECA 201<sup>®</sup>.

Blood sample for laboratory analysis was collected one day before cycling tour activities, in the afternoon on after fasting for 8-10 hours. Five milliliter of venous blood was taken from antecubital vein. The blood samples were stored in a silicon tube without anticoagulant and clotted in room temperature for an hour. Then, the blood samples were centrifuged for 10 to 20 minutes at 2.500-3.600 rpm. The serum were aliquoted and transferred into a labeled plastic tube. The levels of total cholesterol and HDL cholesterol were analyzed using biochemical analysis method by an automated analyzer (Cobas 501, Roche Diagnostics, USA).

Statistical analyses were performed using the Statistical Package for the Social Sciences (IBM SPSS Statistics for Windows, Version 23.0, IBM Corp., NY, USA). Numerical data presented in mean±SD for Gaussian data, whereas non-Gaussian data were presented in the interquartile value and categorical data were



presented in proportion. Differences between groups were tested using either Independent t test, one way Anova test or chi-square test. Multivariate analyses was conducted by binary logistic regression.

### Results and Discussion

The subject was classified into three groups based on cycling distances that they participated. The heavier the tour, the greater the required preparation and performance for cyclists. As such, it represent their daily cycling training habits. Group NC 240K, the heaviest tour in this study, rode 240 km from Semarang to Cirebon through Java North Coast main road. NC 240K was held in October 20th 2017 in clear weather, with mean temperature of 33°C. The minimum temperature was 28°C and maximum temperature was 41°C. Total elevation gained during NC 240K through Java North Coast road was 826 m.

TdB was divided into two groups i.e. TdB 140K group and TdB 100K. TdB 140K rode 140 km from Semarang to Magelang via Boja/Limbangan road. TdB 140K took place on November 11th 2017 in almost rainy weather, with mean temperature of 26°C. The minimum temperature was 21°C and maximum

temperature was 37°C. Total elevation gained during TdB 140K was 2,754 m and is often called Climbing Challenge tour.

TdB 100K, the lightest tour of this study, rode 100 km from Semarang to Borobudur via the main road of Semarang-Yogyakarta. TdB 100K was held on November 12th 2017, one day after TdB 140K in very clear weather, with mean temperature of 30°C. The minimum temperature was 28°C and maximum temperature was 34°C. Total elevation gained during TdB 140K was 1,117 m.

Characteristics of participants based on age and anthropometry parameters was presented in Table 1 which showed no significant difference in all parameters, i.e. age, weight, height, body mass index, waist circumference, hip circumference, and waist to hip circumference ratio. These data showed that all parameters were not influenced by the distance of the tour.

Mean age of all participants were 45.3±11.4 years old. The mean age of participants from NC 240K, TdB 140K and TdB 100K were as follows: 43.3±9.33 years old, 43.5±10.18 years old and 49±13.70 years old, respectively. Minimum and maximum age of all participants

Table 1. Subjects Characteristics Based on Tour Groups

No	Parameters	NC 240K n=28	TdB 140K n=30	TdB 100K n=30	Total n=88	p value
<b>Characteristic of Age and Anthropometry</b>						
1.	Age (years, mean±SD)	43.3±9.33	43.5±10.18	49±13.70	45.3±11.47	0.090 <sup>*</sup>
2.	Age (years, min-max)	25-65	24-59	21-71	21-71	
3.	Weight (kg, mean±SD)	67.9±7.51	68.3±9.10	69.8±11.21	67.7±9.36	0.779 <sup>*</sup>
4.	Height (cm, mean±SD)	167.2±5.02	167.1±4.35	167.3±4.5	167.2±4.58	0.969 <sup>*</sup>
5.	BMI (kg/m <sup>2</sup> , mean±SD)	24.2±2.26	24.5±3.11	23.7±3.60	24.2±3.03	0.653 <sup>*</sup>
6.	Waist circumference (cm, mean±SD)	84.2±6.69	86.4±8.50	85.1±9.78	85.2±8.41	0.539 <sup>*</sup>
7.	Hip circumference (cm, mean±SD)	89.8±5.43	89.3±7.09	88.2±7.89	88.2±6.87	0.617 <sup>*</sup>
8.	Waist-hip ratio (mean±SD)	0.94±0.069	0.96±0.044	0.96±0.053	0.95±0.057	0.157 <sup>*</sup>
<b>Characteristic of Workout and Exercise Frequency</b>						
9.	Duration of cycling experience (years, min-max)	1-4-42	5-1-26	4-75-1-25		0.019 <sup>**</sup>
10.	Long distance tour experience		9/30	6/20		0.080 <sup>**</sup>
	No (n%)	2/7.1	21/21	24/20		
	Yes (n%)	26/92.9				
11.	Frequency of tour participating					0.020 <sup>**</sup>
	Never (n, %)	2/7.1	9/30	6/20		
	1-3 times (n, %)	8/32.1	11/36.7	6/20		
	4-6 times (n, %)	2/7.1	7/23.3	11/36.7		
	≥ 7 times	15/52.6	5/16	7/23.3		
12.	Cycling distance in a month (km, mean±SD)	779±380.1	600±222.2	642±318.0		0.025 <sup>**</sup>
<b>Characteristic of Lipid Profile</b>						
13.	TC level (mg/dL, mean±SD)	216.6±55.94	208.1±27.13	243.1±31.95		0.427 <sup>*</sup>
14.	HDL-C level (mg/dL, mean±SD)	88.9±19.09	52.1±13.9	53.6±12.45		<0.001 <sup>**</sup>
15.	TC/HDL-C (ratio, mean±SD)	3.3±1.12	4.2±1.07	4.0±1.06		0.007 <sup>**</sup>

BMI: Body Mass Index \*One way Anova test \*\*Kruskal-Wallis test \*\*\*Chi Square test

was 22 years old and 71 years old; NC 240K: 25 years old and 63 years old; TdB 140K: 24 years old and 59 years old; and TdB 100K: 22 years old and 71 years old. Characteristic of age in all group were not different significantly ( $p>0.05$ ).

Similar to age, anthropometry parameters were not different in all groups. Mean weight of all participants were  $67.7\pm 9.36$  kg. The mean weight in NC 240K, TdB 140K and TdB 100K were as follows:  $67.9\pm 7.51$  kg,  $68.5\pm 9.10$  kg and  $66.8\pm 11.21$  kg, respectively. Mean BMI of all participants were  $24.2\pm 3.03$  kg/m<sup>2</sup>. Mean BMI in NC 240K, TdB 140K and TdB 100K were as follows:  $24.2\pm 2.26$  kg/m<sup>2</sup>,  $24.5\pm 3.11$  kg/m<sup>2</sup> and  $23.7\pm 3.60$  kg/m<sup>2</sup>, respectively. Mean waist circumference of all participants were  $85.2\pm 8.41$  cm. Mean waist circumference in NC 240K, TdB 140K and TdB 100K were as follows:  $84.2\pm 6.69$  cm,  $86.4\pm 8.50$  cm and  $85.1\pm 9.78$  cm, respectively. Mean hip circumference of all participants were  $88.2\pm 6.87$  cm. Mean hip circumference in NC 240K, TdB 140K and TdB 100K were as follows:  $89.8\pm 5.43$  cm,  $89.5\pm 7.09$  cm and  $88.2\pm 7.89$  cm, respectively. Finally, mean waist-hip ratio of all participants were  $0.95\pm 0.057$  cm. Mean hip circumference in NC 240K, TdB 140K and TdB 100K were as follows:  $0.94\pm 0.069$  cm,  $0.96\pm 0.044$  cm and  $0.96\pm 0.053$  cm, respectively.

Distribution of workouts characteristic was also showed in Table 1. It showed a significantly different duration for cycling experience ( $p=0.019$ ), with the median value and the value of minimum-maximum at NC 240K, TdB 140K, TdB 100K as follows 3;0-42, 5;1-26, 4;75;1-25, respectively. The difference was not significant ( $p=0.088$ ) on long distance cycling tour experience categorized as no experienced at all and experienced, with frequency at NC 240K, TdB 140K, TdB 100K as follows 7.1%; 92.9%, 30%; 21% and 20%; 80%, respectively.

Significant difference ( $p<0.05$ ) was also found in frequency of tour participation categorized as never, 1-3 times, 4-6 times and more than seven times, with the frequency at NC240K, TdB140K, TdB100K as follows 7.1%; 32.1% 7.1%; 53.6%, 30%; 36.7%; 23.3%, 20%; 20%; 36.7%; 23.3%, respectively. The cycling distances (km) in a month, showed as mean  $\pm$  SD on the NC240K, TdB140K, TdB100K

were as follows  $779\pm 388.1$  km,  $600\pm 322.2$  km,  $642\pm 318.0$  km, respectively.

Test results of statistical analysis in Table 1 also showed that there are significant differences between the levels of HDL cholesterol (mg/dL) and distance ( $p$  value  $<0.001$ ), with mean and standard deviation of NC240K group  $68.9\pm 19.09$ , TdB140K group  $52.1\pm 13.9$  and TdB100K  $53.6\pm 12.45$ . Then, the ratio of total cholesterol to HDL cholesterol was significantly different as well ( $p=0.007$ ), with mean and standard deviation on NC240K group  $3.3\pm 1.12$ , TdB140K group  $4.0\pm 1.06$ , and TdB100K group  $4.0\pm 1.06$ . Meanwhile, a significant relationship was not found in total cholesterol ( $p=0.427$ ), with mean and standard deviation on NC240K group  $216.6\pm 55$ , TdB140K group  $208.1\pm 27.13$ , and TdB100K group  $203.1\pm 31.95$ .

Levels of TC indicates that different distance did not affect this parameter. These results are in line with the study that stated the load of sports is not associated with levels of TC (Scheers *et al.*, 2008). High levels of HDL-C were found in the group of cyclists with a longer distance, this finding in line with previous studies which showed high levels of HDL-C after performing high intensity sport (Olson, 2000; Skoumas, 2003; Lecheminant, 2005). Discovery of high levels of TC/HDL in cyclists with longer distance on this study is in accordance with the study which stated that the higher the intensity of physical activity, the higher the levels of TC/HDL (Tran, 2003; De Groot, 2003).

This study used two category of TC/HDL ratio with a cut-off point of  $\geq 4.0$  as first category and the second category was  $< 4.0$ , based on cut-off point used to predict CAD. Higher levels of TC/HDL ratio progressively increases the risk of CAD (Robert F. Dons, Frank H. Wians, 2009). Difference parameter cut-off point TC/HDL ratio also used for 4.5 as the target of primary prevention of CAD (Millan, 2009), but since the mean of this ratio in this study were relatively lower in all subjects, less than 4.5 as showed on Table 1, the cut-off point of 4 was used to determine the groups. Then, based on this group, the data was described in Table 2.

Mean age of all participants based on group with TC/HDL ratio  $\geq 4.0$  and

Table 2. Subjects Characteristics Based on TC/HDL Ratio Groups

Variables	TC/HDL		p value
	≥ 4.0	< 4.0	
1. Age and Anthropometry			
2. Age (years, mean±SD)	47.0±9.70	44.6±12.16	0.352
3. Weight (kg, mean±SD)	72.7±7.28	65.5±9.38	0.001
4. Height (cm, mean±SD)	167.7±4.19	166.9±4.85	0.225
5. BMI (kg/m <sup>2</sup> , mean±SD)	26.0±2.63	23.3±2.85	0.001
6. Waist circumference (cm, mean±SD)	89.5±7.50	83.4±8.17	0.001
7. Hip circumference (cm, mean±SD)	92.8±6.13	87.6±6.61	0.001
8. Waist-hip ratio (mean±SD)	4.5±3.03	6.3±7.09	0.349
9. Cycling touring characteristics and training habits			
Duration of cycling experience (years, mean±SD)	4.5±3.03	6.3±7.09	0.225
10. Long distance tour experience			0.899 <sup>†</sup>
No (n;%)	5;29.4	12;70.6	
Yes (n;%)	22;31	49;69	
11. Frequency of participating in cycling tour			0.998 <sup>†</sup>
Newer (n;%)	5;29.4	12;70.6	
1-3 times (n;%)	8;30.8	18;69.2	
4-6 times (n;%)	6;30	11;70	
≥7 times (n;%)	8;32	17;68	
12. Cycling distance in a month (km, mean±SD)	648±336.6	723±372.6	0.354 <sup>†</sup>
13. Type of Tour participated			0.047 <sup>†</sup>
NC (n;%)	7;25.0	21;75.0	
TdTB140K (n;%)	16;53.3	14;46.7	
TdTB 100K (n;%)	14;46.7	16;53.3	

†Independent t-test ††Chi Square test

<4.0 were 47.0±9.70 and 44.6±12.16, and were not significantly different (p=0.352). Anthropometry measurement were significantly different in the groups. Mean weight of participants based on group with TC/HDL ratio ≥ 4.0 and < 4.0 was significantly different (p= 0.001), with the mean of 72.7±7.28 kg vs 65.5±9.38 kg. Mean BMI of participants based on group with TC/HDL ratio ≥4.0 and <4.0 was also significantly different (p= 0.001), with the mean of 26.0±2.63 kg/m<sup>2</sup> vs 23.3±2.85 kg/m<sup>2</sup>. Mean height was not significantly different (p=0.225). These finding is consistent with other studies that concluded BMI is related to serum cholesterol levels (Al-Ajlan, 2011; Kumar, 2014; Nwaiwu, 2015), TC/HDL ratio (Lemieux, 2015), and other conditions due to overall lipid profile (Ali, 2011; Yildiran, 2011; Febrianti, 2013). Quebec study (Lemieux, 2011) showed that mean BMI of 25.3±3.6 tend to be normolipidemic than mean BMI 27.6±3.4 which tend to be dyslipidemic.

Mean waist circumference of all participants based on group with TC/HDL

ratio ≥ 4.0 and < 4.0 were 89.5±7.50 cm and 83.4±8.17, and were significantly different (p=0.001). Similarly, mean hip circumference of participants based on group with TC/HDL ratio ≥ 4.0 and < 4.0 was significantly different (p= 0.001), with mean of 92.8±6.13 cm vs 87.6±6.61 cm. Nevertheless, mean waist to hip ratio which was not significantly different (p=0.349). This finding was in line with many other studies which concluded that waist circumference and hip circumference are related to lipid profile, including TC/HDL ratio (Mohanna, 2006; Alhamdan, 2008; Rizk, 2012; Devi, 2017).

To the best of our knowledge, the present study is the first report concluding characteristic of cycling training related to lipid profile, especially TC/HDL ratio. Based on the test result analysis for characteristic of exercises frequency on Table 2. It was found that all variables, i.e. duration of cycling experience, long distance tour experience, frequency of participating in cycling tour, and cycling distance in a month. Only type of participation, that represent overall characteristics of cycling



Table 3. Multivariate binary logistic regression

Variable	B	Wald	P	OR	95% CI
BMI	-0.364	13.35	0.001	0.695	0.572 - 0.845
Type of tour	-0.580	3.563	0.049	0.827	0.307 - 0.938
Constants	9.772	15.157	0.001	17.338	

R<sup>2</sup>=0.770 (Cox and Snell)

training habits, that have significantly different proportion in both groups ( $p=0.047$ ). The heaviest tour participated, i.e. NC had more proportion of TC/HDL ratio  $< 4$  (75%) than ratio  $\geq 4$  (25%); on the other hand, TdB 140K and TdB 100K have similar proportion of around 50% in TC/HDL ratio  $< 4$  and  $\geq 4$ .

Descriptively, TC/HDL ratio  $\geq 4$  group had mean duration of cycling experience of  $4.5 \pm 3.03$  years and in TC/HDL ratio  $< 4$  group it was  $6.3 \pm 7.09$  years, but statistically was not different ( $p=0.225$ ). Similarly, long-distance cycling tour experience, which were divided into two categories, i.e. no experienced participating at all and experienced, had no significantly different proportion in both groups based on TC/HDL ratio  $\geq 4$  and TC/HDL ratio  $< 4$ . The proportion were about 30% in TC/HDL ratio  $\geq 4$  and about 70% in TC/HDL ratio  $< 4$ , both for inexperienced and experienced groups. Statistically, there was no different proportion ( $p=0.899$ ).

The same goes with frequency of participation in cycling tour, which were divided into four categories, i.e. Never, 1-3 times, 4-6 times, and more than 7 times. In all groups, proportion were about 30% in TC/HDL ratio  $\geq 4$  and about 70% in TC/HDL ratio  $< 4$ . Statistical analysis also found that there was no significant different proportion ( $p=0.998$ ). Mean distance of cycling over the past month which was done by participants indicated that there is no significance as well ( $p=0.354$ ). Descriptively, participants with TC/HDL ratio  $\geq 4$  had mean distance of  $648 \pm 336.6$  km vs participants with TC/HDL ratio  $< 4$  who had higher mean distance of  $723 \pm 372.6$  km.

Previous study in 2010, which was conducted to 36 non-trained men, found that high intensive exercise of interval running for 40 minutes within 12 weeks lowered the ratio of TC/HDL by 0.08 (Nybo, 2010). Other study also reported that physical training for 6 weeks to 30

men were able to lower their ratio of TC/HDL significantly (Sheikholeslami Vatani, 2011). It proves that people who have experience of exercise or cycling and doing longer physical activity will have lower levels of TC/HDL ratio. Many studies also concluded that exercise habits is related to body composition and lipid profile (Trejo-Gutierrez, 2007; Albuquerque Filho, 2014).

Multivariate analysis was conducted to find the best model of parameters that influence TC/HDL ratio in long distance cyclists. Multivariate analysis involved all independent variables having  $p < 0.25$  in bivariate analysis influenced TC/HDL ratio status. Those variables were: weight, BMI, waist circumference, hip circumference, duration of cycling experience, and type of cycling tour participated. A backward stepwise binary logistic regression was used to determine variables that influenced TC/HDL ratio. Final regression model was shown in Table 2.

Multivariate analysis showed that BMI and type of tour participated were the best model that contribute to TC/HDL ratio category of  $\geq 4.0$  and  $< 4.0$ . Present study identified that lower BMI ( $\beta=-0.364$ ,  $p=0.001$ ) with OR 0.695 (95% CI=0.572 - 0.845) and heaviest of cycling tour participated ( $\beta=-0.580$ ,  $p=0.049$ ) with OR 0.827 (95% CI=0.307 - 0.938) were the most influential variables toward TC/HDL ratio status in long distance cyclists. Our final regression model have R<sup>2</sup> (Cox and Snell) = 0.770, showing that 77.0 % of TC/HDL ratio status  $< 4$  in long distance cyclists was influenced by this model variant (BMI and type of cycling tour) and the other 23.0% were influenced by other factors.

This study have limitations. This study did not measure complete lipid profile, such as, triglyceride, low density lipoprotein cholesterol, apolipoproteins, and other parameters which are the more valuable predictor of CAD events.

Lipid profile is predicted to have high levels after doing long distance cycling activities. In addition, total cholesterol, HDL, TC/HDL ratio were not measured before and after long-distance cycling activity.

#### Conclusion

There is significant differences in the levels of HDL-C based on the characteristics of cycling distance and cycling training habits. Present study also showed that there is difference in TC/HDL ratio in type of cycling tour which represent cyclist's daily exercise habits. Study also found that anthropometry measurement i.e. weight, BMI, waist circumference, and hip circumference have strong association with TC/HDL ratio status as well as the characteristics of the cycling tour. Final regression model involved BMI and type of tour to predict TC/HDL ratio status.

#### Practical Implications:

Long distance cyclists have a synergistic effect of lipid profile (TC/HDL ratio) and anthropometry measurements (weight, BMI, waist and hip circumference)

Long distance cyclists are shown to have an average TC/HDL ratio less than 4.5, in which cut-off point of CAD risk factor determined.

Experienced cyclists who participated in heavier cycling tour (NC cyclists), which represent their cycling training habits, tend to have lower TC/HDL ratio (< 4).

#### Acknowledgment

The study was funded by the Directorate of Research and Community Services, Ministry of Research, Technology and Higher Education of Republic Indonesia (DIPA 042.06.1.401516). The authors thank all investigators and organizers of NC 240K, TdB 140K and TdB 100K cycling tour and participants involved in this study.

#### References

- Aguiló, A., Tauler, P., Pilar, G.M., Villa, J., Córdova, A., Tur, J.A., & Pons, A., 2003. Effect of Exercise Intensity and Training on Antioxidants and Cholesterol Profile in Cyclists. *Journal of Nutritional Biochemistry*, 14(6), pp.319–325.
- Ahn, N., & Kim, K., 2016. High-Density Lipoprotein Cholesterol (Hdl-C) in Cardiovascular Disease: Effect of Exercise Training. *Integrative Medicine Research*, 5(3), pp.212–215.
- Al-Ajlan, A.R., 2011. Lipid Profile in Relation to Anthropometric Measurements among College Male Students in Riyadh, Saudi Arabia: A Cross-sectional Study. *International Journal of Biomedical Science*, 7(2), pp.112–119.
- Albuquerque, F.N., Rebouças, G., Matos, V., Salgueiro, C., Knackfuss, M.L., & de Medeiros, H., 2014. Effect of Concurrent Training on Body Composition and Lipid Profile in Overweight Adolescents. *Journal of Exercise Physiology Online*, 17(6), pp.33–44.
- Alhamsan, A.A., 2008. Body Mass Index, Waist, Waist to Hip Ratio and Lipid Profile in Elderly Subjects Living in a Nursing Home. *Journal of Medical Sciences*, 8(2), pp.177–181.
- Ali, Z.A.U.A., & Al-Zaidi, M.S., 2011. The Association between Body Mass Index, Lipid Profile and Serum Estradiol Levels in a Sample of Iraqi Diabetic Premenopausal Women. *Oman Medical Journal*, 26(4), pp.263–266.
- Arquer, A., Elosua, R., Covas, M.L., Molina, L., & Marrugat, J., 2006. Amount and Intensity of Physical Activity, Fitness, and Serum Lipids in Pre-Menopausal Women. *International Journal of Sports Medicine*, 27(11), pp.911–918.
- Azam, M., Nida, I., Harydayani, O.W.K., & Fibrina, A.I., 2017. Lipid Profiles and Postprandial Glucose Status of Hypertensive Individuals with Stroke in Indonesia. *Asian Journal of Epidemiology*, 10(3), pp.116–122.
- Devi, S., Choudhary, A.K., Verma, P., Jain, N., & Garg, N., 2017. Association of Lipid Profile, Body Mass Index, and Waist Circumference as Cardiovascular Risk Factors for Obese Male Adults of North India. *International Journal of Scientific Study*, 4(10), pp.148–153.
- Endarti, A.T., & Handito, A., 2016. The Relationship between Non-Communicable Diseases History and Health-Related Quality of Life. *Kemas*, 12(1), pp.120–126.
- Febrianti, E.S.Z., Asviandri, Farina, L., Lestari, R., Cahyohadi, S., & Rini, E.A., 2013. Correlation between Lipid Profiles and Body Mass Index of Adolescents Obesity in Padang. *International Journal of Pediatric Endocrinology*, 2013.
- De Groot, P.C., Hjeltnes, N., Heijboer, A.C., Stul, W., Birkeland, K., 2003. Effect of Training Intensity on Physical Capacity, Lipid Profile and Insulin Sensitivity in Early Rehabilitation of Spinal Cord Injured Individuals. *Spinal Cord*, 41(12), pp.673–679.
- Health and Human Service, D., 2005. Lowering Your Cholesterol with TLC (Therapeutic Lifestyle Changes). *National Institute of Health*, pp.1–



- Kannan, U., Vasudevan, K., Balasubramaniam, K., Yerrabelli, D., Shanmugavel, K., & John, N. A., 2014. Effect of Exercise Intensity on Lipid Profile in Sedentary Obese Adults. *Journal of Clinical and Diagnostic Research*, 8(7).
- Kumar, A., 2014. Correlation of Serum Paraoxonase Activities in Known Cases of 130 Elderly Hypertensive South Asian Aged 56-64 Years - a Hospital Based Study. *Asian Pacific Journal of Tropical Disease*, 4(51).
- Lecheminant, J.D., Tucker, L.A., Bailey, B.W., & Peterson, T., 2005. The Relationship between Intensity of Physical Activity and HDL Cholesterol in 272 Women. *Journal of Physical Activity and Health*, 3, pp.333-344.
- Lemieux, I., 2011. Total Cholesterol/HDL Cholesterol Ratio vs LDL Cholesterol/HDL Cholesterol Ratio as Indices of Ischemic Heart Disease Risk in Men. *Archives of Internal Medicine*, 161(22), pp.2685.
- Lemieux, I., Lamarche, B., Couillard, C., Pascot, A., Carlin, B., Bergeron, J., Dagenais, G.R., & Després, J.P., 2015. Total Cholesterol/HDL Cholesterol Ratio vs LDL Cholesterol/HDL Cholesterol Ratio as Indices of Ischemic Heart Disease Risk in Men: the Quebec Cardiovascular Study. *Archives of Internal Medicine*, 161, pp.2685-2692.
- Millán, J., Pintó, X., Muñoz, A., Zúñiga, M., Rubiés-Prat, J., Pallardo, L.F., Masana, L., Mangas, A., Hernández-Mijares, A., González-Santos, P., Ascaso, J.F., Pedro-Botet, J., 2009. Lipoprotein Ratios: Physiological Significance and Clinical Usefulness in Cardiovascular Prevention. *Vascular Health and Risk Management*, 5, pp.757-765.
- Mohanna, S., Baracco, R., & Seclén, S., 2006. Lipid Profile, Waist Circumference, and Body Mass Index in a High Altitude Population. *High Altitude Medicine & Biology*, 7(3), pp.245-255.
- Natarajan, P., Ray, K.K., & Cannon, C.P., 2010. High-Density Lipoprotein and Coronary Heart Disease. Current and Future Therapies. *Journal of the American College of Cardiology*, 55(13), pp.1283-1299.
- Nwaiwu, O., & Ibe, B.C., 2015. Relationship between Serum Cholesterol and Body Mass Index in Nigeria Schoolchildren Aged 2-15 Years. *Journal of Tropical Pediatrics*, 61(2), pp.126-130.
- Nybo, L., Sundstrup, E., Jakobsen, M.D., Mohr, M., Hornstrup, T., Simonsen, L., Bülow, J., Randers, M.B., Nielsen, J.J., Aagaard, P., Krstrup, P., 2010. High-Intensity Training Versus Traditional Exercise Interventions for Promoting Health. *Medicine and Science in Sports and Exercise*, 42(10), pp.1951-1958.
- Olson, M.B., Kelsey, S.F., Bittner, V., Reis, S.E., Reichel, N., Handberg, E.M., & Merz, C.N., 2000. Weight Cycling and High-Density Lipoprotein Cholesterol: Women: Evidence of an Adverse Effect. *Journal of the American College of Cardiology*, 36(5), pp.1565-1571.
- Penalva, R.A., Huoyn, M.Q., Correia, J.C., Feitosa, G.S., & Ladeta, A.M., 2003. Original Article Lipid Profile and Severity of Atherosclerotic Disease in Acute Coronary Syndrome. pp.24-29.
- Riwanto, M., & Landmesser, U., 2013. High Density Lipoproteins and Endothelial Functions: Mechanistic Insights and Alterations in Cardiovascular Disease. *Journal of Lipid Research*, 54(12), pp.3227-3243.
- Rizk, N.M., & Yousef, M., 2012. Association of Lipid Profile and Waist Circumference as Cardiovascular Risk Factors for Overweight and Obesity among School Children in Qatar. *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 5, pp.425-432.
- Robert, F.D., Frank, H., & Wiats, J., 2009. *Endocrine and Metabolic Disorders: Clinical Lab Testing Manual*. 4th edn. CRC Press Taylor & Francis Group.
- Rosjidi, C.H., Isroin, L., & Wahyuni, N.S., 2017. Differences in Risk Factor of Cardiovascular Disease Risk on Rural and Urban Population. *Kemas*, 13(1), pp.69-76.
- Scheers, T., Philippaerts, R., Langendonck, L., Van Duquet, W., Duvigneaud, N., Matton, L., Thomis, M., Wijndaele, K., & Lefevre, J., 2008. Lipid Profile in Men and Women with Different Levels of Sports Participation and Physical Activity. *Public Health Nutrition*, 11(11), pp.109-1106.
- Sheikholeslami, V.D., Ahmadi, S., Ahmadi D.K., & Gharibi, F., 2011. Changes in Cardiovascular Risk Factors and Inflammatory Markers of Young, Healthy, Men after Six Weeks of Moderate or High Intensity Resistance Training. *J Sports Med Phys Fitness*, 51(4), pp.695-700.
- Skouras, J., Pitsavos, C., Panagiotakos, D.B., Chrysohoou, C., Zembekis, A., Papaioannou, I., Toutouza, M., Toutouzas, P., & Stefanadis, C., 2003. Physical Activity, High Density Lipoprotein Cholesterol and other Lipids Levels, in Men and Women from the ATTICA Study. *Lipids in Health and Disease*, 2, pp.1-7.

- Taralov, Z., Boyadjiev, N., & Georgieva, K., 2000. Serum Lipid Profile in Pubescent Athletes. *Acta Physiologica et Pharmacologica Bulgarica*, 25(1), pp.3–8.
- Tran, Z.V., Weltman, A., Glass, G.V., & Mood, D.P., 2003. The Effects of Exercise on Blood Lipids and Lipoproteins: A Meta-Analysis of Studies. *Medicine and Science in Sports and Exercise*, 15(5), pp.393–402.
- Trejo-Gutierrez, J.F., & Fletcher, G., 2007. Impact of Exercise on Blood Lipids and Lipoproteins. *Journal of Clinical Lipidology*, pp.175–181.
- Yildiran, H., Acar, T.N., Keksalp, E., Geremen, K.M., Akbulut, G., Illici, S., & Sanlier, N., 2011. The Association of Anthropometric Measurements and Lipid Profiles in Turkish Hypertensive Adults. *African Health Sciences*, 11(3), pp.407–413.

# Anthropometric-Parameters and Total-Cholesterol to HDL-Cholesterol Ratio are Better in Long-Distance Cyclists

## ORIGINALITY REPORT

7%

SIMILARITY INDEX

7%

INTERNET SOURCES

4%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1

[garuda.ristekdikti.go.id](http://garuda.ristekdikti.go.id)

Internet Source

3%

2

[link.springer.com](http://link.springer.com)

Internet Source

2%

3

Submitted to Wakefield College

Student Paper

2%

Exclude quotes Off

Exclude matches < 2%

Exclude bibliography Off



# Anthropometric-Parameters and Total-Cholesterol to HDL-Cholesterol Ratio are Better in Long-Distance Cyclists

---

GRADEMARK REPORT

---

FINAL GRADE

**/0**

GENERAL COMMENTS

**Instructor**

---

PAGE 1

---

PAGE 2

---

PAGE 3

---

PAGE 4

---

PAGE 5

---

PAGE 6

---

PAGE 7

---

PAGE 8

---

PAGE 9

---