

TURNITIN-Correlation-Between-Portal-Venous-Dimensions

by A. Gunawan Santoso

Submission date: 15-May-2023 11:05AM (UTC+0700)

Submission ID: 2093340910

File name: ARTIKEL_dr._GUNAWAN_MEDICA_HOSPITALIA.docx (76.42K)

Word count: 3225

Character count: 18090



Original Article

Correlation Between Portal Venous Dimensions and Liver Stiffness in Patients of Child Pugh A Cirrhosis

Antonius Gunawan Santoso¹, Nala Khodlil Mubarok¹,
Maya Nuriya Widayarsi¹, Hery Djagat Purnomo²

¹Departement of Radiology Faculty of Medicine Diponegoro University/ Dr. Kariadi Hospital Semarang, Indonesia

²Sub Division Gastroenterohepatology Faculty of Medicine Diponegoro University/
Dr. Kariadi Hospital Semarang, Indonesia

Abstract

p-ISSN: 2301-4369 e-ISSN: 2685-7898
<https://doi.org/10.36408/mhjcm.v10i1.899>

Accepted: January 16th, 2022

Approved: March 13th, 2023

Author Affiliation:

Departement of Radiology
Faculty of Medicine Diponegoro University/
Dr. Kariadi Hospital Semarang,
Indonesia

Author Correspondence:

Antonius Gunawan Santoso
Dr. Sutomo Street No. 16 Semarang,
Central Java 50244, Indonesia

E-mail:

santoso_gun@yahoo.com

Background: Liver cirrhosis is an end-stage liver disease characterized by pathologic fibrosis and regenerative nodules with resultant liver dysfunction. The diagnostic hallmark of portal hypertension is slow flow velocities in addition to the increased caliber of the mean portal vein. That is, the diagnosis of portal hypertension requires the measurement of mean portal velocity and portal diameter, and the correlation between cirrhosis and mean portal velocity and portal diameter is the correlation between liver cirrhosis and its complications. The objectives of this study was to analyze the correlation between portal vein dimensions and liver stiffness in patients of Child-Pugh A cirrhosis.

Methods: This study was a cross-sectional study on 30 subjects with Child-Pugh A liver cirrhosis. The subjects of this research are patients who come to the Radiology Department of the Dr. Kariadi Hospital in Semarang for point shear wave elastography and abdominal ultrasound examination from January to December 2022. Spearman test correlation was used for the analysis.

Results: Spearman test showed no correlation between liver stiffness and portal vein diameter ($p=0.250$, $r=-0.217$), liver stiffness and mean portal vein velocity ($p=0.883$, $r=-0.028$), and portal vein diameter with mean venous velocity in Child-Pugh A liver cirrhosis ($p=0.979$, $r=0.005$)

Keywords: Liver stiffness, portal vein diameter, portal vein velocity, child-pugh A

INTRODUCTION

Liver fibrosis develops as a reversible consequence of a sustained or repeated wound-healing response to liver injury caused by viral, toxic, and/or metabolic insults, and represents an imbalance between the synthesis and degeneration of the extracellular matrix. Accompanied by the distortion of hepatic structure and function, cirrhosis is the result of the progression of liver fibrosis. Liver cirrhosis is a fatal condition with a high fatality rate. The 12th leading cause of death in the US and the fifth leading cause of death for adults between the ages of 45 and 54 is cirrhosis and its consequences. According to the latest Global Burden of Disease Study, the global incidence of cirrhosis and other chronic liver diseases in 2017 was 5,154,900.³ Moreover, from 2007 to 2017, the years lived with disability (or 'YLDs') and all-age deaths from cirrhosis have increased by 34.8% and 15.0% respectively. Hence, the clinical burden of cirrhosis is substantial. Even so, the advancement of liver cirrhosis can be stopped or slowed down with prompt and accurate screening and diagnosis.^{1,14,15}

Biopsy and histopathological evaluation remain the golden standard for assessing liver fibrosis. However, this procedure is painful and has many complications including bleeding (0.3%) and death (0.01%). Hence, early recognition of patients with chronic liver disease (CLD) at high risk for developing its complication in a noninvasive manner is warranted to allow the implementation of optimal preventative management strategies that may modify the natural course of the disease. Cirrhosis itself, on the other hand, causes stiffening, prompting standardized cut-off value for its measurement. Cirrhosis is also known to lead to portal hypertension and metabolic liver failure. Cirrhosis causes intrahepatic portal hypertension secondary to increased hepatic venular resistance caused by intrahepatic fibrosis. The diagnostic hallmark of portal hypertension is slow flow velocities in addition to the increased caliber of the mean portal vein. That is, the diagnosis of portal hypertension requires the measurement of mean portal velocity and portal diameter, and the correlation between cirrhosis and mean portal velocity and port diameter is the correlation between liver cirrhosis and its complications.^{14-17,19}

The purpose of this study is to evaluate the relationship between liver stiffness in patients with Child-Pugh A cirrhosis and the dimensions of the portal vein. The findings of this study are anticipated to serve as a foundation for the management and predictors of mortality in liver cirrhosis.

METHODS

This study employs a cross-sectional methodology. The liver stiffness variable was measured using point SWE using Acuson S2000 from Siemens at RSUP Dr. Kariadi at a specific time throughout the research's data collection period, which ran from July 2022 to August 2022. The ethical council of the Faculty of Medicine at Diponegoro University, Semarang approved the procedure (No.1097/EC/KEPK-RSDK/2022). The cross-sectional study design was used to determine the sample size. In the study of Hong *et al.*, the correlation coefficient between the diameters of the portal vein and the stiffness of the liver is 0.5. The minimal sample size is 30 subjects according to the sample size calculation.⁵

The inclusion criteria in this study are age ≥ 18 years old, Child-Pugh A classification of liver cirrhosis, clinical cirrhosis of the liver, previous sonography showing features of liver cirrhosis and not with acute chronic liver failure, history of iron overload, biliary obstruction (primary sclerosing cholangitis and cholestasis), and passive congestion leading to congestive hepatopathy (congestive heart failure, congenital heart disease with pulmonary stenosis and tricuspid regurgitation). The exclusion criteria for study subjects are invalid readings from point SWE or portal hypertension which is unlikely for technical reasons.¹⁸

The patient fasted for 4-6 hours before the examination. The examination was performed with the patient in a supine or left lateral position with the arm. The measurement was made through the intercostal space as the best location for the acoustic window. The transducer was then positioned perpendicular to the liver capsule about 1-2 cm below the liver capsule to avoid reverberation artifacts. The optimal measurement depth is 4-5 cm from the skin layer. Region of Interest (ROI) should avoid major vessels, bile ducts, costal/rib shadows, and masses. The ROI area is about 1 cm³. The examination was carried out in a neutral position. The patient was asked to take a deep breath and hold it. The examination was repeated ten times. The median value of the examinations was then used as the result of the elastography examination. The diameter of the inner wall (lumen) of the portal vein was measured in millimeters using sonography. Doppler sonography was used to calculate the mean highest and lowest velocity of portal vein blood flow in centimeters per second.

To prove the relationship between portal vein dimensions and liver stiffness, the authors also collected data regarding the subjects' age, sex, BMI, duration of illness, history of previous illness, history of alcohol consumption, and the subjects' previous laboratory test results. The formula used to prove the correlation will be Pearson if the data distribution is normal, or Rank-Spearman if one or both of them are not normally distributed.

RESULTS

In this study, 30 patients already classified with the Child-Pugh classification were chosen; consisting of 17 men (56.67 %) and 13 women (43.33 %). Two-thirds of the subjects were more than fifty years old (66.67%). Twenty-six of the samples (86.67%) have been ill for more than 24 months, two (6.67%) admitted to being ill for 1–6

TABLE 1
Distribution of study sample based on gender, age, length of illness, BMI, liver stiffness, portal vein diameter and mean portal vein velocity

Variabel	Klasifikasi	N	%
Gender	Male	17	56.67
	Female	13	43.33
Age (years)	≤30	3	10
	31–50	8	26.7
	>50	19	63.3
Duration of illness (month)	1–6	2	6.67
	7–12	2	6.67
	13–18	–	–
	19–24	–	–
	>24	26	86.67
BMI (kg/m ²)	<18.5	5	16.67
	18.5–24.9	16	53.30
	25.0–29.9	6	20.00
	30.0–34.9	3	10.00
Liver stiffness (kPa)	10–15	10	33.3
	15–20	8	26.7
	20–25	3	10.0
	25–30	5	16.7
	>30	4	13.3
Portal vein diameter (mm)	<8	3	10.0
	8–10	6	20.0
	10–12	3	10.0
	12–14	14	46.7
	>14	4	13.3
Mean portal vein velocity (cm/s)	<15	4	13.33
	15–30	18	60.00
	30–45	5	16.67
	>45	3	10

months, and another two (6.67%) admitted to being ill for 7–12 months. Sixteen (53.3%) subjects were considered within the normal weight range (18.5–24.9 kg/m²), and only three of the subjects were obese (30.0–34.9 kg/m²).

Ten subjects (33.3%) had the result between 10–15 kPa when examined for liver stiffness, while 5 (16.67%) of them had the result of >30 kPa. Fourteen subjects' portal vein diameters were between 12–14 mm, while 3 subjects (10%) were <8 mm and another 3 were 10–12mm. Eighteen (60%) subjects were found with mean portal vein velocity between 15–30 cm/s, and only three (1%) were found with >45 cm/s.

The results of the data normality test using the Shapiro-Wilk test showed that the liver stiffness variable, portal vein diameter, and mean portal vein velocity were not normally distributed (sig. > 0.05), leading Spearman to be used to analyze the correlation between the two variables.

The results of the Spearman test prove that there is no correlation between liver stiffness and portal vein diameter ($p = 0.250$, $r = -0.217$) and the mean portal vein velocity ($p = 0.883$, $r = -0.028$). There is no correlation between portal vein diameter and mean portal vein velocity with $p = 0.979$ and $r = -0.005$.

DISCUSSION

All chronic liver diseases, whether toxic, hereditary, autoimmune, or infectious, undergo typical histologic changes, ultimately leading to fibrosis/cirrhosis and excessive matrix deposition. Cirrhosis rapidly decompensates and has a high mortality rate. Patients with cirrhosis have reduced liver capacity to metabolize and synthesize proteins, peptides, and hormones. In addition, fibrosis progression and nodal regeneration cause an increase in portal vascular resistance with portal hypertension and an increase in the hepatic venous pressure gradient (HVPG) of 0.10 mmHg. Portal hypertension eventually leads to ascites and the development of vascular collaterals such as esophageal varices. Portal hypertension finally leads to ascites, and vascular collaterals will develop such as esophageal varices. Signs of these clinical features are what we look for when classifying liver disease patients with Child-Pugh classification. Even so, the patients in our study were of Child-Pugh A, whose liver dysfunction is still compensated. This may be why none of our results shows a correlation with one another.¹⁴

Liver stiffness can be affected by confounding factors such as hepatitis, mechanic cholestasis, liver congestion, cellular infiltrations, and deposition of amyloid irrespective of fibrosis stage. Patients with cholestatic liver diseases, such as primary biliary cirrhosis and primary sclerosing cholangitis, seem to have higher stiffness than those with viral hepatitis. Therefore, for each stage of fibrosis, cutoffs are higher than in chronic

TABLE 2
Spearman's test of liver stiffness, portal vein diameter, mean portal vein velocity

Spearman's rho		Liver stiffness	Portal vein diameter	Portal vein velocity
Liver stiffness	Correlation Coefficient	1.000	-.217	-.028
	Sig. (2-tailed)	.	.250	.883
	N	30	30	30
Portal vein diameter	Correlation Coefficient	-.217	1.000	.005
	Sig. (2-tailed)	.250	.	.979
	N	30	30	30
Portal vein velocity	Correlation Coefficient	-.028	.005	1.000
	Sig. (2-tailed)	.883	.979	.
	N	30	30	30

viral hepatitis either because of the nature of the liver disease or because of cholestasis. Similar higher cutoffs for each fibrosis stage were described in alcoholic liver disease. It should be mentioned that steatosis does not increase liver stiffness although it is often regarded as an essential initial state in chronic liver disease. Rather, steatosis may slightly decrease liver stiffness. As seen from our measurement data, our patients have varied levels of liver stiffness. This variety might affect the result of this research.^{14,20}

Correlation between liver stiffness and the mean portal vein velocity

According to the Spearman test results, which were $p=0.883$ ($p > 0.05$) and $r = -0.028$, there is no link between liver stiffness and the mean portal vein velocity. This is consistent with earlier research that linked liver stiffness to mean portal vein velocity such as Gunawan Y *et al.* (2022) which found no connection between liver stiffness and mean portal vein velocity and SWE. However, it disagreed with Sudirman I (2018), who demonstrated a weak correlation ($r=0.271$) between mean portal vein velocity as determined by ultrasonography and the degree of liver stiffness. Meanwhile, there is a significant correlation between the degree of fibrosis and the severity of fibrosis detected with SWE ($p=0.001$, $r=0.672$).^{6,7}

Taking anti-hypertension medication may play a role. In this study, 4 of the subjects (10%) had been using propranolol for more than 3 months, and 15 (37.5%) had been using propranolol plus spironolactone. According to Amin *et al.* (2018), this can affect portal velocity in subjects because propranolol can lower portal pressure by reducing portal vein blood flow. This is associated with reduced cardiac output caused by the blockade of B1 adrenergic receptors.^{8,9}

Correlation between liver stiffness and portal vein diameter

The results of this study showed no correlation between liver stiffness and portal vein diameter, measured at values of $p=0.250$ ($p > 0.05$) and $r = -0.217$. This is consistent with previous studies that correlated the degree of hepatic stiffness with portal vein diameter, including Sudirman I (2018), which showed no significant relationship between portal vein diameter and degree of hepatic stiffness ($r = 0.166$), but in contrast to Zaghoul S.G. (2019) who showed that changes in portal vein diameter are significantly correlated with liver stiffness.^{6,10}

This may be due to several confounding factors beyond the investigator's control, such as the subjects' age and body weight. Ahmed M (2019) showed a significant correlation ($r = 0.234$; $P=0.019$ and $r = 0.22$; $P=0.028$) between portal vein diameter, age, and body weight, and Leao *et al.* (2012) showed that portal vein diameters were correlated significantly with age and BMI ($p=0.02$ and 0.001). Since the portal vein diameter rises with age with age and liver

Other variables that might affect the result are the level of subjects' ALT and the duration of their illness. An increase in portal vein diameter showed a statistically significant association with the duration of illness of more than six months ($P < 0.02$) and increased serum alanine aminotransferase/ALT levels ($P < 0.03$). Several studies have shown that increased aminotransferase levels are associated with higher shear wave velocity by virtual touch tissue quantification (VTQ) compared to that observed in patients with slightly elevated aminotransferases, which means that ALT serum level can also affect liver stiffness. The medication also played a part, as the mean portal vein diameter decreased significantly after 3 months of therapy with propranolol.

12

Propranolol is the most widely used non-selective beta blocker (NSBB) that causes a decrease in portal pressure due to reduced heart rate.^{8,10,11}

14

Correlation between portal vein diameter and portal vein velocity

The results of this study showed that there is no correlation between portal vein diameter and mean portal vein velocity ($p=0.979$ ($p > 0.05$) and $r = 0.005$). This is contrasting previous studies that show a correlation between portal vein diameter and the mean portal vein velocity, including Ahmed M (2019) ($r=-0.628$; $P=0.0$).¹¹

Blood flow should be easier when there is no resistance, and velocity should be higher when the vein diameter is wider. This result might be influenced by previously mentioned factors that affect the portal vein. Further study might be necessary to confirm the hypothesis.

Limitations of the study

There were uncontrolled confounding factors in this study, including subjects' antihypertensive drug use (propranolol, bisoprolol, and spironolactone) over 3 months, age, BMI, disease duration, and serum ALT.

There are no cutoffs for liver stiffness grade, mean portal velocity, and portal diameter in this study because point SWE and ultrasonography are not the gold standards for diagnosis. Liver biopsy is the gold standard for assessing liver stiffness, but MR elastography is the best non-invasive method to assess liver stiffness. The gold standard test for portal hypertension is currently the HVPG.

CONCLUSION

10

This study proves that there is no correlation between portal vein dimensions and liver stiffness in patients with Child-Pugh A liver cirrhosis. Further research may be necessary with more controlled variables to confirm the hypothesis.

REFERENCES

1. Wang J, Wang Q, Yu G *et al.* Correlation Between Liver Stiffness Measured by Shear Wave Elastography And Child-Pugh Classification. *J Ultrasound Med* 2018; 00:00–00
2. Hoodeshenas S, Yin M, Venkatesh SK. Magnetic Resonance Elastography of Liver Current Update. 2018; Volume 27: 319–333
3. Jansen C, Bogs C, Verlinden W, Thiele M, Möller P, Görtzen J, Lehmann J, Vanwollegem T, Vonghia L, Praktijnjo M, Chang J, Krag A, Strassburg CP, Francque S, Trebicka J. Jansen C, *et al.* *Liver Int.* 2017 Mar;37(3):396–405. doi: 10.1111/liv.13243. Epub 2016 Sep 25. *Liver Int.* 2017. PMID: 27569696
4. Bhargava SK. Textbook of Color Doppler Imaging second edition. Jaypee brothers Medical Publishers (P) Ltd. 2010. New Delhi, India
5. Hong WK, Kim MY, Baik SK, Shin SY, *et al.* The usefulness of non-invasive liver stiffness measurements in predicting clinically significant portal hypertension in cirrhotic patients: Korean data. *Clin Mol Hepatol* 2013; 19: 370–375 [PMID: 24459641 DOI: 10.3350/cmh.2013.19.4.370]
6. Sudirman I. Kesesuaian Gambaran Ultrasonografi Transabdominal Duplex Hepar Dan Lien Serta *Transient Elastography (Fibroscan)* Dalam Menilai Fibrosis Hepar Pada Pasien Hepatitis Virus Kronik. Fakultas Kedokteran Universitas Hasanuddin Makassar. 2018
7. Gunawan Y, Murtala B and Asriyani S. Evaluasi *Shear Wave Elastography*, Fibroscan dan Kecepatan Aliran Vena Porta pada Pasien *Non-Alcoholic Fatty Liver Disease*. *Jurnal Biomedik: JBM* 2022; 14(1): 10–16
8. Leao J, Brock M, Castilho M, Scariot A, *et al.* Non-Invasive Assessment of Fibrosis Using Color Doppler Ultrasound in Patients with Hepatitis C Virus in the Amazon Rainforest, Brazil. *The American Society of Tropical Medicine and Hygiene.* 2012. pp. 273–279
9. Amin AI, Zaghoul SG, Lakouz *et al.* Effect Of Non-Selective Beta Blockers On Esophageal Varices And Portal Vein Diameter In Cirrhotic Hcv Patients. *Z.U.M.J.* Vol. 24; No.5 September.;2018: 449–455
10. Zaghoul S.G, Wahab E.A, Seleem W.M *et al.* Impact of non-selective beta blockers on portal hypertension and hepatic elasticity in hepatitis C virus-related liver cirrhosis. *Drug Discoveries & Therapeutics.* 2019; 13(2):108–113.
11. Ahmed, M. Assessment of normal portal vein diameter and velocity using ultrasound. *Clin Radiol Imag.* 2019. J; 3: 000143
12. Luo W, Meng Y, Ji H.L *et al.* Spironolactone Lowers Portal Hypertension by Inhibiting Liver Fibrosis, ROCK-2 Activity and Activating NO/PKG Pathway in the Bile-Duct-Ligated R a t . *P L o S O N E* 7 (3) . 2 0 1 2 : e 3 4 2 3 0 . doi:10.1371/journal.pone.0034230
13. Boursier J, Isselin G, Fouchard- Hubert I, Oberti F, Dib N, Lebigoit J, *et al.* Acoustic radiation force impulse: A new ultrasonographic technology for the widespread noninvasive diagnosis of liver fibrosis. *European Journal of Gastroenterology & Hepatology.* 2010;22(9):1074–1084
14. Sebastian Mueller & Laurent Sandrin. Liver stiffness: a novel parameter for the diagnosis of liver disease, *Hepatic Medicine: Evidence and Research.* 2010;49-67, DOI: 10.2147/hmer.s7394
15. Wu, L, Shen, Y., & Li, F. Non-invasive diagnosis of liver fibrosis: A review of current imaging modalities. *Gastroenterología y Hepatología (English Edition).* 2020;43(4), 211–221. doi:10.1016/j.gastre.2019.11.006
16. Zaki, M., Hazem, M. & Elsamman, M. Shear wave elastography in assessment of liver stiffness and prediction of gastroesophageal varices in patients with liver cirrhosis. *Egypt J Radiol Nucl Med* 50, 16 (2019). <https://doi.org/10.1186/s43055-019-0015-x>
17. Iranpour P, Lall C, Houshyar R, Helmy M, Yang A, Choi JI, *et al.* Altered Doppler flow patterns in cirrhotic patients: an overview. *Ultrasonography.* 2016 Jan;35(1):3–12.
18. Tisoris A, Marlar CA. Use Of The Child Pugh Score In Liver Disease. In: *StatPearls.* StatPearls Publishing, Treasure Island (FL); 2022. PMID: 31194448.
19. Singh, S., Fujii, L. L., Murad, M. H., Wang, Z., Asrani, S. K., Ehman, R. L., Talwalkar, J. A. Liver Stiffness Is Associated With Risk of Decompensation, Liver Cancer, and Death in Patients With Chronic Liver Diseases: A Systematic Review and Meta-analysis. *Clinical Gastroenterology and Hepatology,* 2013; 11(12), 1573–1584.e2. doi:10.1016/j.cgh.2013.07.034
20. Perazzo, H., Veloso, V. G., Grinsztejn, B., Hyde, C., & Castro, R.. Factors That Could Impact on Liver Fibrosis Staging by Transient Elastography. *International Journal of Hepatology,* 2015, 15. doi:10.1155/2015/624596

TURNITIN-Correlation-Between-Portal-Venous-Dimensions

ORIGINALITY REPORT

18%

SIMILARITY INDEX

10%

INTERNET SOURCES

14%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	Siddharth Singh, Larissa L. Fujii, Mohammad Hassan Murad, Zhen Wang et al. "Liver Stiffness Is Associated With Risk of Decompensation, Liver Cancer, and Death in Patients With Chronic Liver Diseases: A Systematic Review and Meta-analysis", Clinical Gastroenterology and Hepatology, 2013 Publication	1%
2	ejrnm.springeropen.com Internet Source	1%
3	scholar.uwindsor.ca Internet Source	1%
4	"Liver Elastography", Springer Science and Business Media LLC, 2020 Publication	1%
5	www.researchgate.net Internet Source	1%
6	www.ddtjournal.com Internet Source	1%

7

J.W. Carter. "The American Association of Suicidology", *Policing: An International Journal of Police Strategies & Management*, 2012

Publication

1 %

8

Jinhuan Wang, Qiao Wang, Guoying Yu, Qilu She, Wenyuan Zhang, Jiagang Zhang. "Correlation Between Liver Stiffness Measured by Shear Wave Elastography and Child-Pugh Classification", *Journal of Ultrasound in Medicine*, 2018

Publication

1 %

9

Middleton, William D., and Kathryn A. Robinson. "Ultrasound Assessment of the Hepatic Vasculature", *Introduction to Vascular Ultrasonography*, 2012.

Publication

1 %

10

ejournal.unsrat.ac.id

Internet Source

1 %

11

Abd ElMohsen ElDesoky, Mohammed Aeriba, Fardous Abd ElFattah, Galal ElHawary, Mahmoud Awad. "Platelet count/spleen diameter ratio as a predictor of high-risk esophageal varices in patients with liver cirrhosis", *Medical Journal of Viral Hepatitis*, 2022

Publication

1 %

12

Submitted to Adtalem Global Education, Inc.

Student Paper

1 %

13

Yuksel Seckin, Murat M. M. Harputluoglu, Kadir Batcioglu, Melih Karıncaoglu et al. "Gastric Tissue Oxidative Changes in Portal Hypertension and Cirrhosis", Digestive Diseases and Sciences, 2007

Publication

1 %

14

Zhang, Chao-Xue, Jian-Ming Xu, Jia-Bin Li, De-Run Kong, Ling Wang, Xiao-Yong Xu, and Dong-Mei Zhao. "Predict esophageal varices via routine trans-abdominal ultrasound: A design of classification analysis model : Predict EV via routine TUS", Journal of Gastroenterology and Hepatology, 2015.

Publication

1 %

15

ir.ymlib.yonsei.ac.kr

Internet Source

1 %

16

www.iiste.org

Internet Source

<1 %

17

Submitted to Southampton Solent University

Student Paper

<1 %

18

Pooya Iranpour, Chandana Lall, Roozbeh Houshyar, Mohammad Helmy, Albert Yang, Joon-II Choi, Garrett Ward, Scott C Goodwin. "Altered Doppler flow patterns in cirrhosis patients: an overview", Ultrasonography, 2016

<1 %

19

Serife Ulsan. "Evaluation of Portal Venous Velocity with Doppler Ultrasound in Patients with Nonalcoholic Fatty Liver Disease", Korean Journal of Radiology, 2011

Publication

<1 %

20

Seung Woo Cha, Woo Kyoung Jeong, Yongsoo Kim, Min Yeong Kim et al. "Nondiseased Liver Stiffness Measured by Shear Wave Elastography", Journal of Ultrasound in Medicine, 2014

Publication

<1 %

21

rcastoragev2.blob.core.windows.net

Internet Source

<1 %

22

doczz.net

Internet Source

<1 %

23

tede2.pucrs.br

Internet Source

<1 %

24

www.cureus.com

Internet Source

<1 %

25

www.mdpi.com

Internet Source

<1 %

26

Stefanescu, Horia. "Noninvasive assessment of portal hypertension in cirrhosis: Liver stiffness and beyond", World Journal of Gastroenterology, 2014.

Publication

<1 %

27 Utaroh Motosugi, Tomoaki Ichikawa, Yoshibumi Niitsuma, Tsutomu Araki. "Acoustic radiation force impulse elastography of the liver: can fat deposition in the liver affect the measurement of liver stiffness?", Japanese Journal of Radiology, 2011

<1 %

Publication

28 F. Piscaglia, G. Donati, L. Cecilioni, N. Celli, B. Stagni, P. Pini, S. Gaiani, F. Gherlinzoni, L. Bolondi. "Influence of the Spleen on Portal Haemodynamics: a Non-invasive Study with Doppler Ultrasound in Chronic Liver Disease and Haematological Disorders", Scandinavian Journal of Gastroenterology, 2009

<1 %

Publication

29 Luis Téllez, Agustín Albillos. "Non - selective beta - blockers in patients with ascites: The complex interplay among the liver, kidney and heart", Liver International, 2022

<1 %

Publication

30 Ma, Jiao-jiao, Hong Ding, Feng Mao, Hui-chuan Sun, Chen Xu, and Wen-ping Wang. "Assessment of Liver Fibrosis with Elastography Point Quantification Technique in Chronic Hepatitis B Virus Patients: a Comparison with Liver Pathological Results : Assessing Liver Fibrosis by ElastPQ

<1 %

Technique", Journal of Gastroenterology and Hepatology, 2013.

Publication

31

Subash Bhattarai, M Gyawali, KR Dewan, G Shrestha, BS Patowary, P Sharma. "Study of Portal Vein Diameter and Spleen Size by Ultrasonography and Their Association with Gastro-esophageal Varices", Nepalese Journal of Radiology, 2017

Publication

<1 %

32

[dokumen.pub](#)

Internet Source

<1 %

33

[spandidos-publications.com](#)

Internet Source

<1 %

34

[vdoc.pub](#)

Internet Source

<1 %

35

Andrew Huang, Grant S. Roberts, Alejandro Roldán-Alzate, Oliver Wieben, Scott B. Reeder, Thekla H. Oechtering. "Reference values for 4D flow magnetic resonance imaging of the portal venous system", Abdominal Radiology, 2023

Publication

<1 %

36

Masoud Baikpour, Arinc Ozturk, Manish Dhyani, Nathaniel D. Mercaldo, Theodore T. Pierce, Joseph R. Grajo, Anthony E. Samir. "Portal Venous Pulsatility Index: A Novel

<1 %

Biomarker for Diagnosis of High-Risk
Nonalcoholic Fatty Liver Disease", American
Journal of Roentgenology, 2020

Publication

37

"Noninvasive Vascular Diagnosis", Springer
Science and Business Media LLC, 2017

Publication

<1 %

38

Muhammad Danish, Hina Ismail, Rani Tulsi,
Nasir Mehmood, Syed Muddasir Laeeq, Nasir
Hassan Luck. "Liver Elastography as a
Predictor of Esophageal Varices in Patients
With Cirrhosis", Cureus, 2021

Publication

<1 %

39

Shi-Hao Xu, Qiao Li, Yuan-Ping Hu, Li Ying.
"Development of a model based on
biochemical, real-time tissue elastography
and ultrasound data for the staging of liver
fibrosis and cirrhosis in patients with chronic
hepatitis B", Molecular Medicine Reports,
2016

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On

TURNITIN-Correlation-Between-Portal-Venous-Dimensions

GRADEMARK REPORT

FINAL GRADE

/0

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5
