

# Correlation between Interleukin-17 Levels with C- Reactive

*by Muchlis Sofro*

---

**Submission date:** 12-May-2023 11:55AM (UTC+0700)

**Submission ID:** 2091055227

**File name:** Correlation\_between\_Interleukin-17\_Levels\_with\_C-Reactive.pdf (249.33K)

**Word count:** 3351

**Character count:** 17732

# Submissions Template for Authors

## Original Research Article

Instructions/Template for Preparing Manuscript for Journal of Biomedicine and Translational Research

12

## Correlation between Interleukin-17 Levels with C-Reactive Protein and Neutrophil Lymphocyte Ratio in Sepsis

Dwi Retnoningrum<sup>1\*</sup>, Muchlis Achsan Udji Sofro<sup>2</sup>, Setyo Gundi Pramudo<sup>2</sup>, Satrio Adi Wicaksono<sup>3</sup>, Andaru Dahesihdewi<sup>4</sup>

48

<sup>1</sup>Department of Clinical Pathology, Faculty of Medicine, Universitas Diponegoro, Semarang

<sup>2</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Diponegoro, Semarang

<sup>3</sup>Department of Anesthesiology and Intensive Care, Faculty of Medicine, Universitas Diponegoro, Semarang

<sup>4</sup>Department of Clinical Pathology and Laboratory Medicine, Faculty of Medicine, Public Health and Nursing, Gadjah Mada University, Yogyakarta

\*Corresponding Author: [dwiretno@fk.undip.ac.id](mailto:dwiretno@fk.undip.ac.id), phone 085225379937

**Short Running Title:** Interleukin-17, CRP, NLR, sepsis

### ABSTRACT

**Background:** Sepsis is a condition in systemic infection associated with organ dysfunction. Interleukin-17 is a pro-inflammatory cytokine produced by Th-17 cells. C-reactive protein and neutrophil-lymphocyte ratio (NLR) have been widely used as markers of inflammation.

**Objective:** This study aims to prove the correlation between IL-17 with CRP and NLR in septic patients.

**Methods:** Analytical observational study with a cross-sectional approach was carried out on 40 septic patients in July–December 2020. IL-17 levels were obtained by the ELISA method while CRP levels were obtained using the immunoturbidimetric method. NLR was the result of dividing the absolute number of neutrophils and lymphocytes from the automatic hematology analyzer Sysmex XN-1000. Correlation between variables was performed using Spearman correlation test.

**Results:** The median levels of IL-17, CRP, and NLR were 363.55 (11.4-1695.80) pg/mL, 13.25 (0.43-53.87) mg/L, and 12.00 (2.26–48.5), respectively. The Spearman correlation test between IL-17 levels and CRP obtained  $p = 0.019$ ,  $r = 0.37$ , and NLR  $p = 0.425$ ,  $r = 0.13$ .

**Conclusion:** There is a weak positive correlation between IL-17 levels and CRP in septic patients. There is no correlation between IL-17 levels and NLR in septic patients.

**Keywords:** Interleukin-17, CRP, NLR, sepsis

For complete information author guidelines please check:

<http://ejournal2.undip.ac.id/index.php/jbtr/about/submissions#authorGuidelines>

# Submissions Template for Authors

## Original Research Article

Instructions/Template for Preparing Manuscript for Journal of Biomedicine and Translational Research

### INTRODUCTION

Sepsis is a collection of symptoms that often occurs in patients with systemic infections associated with organ dysfunction. The state of sepsis still has a sizable proportion of the critically ill patient population. Sepsis is defined as the presence or suspicion of infection associated with a systemic response, a life-threatening condition caused by dysregulation of the body's response to infection. There is no gold standard in the definition of sepsis, so the diagnosis of sepsis is based on a combination of physiology and laboratory abnormalities. The definition of sepsis itself changed at international meetings in 1991, 2001, and, most recently, in 2016. The current recommendations and definitions of sepsis are based on organ dysfunction where a total SOFA score of 2 points is an indication of infection<sup>1</sup>.

The incidence of sepsis in the United States is about 300 cases per 100,000 population, 50% of whom undergo treatment in the Intensive Care Unit (ICU)<sup>2</sup>. The epidemiological trend of sepsis in Spain shows that the incidence of sepsis is increasing from 3.3 per 1,000 population in 2000 to 4.45 in 2013, with an increase in mortality from 6.34 to 7.89 per 1,000 population<sup>3</sup>. The pathogenesis of sepsis begins with the presence of infectious agents that enter a person's blood circulation causing a systemic inflammatory state, but the presence of pathogens in the systemic circulation is not always present in cases of sepsis. The presence of inflammatory mediators released systemically can induce sepsis while the cause of sepsis itself can be various infectious agents, including bacteria, fungi, parasites, and viruses<sup>4</sup>.

Interleukin-17 is a proinflammatory cytokine produced by T helper-17, Natural Killer (NK) cells, CD-8 T cells, neutrophils, and several other cells. IL-17 increases the production of chemokines so that it plays a significant role in the recruitment of monocytes and neutrophils to the site of inflammation. This shows the vital role of IL-17 as a proinflammatory cytokine that plays a role in infectious conditions. Early studies in animal models induced sepsis showed that removing IL-17 was associated with a higher survival rate, but another study using mice with IL-17 receptor deficiency showed the opposite result<sup>5</sup>. Ali et al. reported that serum levels of IL-17 were associated with sepsis, where, in sepsis, IL-17 levels were 72 pg/mL compared to those without sepsis, namely 37 pg/mL with  $p = 0.0001$ . IL-17 at levels more than 45 pg/mL is reported to have a 28-day mortality risk in septic patients ( $p = 0.005$ ), so it can be a predictor of mortality in sepsis<sup>6</sup>.

The state of sepsis is characterized by changes in acute phase protein levels. One of the acute phase proteins is C-reactive protein (CRP). Elevated levels of CRP are produced in response to tissue damage, infection, inflammation, and malignancy<sup>7</sup>. Yang et al. stated that the sensitivity and specificity of CRP in the diagnosis of sepsis were 90.7% and 49.7% while Pradhan et al. in their study using a CRP cut-off value of 50 mg/L obtained a sensitivity of 84.3% and a specificity of 46.15%<sup>8,9</sup>. Ranzani et al. showed that CRP is a prognostic factor for septic patients admitted to the ICU, where a CRP level of 20 mg/dl has a poor prognosis ( $p = 0.024$ )<sup>10</sup>.

Neutrophil lymphocyte ratio (NLR) is an examination that is easily obtained from routine hematological examinations. Recently increased NLR is promising as a diagnostic and prognostic marker in patients with sepsis. Research by Marik et al. showed the role of NLR in sepsis. NLR values  $> 10$  had a sensitivity of 71% and specificity of 56% with a higher mean NLR value in positive bacterial cultures than in negative cultures<sup>11</sup>.

For complete information author guidelines please check:

<http://ejournal2.undip.ac.id/index.php/jbtr/about/submissions#authorGuidelines>

# Submissions Template for Authors

## Original Research Article

Instructions/Template for Preparing Manuscript for Journal of Biomedicine and Translational Research

In sepsis, there is an increase in pro-inflammatory cytokines, one of which is IL-17 which plays a role in increasing natural, adaptive, and humoral immunity, in inflammation, an increase in acute-phase protein, namely CRP, and a response from leukocyte cells in the inflammatory process, so it is necessary to research the correlation of serum IL-17 levels with CRP levels and NLR values in septic patients.

### MATERIALS AND METHODS ← Capitalized, Bold

This research is an analytic observational study with a cross-sectional approach which was conducted in the ICU of Dr. Kariadi Hospital and Diponegoro National Hospital Semarang from July to December 2020. The research subjects were patients with sepsis criteria (SOFA score) with an age range of 18–70 years. Subjects were taken by consecutive sampling that met the inclusion and exclusion criteria. Patients with a history of liver disease and chronic renal failure were not included in this study.

The research variables consisted of levels of IL-17, CRP, and NLR. Examination of IL-17 levels obtained using the ELISA method was carried out at the GAKI Laboratory of Faculty of Medicine, Universitas Diponegoro. From the examination of CRP levels using the phototurbidimetric method, the calculation of the NLR value was done manually by dividing the absolute neutrophil count by the absolute lymphocyte count obtained from the automatic hematology analyzer. Based on the normality test using the Shapiro Wilk test, IL-17, CRP, and NLR data were not normally distributed after data transformation. Then, Spearman Rank correlation test was performed. The p-value is considered significant if  $p < 0.05$ . Ethical clearance was obtained from the Health Research Ethics Committee of Dr. Kariadi General Hospital, Semarang No. 484/EC/KEPK/2020.

### RESULTS

Forty patients with sepsis criteria according to SOFA criteria were included in this study, consisting of 19 (47.5%) males and 21 (52.5%) females. The mean age of the patients was  $54.18 \pm 13.25$  years with a normal mean body mass index ( $23.24 \pm 4.82$ ). The mean hemoglobin level is 11.5 g/dL, the mean leukocyte count is  $16.45 \times 10^3/\mu\text{L}$ , and the mean platelet count is  $287.42 \times 10^3/\mu\text{L}$ . The results of blood cultures from 40 subjects obtained positive blood cultures in 15 subjects (37.5%), whereas the results of gram-positive bacteria were 6 subjects (40%) and gram-negative 9 subjects (60%). The characteristics of the data are presented in Table 1.

The normality test using Shapiro Wilk found that the three variables were not normally distributed, so the Spearman Rank correlation test was carried out. The correlation between IL-17 level with CRP and NLR is presented in Table 2.

There is a weak positive correlation between levels of IL-17 and CRP ( $p=0.019$ ,  $r=0.37$ ). The data distribution of IL-17 and CRP levels is shown in Figure 1. The correlation between IL-17 and NLR levels showed that the results were not statistically significant.

### DISCUSSION

The results showed that there was a positive correlation between IL-17 levels and CRP levels. This is in accordance with the study of Akin et al. who reported a positive correlation between IL-17 and CRP ( $p = 0.014$ ,  $r = 0.225$ ) in patients with polymicrobial sepsis<sup>12</sup>. Comparable results were reported by the study of Zou et al. in patients with

For complete information author guidelines please check:

<http://ejournal2.undip.ac.id/index.php/jbtr/about/submissions#authorGuidelines>

## Submissions Template for Authors

### Original Research Article

Instructions/Template for Preparing Manuscript for Journal of Biomedicine and Translational Research

chronic obstructive pulmonary disease (COPD) and by Al-Saadani et al. in rheumatoid arthritis patients where there was a positive correlation between IL-17 levels and CRP ( $p < 0.001$ ,  $r = 0.506$ , and  $p < 0.001$ ,  $r = 0.696$ )<sup>13,14</sup>. Interleukin-17 is a cytokine produced by several cells, namely Th17 cells, T cells, NK cells, group 3 innate lymphoid cells (ILC3s), CD8+ cells (Tc17), microglia, and mast cells. The general function of IL-17 is to stimulate the production of inflammatory molecules, chemokines, and antimicrobial peptides (AMPs) such as  $\alpha$ -defensin, calgranulin, and lipocalin-2. IL-17 also functions in maintaining mucosal barrier integrity and induces acute-phase proteins. In sepsis, IL-17 stimulates various immune cells (neutrophils, lymphocytes, macrophages, monocytes, and B cells) to enhance the role of natural, adaptive, and humoral immunity. IL-17 signals play a protective role against infections caused by bacteria, fungi, viruses, and parasites through the body's defense and release of cytokines, but IL-17 can also induce pathological immune responses and the occurrence of organ failures such as acute lung injury (ALI), acute respiratory distress syndrome (ARDS), acute kidney injury (AKI), hepatic dysfunction, immunity, and cardiomyocytes<sup>15</sup>.

An experimental study by Marchesi et al. showed that sepsis caused by Klebsiella pneumonia increased the production of IL-17 ( $p = 0.002$ ), and the highest level was reached at 48 hours after infection<sup>16</sup>. Ogiku et al. reported that mice with IL-17 deficiency showed a significant increase in mortality associated with bacteremia at 12 hours after the cecal ligation and puncture (CLP) procedure. This shows a partial protective role of IL-17 in sepsis, resulting in the positive control mice having a higher survival rate, compared to model mice that were knocked out of the IL-17 gene<sup>17</sup>. CRP is an acute-phase protein that can be used as a marker of inflammation. Li et al.'s study reported higher CRP levels in septic patients than in non-septic patients ( $p < 0.05$ ). Patients with poor clinical outcomes (non-survivors) also had higher CRP levels ( $p = 0.047$ )<sup>11</sup>.

The results of this study showed that there was no correlation between IL-17 levels and NLR. This is not following previous studies where the NLR value can be used as a marker for sepsis although, in this study, a high NLR value ( $> 5$ ) was not associated with IL-17 levels<sup>18</sup>. The results of this study are similar to those of Sag et al. which showed that there was no correlation between IL-17 levels and NLR in rheumatoid arthritis patients ( $p > 0.05$ ,  $r = 0.135$ )<sup>19</sup>. In sepsis, there can be a failure of recruitment and migration of neutrophils to the site of infection, which is related to the severity of the disease. Interleukin-17 plays a role in the migration and microbicidal activity of neutrophils, and the mobilization of T lymphocytes<sup>20</sup>. The increase in NLR indicates the presence of an inflammation that, in addition to sepsis, can occur in other inflammatory conditions.

Positive blood culture results in study subjects were obtained in 37.5% of patients, which is almost similar to the study by Liestadi et al. where positive culture results were obtained in 32.6% of patients, and the prevalence of gram-negative bacteria was higher than gram-positive<sup>21</sup>. In the study, the results showed that there were differences in the levels of IL-17, CRP, and NLR in the subjects between the cultures of gram-positive bacteria and gram-negative bacteria.

The limitation of this study was that it did not take into account comorbid factors from the patient that could affect IL-17, CRP, and NLR levels. Patients on immunosuppressant therapy and the presence of hematological abnormalities were not excluded from the study as it could affect the results of complete blood counts.

For complete information author guidelines please check:

<http://ejournal2.undip.ac.id/index.php/jbtr/about/submissions#authorGuidelines>

# Submissions Template for Authors

## Original Research Article

Instructions/Template for Preparing Manuscript for Journal of Biomedicine and Translational Research

### CONCLUSION

In this study, there was a statistically weak positive correlation between IL-17 levels and CRP in septic patients. There was no correlation between IL-17 levels and NLR in septic patients. Further research needs to be done by paying attention to the factors that affect the levels of IL-17, CRP, and NLR values, as well as to the comorbid factors of the patient.

### ACKNOWLEDGMENTS

The authors would like to thank to Universitas Diponegoro, Dr. Kariadi General Hospital, Diponegoro National Hospital, and GAKI laboratory for their support and assistance throughout this study. This study was financially supported by Faculty of Medicine Universitas Diponegoro, Research Grant No. 85/UN7.5.4.2/HK/2020.

### REFERENCES

1. Gul F, Arslantas MK, Cinel I, Kumar A. Changing Definitions of Sepsis. *Turk J Anesth Reanim.* 2017;45:129–38.
2. Mayr FB, Yende S, Angus DC. Epidemiology of severe sepsis. *Virulence.* 2014;5(1):1–11.
3. Álvaro-Meca A, Jiménez-Sousa MA, Micheloud D, Sánchez-Lopez A, Heredia-Rodríguez M, Tamayo E, et al. Epidemiological trends of sepsis in the twenty-first century (2000-2013): An analysis of incidence, mortality, and associated costs in Spain. *Popul Health Metr.* 2018;16(1):1–11.
4. Burkovskiy I, Sardinha J, Zhou J, Lehmann C. Cytokine release in sepsis. *Adv Biosci Biotechnol.* 2013;2013(September):860–5.
5. Flierl MA, Rittirsch D, Gao H, Hoesel LM, Nadeau BA, Day DE, et al. Adverse functions of IL-17A in experimental sepsis. *FASEB J.* 2008;22(July):2199–205.
6. Ali MA, Abdelkader ESMA, El LMR. Interleukin-17 as a predictor of sepsis in polytrauma patients : a prospective cohort study. *Eur J Trauma Emerg Surg.* 2017
7. Faraj M, Salem N. C-Reactive Protein. In: *Blood Cell- An Overview of Studies in Hematology.* 2012. p. 89–100.
8. Yang Y, Xie J, Guo F, Longhini F, Gao Z, Huang Y, et al. Combination of C - reactive protein , procalcitonin and sepsis - related organ failure score for the diagnosis of sepsis in critical patients. *Ann Intensive Care.* 2016;6(51).
9. Pradhan S, Ghimire A, Bhattarai B, Khanal B, Pokharel K. The role of C - reactive protein as a diagnostic predictor of sepsis in a multidisciplinary Intensive Care Unit of a tertiary care center in Nepal. *Indian J Crit Care Med.* 2016;(41):417–20.
10. Ranzani OT, Zampieri FG, Forte DN, Azevedo LCP, Park M. C-Reactive Protein/Albumin Ratio Predicts 90-Day Mortality of Septic Patients. *PLoS One.*

For complete information author guidelines please check:

<http://ejournal2.undip.ac.id/index.php/jbtr/about/submissions#authorGuidelines>

## Submissions Template for Authors

### Original Research Article

Instructions/Template for Preparing Manuscript for Journal of Biomedicine and Translational Research

- 2013;8(3):e59321.
11. Li Q, Gong X. Clinical significance of the detection of procalcitonin and C-reactive protein in the intensive care unit. *Exp Ther Med*. 2018;15(5):4265–70.
  12. Akin H, Akalin H, Budak F, Ener B, Ocakoğlu G, Gürcüoğlu E, et al. Alterations of serum cytokine levels and their relation with inflammatory markers in candidemia. *Med Myco*. 2015;53(3):258–68.
  13. Zou Y, Chen X, Liu J, Zhou D bo, Kuang X, Xiao J, et al. Serum IL-1 $\beta$  and IL-17 levels in patients with COPD: association with clinical parameter. *Int J COPD*. 2017;12:1247–54.
  14. Al-Saadany HM, Hussein MS, Gaber RA, Zaytoun HA. Th-17 cells and serum IL-17 in rheumatoid arthritis patients: Correlation with disease activity and severity. *Egypt Rheumatol*. 2016;38(1):1–7.
  15. Ge Y, Huang M, Yao YM. Biology of Interleukin-17 and Its Pathophysiological Significance in Sepsis. *Front Immunol*. 2020;11(July):1–13.
  16. Marchesi V V, Rukavina T. Interleukin-17 in experimental Klebsiella sepsis. *Period Biol*. 2011;113(1):99–102.
  17. Ogiku M, Kono H, Hara M, Tsuchiya M, Fujii H. Interleukin-17A Plays a Pivotal Role in Polymicrobial Sepsis According to Studies Using IL-17A Knockout Mice. *J Surg Res*. 2012;174(1):142–9.
  18. Martins EC, Da Fe Silveira L, Viegas K, Beck AD, Júnior GF, Cremonese RV, et al. Neutrophil-lymphocyte ratio in the early diagnosis of sepsis in an intensive care unit: A case-control study. *Rev Bras Ter Intensiva*. 2019;31(1):63–70.
  19. Sağ S, Sağ MS, Tekeoğlu I, Kamanlı A, Nas K, Acar BA. Relationship of hematologic markers with IL-17 and IL-1 beta in patients with rheumatoid arthritis. *J Back Musculoskelet Rehabil*. 2018;31(4):703–7.
  20. Delano MJ, Ward PA. Sepsis-induced immune dysfunction: Can immune therapies reduce mortality? *J Clin Invest*. 2016;126(1):23–31.
  21. Liestiadi DEF, Azlin E, Nafianti S. A hematologic scoring system and C-reactive protein compared to blood cultures for diagnosing bacterial neonatal sepsis. *Paediatr Indones*. 2017;57(2):70–5.

For complete information author guidelines please check:

<http://ejournal2.undip.ac.id/index.php/jbtr/about/submissions#authorGuidelines>


# Submissions Template for Authors

## Original Research Article

Instructions/Template for Preparing Manuscript for Journal of Biomedicine and Translational Research

### FIGURE CAPTIONS

Figure 1. Correlation between IL-17 and CRP in Sepsis



### TABLE CAPTIONS

Table 1. Subjects Characteristic Data <sup>44</sup>

Parameter	Mean±SD	Median (min;max)	p*
Age (year)	54.18±13.25	56 (19;70)	0.316
BMI	23.24±4.82	22.8 (12.8;34.52)	0.244
Hemoglobin (g/dL)	11.01±2.97	11.5 (2.9;14.9)	0.006
Leukocyte (10 <sup>3</sup> /uL)	16.45±9.08	14.75 (2;38)	0.304
Platelets (10 <sup>3</sup> /uL)	287.42±166.5	277 (35;626)	0.186
Neutrophil absolute count (10 <sup>3</sup> /uL)	13.7±7.85	12.2 (1.34;32.97)	0.087
Lymphocyte absolute count (10 <sup>3</sup> /uL)	1.39±1.29	1.09 (0.08;5.48)	0.001
NLR	16.57±13.70	13.51 (2.26;46.5)	0.001
IL-17 (pg/mL)	549.60±489.84	337.95 (11.4;1695.80)	0.001
CRP (mg/L)	17.6±14.97	13.25 (0.43;53.87)	0.001
SOFA score	9.74±3.10	9 (4;17)	0.318

Note: BMI: Body Mass Index. SD: Standard Deviation. NLR: Neutrophil Lymphocyte Ratio. CRP: C-Reactive Protein. SOFA: Sequential Organ Failure Assessment. \* *Saphiro Wilk*

For complete information author guidelines please check:

<http://ejournal2.undip.ac.id/index.php/jbtr/about/submissions#authorGuidelines>



## Submissions Template for Authors

---

### Original Research Article

Instructions/Template for Preparing Manuscript for Journal of Biomedicine and Translational Research

**Table 2. Correlation between IL-17 with CRP and NLR**

Variable	IL-17 (pg/mL)	
	<i>p</i>	<i>r</i> <sup>§</sup>
CRP (mg/L)	0.019*	0.37
NLR	0.425	0.13

IL-17: Interleukin-17, CRP: C-Reactive Protein, NLR: Neutrophil Lymphocyte Ratio;

<sup>§</sup>Spearman Rank correlation test; \**p* < 0.05

For complete information author guidelines please check:

<http://ejournal2.undip.ac.id/index.php/jbtr/about/submissions#authorGuidelines>

# Correlation between Interleukin-17 Levels with C-Reactive

## ORIGINALITY REPORT

24%

SIMILARITY INDEX

20%

INTERNET SOURCES

16%

PUBLICATIONS

11%

STUDENT PAPERS

## PRIMARY SOURCES

1

[pure.uva.nl](https://pure.uva.nl)

Internet Source

1%

2

[sciendo.com](https://sciendo.com)

Internet Source

1%

3

Fakhriati Fakhriati, Mu'jizah Mu'jizah, Yeni Mulyani Supriatin, Asep Supriadi et al.

"Cultural Coping Strategies to Mitigate Local Impact of Baribis Fault Disasters", Research Square Platform LLC, 2023

Publication

1%

4

Castro, Myrella L, Gilson C N Franco, Luciana S Branco-de-Almeida, Ana L Anbinder, Karina Cogo-Müller, Sheila C Cortelli, Simone Duarte, Deepak Saxena, and Pedro L Rosalen. "Down-Regulation of Protease Activated Receptor 2, Interleukin-17 and Other Pro-Inflammatory Genes by Subantimicrobial Doxycycline Dose in a Rat Periodontitis Model", Journal of Periodontology, 2015.

Publication

1%

[online.medunigraz.at](https://online.medunigraz.at)

5	Internet Source	1 %
6	<a href="http://ntnuopen.ntnu.no">ntnuopen.ntnu.no</a> Internet Source	1 %
7	<a href="http://annalsofintensivecare.springeropen.com">annalsofintensivecare.springeropen.com</a> Internet Source	1 %
8	<a href="http://journal-inflammation.biomedcentral.com">journal-inflammation.biomedcentral.com</a> Internet Source	1 %
9	Submitted to Marmara University Student Paper	1 %
10	<a href="http://bmcnephrol.biomedcentral.com">bmcnephrol.biomedcentral.com</a> Internet Source	1 %
11	<a href="http://medic.upm.edu.my">medic.upm.edu.my</a> Internet Source	1 %
12	<a href="http://ejournal2.undip.ac.id">ejournal2.undip.ac.id</a> Internet Source	1 %
13	Submitted to Universitas Islam Indonesia Student Paper	1 %
14	<a href="http://topsecretapiaccess.dovepress.com">topsecretapiaccess.dovepress.com</a> Internet Source	1 %
15	<a href="http://publikationen.ub.uni-frankfurt.de">publikationen.ub.uni-frankfurt.de</a> Internet Source	1 %
16	<a href="http://repository.stikim.ac.id">repository.stikim.ac.id</a> Internet Source	1 %

17	<a href="https://journals.sagepub.com">journals.sagepub.com</a> Internet Source	1 %
18	Na Cui, Hongwei Zhang, Zhi Chen, Zhanbiao Yu. "Prognostic significance of PCT and CRP evaluation for adult ICU patients with sepsis and septic shock: retrospective analysis of 59 cases", Journal of International Medical Research, 2019 Publication	<1 %
19	<a href="http://www.tara.tcd.ie">www.tara.tcd.ie</a> Internet Source	<1 %
20	<a href="http://tede.bc.uepb.edu.br">tede.bc.uepb.edu.br</a> Internet Source	<1 %
21	<a href="http://www.frontiersin.org">www.frontiersin.org</a> Internet Source	<1 %
22	Submitted to Tufts University Student Paper	<1 %
23	<a href="http://academic.oup.com">academic.oup.com</a> Internet Source	<1 %
24	Submitted to Amman Arab University for Graduate Studies Student Paper	<1 %
25	<a href="http://www.ccsenet.org">www.ccsenet.org</a> Internet Source	<1 %

26 Submitted to Queen Mary and Westfield College  
Student Paper <1 %

---

27 [www.oncotarget.com](http://www.oncotarget.com)  
Internet Source <1 %

---

28 Pablo Palazon-Riquelme, Gloria Lopez-Castejon. "The inflammasomes, Immune guardians at defence barriers", Immunology, 2018  
Publication <1 %

---

29 [core.ac.uk](http://core.ac.uk)  
Internet Source <1 %

---

30 [www.science.gov](http://www.science.gov)  
Internet Source <1 %

---

31 [balimedicaljournal.org](http://balimedicaljournal.org)  
Internet Source <1 %

---

32 [kupdf.net](http://kupdf.net)  
Internet Source <1 %

---

33 [pubmed.ncbi.nlm.nih.gov](http://pubmed.ncbi.nlm.nih.gov)  
Internet Source <1 %

---

34 [www.pneumon.org](http://www.pneumon.org)  
Internet Source <1 %

---

35 [www.wjgnet.com](http://www.wjgnet.com)  
Internet Source <1 %

---

36 Basak, P.Y.. "The role of helper and regulatory T cells in the pathogenesis of vitiligo", Journal of the American Academy of Dermatology, 200902 <1 %  
Publication

---

37 Maira G.R. Pitta. "IL-17 and IL-22 are associated with protection against human kala azar caused by Leishmania donovani", Journal of Clinical Investigation, 07/13/2009 <1 %  
Publication

---

38 bioscmed.com <1 %  
Internet Source

---

39 dspace.knust.edu.gh <1 %  
Internet Source

---

40 hqlo.biomedcentral.com <1 %  
Internet Source

---

41 repository.up.ac.za <1 %  
Internet Source

---

42 scholarworks.bridgeport.edu <1 %  
Internet Source

---

43 www.ajp.eg.net <1 %  
Internet Source

---

44 www.jointdrs.org <1 %  
Internet Source

---

45 www.mdpi.com  
Internet Source

<1 %

46

Hai-Feng Li. "Does increased serum IL-17 level imply its pathogenic role in myasthenia gravis?", Muscle & Nerve, 01/2012

Publication

<1 %

47

Kolls, Jay K.. "CD4<sup>+</sup> T-cell subsets and host defense in the lung", Immunological Reviews, 2013.

Publication

<1 %

48

Songane, Mario, Johanneke Kleinnijenhuis, Bacht Alisjahbana, Edhyana Sahiratmadja, Ida Parwati, Marije Oosting, Theo S. Plantinga, Leo A. B. Joosten, Mihai G. Netea, Tom H. M. Ottenhoff, Esther van de Vosse, and Reinout van Crevel. "Polymorphisms in Autophagy Genes and Susceptibility to Tuberculosis", PLoS ONE, 2012.

Publication

<1 %

49

Francisco Ujueta, Roshanak Azimi, Matthew R. Lozier, Robert Poppiti, Ari Ciment. "Lymphohistocytic myocarditis after Ad26.COV2.S viral vector COVID-19 vaccination", IJC Heart & Vasculature, 2021

Publication

<1 %

50

Mustafa Özgür Cirik, Derya Yenibertiz. "What are the prognostic factors affecting 30-day

<1 %

mortality in geriatric patients with respiratory failure in the Intensive Care Unit?", Pakistan Journal of Medical Sciences, 2020

Publication

---

51

Shi-tong Yu, Zhiwei Zhou, Qian Cai, Faya Liang, Ping Han, Renhui Chen, Xiaoming Huang.

<1 %

"Prognostic value of the C-reactive protein/albumin ratio in patients with laryngeal squamous cell carcinoma", OncoTargets and Therapy, 2017

Publication

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off