

The Effects of Pancreatic Omentoplasty on Weight Loss

by Heri Nugroho


Submission date: 10-May-2023 03:33PM (UTC+0700)

Submission ID: 2089329201

File name: The_Effects_of_Pancreatic_Omentoplasty_on_Weight_Los.pdf (262.35K)

Word count: 2715

Character count: 15126



Bioscientia Medicina: Journal of Biomedicine & Translational Research

Journal Homepage: www.bioscmed.com

The Effects of Pancreatic Omentoplasty on Weight Loss, Improvement of Triglyceride and Interleukin-1 Levels in Obese Rats with Type II Diabetes Mellitus Performed Sleeve Gastrectomy

Anggoro Teguh Prasetya^{1*}, K. Heri Nugroho², Abdul Mughni³

¹Biomedicine Postgraduate Student, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

²Subdivision of Endocrinology and Diabetes Metabolic, Faculty of Medicine, Universitas Diponegoro/Dr. Kariadi General Hospital, Semarang, Indonesia

³Subdivision of Digestive Surgery, Department of Surgery, Faculty of Medicine, Universitas Diponegoro/Dr. Kariadi General Hospital, Semarang, Indonesia

ARTICLE INFO

Keywords:

Diabetes mellitus
Obesity
Omentoplasty
Pancreas
Sleeve gastrectomy

*Corresponding author:

Anggoro Teguh Prasetya

2

E-mail address:

Anggoroprasetya19@gmail.com

All authors have reviewed and approved the final version of the manuscript.

<https://doi.org/10.37275/bsm.v7i2.774>

21

1. Introduction

Obesity is one of the main risk factors for non-communicable diseases, where non-communicable diseases contribute >70% of premature deaths worldwide and are a major cause of premature mortality and disability. Obesity is a major health challenge because it increases risk factors for diseases such as type II diabetes mellitus, fatty liver, obstructive sleep apnea, hypertension, stroke,

ABSTRACT

Background: Obesity can trigger inflammation or inflammatory disease, which is one of the main features of adipose tissue dysfunction. Adipose secretes several different peptides called adipocytokines, which can induce several pro-inflammatory cytokines such as IL-6, TNF- α , and IL-1. Omentoplasty pancreas dan sleeve gastrectomy is a surgical technique that can treat overweight and metabolic syndrome in obesity and type II DM. This study aims to determine the effect of pancreatic omentoplasty on weight loss and improvement of triglyceride and interleukin-1 levels in rats obesity with type II diabetes mellitus performed sleeve gastrectomy. **Methods:** This study is an experimental study in vivo. Eighteen rats participated in this study and were grouped into the control group (K1) and the KII and KIII treatment groups. Triglyceride levels were assessed using the spectrophotometric method, and IL-1 levels were assessed using the ELISA method. Data analysis was carried out using SPSS univariate and bivariate. **Results:** The KII and KIII treatment groups experienced a decrease in body weight, triglyceride levels, and Interleukin-1 levels of rats after the treatment intervention, $p < 0.05$. The KIII treatment group showed the most significant reduction in body weight, triglyceride levels, and Interleukin-1 levels compared to the KII treatment. **Conclusion:** Pancreatic omentoplasty is effective in reducing body weight, reducing triglyceride levels, and reducing IL-1 levels in obese rats with type II diabetes mellitus undergoing sleeve gastrectomy surgery.

myocardial infarction, and several cancers. Obesity also reduces the quality of life, causes unemployment, unfavorable social impact, and causes a decline in socioeconomic productivity. The magnitude of the impact of obesity is a health problem that must be overcome. Obesity is strongly associated with dyslipidemia, mainly driven by the effects of insulin resistance and adiposopathy, which is characterized by increased triglyceride levels and decreased HDL

levels. Insulin resistance causes enhancement lipolysis in adipose tissue due to the activation of hormone sensitive lipase (HSL) so that many FFA are released into the circulation and increase the secretion of very low-density lipoprotein (VLDL) in the liver. Circulating very low-density lipoprotein (VLDL) will be hydrolyzed into triglycerides (TG) with the help of lipoprotein lipase (LPL) so that insulin resistance will cause plasma clearance of TG-rich lipoproteins to be inhibited, resulting in hypertriglyceridemia. Obesity can trigger inflammation or inflammatory disease, which is one of the main features of adipose tissue dysfunction. Adipose secretes several different peptide called adipocytokines, which can induce several pro-inflammatory cytokines such as IL-6, TNF- α , and IL-1.¹⁻⁵

The management of obesity consists of various therapies such as lifestyle modification, restricted diet, pharmacokinetics, and bariatric surgery. Despite acting as first-line therapy, nonoperative management is not a profitable method for obese patients because of poor adherence. Sleeve gastrectomy is one of the fastest-growing surgical techniques and is widely used because of its benefits for syndrome metabolic rate of severe obesity and a better risk of perioperative morbidity and mortality than other techniques. Technique Sleeve gastrectomy with the aim of cutting nearly 80% of the stomach, resection begins 3-6 cm from the pylorus cutting along the greater curvature to the gastric fundus to form a tube or banana. Mechanism Metabolically, sleeve gastrectomy is not known with certainty, but some explain that due to gastric mechanical restriction, decreased ghrelin levels due to fundus resection, changes in bile acid signals, hindgut effect causes weight loss due to reduced caloric intake, which ultimately improves adipose tissue and insulin sensitivity. The omentum is a broad layer of adipose tissue that is covered by the visceral peritoneum and hangs over the greater curvature of the stomach to the intraperitoneal organs. Omentum acts as an organ that has a role in controlling inflammation and promoting revascularization and tissue regeneration. Omental adipose tissue plays a

role in local immunological responses and also contains a population of multipotent mesenchymal stem cells (MSCs) that facilitate tissue regeneration. Omentoplasty is a surgical procedure that attaches the omentum to certain organs to close the defect, repair tissue and improve portal arterial and venous circulation.⁶⁻¹⁰ This study aimed to determine the effect of pancreatic omentoplasty on weight loss and improvement of triglyceride and interleukin-1 levels in obese rats with type II diabetes mellitus performed sleeve gastrectomy.

2. Methods

This study is an in vivo experimental study with a pre-post test approach with a control group design. The research subjects in this study were male rats (*Rattus norvegicus*) Wistar strain and around 6 weeks old. A total of 18 rats (each 6 rats/group) were included in this study, where the White Rats were grouped into 3 groups, namely group I: the control group where the White Rats were induced by obesity and type II diabetes mellitus; Group II: Obese and type II diabetes mellitus rats have used sleeve gastrectomy; Group III: Obese and type II diabetes mellitus rats underwent sleeve gastrectomy and omentoplasty. Each rat was placed in its own cage with the temperature of the cage made at 25°C, 55% humidity, and a 12:12 hour light-dark cycle. This study has received approval from the medical and health research ethics committee of the Faculty of Medicine, Universitas Diponegoro (No.118/EC/H/FK-UNDIP/XII/2020).

Each group was given a high-fat diet (comfeed pars 60%, flour 27.8%, cholesterol 2%, cholic acid 0.2%, lard 10%, and fructose 2 ml/head/day) for 4 weeks and then administration of injection of streptozotocin in sodium citrate buffer pH 4.5 intraperitoneal a single dose of 45 mg/kg BW. After 3 days, data collection was carried out pre-test and sleeve gastrectomy and omentoplasty surgery and maintenance for 10 days thereafter in analysis data post-test. The process of sleeve gastrectomy surgery is that rats have fasted for 10 hours before surgery, injection of ketamine at a

dose of 0.5 mg/kgBB is carried out intramuscularly, abdominal hair is cleaned using hair clippers until the rat's skin is visible, asepsis and antisepsis of the operating area are carried out. Incision transversely starting from the xiphoid processus latera abdomen, deepen layer by layer cutis, subcutis, internal rectus abdominis muscle, external abdominis muscle, and peritoneum, identify gastrically, perform gastric removal along the curvatura major and treat bleeding, suture the stomach with polypropylene 3.0, clean the cavity abdomen with warm 0.9% NaCl, cover the surgical wound layer by layer. The surgical procedure for pancreatic omentoplasty was that the rats were fasted for 10 hours prior to the operation, injected ketamine at a dose of 0.5 mg/kgBB intramuscularly, the abdominal hair was cleaned using a hair shave until the rat's skin was visible, asepsis and antisepsis were carried out in the operating area, incision transversely from the xiphoid process to the lateral abdomen, deepen layer by layer cutis, subcutis, internal rectus abdominis, external abdominis muscles, and peritoneum, identify omentum and

suture the omentum to the pancreas with polypropylene 3.0, clean the abdominal cavity with warm 0.9% NaCl, cover the surgical wound layer by layer. Measurement of triglyceride levels was carried out using the spectrophotometric method with the triglyceride FS kit®. IL-1 levels were assessed using the ELISA method using the Rat ELISA Kit Abclonal®. Data analysis was carried out using SPSS software version 21. Data analysis was performed univariately to describe the frequency distribution of each test variable. Bivariate analysis was carried out to determine the relationship between the test variables with a p-value <0.05.

3. Results

Table 1 presents a comparison of pre and post-test rats' body weight. The results of the study stated that in the KII and KIII treatment groups, there was a decrease in the weight of the rats after the treatment intervention, p<0.05. The KIII treatment group showed the most significant weight loss compared to the KII treatment.

Table 1. Comparison of pre-test and post-test rats' body weight.

Body weight	Treatment		p-value*
	Pre-test	Post-test	
KI	252,90 ± 6,57	241,40 ± 6,75	0,001
KII	254,90 ± 5,44	207,80 ± 5,27	0,001
KIII	256,50 ± 5,25	194,50 ± 4,80	0,001

*Dependent T-test, p< 0,05.

Table 2 presents a comparison of pre and post-test rat triglyceride levels. The results of the study stated that in the KII and KIII treatment groups, there was a decrease in rat triglycerides after the treatment

intervention, p<0.05. The KIII treatment group showed the most significant decrease in triglyceride levels compared to the KII treatment.

Table 2. Comparison of pre-test and post-test triglyceride levels.

Triglycerides	Treatment		p-value*
	Pre-test	Post-test	
KI	134,29 ± 2,76	153,98 ± 3,17	0,001
KII	132,24 ± 1,93	114,86 ± 2,57	0,001
KIII	132,46 ± 3,75	94,25 ± 3,62	0,001

*Dependent T-test, p< 0,05.

Table 3 presents a comparison of Interleukin-1 levels in pre and post-test rats. The results of the study stated that in the KII and KIII treatment groups, there was a decrease in Interleukin-1 levels of rats after the

intervention treatment, $p < 0.05$. The KIII treatment group showed the most significant decrease in Interleukin-1 levels compared to the KII treatment.

Table 3. Comparison of Interleukin-1 pre-test and post-test levels.

IL-1	Treatment		p-value*
	Pre-test	Post-test	
KI	91,38 ± 3,61	91,27 ± 4,06	0,897
KII	89,98 ± 3,12	73,67 ± 3,81	0,001
KIII	88,90 ± 2,73	60,20 ± 2,84	0,001

*Dependent T-test, $p < 0,05$.

4. Discussion

Sleeve gastrectomy surgery can reduce body weight significantly through restriction mechanisms due to reduced gastric volume and reduced ghrelin secretion, and increased secretion of intestinal hormones such as GLP-1 and PYY. In group KIII, where obese and type II DM rats underwent sleeve gastrectomy and pancreatic omentoplasty surgery, they showed greater weight loss than the other groups. This can be explained because the omentum can play a role in weight loss through the mechanism of restriction of leptin secretion mediated by CRP, IL-6, and activity-high lipolysis. Pancreatic omentoplasty will have the effect of reducing CRP and IL-6 levels which will increase leptin secretion and indirectly reduce body weight.¹¹⁻¹⁶

Improvements in triglyceride levels in the group that underwent sleeve gastrectomy and pancreatic omentoplasty are related to the regeneration of pancreatic tissue by the effect of the omentum, which can increase activity insulin signaling in the pancreas thereby inhibiting lipolysis and reducing FFA secretion in plasma which is associated with reduced triglyceride hydrolysis. Improvements in interleukin-1 levels in the sleeve gastrectomy and pancreatic omentoplasty surgery group are related to weight loss which causes an increase in the hormone adiponectin so that adipocytokine secretion decreases. Reduced adipocytokine secretion leads to decreased induction of pro-inflammatory cytokines such as IL-6, TNF α , and IL-1 associated with a decrease in pro-

inflammatory cytokines. This is in accordance with previous studies that omental stromal cells can reduce levels of pro-inflammatory cytokines such as IL-6, IL-1, and IL-12.¹⁷⁻²¹

5. Conclusion

Pancreatic omentoplasty is effective in reducing body weight, reducing triglyceride levels, and reducing IL-1 levels in obese white rats with type II diabetes mellitus undergoing sleeve gastrectomy surgery.

6. References

1. Vekic J, Zeljkovic A, Stefanovic A, Jelic-Ivanovic Z, Spasojevic-Kalimanovska V. Obesity and dyslipidemia. *Metabolism*. 2019; 92: 71-81.
2. Sánchez-García A, Rodríguez-Gutiérrez R, Mancillas-Adame L, González-Nava V, Díaz González-Colmenero A, Solis RC, et al. Diagnostic accuracy of the triglyceride and glucose index for insulin resistance: A systematic review. *Int J Endocrinol*. 2020; 2020.
3. Amin MN, Hussain MS, Sarwar MS, Rahman Moghal MM, Das A, Hossain MZ, et al. How the association between obesity and inflammation may lead to insulin resistance and cancer. *Diabetes Metab Syndr Clin Res Rev*. 2019; 13(2): 1213-24.
4. Febbraio MA. Role of interleukins in obesity: Implications for metabolic disease. *Trends Endocrinol Metab*. 2014; 25(6): 312-9.

5. Lee MKS, Yvan-Charvet L, Masters SL, Murphy AJ. The modern interleukin-1 superfamily: Divergent roles in obesity. *Semin Immunol.* 2016; 28(5): 441–9.
6. Cheng J, Gao J, Shuai X, Wang G, Tao K. The comprehensive summary of surgical versus non-surgical treatment for obesity: A systematic review and meta-analysis of randomized controlled trials. *Oncotarget.* 2016; 7(26): 20016–30.
7. Nguyen NT, Blackstone RP, Morton JM, Ponce J, Rosenthal PJ. The ASMBS textbook of bariatric surgery: History of bariatric Procedure. *Springer.* 2015; 1(3): 37
8. Arondel J, Singer M, Matsukawa A, Zychlinsky A, Sansonetti PJ. Increased interleukin-1 (IL-1) and imbalance between IL-1 and IL-1 receptor antagonist during acute inflammation in experimental shigellosis. *Infect Immun.* 1999; 67(11): 6056–66.
9. Kafka D, Ling E, Feldman G, Benharroch D, Voronov E, Givon-Lavi N, et al. Contribution of IL-1 to resistance to *Streptococcus pneumoniae* infection. *Int Immunol.* 2008; 20(9): 1139–46.
10. Romero M. del M, Fernández-López J, Esteve, M, Alemany M. Different modulation by dietary restriction of adipokine expression in white adipose tissue sites in the rat. *Cardiovascular Diabetology.* 2009; 8(1): 42.
11. Ahmad R, Shihab PK, Thomas R, Alghanim M, Hasan A, Sindhu S, et al. Increased expression of the interleukin-1 receptor-associated kinase (IRAK)-1 is associated with adipose tissue inflammatory state in obesity. *Diabetology & Metabolic Syndrome.* 2015; 7(1).
12. McArdle MA, Finucane OM, Connaughton RM, McMorrow AM, Roche HM. Mechanisms of obesity-induced inflammation and insulin resistance: Insights into the emerging role of nutritional strategies. *Front Endocrinol (Lausanne).* 2013; 4(5): 1–24.
13. Askarpour M, Khani D, Sheikhi A, Ghaedi E, Alizadeh S. Effect of bariatric surgery on serum inflammatory factors of obese patients: A systematic review and meta-analysis. *Obes Surg.* 2019; 29(8): 2631–47.
14. Salman MA, Salman AA, Nafea MA, Sultan AAEA, Anwar HW, Ibrahim AH, et al. Study of changes of obesity-related inflammatory cytokines after laparoscopic sleeve gastrectomy. *ANZ J Surg.* 2019; 89(10): 1265–9.
15. Collins D, Hogan AM, O'Shea D, Winter DC. The omentum: anatomical, metabolic, and surgical aspects. *J Gastrointest Surg* 2009; 13: 1138e46.
16. Shah S, Lowery E, Braun RK, Martin A, Huang N, Medina M, et al. Cellular basis of tissue regeneration by omentum. *PLoS One.* 2012; 7(6): 1–11.
17. Maedler K, Sergeev P, Ris F, Oberholzer J, Joller-Jemelka HI, Spinas GA, et al. Glucose-induced β cell production of IL-1 β contributes to glucotoxicity in human pancreatic islets. *Journal of Clinical Investigation.* 2002; 110: 851–60.
18. Bartholomeus K, Jacobs-Tulleneers-Thevissen D, Shouyue S, Suenens K, In'T Veld PA, Pipeleers-Marichal M, et al. Omentum is better site than kidney capsule for growth, differentiation, and vascularization of immature porcine β -cell implants in immunodeficient rats. *Transplantation.* 2013; 96(12): 1026–33.
19. Stice MJ, Dunn TB, Bellin MD, Skube ME, Beilman GJ. Omental pouch technique for combined site islet autotransplantation following total pancreatectomy. *Cell Transplant.* 2018; 27(10): 1561–8.
20. Holmes A, Coppey LJ, Davidson EP, Yorek MA. Rat models of diet-induced obesity and high fat/low dose streptozotocin type 2 diabetes: effect of reversal of high fat diet compared to treatment with enalapril or menhaden oil on glucose utilization and neuropathic endpoints. *J Diabetes Res.* 2015; 2015.

21. Marques C, Meireles M, Norberto S, Leite J, Freitas J, Pestana D, et al. High-fat diet-induced obesity Rat model: a comparison between Wistar and Sprague-Dawley Rat. *Adipocyte*. 2016; 5(1): 11-21.

The Effects of Pancreatic Omentoplasty on Weight Loss

ORIGINALITY REPORT

24%

SIMILARITY INDEX

22%

INTERNET SOURCES

20%

PUBLICATIONS

11%

STUDENT PAPERS

PRIMARY SOURCES

1	oamjms.eu Internet Source	2%
2	Submitted to Sriwijaya University Student Paper	1%
3	citeseerx.ist.psu.edu Internet Source	1%
4	assets.researchsquare.com Internet Source	1%
5	www.jmb.or.kr Internet Source	1%
6	bmcendocrdisord.biomedcentral.com Internet Source	1%
7	www.sochob.cl Internet Source	1%
8	papyrus.bib.umontreal.ca Internet Source	1%
9	www.wjgnet.com Internet Source	1%

10	academic.oup.com Internet Source	1 %
11	docksci.com Internet Source	1 %
12	doaj.org Internet Source	1 %
13	bmccardiovascdisord.biomedcentral.com Internet Source	1 %
14	www.bds.org.rs Internet Source	1 %
15	Wardiansah, Rachmat Hidayat, Msy Rulan Adnindya. "The Potential of CO2 Incubator "Sriwijaya CO2 Incubator" Against Cell Culture Proliferation In Invitro Study As Smart Controlling-Based CO2 Incubator For Cell Culture", Bioscientia Medicina : Journal of Biomedicine and Translational Research, 2020 Publication	1 %
16	balimedicaljournal.org Internet Source	1 %
17	www.sid.ir Internet Source	1 %
18	eprints.qut.edu.au Internet Source	1 %
19	scholarshare.temple.edu	

Internet Source

1 %

20

Submitted to Universiti Teknologi MARA

Student Paper

1 %

21

www.medrxiv.org

Internet Source

<1 %

22

edisciplinas.usp.br

Internet Source

<1 %

23

Submitted to Universitas Diponegoro

Student Paper

<1 %

24

ejournal.radenintan.ac.id

Internet Source

<1 %

25

Agustini Utari, Muhammad Saifulhaq Maududi, Ninung Rose Diana Kusumawati, Maria Mexitalia. "Effects of low glycemic index diet on insulin resistance among obese adolescent with non-alcoholic fatty liver disease: a randomized controlled trial", *Medical Journal of Indonesia*, 2019

Publication

<1 %

26

doi.org

Internet Source

<1 %

27

link.springer.com

Internet Source

<1 %

28

"Abstracts from 38th Congress of the Société Internationale d'Urologie Seoul Dragon City, October 4-7, 2018", World Journal of Urology, 2018

Publication

<1 %

29

Edward Schiappa, Peter B. Gregg, Dean E. Hewes. "The Parasocial Contact Hypothesis", Communication Monographs, 2005

Publication

<1 %

30

Guangzhong Xu, Ming Song. "Recent advances in the mechanisms underlying the beneficial effects of bariatric and metabolic surgery", Surgery for Obesity and Related Diseases, 2020

Publication

<1 %

31

hqlo.biomedcentral.com

Internet Source

<1 %

32

www.scielo.br

Internet Source

<1 %

33

G D Kolovou. "Pathophysiology of dyslipidaemia in the metabolic syndrome", Postgraduate Medical Journal, 2005

Publication

<1 %

34

www.ncbi.nlm.nih.gov

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off