

# Correlation between Interleukin17 Levels with C- Reactive Protein and Neutrophil Lymphocyte Ratio in Sepsis - JBTR.pdf *by* Check Turnitin

---

**Submission date:** 11-May-2023 07:44AM (UTC-0400)

**Submission ID:** 2090327174

**File name:** Correlation between Interleukin17 Levels with C-Reactive Protein and Neutrophil Lymphocyte Ratio in Sepsis - JBTR.pdf (212.35K)

**Word count:** 3681

**Character count:** 19049

# JOURNAL OF BIOMEDICINE AND TRANSLATIONAL RESEARCH

Available online at JBTR website: <https://jbtr.fk.undip.ac.id>

Copyright © 2022 by Faculty of Medicine Universitas Diponegoro, Indonesian Society of Human Genetics and Indonesian Society of Internal Medicine

## Original Research Article

### Correlation between Interleukin-17 Levels with C-Reactive Protein and Neutrophil Lymphocyte Ratio in Sepsis

Dwi Retnoningrum<sup>1\*</sup>, Muchlis Achsan Udji Sofro<sup>2</sup>, Setyo Gundi Pramudo<sup>2</sup>, Satrio Adi Wicaksono<sup>3</sup>, Andaru Dahesihdewi<sup>4</sup>

<sup>1</sup>Department of Clinical Pathology, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

<sup>2</sup>Department of Internal Medicine, Faculty of Medicine, Universitas Diponegoro, Semarang, Indonesia

<sup>3</sup>Department of Anesthesiology and Intensive Care, Universitas Diponegoro, Semarang, Indonesia

<sup>4</sup>Department of Clinical Pathology and Laboratory Medicine, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta, Indonesia

#### Article Info

History

Received: 26 Jul 2022

Accepted: 25 Aug 2022

Available: 31 Aug 2022

#### Abstract

**Background:** Sepsis is a condition in systemic infection associated with organ function. Interleukin-17 is a pro-inflammatory cytokine produced by Th-17 cells. C-reactive protein and neutrophil-lymphocyte ratio (NLR) have been widely used as markers of inflammation. The relationship between IL-17 as a proinflammatory cytokine with CRP and NLR has not been reported.

**Objective:** This study aims to prove the correlation between IL-17 with CRP and NLR in septic patients.

**Methods:** Analytical observational study with a cross-sectional approach was carried out on 40 septic patients in July–December 2020. IL-17 levels were obtained by the ELISA method while CRP levels were obtained using the immunoturbidimetric method. NLR was the result of dividing the absolute number of neutrophils and lymphocytes from the automatic hematology analyzer. Correlation between variables was performed using Spearman correlation test.

**Results:** The median levels of IL-17, CRP, and NLR were 363.55 (11.4-1695.80) pg/mL, 13.25 (0.43-53.87) mg/L, and 12.00 (2.26–48.5), respectively. The Spearman correlation test between IL-17 levels and CRP obtained  $p = 0.019$ ,  $r = 0.37$ , and NLR  $p = 0.425$ ,  $r = 0.13$ .

**Conclusion:** There is a weak positive correlation between IL-17 levels and CRP in septic patients. There is no correlation between IL-17 levels and NLR in septic patients.

**Keywords:** Interleukin-17; CRP; NLR; sepsis

**Permalink/ DOI:** <https://doi.org/10.14710/jbtr.v8i2.15323>

#### INTRODUCTION

Sepsis is a collection of symptoms that often occurs in patients with systemic infections associated with organ dysfunction. The state of sepsis still has a sizable proportion of the critically ill patient population. Sepsis is defined as the presence or suspicion of infection associated with a systemic response, a life-threatening condition caused by dysregulation of the body's response to infection. There is no gold standard in the definition of sepsis, so the diagnosis of sepsis is based on a combination of physiology and laboratory abnormalities. The definition of sepsis itself changed at

international meetings in 1991, 2001, and, most recently, in 2016. The current recommendations and definitions of sepsis are based on organ dysfunction where a total SOFA score of 2 points is an indication of infection<sup>1</sup>.

The incidence of sepsis in the United States is about 300 cases per 100,000 population, 50% of whom undergo treatment in the Intensive Care Unit (ICU)<sup>2</sup>.

42

\* Corresponding author:

E-mail: [dwiretno@fk.undip.ac.id](mailto:dwiretno@fk.undip.ac.id)  
(Dwi Retnoningrum)

**Table 1.** Subjects Characteristic Data

Parameter	n (%)	Mean±SD	Median (min;max)	p*
Gender				
Male	19 (47.5)			
Female	21 (52.5)			
Age (year)		54.18±13.25	56 (19;70)	0.316
BMI		23.24±4.82	22.8 (12.8;34.52)	0.244
Hemoglobin (g/dL)		11.01±2.97	11.5 (2.9;14.9)	0.006
Leukocyte (10 <sup>3</sup> /uL)		16.45±9.08	14.75 (2;38)	0.304
Platelets (10 <sup>3</sup> /uL)		287.42±166.5	277 (35;626)	0.186
Neutrophil absolute count (10 <sup>3</sup> /uL)		13.7±7.85	12.2 (1.34;32.97)	0.087
Lymphocyte absolute count (10 <sup>3</sup> /uL)		1.39±1.29	1.09 (0.08;5.48)	0.001
NLR		16.57±13.70	13.51 (2.26;46.5)	0.001
IL-17 (pg/mL)		549.60±489.84	337.95 (11.4;1695.80)	0.001
CRP (mg/L)		17.6±14.97	13.25 (0.43;53.87)	0.001
SOFA score		9.74±3.10	9 (4;17)	0.318
Comorbidity				
Diabetes mellitus	8 (20)			
Hypertension	5 (12.5)			
Positive culture	15 (37.5)			
Gram (+) bacteria	6 (40)			
Gram (-) bacteria	8 9 (60)			

Note: BMI: Body Mass Index. SD: Standard Deviation. NLR: Neutrophil Lymphocyte Ratio. CRP: C-Reactive Protein. SOFA: Sequential Organ Failure Assessment. \* *Saphiro Wilk*

**Table 2.** Correlation between IL-17 with CRP and NLR

Variable	IL-17 (pg/mL)	
	p	r <sup>s</sup>
CRP (mg/L)	0.019*	0.37
NLR	0.425	0.13

IL-17: Interleukin-17, CRP: C-Reactive Protein, NLR: Neutrophil Lymphocyte Ratio; <sup>s</sup>Spearman Rank correlation test; \*p < 0.05

The epidemiological trend of sepsis in Spain shows that the incidence of sepsis is increasing from 3.3 per 1,000 population in 2000 to 4.45 in 2013, with an increase in mortality from 6.34 to 7.8 per 1,000 population<sup>3</sup>. Research in Asia (2009) from 150 intensive care patients in 16 countries (including Indonesia) showed severe sepsis and septic shock accounted for 10.9% of treatment diagnoses intensive care, with a mortality rate of 44.5%. In Indonesia, data from Cipto Mangunkusumo Hospital (2012), sepsis and septic shock occurred in 27% of patients in the ICU, with a mortality rate of 47.8%. The pathogenesis of sepsis begins with the presence of infectious agents that enter a person's blood circulation causing a systemic inflammatory state, but the presence of pathogens in the systemic circulation is not always present in cases of sepsis. The presence of inflammatory mediators released systemically can induce sepsis while the cause of sepsis itself can be various infectious agents, including bacteria, fungi, parasites, and viruses<sup>4</sup>.

Interleukin-17 is a proinflammatory cytokine produced by T helper-17, Natural Killer (NK) cells, D-8 T cells, neutrophils, and several other cells. IL-17 increases the production of chemokines so that it plays a significant role in the recruitment of monocytes and neutrophils to the site of inflammation. This shows the vital role of IL-17 as a proinflammatory cytokine that plays a role in infectious conditions. Neutrophil

lymphocyte ratio (NLR) has been widely used as a marker of the severity of the bacterial infection and to determine the prognosis of patients with infection. Recruitment of neutrophils to the site of infection will affect the value of NLR. Early studies in animal models induced sepsis showed that removing IL-17 was associated with a higher survival rate, but another study using mice with IL-17 receptor deficiency showed the opposite result<sup>5</sup>. Ali et al. reported that serum levels of IL-17 were associated with sepsis, where, in sepsis, IL-17 levels were 72 pg/mL compared to those without sepsis, namely 37 pg/mL with p = 0.0001. IL-17 levels more than 45 pg/mL is reported to have a 28-day mortality risk in septic patients (p = 0.005), so it can be a predictor of mortality in sepsis<sup>6</sup>.

The state of sepsis is characterized by changes in acute phase protein levels. One of the acute phase proteins is C-reactive protein (CRP). Elevated levels of CRP are produced in response to tissue damage, infection, inflammation, and malignancy<sup>7</sup>. Yang et al. stated that the sensitivity and specificity of CRP in the diagnosis of sepsis were 90.7% and 49.7% while Pradhan et al. in their study using a CRP cut-off value of 50 mg/L obtained a sensitivity of 84.3% and a specificity of 46.15%<sup>8,9</sup>. Ranzani et al. showed that CRP is a prognostic factor for septic patients admitted to the ICU, where a CRP level of 20 mg/dl has a poor prognosis (p = 0.024)<sup>10</sup>.

Neutrophil lymphocyte ratio (NLR) is an examination that is easily obtained from routine hematological examinations. Recently increased NLR is promising as a diagnostic and prognostic marker in patients with sepsis. Research by Marik et al showed the role of NLR in sepsis. NLR values > 10 had a sensitivity of 71% and specificity of 56% with a higher mean NLR value in positive bacterial cultures than in negative cultures<sup>11</sup>.

In sepsis, there is an increase in pro-inflammatory cytokines, one of which is IL-17 which plays a role in increasing natural, adaptive, and humoral immunity, in inflammation, an increase in acute-phase protein, namely CRP, and a response from leukocyte cells in inflammatory process, so it is necessary to research the correlation of serum IL-17 levels with CRP levels and NLR values in septic patients.

12

## MATERIALS AND METHODS

This research is an analytic observational study with a cross-sectional approach which was conducted in the ICU of Dr. Kariadi Hospital and Diponegoro National Hospital Semarang from July to December 2020. The research subjects were patients with sepsis criteria (SOFA score 2) with an age range of 18–75 years. Subjects were taken by consecutive sampling that met the inclusion and exclusion criteria. Patients with a history of liver disease and chronic renal failure were not included in this study.

The research variables consisted of levels of IL-17, CRP, and NLR. Examination of IL-17 levels measured using the sandwich enzyme-linked immunoassay (ELISA) principle from Elabscience® USA, was carried out at the GAKI Laboratory of Faculty of Medicine, Universitas Diponegoro. Sample allow to clot for 1 hour before centrifugation, 100 µL serum used for this assay. From the examination of CRP levels using the immunoturbidimetric method, the calculation of the NLR value was done manually by dividing the absolute neutrophil count by the absolute lymphocyte count obtained from the automatic hematology analyzer. Based on the normality test using the Saphiro Wilk test, IL-17, CRP, and NLR data were not normally distributed after data transformation. Then, Spearman Rank correlation test was performed. The p-value is considered significant if  $p < 0.05$ . Ethical clearance was obtained from the Health Research Ethics Committee of Dr. Kariadi General Hospital, Semarang No. 484/EC/KEPK/2020.

## RESULTS

In this study, forty patients with sepsis criteria according to SOFA criteria were included in this study, consisting of 19 (47.5%) males and 21 (52.5%) females. The mean age of the patients was  $54.18 \pm 13.25$  years with a normal mean body mass index ( $23.24 \pm 4.82$ ). The mean hemoglobin level is 11.5 g/dL, the mean leukocyte count is  $16.45 \times 10^3/\mu\text{L}$ , and the mean platelet count is  $287.42 \times 10^3/\mu\text{L}$ . The results of blood cultures from 40 subjects obtained positive blood cultures in 15 subjects (37.5%), whereas the results of gram-positive bacteria were 6 subjects (40%) and gram-negative 9 subjects (60%). The characteristics of the data are presented in Table 1.

The normality test using Saphiro Wilk found that three variables were not normally distributed, so the Spearman Rank correlation test was carried out. The correlation between IL-17 levels with CRP and NLR is presented in Table 2.




Figure 1. Correlation between IL-17 and CRP in Sepsis

9

There is a weak positive correlation between levels of IL-17 and CRP ( $p=0.019$ ,  $r=0.37$ ). The data distribution of IL-17 and CRP levels is shown in Figure 1. The correlation between IL-17 and NLR levels showed that the results were not statistically significant.

## DISCUSSION

The results showed that there was a positive correlation between IL-17 levels and CRP levels. C-reactive protein is a marker of inflammation routinely examined in clinical practice. The increase in CRP levels illustrates the increase in IL-17 levels in this study, which indicates an increase in proinflammatory cytokines. This is in accordance with the study of Akin et al. who reported a positive correlation between IL-17 and CRP ( $p = 0.014$ ,  $r = 0.225$ ) in patients with polymicrobial sepsis<sup>12</sup>. Comparable results were reported by the study of Zou et al. in patients with chronic obstructive pulmonary disease (COPD) and by Al-Saadani et al. in rheumatoid arthritis patients where there was a positive correlation between IL-17 levels and CRP ( $p < 0.001$ ,  $r = 0.506$ , and  $p < 0.001$ ,  $r = 0.696$ )<sup>13,14</sup>. Interleukin-17 is a cytokine produced by several cells, namely Th17 cells, T cells, NK cells, group 3 innate lymphoid cells (ILC3s), CD8+ cells (Tc17), microglia, and mast cells. The general function of IL-17 is to stimulate the production of inflammatory molecules, chemokines, and antimicrobial peptides (AMPs) such as defensin, calgranulin, and lipocalin-2. IL-17 also functions in maintaining mucosal barrier integrity and induce acute-phase proteins. In sepsis, IL-17 stimulates various immune cells (neutrophils, lymphocytes, macrophages, monocytes, and B cells) to enhance the role of natural, adaptive, and humoral immunity. IL-17 signals play a protective role against infections caused by bacteria, fungi, viruses, and parasites through the body's defense



and release of cytokines, but IL-17 can also induce pathological immune responses and the occurrence of organ failures such as acute lung injury (ALI), acute respiratory distress syndrome (ARDS), acute kidney injury (AKI), hepatic dysfunction, immunity, and cardiomyocytes<sup>15</sup>. The results of the observation of IL-17 in this study indicate pathological immune responses, that showed a high level of IL-17. An experimental study by Marchesi et al.<sup>55</sup> showed that sepsis caused by *Klebsiella pneumoniae* increased the production of IL-17 ( $p = 0.002$ ), and the highest level was reached at 48 hours after infection<sup>16</sup>. Ogiku et al. reported that mice with IL-17 deficiency showed a significant increase in mortality associated with bacteremia at 12 hours after the cecal ligation and puncture (CLP) procedure. This shows a partial protective role of IL-17 in sepsis, resulting in the positive control mice having a higher survival rate, compared to  $\Delta$ IL-17 mice that were knocked out of the IL-17 gene<sup>17</sup>. CRP is an acute-phase protein that can be used as a marker of inflammation. Li et al.'s study reported higher CRP levels in septic patients than in non-septic patients ( $p < 0.05$ ). Patients with poor clinical outcomes (non-survivors) also had higher CRP levels ( $p = 0.047$ )<sup>11</sup>.

The results of this study showed that there was no correlation between IL-17 levels and NLR. This is not following previous studies where the NLR value can be used as a marker for sepsis although, in this study, a high NLR value ( $> 5$ ) was not associated with IL-17 level<sup>23</sup>. The results of this study are similar to those of Sag et al. which showed that there was no correlation between IL-17 levels and NLR in rheumatoid arthritis patients ( $p > 0.05$ ,  $r = 0.135$ )<sup>19</sup>. In sepsis<sup>54</sup> there can be a failure of recruitment and migration of neutrophils to the site of infection, which is related to the severity of the disease. Interleukin-17 plays a role in the migration and microbicidal activity of neutrophils, and the mobilization of T lymphocytes<sup>20</sup>. Brunialti et al. show profound lymphopenia and decreased CD4+T cell counts in septic patients, and increased percentage of IL-17-producing T helper lymphocytes in the peripheral blood of septic patients<sup>21</sup>. The increase in NLR indicates the presence of an inflammation that, in addition to sepsis, can occur in other inflammatory conditions. The comorbidity of the patients can affect the value of NLR.

Positive blood culture results in study subjects were obtained in 37.5% of patients, which is almost similar to the study by Liestiadi et al. where positive culture results were obtained in 32.6% of patients, and the prevalence of gram-negative bacteria was higher than gram-positive<sup>22</sup>. In the study, the results showed that there were no differences in the levels of IL-35, CRP, and NLR in the subjects between the cultures of gram-positive bacteria and gram-negative bacteria.

The limitation of this study was that it did not take into account comorbid factors from the patient that could affect IL-17, CRP, and NLR levels. Patients on immunosuppressant therapy and the presence of hematological abnormalities were not excluded from the study as it could affect the results of complete blood counts.

## CONCLUSION

In summary, there was a weak correlation between IL-17 levels and CRP in the septic patients. Potential biases of the study could not be avoided due to small sample size, selection bias and variables in the clinical data of septic patients.

## ACKNOWLEDGMENTS

The authors would like to thank to Universitas Diponegoro, Dr. Kariadi General Hospital, Diponegoro National Hospital, and GAKI laboratory for their support and assistance throughout this study. This study was financially supported by Faculty of Medicine Universitas Diponegoro, Research Grant No. 85/UN7.5.4.2/HK/2020

## REFERENCES

- Gul F, Arslantas MK, Cinel I, Kumar A. Changing Definitions of Sepsis. *Turk J Anesth Reanim* [Internet]. 2017;45:129–38. Available from: <https://doi.org/10.5152/TJAR.2017.93753>
- Mayr FB, Yende S, Angus DC. Epidemiology of severe sepsis. *Virulence* [Internet]. 2014;5(1):1–11. Available from: <https://doi.org/10.4161/viru.27372>
- Álvaro-Meca A, Jiménez-Sousa MA, Micheloud D, Sánchez-Lopez A, Heredia-Rodríguez M, Tamayo E, et al. Epidemiological trends of sepsis in the twenty-first century (2000-2013): An analysis of incidence, mortality, and associated costs in Spain. *Popul Health Metr* [Internet]. 2018;16(1):1–11. Available from: <https://doi.org/10.1186/s12963-018-0160-x>
- Burkovskiy I, Sardinha J, Zhou J, Lehmann C. Cytokine release in sepsis. *Adv Biosci Biotechnol* [Internet]. 2013;2013(September):860–5. Available from: <http://dx.doi.org/10.4236/abb.2013.49114>
- Flierl MA, Rittirsch D, Gao H, Hoesel LM, Nadeau BA, Day DE, et al. Adverse functions of IL-17A in experimental sepsis. *FASEB J* [Internet]. 2008;22(July):2199–205. Available from: <https://doi.org/10.1096/fj.07-105221>
- Ali MA, Abdelkader ESMA, El LMR. Interleukin-17 as a predictor of sepsis in polytrauma patients: a prospective cohort study. *Eur J Trauma Emerg Surg* [Internet]. 2017; Available from: <http://dx.doi.org/10.1007/s00068-017-0841-3>
- Faraj M, Salem N. C-Reactive Protein. In: *Blood Cell- An Overview of Studies in Hematology*. 2012. p. 89–100. Available from: <https://www.intechopen.com/chapters/39106> doi: 10.5772/47735
- Yang Y, Xie J, Guo F, Longhini F, Gao Z, Huang Y, et al. Combination of C - reactive protein , procalcitonin and sepsis - related organ failure score for the diagnosis of sepsis in critical patients. *Ann Intensive Care* [Internet]. 2016;6(51). Available from: <https://doi.org/10.1186/s13613-016-0153-5>
- Pradhan S, Ghimire A, Bhattarai B, Khanal B, Pokharel K. The role of C - reactive protein as a diagnostic predictor of sepsis in a multidisciplinary Intensive Care Unit of a tertiary care center in Nepal. *Indian J Crit Care Med* [Internet]. 2016;(41):417–20. Available from: <https://doi.org/10.4103/0972-5229.186226>

10. Ranzani OT, Zampieri FG, Forte DN, Azevedo LCP, Park M. C-Reactive Protein/Albumin Ratio Predicts 90-Day Mortality of Septic Patients. *PLoS One* [Internet]. 2013;8(3):e59321. Available from: <https://doi.org/10.1371/journal.pone.0059321>
11. Li Q, Gong X. Clinical significance of the detection of procalcitonin and C-reactive protein in the intensive care unit. *Exp Ther Med* [Internet]. 2018;15(5):4265–70. Available from: <https://doi.org/10.3892/etm.2018.5960>
12. Akin H, Akalin H, Budak F, Ener B, Ocakoğlu G, Gürçüoğlu E, et al. Alterations of serum cytokine levels and their relation with inflammatory markers in candidemia. *Med Mycol* [Internet]. 2015;53(3):258–68. Available from: <https://doi.org/10.1093/mmy/myu084>
13. Zou Y, Chen X, Liu J, Zhou D bo, Kuang X, Xiao J, et al. Serum IL-1 $\beta$  and IL-17 levels in patients with COPD: association with clinical parameter. *Int J Chron Obstruct Pulmon Dis* [Internet]. 2017;12:1247–54. Available from: <https://doi.org/10.2147/COPD.S131877>
14. Al-Saadany HM, Hussein MS, Gaber RA, Zaytoun HA. Th-17 cells and serum IL-17 in rheumatoid arthritis patients: Correlation with disease activity and severity. *Egypt Rheumatol* [Internet]. 2016;38(1):1–7. Available from: <http://dx.doi.org/10.1016/j.ejr.2015.01.001>
15. Ge Y, Huang M, Yao YM. Biology of Interleukin-17 and Its Pathophysiological Significance in Sepsis. *Front Immunol* [Internet]. 2020;11(July):1–13. Available from: <https://doi.org/10.3389/fimmu.2020.01558>
16. Marchesi V V, Rukavina T. Interleukin-17 in experimental Klebsiella sepsis. *Period Biol* [Internet]. 2011;113(1):99–102. Available from: <https://hrcaak.srce.hr/67272>
17. Ogiku M, Kono H, Hara M, Tsuchiya M, Fujii H. Interleukin-17A Plays a Pivotal Role in Polymicrobial Sepsis According to Studies Using IL-17A Knockout Mice. *J Surg Res* [Internet]. 2012;174(1):142–9. Available from: <http://dx.doi.org/10.1016/j.jss.2010.11.901>
18. Martins EC, Da Fe Silveira L, Viegas K, Beck AD, Júnior GF, Cremonese RV, et al. Neutrophil-lymphocyte ratio in the early diagnosis of sepsis in an intensive care unit: A case-control study. *Rev Bras Ter Intensiva* [Internet]. 2019;31(1):63–70. Available from: <https://doi.org/10.5935/0103-507X.20190010>
19. Sağ S, Sağ MS, Tekeoğlu I, Kamanlı A, Nas K, Acar BA. Relationship of hematologic markers with IL-17 and IL-1 beta in patients with rheumatoid arthritis. *J Back Musculoskelet Rehabil* [Internet]. 2018;31(4):703–7. Available from: <https://doi.org/10.3233/BMR-170903>
20. Delano MJ, Ward PA. Sepsis-induced immune dysfunction: Can immune therapies reduce mortality? *J Clin Invest* [Internet]. 2016;126(1):23–31. Available from: <https://doi.org/10.1172/JCI82224>
21. Colo-Brunialti MK, Santos MC, Rigato O, Machado FR, Silva E, Salomao R. Increased Percentages of T Helper Cells Producing IL-17 and Monocytes Expressing Markers of Alternative Activation in Patients with Sepsis. *PLoS ONE* [Internet]. 7(5): e37393. Available from: <https://doi.org/10.1371/journal.pone.0037393>
22. Liestyadi DEF, Azlin E, Nafianti S. A hematologic scoring system and C-reactive protein compared to blood cultures for diagnosing bacterial neonatal sepsis. *Paediatr Indones* [Internet]. 2017;57(2):70–5. Available from: <https://doi.org/10.14238/pi57.2.2017.70-5>

# Correlation between Interleukin17 Levels with C-Reactive Protein and Neutrophil Lymphocyte Ratio in Sepsis - JBTR.pdf

## ORIGINALITY REPORT

**33%**  
SIMILARITY INDEX

**25%**  
INTERNET SOURCES

**22%**  
PUBLICATIONS

**9%**  
STUDENT PAPERS

## PRIMARY SOURCES

**1** Submitted to Universitas Diponegoro **4%**  
Student Paper

**2** [www.frontiersin.org](http://www.frontiersin.org) **3%**  
Internet Source

**3** Sianny Herawati, Yenny Kandarini, Ni Kadek Mulyantari, Putu Yuda Prabawa. "Correlation of Neutrophil to Lymphocyte Ratio with Interleukin-10 in Diagnosis and Monitoring of Coronavirus Disease-19 Patients", Open Access Macedonian Journal of Medical Sciences, 2022 **1%**  
Publication

**4** [icm-experimental.springeropen.com](http://icm-experimental.springeropen.com) **1%**  
Internet Source

**5** [balimedicaljournal.org](http://balimedicaljournal.org) **1%**  
Internet Source

**6** [www.pubmedcentral.nih.gov](http://www.pubmedcentral.nih.gov) **1%**  
Internet Source

[medic.upm.edu.my](http://medic.upm.edu.my)

7	Internet Source	1 %
8	topsecretapiaccess.dovepress.com Internet Source	1 %
9	inabj.org Internet Source	1 %
10	www.science.gov Internet Source	1 %
11	core.ac.uk Internet Source	1 %
12	jddtonline.info Internet Source	1 %
13	journals.sagepub.com Internet Source	1 %
14	Andi Itha Maghfirah, Rachmawati Muhiddin, Irda Handayani, Nursin Abdul Kadir. "Analysis of procalcitonin and urine nitrite to predictor sepsis patients", MEDISAINS, 2021 Publication	1 %
15	Submitted to University of Sydney Student Paper	1 %
16	Submitted to Curtin University of Technology Student Paper	1 %
17	Submitted to Tufts University Student Paper	1 %



18 [paper.researchbib.com](http://paper.researchbib.com) <1 %  
Internet Source

---

19 [www.ccsenet.org](http://www.ccsenet.org) <1 %  
Internet Source

---

20 [www.ijretina.com](http://www.ijretina.com) <1 %  
Internet Source

---

21 [scholararchive.ohsu.edu](http://scholararchive.ohsu.edu) <1 %  
Internet Source

---

22 "Abstracts from 38th Congress of the Société  
Internationale d'Urologie Seoul Dragon City,  
October 4-7, 2018", World Journal of Urology,  
2018 <1 %  
Publication

---

23 Submitted to Bilkent University <1 %  
Student Paper

---

24 Submitted to Queen Mary and Westfield  
College <1 %  
Student Paper

---

25 Submitted to Universitas Jenderal Soedirman <1 %  
Student Paper

---

26 [www.oncotarget.com](http://www.oncotarget.com) <1 %  
Internet Source

---

27 [link.springer.com](http://link.springer.com) <1 %  
Internet Source

---

[repository.stikim.ac.id](http://repository.stikim.ac.id)

28

Internet Source

&lt;1 %

29

[synapse.koreamed.org](https://synapse.koreamed.org)

Internet Source

&lt;1 %

30

Pablo Palazon-Riquelme, Gloria Lopez-Castejon. "The inflammasomes, Immune guardians at defence barriers", Immunology, 2018

Publication

&lt;1 %

31

Schuerwegh, A.J.. "FLOW CYTOMETRIC DETECTION OF TYPE 1 (IL-2, IFN-@c) AND TYPE 2 (IL-4, IL-5) CYTOKINES IN T-HELPER AND T-SUPPRESSOR/CYTOTOXIC CELLS IN RHEUMATOID ARTHRITIS, ALLERGIC ASTHMA AND ATOPIC DERMATITIS", Cytokine, 199910

Publication

&lt;1 %

32

[journals.plos.org](https://journals.plos.org)

Internet Source

&lt;1 %

33

Meichun Tan, Yunxia Lu, Hao Jiang, Liandong Zhang. "The diagnostic accuracy of procalcitonin and C-reactive protein for sepsis: A systematic review and meta-analysis", Journal of Cellular Biochemistry, 2018

Publication

&lt;1 %

34

[journals.lww.com](https://journals.lww.com)

Internet Source

&lt;1 %

35	<a href="http://kupdf.net">kupdf.net</a> Internet Source	<1 %
36	<a href="http://pubmed.ncbi.nlm.nih.gov">pubmed.ncbi.nlm.nih.gov</a> Internet Source	<1 %
37	<a href="http://worldwidescience.org">worldwidescience.org</a> Internet Source	<1 %
38	<a href="http://www.pneumon.org">www.pneumon.org</a> Internet Source	<1 %
39	<a href="http://www.wjgnet.com">www.wjgnet.com</a> Internet Source	<1 %
40	Basak, P.Y.. "The role of helper and regulatory T cells in the pathogenesis of vitiligo", Journal of the American Academy of Dermatology, 200902 Publication	<1 %
41	Dreesen, L., K. De Bosscher, G. Grit, B. Staels, E. Lubberts, E. Bauge, and P. Geldhof. "Giardia muris Infection in Mice Is Associated with a Protective Interleukin 17A Response and Induction of Peroxisome Proliferator-Activated Receptor Alpha", Infection and Immunity, 2014. Publication	<1 %
42	Meira Erawati, Nyoman Suci Widyastiti, Tri Indah Winarni, Edi Dharmana. "beta-Glucan Increases IFN-gamma and IL-12 Production of	<1 %

Peripheral Blood Mononuclear Cells  
with/without Induction of Mycobacterium  
tuberculosis Wild-type/Mutant DNA", The  
Indonesian Biomedical Journal, 2019

Publication

43

Na Cui, Hongwei Zhang, Zhi Chen, Zhanbiao Yu. "Prognostic significance of PCT and CRP evaluation for adult ICU patients with sepsis and septic shock: retrospective analysis of 59 cases", Journal of International Medical Research, 2019

Publication

<1 %

44

[bioscmed.com](http://bioscmed.com)

Internet Source

<1 %

45

[dspace.knust.edu.gh](http://dspace.knust.edu.gh)

Internet Source

<1 %

46

[eujournal.org](http://eujournal.org)

Internet Source

<1 %

47

[journal-inflammation.biomedcentral.com](http://journal-inflammation.biomedcentral.com)

Internet Source

<1 %

48

[nutritionj.biomedcentral.com](http://nutritionj.biomedcentral.com)

Internet Source

<1 %

49

[ouci.dntb.gov.ua](http://ouci.dntb.gov.ua)

Internet Source

<1 %

50

[pjmhsonline.com](http://pjmhsonline.com)

Internet Source

<1 %

51	<a href="https://repository.up.ac.za">repository.up.ac.za</a> Internet Source	<1 %
52	<a href="https://scholarworks.bridgeport.edu">scholarworks.bridgeport.edu</a> Internet Source	<1 %
53	<a href="https://static-site-aging-prod2.impactaging.com">static-site-aging-prod2.impactaging.com</a> Internet Source	<1 %
54	<a href="https://www.dovepress.com">www.dovepress.com</a> Internet Source	<1 %
55	<a href="https://www.jrheum.org">www.jrheum.org</a> Internet Source	<1 %
56	<a href="https://www.nature.com">www.nature.com</a> Internet Source	<1 %
57	<a href="https://www.tara.tcd.ie">www.tara.tcd.ie</a> Internet Source	<1 %
58	Hongxia Zhao, Cheol-In Kang, Mark S. Rouse, Robin Patel, Hirohito Kita, Young J. Juhn. "The Role of IL-17 in the Association between Pneumococcal Pneumonia and Allergic Sensitization", International Journal of Microbiology, 2011 Publication	<1 %
59	Kristen N. Morrow, Craig M. Coopersmith, Mandy L. Ford. "IL-17, IL-27, and IL-33: A Novel Axis Linked to Immunological Dysfunction During Sepsis", Frontiers in Immunology, 2019	<1 %



60

Anne C. Bay-Jensen, Anne Sofie Siebuhr, Dres Damgaard, Patryk Drobinski, Christian Thudium, Joachim Mortensen, Claus H Nielsen. "Objective and noninvasive biochemical markers in rheumatoid arthritis: where are we and where are we going?", Expert Review of Proteomics, 2021

Publication

---

<1 %

61

Hongmin Zhang, Xiaoting Wang, Qing Zhang, Ying Xia, Dawei Liu. "Comparison of procalcitonin and high-sensitivity C-reactive protein for the diagnosis of sepsis and septic shock in the oldest old patients", BMC Geriatrics, 2017

Publication

---

<1 %

62

Senem Buyukkara Yilmaz, Nilüfer Cicek, Mesut Coskun, Olcay Yegin, Erkan Alpsoy. "Serum and tissue levels of IL-17 in different clinical subtypes of psoriasis", Archives of Dermatological Research, 2012

Publication

---

<1 %

---

Exclude quotes Off

Exclude matches Off

Exclude bibliography On