

# 2019 Al-Baarri Production of ozone

*by* Setya Budi Muhammad Abduh

---

**Submission date:** 12-May-2023 01:45AM (UTC+0700)

**Submission ID:** 2090631641

**File name:** 2019\_Al-Baarri\_Production\_of\_ozone.pdf (432.08K)

**Word count:** 1349

**Character count:** 6745

PAPER · OPEN ACCESS

## Production of Ozone and the Simple Detection using Potassium Iodide Titration Method

<sup>3</sup> To cite this article: A N Al-Baarri *et al* 2019 *IOP Conf. Ser.: Earth Environ. Sci.* **292** 012062

### Recent citations

<sup>4</sup> - [Cleaner approach for medium consistency eucalyptus slab pulp production using ozone bleaching under turbulent mixing](#)  
Xi Zhang *et al*

View the [article online](#) for updates and enhancements.



The Electrochemical Society  
Advancing solid state & electrochemical science & technology  
2021 Virtual Education

**Fundamentals of Electrochemistry:**  
Basic Theory and Kinetic Methods

Instructed by: **Dr. James Noël**

Sun, Sept 19 & Mon, Sept 20 at 12h–15h ET

Register early and save!



## Production of Ozone and the Simple Detection using Potassium Iodide Titration Method

A N Al-Baarri <sup>1,2\*</sup>, A M Legowo <sup>1</sup>, S B M Abduh <sup>1</sup>, A A Mawarid <sup>2</sup>, K M Farizha <sup>2</sup>, M Silvia <sup>2</sup>

<sup>1</sup>Food Technology Department, Faculty of Animal and Agricultural Sciences, Diponegoro University, Indonesia

<sup>2</sup>Center of Research and Services-Diponegoro University, Semarang 50275

\*Corresponding author: albari@live.undip.ac.id

**Abstract.** Ozone is a molecule consisting of three oxygen atoms ( $O_3$ ) and a powerful oxidizing agent that can also act as a non-chemical disinfectant. Ozone can be detected using potassium iodide and titrated using sodium thiosulfate to know the concentration. The purpose of this study is to know the output of ozone concentration from the ozone generator. Ozone measurement was done by making KI solutions and the concentration are 0,2; 0,4; 0,6; 0,8; and 1 M then flew by ozone for 2 minutes with flow rate 1 L/min and the color of the solution will change from clear to brown. The brown solution titrated using  $Na_2S_2O_3$  0,4M solution, until the color of the KI solution becomes clear again. In conclusion, potassium iodide (KI) can detect the output of ozone generator which means potassium iodide (KI) concentrations linier with ozone concentration.

**Keywords:** ozone, simple detection, potassium iodide, titration method

### I. Introduction

Ozone is a molecule consisting of three oxygen atoms ( $O_3$ ) and is a powerful oxidizing agent which can be also acted as a non-chemical disinfectant [6]. Ozone was specified as non-toxic in low concentrations, environmentally friendly, relatively harmless, and almost similar to oxygen. The production of ozone can be made by flowing oxygen into the ozone generator. The initial method for ozone generation is utilization of splash from an electrode that may break up oxygen to convert into three bonds [5]. The high concentration and purity of ozone may be adjusted using controlled flow rate and the pure of oxygen input, therefore for the achievement of good validity of ozone measurement, the pure oxygen is advised to be applied in the system [5].

Ozone can be detected via the utilization of potassium iodide or KI and sodium thiosulphate or  $Na_2S_2O_3$  using titration methods. This can be applied since reaction of the system produces the visible color at the final reaction, thus this visible color appearance can be detected with ease and may provide the linear graph as the volume of applied solution. The amount of ozone may be detected by flowing ozone into the KI solution, then analyzed using titration method using sodium thiosulfate [3].

The purpose of this study was to expose the color appearance of titrated sample and concentration of ozone caught by KI with the high volume of KI.



## <sup>7</sup> **2. Materials and Methods**

### **2.1. Materials**

#### **2.1.1. Chemical materials**

<sup>3</sup> KI (0.2 mM) from Roche (Germany), ozone generator, oxygen container from <sup>1</sup> modern health shop in Semarang, Indonesia, aquadest, and sodium thiosulfate were obtained from Center of Research and Services-Diponegoro University, Semarang, Indonesia.

#### **2.2. Methods**

##### **2.2.1. KI solution preparation**

KI in aquadest solution was prepared from the initial KI at the concentration of 0.2 mM using KI powder. Other concentration was also made to produce KI at concentration of 0.4, 0.6, 0.8, and 1 mM.

##### **2.2.2. Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution preparation**

Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> in aquadest solution was made from the powder at the concentration of 0.4 mM.

##### **2.2.3. Ozone treatment**

Ozone treatment was applied using following the method with small modification [ 4 ]. Potassium iodide at the concentration of 0.2, 0.4, 0.6, 0.8, and 1 mM were flew by ozone using ozone generator for 2 minutes with the adjusted flow rate at 1 L/min. The color of the solution changed from clear to brick red.

##### **2.2.4. Titration**

The brick-red-colored solution then was titrated using 0.4 M Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution until the color of the KI solution turned back to clear.


##### **2.2.5. Titration analysis**

Testing was done by calculating the dissolving ozone based at the input oxygen rate. For example 10 L/min (Vair = 10 L min) was applied into KI solution, for example 2 minutes (t = 2 minutes). KI solutions then was exposed to ozone. KI and ozone solutions were then titrated using Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> solution until the KI solution turned back to colorless. The volume Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> was then recorded and substituted into the following formula:

$$\text{Ozone Concentration} = \frac{Mr \ O_3 \cdot V \cdot N}{Vair \cdot e \cdot t}$$

while Mr O<sub>3</sub> is 16.3 = 48, V is Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> volume, N is Na<sub>2</sub>S<sub>2</sub>O<sub>3</sub> concentration, Vair is air flow rate (oxygen), e is electron mass X 2, t is time (min)

### 3. Result and Discussions



**Figure 1.** The ozone concentration changes using the KI at the of 0.2, 0.4, 0.6, 0.8, and 1 mM

Determining the product of formed ozone is based on the reaction of  $I^-$  with  $O_3$  which produced  $I_2$  by reaction  $O_3 + 2I^- + H_2O \Leftrightarrow I_2 + 2OH^- + O_2$ . The equivalent amount of  $I_2$  formed in the KI solution was then titrated with sodium thiosulfate by reaction  $I_2 + 2 Na_2S_2O_3 \Leftrightarrow 2NaI + Na_2S_4O_6$  [ 1 ]. The data shows that the concentration of 0.2 mmol potassium iodide could produce ozone at concentration of 2.5 mM while 0.4 mmol potassium iodide generated 3.3 mM ozone. Potassium iodide at concentration of 0.6, 0.8, and 1 mM produced 5.9, 5.8, and 6.2 mM ozone, respectively. This shows that **the higher the concentration of potassium iodide (KI), the higher the ozone** could be trapped. This is in accordance with the statement that KI binds ozone and more KI concentration the more ozone may be bound [ 2 ] .

### 4. Conclusion

Based on this study, potassium iodide could bind ozone and the more potassium iodide concentration the more ozone might be bound.

### 5. References

- [ 1 ] Atkinson, R., and S.M. Aschmann. 1993. Hydroxyl radical production from the gas-phase reactions of ozone with a series of alkenes under atmospheric conditions. *Environmental Science & Technology*. 27(7): 1357-1363.
- [ 2 ] Boelter, E. D., G. L. Putnam and E. I. Lash. 1950. Iodometric determination of ozone of high concentration. *Analytica*.

- [ 3 ] Ding, W., Wang, Y., Zhang, W., Shi, Y., & Wang, D. 2015. Effect of ozone treatment on physicochemical properties of waxy rice flour and waxy rice starch. *International Journal of Food Science & Technology*. 50(3): 744-749.
- [ 4 ] Kusvuran, E., O. Gulnaz, A. Samil and M. Erbil. 2010. Detection of double bond-ozone stoichiometry by an iodimetric method during ozonation processes. *Journal of hazardous materials*. 175(1-3): 410-416.
- [ 5 ] Rodden, R.M., 2001. Ozone generator. U.S. Patent 6,270,733.
- [ 6 ] Schork, B. M., J. Besore, Y. K. Allimalli and L. Miller. 2011. U.S. Patent Application No. 12/707,779.

# 2019 Al-Baarri Production of ozone

---

## ORIGINALITY REPORT

---

19%

SIMILARITY INDEX

%

INTERNET SOURCES

19%

PUBLICATIONS

%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

1

N Yaqin, A N Al-Baarri, A M Legowo, Widayat, M A Budihardjo. "Physical Characteristics of Modified Cassava Flour Wastewater at Room Temperature", IOP Conference Series: Earth and Environmental Science, 2019

Publication

6%

2

N D Indrawati, E Kusumawati, R Maria. "Nutrition and Contraception: The Most Influence Factors in Extending the Age of menopause", IOP Conference Series: Earth and Environmental Science, 2019

Publication

5%

3

A N Al-Baarri, A M Legowo, E Auliana, A D Puspitoasih, A C D Wratsongko, L Izzati, Michael, M Hadipernata, W Broto. "Stability of Antibacterial Agent Hypoiodous Acid Against Time and Temperature of Storage", IOP Conference Series: Earth and Environmental Science, 2019

Publication

3%

---

4

Xi Zhang, Jun Li, Jie Gong, Yishan Kuang, Shuaiming He, Jun Xu, Lihuan Mo, Yi Man, Wenyan Zhu, Junlong Song, Tao Song. "Cleaner approach for medium consistency eucalyptus slab pulp production using ozone bleaching under turbulent mixing", *Journal of Cleaner Production*, 2020

Publication

2%

5

Jéssica Luiza Bueno Trevizani, André Nagalli, Fernando Hermes Passig, Karina Querne de Carvalho et al. **"Influence of pH and concentration on the decolorization and degradation of BR red azo dye by ozonization"**, *Acta Scientiarum. Technology*, 2018

Publication

1%

6

Min Li, Jiaqi Tang, Yan Luo, Jinglun Yang, Kaiqiang Liu, Zhongshan Liu, Yu Fang. **"Controllable Preparation of Imine-based Nanofilms Towards Ozone Detection and Removal"**, *Sensors and Actuators B: Chemical*, 2023

Publication

1%

7

Mohsen Faghinezhad, Majid Baghdadi, Mohammad Sajjad Shahin, Ali Torabian. **"Catalytic ozonation of real textile wastewater by magnetic oxidized g-C<sub>3</sub>N<sub>4</sub> modified with Al<sub>2</sub>O<sub>3</sub> nanoparticles as a**

1%



# novel catalyst", Separation and Purification Technology, 2022

Publication

---

---

Exclude quotes      On

Exclude matches      Off

Exclude bibliography      On