

African leaf (*Vernonia amygdalina*) extracts improve Japanese quail (*Coturnix coturnix japonica*) carcass traits

by Teysar Adi Sarjana

Submission date: 25-May-2023 02:49PM (UTC+0700)

Submission ID: 2101469926

File name: ove_Japanese_quail_Coturnix_coturnix_japonica_carcass_traits.pdf (482.08K)

Word count: 5340

Character count: 28405

African leaf (*Vernonia amygdalina*) extracts improve Japanese quail (*Coturnix coturnix japonica*) carcass traits

Sri Kismiati¹, Teysar Adi Sarjana², Luthfi Djauhari Mahfudz³, and Dwi Sunarti Prayitno⁴

¹Department of Animal Science, Faculty of Animal and Agricultural Sciences, Diponegoro University, Tembalang Campus, Semarang, Central Java, Indonesia.

Corresponding author: Sri Kismiati, e-mail: kismiati59@gmail.com

Co-authors: TAS: teysaradisarjana@lecturer.undip.ac.id, LDM: lmahfudz@gmail.com, DSP: dwisunarti@gmail.com

Received: 28-11-2022, **Accepted:** 09-03-2023, **Published online:** 15-04-2023

doi: www.doi.org/10.14202/vetworld.2023.773-778 **How to cite this article:** Kismiati S, Sarjana TA, Mahfudz LD, and Prayitno DS (2023) African leaf (*Vernonia amygdalina*) extracts improve Japanese quail (*Coturnix coturnix japonica*) carcass traits, *Veterinary World*, 16(4): 773-778.

Abstract

Background and Aim: The *Vernonia amygdalina* Del. leaf extract (VALE) contains several natural antioxidants including flavonoids, which effectively ameliorate cholesterol levels while improving quail carcass traits and meat quality. This study aimed to evaluate the effects of VALE on Japanese quail (*Coturnix coturnix japonica*) carcass traits and meat.

Materials and Methods: In total, 260 Japanese quails (aged 5 weeks and body weight = 129.1 ± 2.2 g) were raised in an open-sided house and randomized to four VALE treatments: T0: Control, T1: 10 mL/L, T2: 20 mL/L, and T3: 10 mL/L in drinking water. After 12 weeks, carcass traits and chemical and physical meat qualities were evaluated.

Results: *Vernonia amygdalina* leaf extract in drinking water exerted significant effects ($p < 0.05$) on carcass weight, cholesterol levels, and meat water holding capacity (WHC) without significantly affecting carcass and non-carcass percentages, moisture, protein, fat, and meat color qualities. The highest carcass weights and lowest cholesterol levels were identified in the T2 group, while WHC improved in the T3.

Conclusion: Thus, VALE supplementation (20 mL/L) to quails improved carcass traits, especially cholesterol levels and carcass weights.

Keywords: chemical and physical, cholesterol, fat and meat color, moisture, non-carcass percentage, protein.

Introduction

Quail is a domesticated bird of important economic value. Its meat and eggs have specific flavors, with Japanese quails (*Coturnix coturnix japonica*) producing one of the best meats among domesticated quails. The average adult weight is approximately 200 g, with birds easily stressed due to environmental conditions. Quail meat contains nutrients which provide valuable sources of animal protein. In Indonesia, the quail agri-business is unique, in some case female quail of laying type are harvested even during egg production phases. Thus, apart from egg production, carcass production traits and meat quality have become serious issues during the rearing processes. Studies have indicated that 8-week-old Japanese quail meat contains high protein and fat levels (23.32–24.55 g/100 g and 1.1–1.30 g/100 g, respectively) [1], but high cholesterol levels (71.07–87.0 mg/100 g) [1, 2]. However, quail farming in Indonesia faces highly fluctuating temperature and humidity conditions; too high or low temperatures

affect bird health, growth, product quality [3]; carcass quality [4, 5]; body weight, carcass weight, carcass percentages [6, 7]; immunity, nutrient digestibility, and product quality [8, 9]. Environmental farming factors cannot be ignored and have key roles in determining performance and meat quality, especially when oxidative stress mechanisms and potential negative impacts are considered. Long-term bird stress also disturbs protein synthesis in different muscle groups and reduces protein breakdown and disposition in tissue [5]. To cope with stress effects, providing plant-based feed additives is a safe and viable solution [8]; using natural ingredients in Japanese quail diets improves carcass quality characteristics and meat quality [10].

Vernonia amygdalina Del. leaf extracts (VALE) contain phenolic acid and flavonoid compounds [11]. Phenolic acid is an immunomodulator, anti-mutagenic, and anti-inflammatory agent [12]. *Vernonia amygdalina* is a natural source of antioxidants, including phenols, flavonoids, anthocyanin, proanthocyanin, and tannins [13], which combat oxidative stress [14]. Adding flavonoids to broiler diets increased villus height and ratios, duodenum and jejunum crypt depth, growth, and carcass composition [15]. Natural antioxidant supplementation to quail feed reduced heat stress, significantly increased body and carcass weight [16], and increased meal quality [17]. In broilers, Saracila *et al.* [18] used antioxidants to reduce stress effects

Copyright: Kismiati, *et al.* Open Access. This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated.

and successfully increase protein content. Japhet and Godgift [19] reported that VALE supplementation (50 mL/L) to drinking water increased broiler carcass performance but did not increase the laying hens pullet performance. Supplemented *V. amygdalina* (30 mL/L) juice in drinking water increased carcass weight but percentages were not significantly different [20]. Suhaemi and Hidayati [21] showed that African leaves reduced cholesterol levels in broiler meat. Efforts to overcome stress in quail using VALE also improved egg production and quality [22].

Research of African leaves extract (VALE) in broiler chickens has been widely conducted, while there has been no publication on quail during the laying period related to carcass trait. Therefore, this study aimed to assess the impact of VALE on quail characteristics and carcass quality and observed that appropriate VALE supplementation increased carcass traits and improved final products.

Materials and Methods

Ethical approval

The study was approved by the Animal Ethics Committee in the Faculty of Animal Sciences, Diponegoro University, Semarang, Indonesia (Approval number 58-09/A-8/KEP-FPP/2022).

Study period and location

This study was conducted from September to December 2022 in the Poultry Production Laboratory of the Faculty of Animal and Agricultural Sciences, Diponegoro University.

Vernonia amygdalina leaf extract preparation

This study used African leaves (*V. amygdalina*) that are comprised of young and old leaves bought from a Small Medium Enterprise herb producer (Bibitherb, Jember City, Indonesia). *Vernonia amygdalina* leaf extract was generated using a previous decoction method [23], where leaves were aerated until dry and then ground. Approximately 50 g *V. amygdalina* powder was boiled for 30 min in water, left cool for 5 min at room temperature (25°C) and filtered using Whatman No. 41. The filtrate was used for treatments.

Animals

We used 260 5-week-old quails (average body weight = 129.1 ± 2.2 g) who were raised in an open-sided house. Quails were obtained from Kalisidi Village, West Ungaran District, Semarang Regency, Indonesia. Feed composition: Yellow corn, soybean meal, meat and bone meal, limestone, top mix product from PT Medion (Indonesia), and NaCl. The nutrient ratio was: protein: 21.23%, metabolic energy: 2741.57 kcal/kg/ration, fat: 1.2%, and crude fiber: 2.3%.

Study design

Our study incorporated a completely randomized design with four treatments and five repetitions (13 quails/repetition) with 4 treatments. *Vernonia amygdalina* leaf extract was added to drinking water,

that is, T0: Control, T1: 10 mL/L, T2: 20 mL/L, and T3: 30 mL/L.

Parameters

Life weight, carcass weight, and meat chemical analyses were performed at the study end (17-week-old), with two quails/experimental units used after 8 h of fasting. Quails were slaughtered according to halal procedures, then scalded, defeathered, and eviscerated [24]. Meat chemical composition analyses were performed on breasts and thighs. Samples were taken separately and stored at -20°C. Meat was removed from bones and homogenized and analyzed for moisture, crude protein, and crude fat content according to Association of Official Agricultural Chemists method [25]. Cholesterol content was determined using an enzymatic colorimetric method; samples were saponified using methanolic potassium hydroxide and Fluitest cholesterol kit (DSI, Indonesia) was used for assay. Results were recorded at 500 nm using a spectrophotometer (Shimadzu, UV1900I, EUROPA GmbH, Germany).

Statistical analysis

Data were analyzed in a completely randomized design using a one-way analysis of variance and Duncan's Multiple Range Tests with 95% confidence level ($\alpha = 0.05$). Analyses were performed using JMP® Pro 13.0.0 (64-bit) software (JMP Statistical Discovery LLC, Netherlands). Orthogonal polynomial analyses were performed using MS Excel 16 (Microsoft Corporation, USA).

Results

Carcass performance

Vernonia amygdalina leaf extract significantly affected carcass weight ($p < 0.05$), although live weight and carcass and non-carcass percentages were not significantly affected ($p > 0.05$) (Table-1).

Meat quality

Vernonia amygdalina leaf extract (T1, T2, and T3) did not significantly ($p > 0.05$) affect fat and protein content but significantly ($p < 0.05$) affected cholesterol levels in meat (Table-2). Carcass cholesterol levels were decreased significantly ($p < 0.05$) in the T1 group but had decreased more in T2. T3 and T2 groups were not significantly different ($p > 0.05$).

Carcass physical qualities

Vernonia amygdalina leaf extract significantly ($p < 0.05$) affected water holding capacity (WHC) in breast meat, but did not significantly affect ($p > 0.05$) pH, drip loss, breast and thigh meat color, and WHC of thigh meat (Table-3).

Discussion

Live weight was not significantly different between groups ($p > 0.05$), but VALE supplementation significantly increased carcass weight ($p < 0.05$) (Table-1). This result was similar to Osho *et al.* [26], who showed that VALE (aqueous) supplementation

Table-1: Carcass performance of Japanese Quail when *Vernonia amygdalina* leaf extract was added to their drinking water.

Parameter	Treatment				p-value
	T0	T1	T2	T3	
Live weight (g)	166.50 ± 4.18	178.50 ± 9.62	174.00 ± 15.47	162 ± 7.37	0.08
Carcass weight (g)	94.45 ± 4.83 ^b	103.27 ± 4.23 ^a	103.21 ± 9.92 ^a	90.84 ± 7.50 ^b	<0.01
Percentage of carcass (%)	56.75 ± 3.23	57.93 ± 2.51	59.31 ± 2.08	56 ± 2.10	0.22
Percentage of non-carcass (%)	24.50 ± 1.95	24.79 ± 1.38	25.30 ± 1.64	24.14 ± 1.54	0.73

^{a,b}Means with different superscripts along the same row are significantly different ($p < 0.01$)

Table-2: The chemical quality of quail meat when *Vernonia amygdalina* leaf extract was added to the drinking water.

Parameter	Treatment				p-value
	T0	T1	T2	T3	
Moisture (%)	73.74 ± 0.45	72.08 ± 0.45	72.23 ± 0.55	72.70 ± 0.84	0.08
Fat (%)	1.57 ± 0.22	1.79 ± 0.24	1.80 ± 0.44	1.84 ± 0.21	0.49
Protein (%)	21.22 ± 0.56	21.91 ± 0.48	21.98 ± 1.24	22.73 ± 0.87	0.22
Cholesterol (mg/100 g)	294.31 ± 47.09 ^a	226.12 ± 24.02 ^b	167.85 ± 30.1 ^c	164.46 ± 41.1 ^c	0.01

^{a,b,c}Means with different superscript along the same row are significantly different ($p < 0.01$)

Table-3: Physical properties of meat quail when *Vernonia amygdalina* extract leaf extract was added to the drinking water.

Parameter	Treatment				p-value
	T0	T1	T2	T3	
pH	6.36 ± 0.54	6.28 ± 0.39	6.30 ± 0.38	6.26 ± 0.46	0.28
Breast	6.36 ± 0.54	6.28 ± 0.39	6.30 ± 0.38	6.26 ± 0.46	0.28
Thigh	6.78 ± 0.29	6.70 ± 0.24	6.82 ± 0.13	6.82 ± 0.18	0.28
WHC					
Breast	23.76 ± 1.17 ^b	24.21 ± 1.70 ^b	23.43 ± 0.90 ^b	30.81 ± 1.84 ^a	<0.01
Thigh	35.15 ± 16.73	41.50 ± 11.98	39.27 ± 9.25	38.31 ± 7.03	0.95
Drip loss					
Breast	5.79 ± 0.84	5.12 ± 0.86	5.98 ± 0.10	5.59 ± 1.30	0.89
Thigh	5.66 ± 1.40	5.36 ± 1.69	4.22 ± 1.99	4.21 ± 1.85	0.95
L* Breast	26.90 ± 5.66	23.38 ± 8.94	27.70 ± 11.10	27.30 ± 9.65	0.91
L* Thigh	37.60 ± 6.82	37.60 ± 11.54	37.50 ± 14.56	38.00 ± 13.38	0.89
a* Breast (a)	15.50 ± 1.66	14.60 ± 1.47	15.80 ± 1.20	15.40 ± 10.74	0.90
a*Thigh (a)	16.20 ± 1.04	16.60 ± 1.08	16.50 ± 2.52	28.75 ± 6.08	0.68
b* Breast	3.13 ± 0.85	2.60 ± 0.96	3.10 ± 1.56	2.20 ± 0.67	0.51
b* Thigh	4.00 ± 0.91	4.10 ± 0.47	3.90 ± 1.08	4.10 ± 2.79	0.28

L* = Lightness, a* = Redness, b* = Yellowness. ^{a,b}Means with different superscripts along the same row are significantly different ($p < 0.01$), WHC = Water holding capacity

(up to 2.5 g/L) did not significantly affect live weight, while Japhet and Godgift [19] reported that VALE supplementation (50 mL/L) did not significantly affect live pullet weight, but increased live broiler weight. However, Daramola *et al.* [27] and Tokofai *et al.* [28] reported that VALE (100 mg/kg body weight) increased 42-day-old broiler live weight. In our study, live weight was not significantly different between groups because our observations were made during laying periods. After entering this period, quails do not experience many body weight changes. This result concurred with Korwin-Kossakowska *et al.* [29], who reported little change in female quail body weight during laying periods and from the same generation unless nutrients were used.

Vernonia amygdalina leaf extract (10 mL/L and 20 mL/L: T1 and T2, respectively) significantly

($p < 0.05$) increased carcass weight, but this was reduced at 30 mL/L (T3), while carcass and non-carcass percentages were not significantly different (Table-1). These observations concurred with the previous studies: Japhet and Godgift [19] reported that carcass percentages did not always have positive correlation with carcass weights. Daramola *et al.* [27] stated that *V. amygdalina* leaf powder (0.3%) significantly increased carcass weight. Rusli *et al.* [30] recorded that *V. amygdalina* leaf powder (2–6%) did not significantly affect carcass percentages. Mandey *et al.* [20] reported that *V. amygdalina* leaf juice (20 mL/L) did not significantly affect broiler carcass weight, while 30 mL/L supplementation significantly increased carcass weight but not carcass weight percentages. Abdel-Moneim *et al.* [8] stated that heat stress reduced intestinal function and protein digestibility. Ugokwe

and Ugokwe [31] reported that VALE significantly increased duodenal, jejunal, ileal villus height, and crypt depth. Thus nutrient absorption and growth indices in broilers were better, while higher doses (up to 100 mg/g) did not induce significant changes. In our study, VALE (10 mL/L and 20 mL/L in T1 and T2 groups, respectively) significantly increased carcass weight ($p < 0.05$), but 30 mL/L did not ($p > 0.05$). Thus, low VALE doses improved carcass weights, while 30 mL/L had no significant effects ($p > 0.05$); this was possibly attributed to reduced nutrient bio-availability from high anti-nutritional substances in extracts. Increased carcass weights without increased non-carcass percentages (Table-1) suggested VALE increased meat production.

Quail carcass moisture and meat protein levels were not significantly different ($p > 0.05$) (Table-2). *Vernonia amygdalina* leaf extract increased nutrient digestibility and absorption only to increase carcass weight, but was insufficient in increasing carcass meat protein levels. Quails were maintained in open cages at 23.9°C–33.5°C and humidity = 85–90%, so they were potentially stressed. Research data on stress reduction by VALE are often inconsistent. Oyesola *et al.* [14] stated that *V. amygdalina* contained tannin, alkaloid, phenol, and flavonoid antioxidants and were effective against oxidative stress. Antioxidant feed supplementation increased feed nutrient digestibility and absorption [32, 33]. Vargas-Sánchez *et al.* [10] reported that natural herb antioxidant actions mostly improved quail carcass quality without affecting carcass weight. In our study, VALE, as an antioxidant herb source, effectively increased carcass weight. The type, conformance, and concentration of active compounds determined its effects; too high concentrations decreased absorption and metabolism indices [10]. Only one study reported effective antioxidant supplementation on increasing protein deposition in carcasses [18].

Fat content was not significantly different ($p > 0.05$), but cholesterol levels were significantly decreased ($p < 0.05$) and were relatively lower when compared with Ilavarasan *et al.* [34], who showed that fat levels were 2.5–2.78%. Omede *et al.* [13], Praptiwi *et al.* [35], and Nowak *et al.* [11] reported that *V. amygdalina* contained high flavonoid levels, which reduced bird meat fat [34, 36–39]. Furthermore, Tan *et al.* [40] showed that flavonoids inhibited adipogenesis and increased lipolysis and apoptosis in adipose tissue cells, with reduced meat fat. In our study,

fat levels were not significantly different between groups.

Vernonia amygdalina leaf extract significantly ($p < 0.05$) affected meat cholesterol. At 10 mL/L VALE (T1), cholesterol levels were decreased and were reduced at 20 mL/L (T2), but T2 and T3 groups were not significantly different. *Vernonia amygdalina* leaf extract contains antioxidants that reduce bile acid secretion and cholesterol deposition in meat, in agreement with Suhaemi and Hidayati [21], who showed that *V. amygdalina* leaf powder (1–2%) reduced cholesterol levels in broiler meat. *Vernonia amygdalina* contains many antioxidants [41, 42] which inhibit HMG CoA reductase and lipase enzymes [43]. Reduced lipase activity is followed by reduced cholesterol synthesis [44, 45] and reduced meat cholesterol levels [34]. Vargas-Sánchez *et al.* [10] indicated that particular herb antinutritive and/or active compound (e.g., tannin) at high concentrations reduced carcass quality. Cholesterol levels in T2 and T3 groups were non-significantly different and showed that high VALE values did not reduce meat cholesterol levels.

Vernonia amygdalina leaf extract at 10–30 mL/L concentrations did not significantly affect ($p > 0.05$) physical carcass parameters, except WHC (Table-3). We hypothesized that bioactive *Vernonia amygdalina* leaf extract improved carcass production (Table-1), improved chemical (Table-2) and physical meat qualities, especially WHC (Table-3). Improved meat quality is related to increased antioxidant levels and activity [46], thus, our observations may reflect increased antioxidant levels. Victoria *et al.* [46] showed that 7.5% African leaf meal significantly increased physical properties in broiler meat, that is, pH, WHC, and drip loss, but meat color was not significantly changed. These observations concurred with Hosseindoust *et al.* [47], who showed that natural antioxidant addition did not affect other meat quality parameters, including meat color.

To determine optimum VALE doses, we used polynomial orthogonal analysis (Table-4), which showed strong R^2 values for cholesterol (0.720) and WHC (0.701), but weak values for carcass weight (0.066). Based on the equation that was formed, the optimum VALE dose to produce low meat cholesterol levels was 28.821 mL/L VALE in drinking water that was predicted to reach an optimized minimum cholesterol level of 161.992 g/100 g. Optimization of quail meat WHC values obtained at VALE dose of 9.141 mL/L.

Table-4: Parameters, equation, R^2 value, VALE dosage, and predicted observed parameter optimized value as polynomial orthogonal analysis results.

Parameter	Equation	R^2	Predicted VALE dosage	Optimized value of the observed parameter
Cholesterol (mg/100 g)	$Y = 0.162x^2 - 9.3381x + 296.56$	0.720	28.821	161.992
Carcass weight (g)	$Y = -0.003x^2 + 0.5804x + 95.279$	0.066	96.730	123.351
WHC (%)	$Y = 0.0191x^2 - 0.3492x + 23.6$	0.701	9.141	22.004

WHC=Water holding capacity, VALE=*Vernonia amygdalina* Del. leaf extract

Conclusion

Vernonia amygdalina leaf extract (20 mL/L) improved quail carcass traits, especially cholesterol levels in quail meat and carcass weight. The optimum efficacy of VALE in reducing quail meat cholesterol might be achieved with a higher dosage, however, it can lower carcass weight.

Authors' Contributions

SK and TAS: Prepared the materials, planned and designed the study, recorded and analyzed the data, and drafted and revised the manuscript. LDM and DSP: Prepared the materials, conducted the study, and drafted and revised the manuscript. All authors have read, reviewed, and approved the final manuscript.

Acknowledgments

The authors are thankful to Edjeng Suprijatna, Rina Muryani, Arif Maulana, Nadira Rizqi Davika, Hedy Suryatama, Maulana Syamsudin, Maulana Umar, Fuad, Anita Damayanti, Ela Marin and Lulu Hanisa for help during data collection. The study was funded by Diponegoro University, Indonesia (Grant number 36/UN7.5.5/PP/2021).

Competing Interests

The authors declare that they have no competing interests.

Publisher's Note

Veterinary World remains neutral with regard to jurisdictional claims in published institutional affiliation.

References

- Pavelková, A., Haščík, P., Kalařová, A., Capcarová, M., Čuboň, J., Bučko, O., Kačániová, M., Hanusová, E., Tkáčová, J. and Bobko, M. (2020) Chemical composition of muscle after bee bread application in the nutrition of Japanese Quails. *J. Microbiol. Biotech. Food Sci.*, 9(4): 831–835.
- Quaresma, M.A.G., Antunes, I.C., Ferreira, B.G., Parada, A., Elias, A., Barros, M., Santos, C., Partidário, A., Mourato, M. and Roseiro, L.C. (2022) The composition of the lipid, protein and mineral fractions of quail breast meat obtained from wild and farmed specimens of Common quail (*Coturnix coturnix*) and farmed Japanese quail (*Coturnix japonica domestica*). *Poult. Sci.*, 101(1): 101505.
- Leishman, E.M., Ellis, J., van Staaveren, N., Barbut, S., Vanderhout, R.J., Osborne, V.R., Wood, B.J., Harlander-Matauschek, A. and Baes, C.F. (2021) Meta-analysis to predict the effects of temperature stress on meat quality of poultry. *Poult. Sci.*, 100(11): 101471.
- Zaboli, G., Xi, H., Xi, F. and Ahn, D.U. (2019) How can heat stress affect chicken meat quality? A review. *Poult. Sci.*, 98(3): 1551–1556.
- Nawaz, A.H., Amoah, K., Leng, Q.Y., Zheng, J.H., Zhang, W.L. and Zhang, L. (2021) Poultry response to heat stress: Its physiological, metabolic, and genetic implications on meat production and quality including strategies to improve broiler production in a warming world. *Front. Vet. Sci.*, 8: 699081.
- Nasr, M.A.F., Mohammed, H., Hassan, R.A., Swelum, A.A. and Saadeldin, I.M. (2019) Does light intensity affect the behavior, welfare, performance, meat quality, amino acid profile, and egg quality of Japanese quails. *Poult. Sci.*, 98(8): 3093–3102.
- Batool, F., Bilal, R.M., Hassan, F.U., Nasir, T.A., Rafique, M., Elnesr, S.S., Farag, M.R., Mahgoub, H.A.M., Naiel, M.A.E. and Alagawany, M. (2021) An updated review on behavior of domestic quail with reference to the negative effect of heat stress. *Anim. Biotechnol.*, 1–14.
- Abdel-Moneim, A.M.E., Shehata, A.M., Khidr, R.E., Paswan, V.K., Ibrahim, N.S., El-Ghoul, A.A., Aldhumri, S.A., Gabr, S.A., Mesalam, N.M., Elbaz, A.M., Elsayed, M.A., Wakwak, M.M. and Ebeid, T.A. (2021) Nutritional manipulation to combat heat stress in poultry—a comprehensive review. *J. Therm. Biol.*, 98: 102915.
- Hosseindoust, A., Kang, H.K. and Kim, J.S. (2022) Quantifying heat stress; The roles on metabolic status and intestinal integrity in poultry, a review. *Domest. Anim. Endocrinol.*, 81: 106745.
- Vargas-Sánchez, R.D., Ibarra-Arias, F.J., del Mar Torres-Martínez, B., Sánchez-Escalante, A. and Torrescano-Urrutia, G.R. (2019) Use of natural ingredients in Japanese quail diet and their effect on carcass and meat quality—a review. *Asian-Aust. J. Anim. Sci.*, 32(11): 1641–1656.
- Nowak, J., Kiss, A.K., Wambebe, C., Katuura, E. and Kuźma, Ł. (2022) Phytochemical analysis of polyphenols in leaf extract from *Vernonia amygdalina* Delile plant growing in Uganda. *Appl. Sci.*, 12(2): 912.
- Abdel-Moneim, A.M.E., Shehata, A.M., Alzahrani, S.O., Shafi, M.E., Mesalam, N.M., Taha, A.E., Swelum, A.A., Arif, M., Fayyaz, M. and Abd El-Hack, M.E. (2020) The role of polyphenols in poultry nutrition. *J. Anim. Physiol. Anim. Nutr. (Berl.)*, 104(6): 1851–1866.
- Omede, A., Suleiman, M.S., Atanu, F.O., Momoh, S., Friday, E.T., Sheneni, V.D. and Jegede, E.R. (2018) Evaluation of antioxidant and cytotoxic properties of *Vernonia amygdalina*. *Int. J. Cell Sci. Mol. Biol.*, 4(4): 555644.
- Oyesola, O.A., Sampson, I.I., Augustine, A.A., Adejoke, O.B. and Taiwo, G.E. (2022) Comparison of phytochemical constituents of ethanol leaf extracts of *Solanum macrocarpon* and *Vernonia amygdalina*. *Asian J. Nat. Prod. Biochem.*, 20(1): 6–10.
- Priambodo, T.R., Sholikin, M.M., Qomariyah, N., Jayanegara, A., Batubara, I., Utomo, D.B. and Nahrowi, N. (2021) Effects of dietary flavonoids on performance, blood constituents, carcass composition and small intestinal morphology of broilers: A meta-analysis. *Anim. Biosci.*, 34(3): 434–442.
- Sarica, S., Islim, P. and Tugay, A. (2019) Supplementation of natural antioxidants to reduced crude protein diets for Japanese quails exposed to heat stress. *Braz. J. Poult. Sci.*, 21(1): 1–14.
- Manassis, G., Kalogianni, A.I., Lazou, T., Moschovas, M., Bossis, I. and Gelasakis, A.I. (2020) Plant-derived natural antioxidants in meat and meat products. *Antioxidants (Basel)*, 9(12): 1215.
- Saracila, M., Panaite, T.D., Mironeasa, S. and Untea, A.E. (2021) Dietary supplementation of some antioxidants as attenuators of heat stress on chicken meat characteristics. *Agriculture*, 11(7): 638.
- Japhet, Y.B. and Godgift, T. (2021) Effect of *Vernonia amygdalina* leaf extract on growth performance, carcass characteristics of pullet and broiler birds. *Int. J. Sci. Res. Arch.*, 3(02): 209–213.
- Mandey, J.S., Sompie, M. and Wolayan, F.R. (2021) Growth Assessment of Broiler Chickens Given Bitter Leaves (*Vernonia amygdalina*) as Phyto-Additive, Potentially Antimicrobial Agents of Lipids and Amino Acids. In: Proceedings of the 9th International Seminar on Tropical Animal Production (ISTAP 2021). p21–25.
- Suhaemi, Z. and Hidayati, S.G. (2020) Improvement of the Quality of Duck's and Chicken's Meat using African Leaf

- (*Vernonia amygdalina*). In: Proceedings of the International Seminar on Promoting Local Resources for Sustainable Agriculture and Development (ISPLRSAD 2020). p163–167.
22. Hasbullah, I., Wulandari, Z. and Suci, D.M. (2020) African leaf (*Vernonia amygdalina*) juice supplement in drinking water on the chemical properties of quail egg (*Coturnix coturnix japonica*). *J. Ilmu Nutr. Teknol. Pakan.*, 18(2): 43–48.
 23. Kaneria, M., Kanani, B. and Chanda, S. (2012) Assessment of effect of hydroalcoholic and decoction methods on extraction of antioxidants from selected Indian medicinal plants. *Asian Pac. J. Trop. Biomed.*, 2(3): 195–202.
 24. Rasid, R.A., Baba, A.R. Yaakub, N.M. and Milan, A.R. (2019) Performance and carcass characteristics of broiler chickens fed various components of Candlenut Kernel. *Trop. Anim. Sci. J.*, 42(3): 203–208.
 25. Association of Official Analytical Chemists. (2007) Official Methods of Analysis of AOAC International 18th ed., 2nd Revised. AOAC International, Maryland.
 26. Osho, I.B., Akindahunsi, A., Igban, F.A. and Adekunle, D.J. (2014) Effect of orally administered bitter leaf (*Vernonia amygdalina*) extract on the growth performance and haematological parameters of broiler chicken. *J. Vet. Med. Anim. Health*, 6(10): 251–256.
 27. Daramola, O.T., Oloruntola, O.D. and Ayodele, S.O. (2018) Nutritional evaluation of bitter leaf meal (*Vernonia amygdalina*): Effects on performance, carcass and serum metabolites of broiler chickens. *Niger. J. Anim. Sci.*, 20(2): 211–221.
 28. Tokofai, B.M., Idoh, K., Oke, O.E. and Agbonon, A. (2021) Hepatoprotective effects of *Vernonia amygdalina* (*Asteraceae*) extract on CCl₄-induced liver injury in broiler chickens. *Animals (Basel)*, 11(12): 3371.
 29. Korwin-Kossakowska, A., Sartowska, K., Linkiewicz, A., Tomczyk, G., Prusak, B. and Sender, G. (2013) Evaluation of the effect of genetically modified RR soya bean and MON 810 maize in the diet of Japanese quail on chosen aspects of their productivity and retention of transgenic DNA in tissues. *Arch. Tierzucht*, 56(1): 60.
 30. Rusli, B., Astuty, A.F., Syamsuddin, Natsir, S. and Muh, M.L. (2022) Percentage of carcass parts of broiler chicken given additional African leaf flour feed (*Vernonia amygdalina*). *Russ. J. Agric. Socioecon. Sci.*, 124(4): 54–61.
 31. Ugokwe, N.I. and Ugokwe, C. (2022) Comparative study on the prebiotic effect of Inulin and aqueous extract of *Vernonia amygdalina* on growth and intestinal morphometry of broiler chicken (Cobb strain). *J. Anim. Sci. Vet Med.*, 7(1): 17–27.
 32. Shilov, V., Khakimova, G., Semina, O., Akhmadullin, R. and Akhmadullina, A. (2020) Effect of the antioxidant "Bisphenol-5" on the digestibility of nutrients in the diet of broilers. *BIO Web Conf.*, 17: 00135.
 33. Abbasi, M.A., Ghazanfari, S., Sharifi, S.D. and Gavligi, A.H. (2020) Influence of dietary plant fats and antioxidant supplementations on performance, apparent metabolizable energy and protein digestibility, lipid oxidation and fatty acid composition of meat in broiler chicken. *Vet. Med. Sci.*, 6(1): 54–68.
 34. Ilavarasan, R., Abraham, R.J.J. and Rao, V.A. (2016) The relationship between meat quality characteristics and nutritional composition of Nandanam quail-III slaughtered at different ages. *J. Anim. Res.* 6(2): 95–100.
 35. Praptiwi, Fathoni, A. and Ilyas, M. (2020) Diversity of endophytic fungi from *Vernonia amygdalina*, their phenolic and flavonoid contents and bioactivities. *Biodiversitas*, 21(2): 436–441.
 36. Radaelli, G., Piccirillo, A., Birolo, M., Bertotto, D., Gratta, F., Ballarin, C., Vascellari, M., Xiccato, G. and Trocino, A. (2017) Effect of age on the occurrence of muscle fiber degeneration associated with myopathies in broiler chickens submitted to feed restriction. *Poult. Sci.*, 96(2): 309–319.
 37. Kamboh, A.A., Leghari, R.A., Khan, M.A., Kaka, U., Naseer, M., Sazili, A.Q. and Malhi, K.K. (2019) Flavonoids supplementation-an ideal approach to improve quality of poultry products. *Worlds Poult. Sci. J.*, 75(1): 115–126.
 38. Nath, S. and Aravindkumar, K. (2021) Role of flavonoids in poultry nutrition. *Acta Sci. Vet. Sci.*, 3(12): 88–91.
 39. Nufer, A.I. and Shatskikh, E.V. (2021) The effect of herbal supplements on development of internal organs and chemical composition of broilers muscles. *Theory Pract. Meat Process.*, 6(2): 141–150.
 40. Tan, Z., Halter, B., Liu, D., Gilbert, E.R. and Cline, M. A. (2022) Dietary flavonoid ass modulators of lipid metabolism in poultry. *Front Physiol.*, 13: 863860
 41. Raimi, C.O., Oyelade, A.R. and Adesola, O.R. (2020) Phytochemical screening and *in-vitro* antioxidant activity on *Vernonia amygdalina* (Ewuro-Bitter leaf). *Eur. J. Agric. Forest Res.*, 8(2): 12–17.
 42. Yazid, F., Hasanah, N.B., Rosmalena, Hanafi, M. and Prasasty, V.D. (2020) Antidiabetic and antioxidant potential of *Vernonia amygdalina* leaf extract in alloxan-induced Sprague-Dawley rat. *J. Biol. Sci.*, 20(4): 190–200.
 43. Yunarto, N., Aini, N., Oktoberia, I.S., Sulistyowati, I. and Kurniatri, A.A. (2019) Antioxidant activity along with inhibition of HMG CoA reductase and lipase from *Anredera cordifolia* leaf-*Curcuma xanthorrhiza* rhizome combination. *J. Kefarmasian Indones.*, 9(2): 89–96.
 44. Shoaib, M., Bhatti, S.A., Nawaz, H. and Saif-Ur-Rehman, M. (2021) Effect of lipase and bile acids on growth performance, nutrient digestibility, and meat quality in broilers on energy-diluted diets. *Turk. J. Vet. Anim. Sci.*, 45: 148–157.
 45. Ge, X.K., Wang, A.A., Ying, Z.X., Zhang, L.G., Su, W.P., Cheng, K., Feng, C.C., Zhou, Y. Zhang, L.L. and Wang, T. (2019) Effects of diets with different energy and bile acids levels on growth performance and lipid metabolism in broilers. *Poult. Sci.*, 98(2): 887–895.
 46. Victoria, A.O., Okpara, J.O., Oshibanjo, Debola, O. and Salihu, A. (2022) Inclusion levels of bitter (*Vernonia amygdalina*) leaf meal on meat and sensory qualities of broiler chickens. *Asian J. Res. Anim. Vet. Sci.*, 9(4): 26–35.
 47. Hosseindoust, A., Oh, S.M., Ko, H.S., Jeon, S.M., Ha, S.H., Jang, A., Son, J.S., Kim, G.Y., Kang, H.K. and Kim, J.S. (2020) Muscle antioxidant activity and meat quality are altered by supplementation of astaxanthin in broilers exposed to high temperature. *Antioxidants (Basel)*, 9(11): 1023.

African leaf (*Vernonia amygdalina*) extracts improve Japanese quail (*Coturnix coturnix japonica*) carcass traits

ORIGINALITY REPORT

18%	15%	15%	8%
SIMILARITY INDEX	INTERNET SOURCES	PUBLICATIONS	STUDENT PAPERS

PRIMARY SOURCES

1	e-sciencecentral.org Internet Source	3%
2	www.ncbi.nlm.nih.gov Internet Source	2%
3	Submitted to Universitas Airlangga Student Paper	2%
4	usefil.eu Internet Source	1%
5	Slamet Hartanto, Agung Budiyanto, Rini Widayanti, Erif Maha Nugraha Setyawan, Imawan Daru Prasetya. "Characterization of polymorphisms in the follicle-stimulating hormone receptor and insulin-like growth factor-1 genes and their association with fertility traits in Jawa-Brebes cows", Veterinary World, 2023 Publication	1%
6	Submitted to SDM Universitas Gadjah Mada Student Paper	1%

7	www.mdpi.com Internet Source	1 %
8	Yusuf Ridwan, Etih Sudarnika, Tri Isyani Tungga Dewi, Novericko Ginger Budiono. "Gastrointestinal helminth parasites of pets: Retrospective study at the veterinary teaching hospital, IPB University, Bogor, Indonesia", Veterinary World, 2023 Publication	1 %
9	Submitted to Universitas Diponegoro Student Paper	1 %
10	Wayan Masa Tenaya, Ida Bagus Ngurah Swacita, Ketut Wirata, Made Damriyasa et al. "A study of African swine fever virus in Regional VI of the Disease Investigation Center of Denpasar Bali in Indonesia", Veterinary World, 2023 Publication	1 %
11	www.animbiosci.org Internet Source	<1 %
12	koreascience.kr Internet Source	<1 %
13	vet.kku.ac.th Internet Source	<1 %
14	Mohamed M. Alghirani, Eric Lim Teik Chung, Nafeesa Abu Kassim, Yee Lyn Ong et al.	<1 %

"Effect of Brachiaria decumbens as a novel supplementation on the production performance of broiler chickens", Tropical Animal Health and Production, 2022

Publication

15

Yuli Purwandari Kristianingrum, Tri Untari, Asmarani Kusumawati. "Severe acute respiratory syndrome coronavirus-2 detection in domestic animals as a reservoir for the virus transmission to humans in Yogyakarta, Indonesia", Veterinary World, 2023

Publication

<1 %

16

academic.oup.com

Internet Source

<1 %

17

talenta.usu.ac.id

Internet Source

<1 %

18

Submitted to Melbourne Institute of Business and Technology

Student Paper

<1 %

19

univendspace.univen.ac.za

Internet Source

<1 %

20

etheses.bham.ac.uk

Internet Source

<1 %

21

Belal S. Obeidat. "Effect of feeding pomegranate seed pulp on Awassi lambs' nutrient digestibility, growth performance, and carcass quality", Veterinary World, 2023

<1 %

22

Emmanuel U. Ejiofor, Samuel O. Onoja, Ernest C. Agwamba, Hitler Louis et al.

<1 %

"Computational, chemical profile, in vitro antioxidant, hypoglycaemic, and anti-inflammatory activity of hexane extract of some selected dark green vegetables", Proceedings of the Indian National Science Academy, 2023

Publication

23

Fernando, Sharma M., and Patricia A. Murphy. "HPLC determination of thiamin and riboflavin in soybeans and tofu", Journal of Agricultural and Food Chemistry, 1990.

<1 %

Publication

24

Parviz Hassanzadeh, Mohammad Ahmadvand, Shadi Aslani, Najmeh Sheikhzadeh et al. "Dietary astaxanthin mitigated paraquat - induced oxidative stress in rainbow trout () fillet ", Aquaculture Research, 2022

<1 %

Publication

25

Yong Ju Kim, Min Ho Song, Ji Hwan Lee, Han Jin Oh et al. "Partial Replacement of Animal Fat with Full-Fat Almond in Broiler Chicken Diets: Performance, Nutrient Digestibility, Blood Profile, Cecal-Fecal Microflora

<1 %

Composition, and Foot-Pad Dermatitis", Animals, 2021

Publication

26

plantarchives.org

Internet Source

<1 %

27

bitok.datastore.pt

Internet Source

<1 %

28

eprints.undip.ac.id

Internet Source

<1 %

29

www.ajol.info

Internet Source

<1 %

30

www.european-poultry-science.com

Internet Source

<1 %

31

www.onehealthjournal.org

Internet Source

<1 %

32

Sugiharto Sugiharto, Turrini Yudiarti, Isroli Isroli, Endang Widiastuti, Fatan Dwi Putra. "Effect of dietary supplementation with *Rhizopus oryzae* or *Chrysonilia crassa* on growth performance, blood profile, intestinal microbial population, and carcass traits in broilers exposed to heat stress", Archives Animal Breeding, 2017

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On