Intake of Sugar-Sweetened Beverage and Metabolic Syndrome Components in Adolescents.

by Fillah Fithra Dieny

Submission date: 02-Sep-2020 01:49PM (UTC+0700) Submission ID: 1378097789 File name: ed_Beverage_and_Metabolic_Syndrome_Components_in_Adolescents.pdf (2.21M) Word count: 4486 Character count: 24101



Intake of Sugar-Sweetened Beverage and Metabolic Syndrome Components in Adolescents

Evi <mark>13 r</mark>tini

Department of Nutrition Science Faculty of Medicine, Diponegoro University Semarang, Indonesia evikartini 10@gmail.com Fillah Fithra Dieny Department of Nutrition Science Faculty of Medicine, Diponegoro University Semarang, Indonesia fillahdieny@gmail.com Etisa Adi Murbawanni Department of Nutrition Science Faculty of Medicine, Diponegoro University Semarang, Indonesia etisatitis@gmail.com A.Fahmy Arif Tsani Department of Nutrition Science Faculty of Medicine, Diponegoro University Semarang, Indonesia fahmi_tsani@yahoo.com

1

Abstract-Adolescents with metabolic syndrome have a higher risk for cardiovascular diseases and diabetes. Risk factors of metabolic syndrome include central obesity, hypertriglycerid, and hyperglycemia. High intake (>50 g/day) of sugar-sweetened beverages (SSBs) is known to be positively associated with increased waist circumference, triglyceride and fasting blood glucose levels. This study aimed to analyze the relationship between SSBs intake with waist circumference, triglyceride and fasting blood glucose levels in adolescents. This study was a cross-sectional design with fifty-nine subjects of adolescents aged 15-18 years old. They were performed by simple random sampling technique. SSBs and energy intake were assessed with Semi Quantitative Food Frequency Questionnaire, while waist circumference used a tape measure. Blood glucose levels were measured with Glucose Oxidation method, triglyceride levels using colorimetric enzymatic method (GPO-PAP). Data were analyzed by Person Product Moment and Rank Spearman correlation test. High triglyceride levels were observed in 62.7% subjects and total of 44.11% subjects had large waist circumference. Excessive intake of SSB was found in 72.9% subjects. Intake of SSBs was correlated with waist circumference (p=0.020) and triglyceride levels (p=0.044), but not fasting blood glucose levels (p=0.060). Consumption of SSBs > 50 g/day can increased waist circumference and triglyceride levels that were component of metabolic syndrome.

Keywords— Sugar-Sweetened Beverage, Metabolic Syndrome, Adolescents

I. INTRODUCTION

Metabolic syndrome is a metabolic disorder which increase cardiovascular diseases and diabetes mellitus. The components of metabolic syndrome are central obesity, the increase of triglyceride level and fasting blood glucose, HDL level decrease and also high blood pressure [1]. The increase of metabolic syndrome prevalence on adolescents in line with the increase of obesity prevalence as one or the risk of metabolic syndrome [2]. Based on study result in Semarang city in 2015 showed that the prevalence of metabolic syndrome in obese adolescents as 68.4% [3]. The research conducted in senior high school students in 2016 showed that from 66 obese adolescents, there were 47.5% has metabolic syndrome [4].

Obesity is the main trigger of metabolic syndrome, because excess calorie intake from carb 15 drate, fat, and protein more than total energy needed [5]. The excess energy will be convert into fat and stored in adipose tissue which will increase waist circumference [6]. Adolescents with central obesity has waist circumference for more than 80th percentile. Obesity can trigger the increase of triglyceride by excess fat storing. Triglyceride accumulation in hepar can cause insulin resistance which will increase fasting blood glucose [3].

Consumption SSBs increases on adolescents because it is affected by several factors, those are: advertisements broadcasted in media, peer group influences, and distribution of SSBs in accessible cafeteria or supermarket [7], [8]. Sugarsweetened beverage is a 2everage of which has been added calorie sweetener into it, such as soft drink, sport drink, fruit drink, energized drink, tea and coffee, milk, fruit juice, and isotonic beverage [9]. Based on the study of cohort conducted on adolescents showed that the consumption of SSBs for 375 ml which contain sugar approximately 37-54 g contribute energy as 150 calories which related to the increase of metabolic syndrome risk [10], [11]. The additional sugar content in SSBs contributes energy as 21.2% on adolescents which can increase obesity Excess energy in the body will be stored in a form of triglyceride in depose tissue which lead to obesity if it is occur continuously [12], [13]. The study on female adolescents aged 12-19 showed the consumption of SSBs 250g/day will lead to the increase of triglyceride as 2,3mg/dL [14]. Another study stated that the consumption of SSBs > 1 portion/ day can increase the prevalence of diabetes mellitus type 2 as 26% [15]. Based on the background the author interested to analyze the correlation of SSBs intake with waist circumference, triglyceride and fasting blood glucose on adolescents.

II. MATERIALS AND METHODS

This study used a cross sectional design with data collecting on July until August 2017. This study was



Advances in Health Science Research, volume 12

2

conducted on students in SMA Negeri 15 semarang. The selection of 59 subjects was performed by simple random sampling technique. Data on age, subjects identity, antropometri, sugar sweetenec everage (SSBs) intake, energy intake, waist circumference, triglyceride levels, and fasting blood glucose levels.

Waist circumference was the circle line of a waist right on navel measured through the middle of lowest interior edge with crista iliaca. The category of central obese waist circumference if the waist circumference was more than 80 cm and normal if it less than 80. Fasting blood glucose levels (GDP) are a parameter that describes the glucose concentration in the blood plasma measured in subjects who were fasting for 8-12 hours. GDP levels were measured using spectophotometry by the Glucose Oxidation (GOD) method. GDP levels are considered to be high if it is $\geq 100 \text{ mg} / \text{dL}$ and normal if it is ≤100 mg / dL. Triglyceride was one of the fats in the bloods 14 am that come from fat intake and were formed in the liver. Triglyceride levels were measured by enzymatic colorimetric using glycerol phosphate oxidase (GPO). Triglyceride levels was considered high if it was ≥110 mg /dL and normal if it is <110 mg/dL [16].

Sugar-sweetened beverage intake was the average of sugar from packaged and non-packaged beverages. Packaged beverages are processed drinks containing additional sugars recorded on the nutritional fact label on the packaging, whereas non-packaged beverages are sugar-added drinks during brewing and are not recorded on the nutritional fact label on the packaging. Sugar was calculated from the sugar content contained on the package and added sugar during brewing. Measurement of SSBs intake was obtained by semi quantitative consumption frequency method with frequency question of consumption in a day, week, month, and year and the amount of each time consumption was recorded in household size (URT) and converted to gram. The intake of SSBs was classified as high 2 f it contains sugar ≥ 50 g / day and as normal if it was<50 g / day [17]. The energy of SSBs intake was categorized to be high if it was $\geq 10\%$ of total energy intake and as a normal if i 2 vas<10% of total energy intake [18]. Data of energy intake obtained by interview using Semi Quantitative Food Frequency Questionnaire (SQ-FFQ). Energy requirements were calculated based on individual needs and then the energy intake was grouped into the adequacy level based on the National Widyakarya Food and Nutrition (WNPG) 2012. The intake of ≥120% of individual needs was call gorized as high, 90-119% is sufficient, and <90% is less. Data were analyzed by Person Product Moment and Rank Spearman correlation test. Data analysis using SPSS with 95% confidence degree ($\alpha = 0.05$).

III. RESULTS AND DISCUSSIONS

The subjects aged 15-18 years old including 35 (60⁽¹³⁾ were female subject and 24 (40%) male subject. The results in Table 1 show that the average age of the subjects was 16.2 years old and there were subjects with z-score of body mass index for age of 4.9 SD who fall into the obesity category.

Table 1: Characteristic of Subject							
Variables Minimum Maximum Mean+DS							
Age (years)	15	18	16,2±0,7				
Height (cm)	145,0	183,0	160,2±7,4				
Weight (kg)	35,9	141,0	63,2±19,1				
BMI/age (SD)	-2,7	4,9	0,7±1,5				
Intake of SSBs (g/day)	11,8	236,2	92,5±60,8				
SSBs packaging (g/day)	4,6	198,2	62,1±46,3				
SSBs non-packaging (g/day)	1,7	89,6	30,3±25,2				
Energy Intake of SSBs (kcal/day)	72,4	1516,0	520,5±328,3				
Energy Intake of SSBs (% Energy)	3	41,2	19,2±8,7				
Energy Intake (kcal/day)	1171,9	6327,7	2707,2±1037,9				
Waist circumference (cm)	56,5	114,7	79,9±12,6				
Triglyceride (mg/dL)	96,0	150,0	113,9±10,3				
Fasting Blood Glucose Levels (mg/dL)	70,0	98,0	81,5±7,6				

The intake of sugar-sweetened beverage subjects ranged from 11.8-236.2 g/day with an average SSBs intake greater than non-packaged beverage, i.e. 62.1 g/day and 30.3 g/day. While the mean SSBs intake of 92.5 g/day, which means the consumption of SSBs in the subject tend 20 be high. The mean triglyceride and fasting blood glucose levels were 113.9 mg/dL and 81.5 mg/dL, respectively, high triglycerides and normal at fasting blood glucose levels.

Table 2 shows that 72,9% subjects were high intake of SSB, but subjects with sufficient energy intakes were only 28.8%, but 91.5% of subjects took energy from a high 5 garsweetened beverage (> 10% of total energy). The high intake of sugar-sweetened beverage is in line with energy intake in the subject, where found in 49.2% of subjects who have higher energy intake. The average total SSBs intake was \pm 19.2% of total energy intake.

Table 2. Characteristics of Sugar-Sweetened Beverage and Energy Intakes

Cotogomy of SSRs and	S	Total	
energy inatake	Male n (%)	Female n (%)	n (%)
Sugar Sweetened			
Beverage (g)	6 (10,2)	10 (16,9)	16(27,1)
Sufficient	18 (30,5)	25 (42,4)	43(72,9)
High			
Energy intake from			
SSBs (%)	1 (1,7)	4 (6,8)	5 (8,5)
Sufficient	23 (39,0)	31 (52,5)	54 (91,5)
High			
Energy intake (%)			
Less	7(119)	6(10,2)	13(22,1)
Sufficient	8 (13,5)	9 (15,3)	17(28,8)
High	9 (15,3)	20 (33,9)	29(49,2)
		11	

This result is greater than the results of a survey conducted 12 Centers for Disease Control and Prevention (CDC) in children and adolescents in the United States in 2005-2008, the average intake of sugar from foods and beverages by 16% of total energy intake [19]. The study conducted on adolescents aged 12-14 years old in 2014 stated that the average sugar consumption in beverages sweetened was 60.43 g (19.04% of total energy) [20]. In fact, the suggestion of 50-60% of energy needs obtained from sources of carbohydrates, both complex and simple. However, according to Tumpeng Gizi Seimbang (TGS) Indonesia, the greatest fulfillment of needs is derived from complex carbohydrate sources, such as whole grains, cereals, various



tubers and their processed products. Whereas simple carbohydrate fulfillment is limited for no more than 50 g/day or <10% of total energy intake [17] [18] [21].

Sugar-sweetened beverage consumption that tends to be high in adolescents is affected by several factors, such as advertisements that are widely broadcasted in the media, the distribution of SSBs in cafeterias or accessible supermarkets, as well as the influenced by peers. The choice of baverage and eating habits on adolescents is influenced by peer because it is considered as the solidarity of the peers [7] [8]. The most commonly consumed type of SSBs was the type of package beverage, which is tea, milk drinks, fruit drinks, and soft drinks which contribute an average of sugar 20-46 g/package as well as non-packaged beverages such as tea and various kinds of ice that has average sugar content of 22-26 g/serving or equivalent to ± 2 tbsp sugar.

Table 3 shows that 14 subjects were classified as obese based on body mass index for age, meanwhile 44.1 % subjects abdominal obesity, 62.7% of subjects had were 19 pertriglyceride and 44.1% subjects had central obesity. The prevalence of obesity in this study was higher than that of the study in 2016, whi 27 was 12.8% [4]. The increased of obesity, especially central obesity is closely related to the 28 dence of metabolic syndrome. Adolescents who fulfill two components of the metabolic syndrome include pre-metabolic syndrome, most of which already occur in the subject, including high waist circumference and high 24 lyceride levels. Adolescents with pre-metabolic syndrome are at greater risk for becoming metabolic syndrome than with normal adolescents [22]. Subjects classified as central obese (≥80th percentile) was 44.1%, with female subjects as (28.8%) was greater than female (15.3%).

Table 3. Nutritional	status, and	i component	of metal	bolic syr	adrome in	subjects
----------------------	-------------	-------------	----------	-----------	-----------	----------

	S	Total	
Metabolic syndrome components	Male	Female	n (%)
	n (%)	n (%)	n (%)
Nutritional status			
Thin	-	2 (3,4)	2(3,4)
Normal	13 (22,0)	21 (35,6)	34 (57,6)
Overweight	3(5,1)	6 (10,2)	9 (15,3)
Obese	8 (13,5)	6 (10,2)	14(23,7)
Waist circumference (cm)			
Normal (< persentil 80)	15(25,4)	18 (30,5)	33 (55,9)
Abdominal obesity (≥persentil 80)	9 (15,3)	17 (28,8)	26(44,1)
Fasting Blood Glucose Levels			
(mg/dL)	24 (40,7)	35 (59,3)	59 (100)
Normal (<100 mg/dL)			
Triglyceride Levels (mg/dL)			
Normal (<110 mg/dL)	8 (13,6)	14 (23,7)	22 (37,3)
High ($\geq 110 \text{ mg/dL}$)	16(27,1)	21 (35,6)	37 (62,7)

This result is in accordance with the study of 13-15 years old adolescents in Semarang which stated that the incidence of central obesity in female is greater than male that were 52.6% and 38.6% [3]. This is because female body fat composition is greater than male, ranging from 21-35% as well as women's physical activity tend to be lower than men, so that the excess energy consumed more easily converted into fat and accumulate faster [23].

Central obesity in individuals due to body fat stores and intra-abdominal fat is closely related to metabolic abnormalities and cardiovascular disease [24]. Adipose tissue is a tissue that plays an active role in the release of free fat and pro and anti-inflammatory cytokines, so that obese individuals tend to experience cholesterol homeostasis disorders, which one of which was elevated triglyceride levels [25]. In this study there were subjects with hypertriglyceride of 62.7%. This number is higher than the research in 2016, which is 37.5% [4]. The increase of components of the metabolic syndrome, one of which is obesity in adolescence occurs due to many factors, one of which is excessive food intake and low physical activity.

The Correlation of SSBs and energy intake with Metabolic Syndrome Components

Table 4 shower that there was significant correlation between SSBs intake with waist circumference (p = 0.015) and triglyceride levels (p = 0.044), as well as energy intake (p = 0.014) with triglyceride levels on adolescents. But there was no significant correlation of SSBs intake with fasting blood glucose levels in adolescents (p = 0.06).

Table 4. Sugar-Sweetened B	everage Intake with Waist circumference
Triglyceride Levels,	and Fasting Blood Glucose Levels

Variabel	Waist circumf	erence	Triglyce Levels	eride	Fasting B Glucose I	lood .evels
	r	р	r	р	r	р
Intake of						
Sugar Sweetened	0,302	0,020 ^a	0,264	0,044 ^a	0,246	0,060ª
Beverage (SSBs) (g)						
Energy (kcal)	0,208	0,113 ^b	0,318	0,014 ^b	0,043	0,745 ^b
^a Rank-Spearman	^b Pearso	n correlatio	n			

This result same with study conducted on a 26 scents aged 12-16 in Taiwan who stated that SSBs intake associated with an increased component of the metabolic syndrome [14].

Sugar-sweetened beverage can increase waist circumference because the excess energy from SSBswill be stored in a form of fat inside of body ar 23 hen it was stored as energy storage in diposa tissue. After that the energy intake from sugar-sweetened beverage is known as liquid form which does not give the effect of full in stomach compare to solid food, therefore will increase excess energy intake. In addition, the fructose content in Sugar-sweetened beverage does not stimulate leptin, hence people who consume high fructose will complain to be hungry faster and eat excessively which will increase the fat accumulation intra abdominal over weight [26] [27]. This is in line with the study conducted to children aged 3-11 years old in England in 2004 and adolescents 12-18 years 17 in Taiwan in 2014 that high intake of SSBs can increased Body mass Index (BMI) and waist circumference [14] [28]. The increase of triglyceride levels may occur due to the fructose content present in SSBs. As much as 60% of the carbohydrates used as sweeteners in SSBs are fructose that will largely be metabolized in the liver. The first working enzyme is fructokinase or ketoheksokinase (KHK-C) using adenosine tri phosphate (ATP) to phosphorylate fructose to fructose -1



phosphate. Then the fructose-1 phosphate was converted to dihydroxyac 6 ne phosphate and glyceraldehyde 3-phosphate which is the material to form glycerol 3-phosphate and acetyl-KoA. Furthermore, acetyl-KoA was converted to acyl-KoA which binds to glycerol-3 phosphate to form triacyl glycerol or triglyceride. The cross-sectional study of adolescents in the UK in 2011 suggests that cons 22 ng high intake SSBs may increase triglyceride levels as 25 as the risk of cardiovascular disease [29]. A c10 rt study in children aged 8-15 years also mentions that the intake of SSBs was positively associa 10 with triglyceride levels [30].

This study showed that no correlation of SSBs intake with fasting blood glucose levels in adolescents (p > 0.05). These results are supported by study conducted on obese adolescets showing no association between intake of sweet beverage with fasting blood glucose levels [31]. The absence of sugar-sweetened beverage intake link with fasting blood glucose levels in this study because all subjects had fasting blood glucose levels in the normal category, so it cannot give 21 idea of the relationship of SSBs intake with fasting blood glucose levels. This result is supported by a Brazilian study in 2013 which states that hyperglycemia in adolescents has the smallest percentage, which is 2% [32]. Moreover, fasting blood glucose indicator is more describing of was conducted on subjects with overweight conditions, has a family history of type 2 diabetes mellitus, and have signs of insulin resistance or conditions associated with insulin resistance [33]. The normal fasting blood glucose levels in adolescents show because the body's compensating system is still good so the body can maintain normal blood glucose levels through the hormone insulin secreted by the pancreas [34]. Based on the correlation test, the intake of SSBs showed a positive correlation with fasting blood glucose, which means higher 23Bs intake then fasting blood glucose level will be higher. This is in line with a cross-sectional study conducted on children and adolescents claiming that sugar intake from beverages can increase fasting blood glucose levels [11] [35]. Sugar-steetened beverage contains more easily absorbed simple carbohydrates such as sucrose fructose corn syrup (HFCS) **65** a high-glycemic index diet which would then increase appetite. and weight gain associated with the development of glucose intolerance and insulin resistance [66]. In addition, the fructose used as a sweetener in SSBsalso contributes to the failure of glucose tolerance and insulin resistance through the mechanism of de novo lipogenesis (DNL). Fructose induces DNL by providing carbon atoms (glycerol 3 phosphate and acyl-KoA) converted into monoacylglycerol and diacylglycerol (DAG) which were subsequently converted to triglycerides. The accumulation of triglycerides in the liver will result in insulin resistance [37].

IV. CONCLUSION

Most of the subjects in this study had a high energy eating habits (49.2%), and consumed a high sugar beverage (72.9%). The average sugar intake in drinks reached 92.5 g/day (19.2% of total energy). So that found 44.1% subjects have experienced central obesity and 62.7% have hyper triglyceride which is a component of metabolic syndrome. Adolescents need to be educated regarding the limit of consuming SSBs maximum 250-350 mL/day or additional sugar to foods and beverage <50 g/day (<4 tbsp/day) and offset by physical activity for achieving an energy balance and lowering the risk of metabolic syndrome.

REFERENCES

- [1] Alberti KGMM, Eckel RH, Grundy SM, Zimmet PZ, Cleeman JI, Donato KA, et al. Harmonizing the Metabolic Syndrome: A Joint Interim Statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Hearth Federation; International Atheroclerosis Society; and International Association for the Study of Obesity. Circulation. 2009;120:1640-1645
- [2] Weiss R, Dziura J, Burgert TS, Tamborlane WV, Taksali SE, Yeckel CW, et al. Obesity and the Metabolic Syndrome in Chldren and Adolescents. New England Journal of Medicine. 2004;350:2362-2374
- [3] Dieny FF, Widyastuti N, Fitranti DY. Metabolic Syndrome among Adolescents Obese: Prevalence and Relation with Diet Quality. Indonesian Journal Clinical of Nutrition. 2015;12 (1): 1–11
- [4] Muhammad D, Dieny FF. Intake of Vitamins A, C, and E with the Incidence of Metabolic Syndrome in Adolescent Obese. Journal of Nutrition College. 2016;5(4):289–297
- [5] Santos A-C, Lopes C, Guimaraes J, Barros H. Central obesity as A Major Determinant of Increased High-Sensitivity C-reactive Protein in Metabolic Syndrome. International Journal of Obesity. 2005;29:1452– 1456
- [6] Oktaviani WD, Saraswati LD, Rahfiludin MZ. Fast Food Consumption Habits, Consumption Patterns, Characteristics of Youth and Parents with Body Mass Index (BMI). Public Health Journal.. 2012;1(2):542-553
- [7] Robert Wood Johnson Foundation. Food and Beverage Marketing to Children and Adolescents: An Environment at Odds with Good Health. Healthy Eating Research. 2011:1-11
- [8] Flood JE, Roe LS, Rolls BJ.The Effect of Increased Beverage Portion Size on Energy Intake at A Meal. Journal of the Academy of Nutrition and Dietetics.2006;106:1984-1990
- [9] Centers for Disease Control and Prevention. The CDC Guide to Strategies for Reducing the Consumption of Sugar Sweetened Beverage (SSBs). 2010: 1-41
- [10] Barrio-Lopez MT, Martinez-Gonzales MA, Fernandez-Montero A, Beunza JJ, Zazpe I, Bes-Rastrollo M. Prospective Study of Changes in Sugar-Sweetened Beverage Consumption and the Incidence of the Metabolic Syndrome and Its Components : the SUN cohort. British Journal of Nutrition. 2013;(110):1722–1731
- Wang J. Consumption of Added Sugars and Development of Metabolic Syndrome Components Among A Sample of Youth At Risk of Obesity. McGill University; 2013:1–244
- [12] Barquera S, Hernandez-Barrera L, Tolentino ML, Espinosa J, Ng SW, Rivera JA, et al. Energy Intake from Beverages is Increasing among Mexican Adolescents and Adults. Journal of Nutrition. 2008; 138: 2454-2461
- [13] Febriyani NMPS, Hardinsyah, Briawan D. Calorie Drinks and Its Contribution to Total Energy Intake in adolescents and adults. Journal of Nutrition and Food.. 2012; 7(1): 35-42
- [14] Chan T-F, Lin W, Chen Y-L, Huang H-L, Yang W-Z, Lee C-Y, et al. Elevated Serum Triglyceride and Retinol-Binding Protein 4 Levels Associated with Fructose-Sweetened Beverages in Adolescents. Plos One Journal. 2014;9(1): 1–9
- [15] Malik VS, Hu FB. Sweeteners and Risk of Obesity and Type 2 Diabetes: The Role of Sugar-Sweetened Beverages. Springer Link Journal. 2012; 12: 195-203.

Advances in Health Science Research, volume 12



- [16] Pulungan AB, Marzuki AN, Julia M, Rosaliana I, Damayanti W, Yanuarso P et al. Diagnosis and Management of Metabolic Syndrome in Children and Adolescents. Consensus of Indonesian Pediatric Association. 2014;1-25
- [17] Ministry of Health of the Republic of Indonesia. Regulation of the Minister of Health of the Republic of Indonesia concerning Inclusion of Sugar, Salt and Fat Content Information and Health Message for Prepared Food and Ready Food. Jakarta; 2013.
- [18] World Health Organization. Guideline: Sugars Intake for Adults and Children. Geneva; 2015 p. 1–46
- [19] Ervin RB, Kit BK, Carroll MD, Ogden CL. Consumption of Added Sugar Among U. S. Children and Adolescents, 2005–2008. U.S Department of Healh and Human Services. 2012;(87):2005–2008
- [20] Akhriani M, Fadhilah E, Kurniasari FN. The relationship of consumption of sweetened beverages with overweight among students in SMP Negeri 1 Bandung. Indonesian Journal of Human Nutrition. 2015;1(1):57–70
- [21] Ministry of Health of the Republic of Indonesia. Balanced Nutrition Guidelines. Jakarta; 2014.
- [22] Widyastuti N, Dieny FF, Fitranti DY. Saturated Fat and Fiber Intake in Adolescent Obesity; Associated with Metabolic Syndrome. Indonesian Journal of Clinical Nutrition. 2016;12(4):131–137
- [23] Piche M-E, Weisnagel SJ, Comeau L, Nadeau A, Bergeron J, Lemieux S. Contribution of Abdominal Visceral Obesity and Insulin Resistance to the Cardiovascular Risk Profile of Postmenopausal Women. American Diabetes Association. 2005;54(March):770–777
- [24] Rodriguez G, Moreno L, Blay M, Blay V, Garagorri J, Bueno M. Body Composition in Adolescents: Measurements and Metabolic Aspects. International Journal of Obesity. 2004;28:54–58
- [25] Taverne F, Richard C, Couture P, Lamarche B. Abdominal Obesity, Insulin Resistance, Metabolic Syndrome and Cholesterol Homeostasis. Pharma Nutrition. 2013;1(4):130–136
- [26] Pereira MA. Sugar-Sweetened and Artificially-Sweetened Beverages in Relation to Obesity Risk. Am SocNutr. 2014;797–808
- [27] Shapiro A, Mu W, Roncal CA, Cheng KY, Johnson RJ SP. Fructose-Induced Leptin Resistance Exacerbates Weight Gain Inresponse to Subsequent High Fat Feeding. American Journal Physiology Regulation Integrative and Comparative Physiology. 2008;1370–1375

- [28] Kosova EC, Auinger P, Bremer AA. The Relationship between Sugar-Sweetened Beverage Intake and Cardiometabolic Markers in Young Children. Journal of Academy of Nutrition Dietetic. 2014;113(2):219– 227
- [29] 4 mpay MI Van, Mckeown NM, Goodman E, Eliasziw M, Chomitz VR, Gordon CM, et al. Sugar-Sweetened Beverage Intake is Positively Associated with Baseline Triglyceride Concentrations, and Changes in Intake Are Inversely Associated with Changes in HDL Cholesterol over 12 Months in a Multi-Ethnic Sample of Children. The Journal of Nutrition. 2015;145:2389–2395
- [30] Welsh JA, Sharma A, Argeseanu S, Vos MB. Consumption of Added Sugars and Cardiometabolic Risk Indicators Among US Adolescents. Circulation Journal National Institutes of Health. 2014;123(3):249–257
- [31] Sudono PP, Sulistyoningrum DC, Tsani AFA. The Relationship between Sweet Drinks to Blood Sugar Levels in Adolescent Obesity in the Yogyakarta. Universitas GadjahMada; 2015
- [32] Park HS, Oh SW, Cho S, Choi WH, Kim YS. The Metabolic Syndrome and Associated Lifestyle Factors Among South Korean Adults. Interhational Journal of Epidemiology. 2004;33(2):328–336
- [33] Mittal S. The Metabolic Syndrome in Clinical Practice. Inggris: Springer; 2008. P. 226
- [34] Buse JB, Kenneth SP, Charles FB. Type 2 Diabetes Mellitus. William Textbook of Endocrinology. 2002;1427–1451
- [35] Wang YC, Bleich SN, Gortmaker SL. Increasing Caloric Contribution from Sugar Sweetened Beverage (SSBs) and 100% Juices among US Children and Adolescents, 1988-2004. Official Journal of The American Academy of Pediatrics. 2008; 121(6): 1604-1614
- [36] Perichart-perera O, Balas-nakash M, Rodríguez-cano A, Muñozmanrique C. Correlates of Dietary Energy Sources with Cardiovascular Disease Risk Markers in Mexican School-Age Children. Journal of American Dietetic Association. 2010;110(2):253–260
- [37] Schulze MB, Manson JE, Ludwig DS, Colditz GA, Stampfer MJ, Willett WC. Sugar Sweetened Beverages, Weight Gain, and Incidence of Type 2 Diabetes in Young and Middle-Aged Women. American Medical Association. 2004;292(8):927–934
- [38] Malik VS, Popkin BM, Bray GA, Despres J-P, Hu FB. Sugar Sweetened Beverage, Obesity, Type 2 Diabetes and Cardiovascular Disease Risk. National Institutes of Healh. 2010;83(2):101–108

Intake of Sugar-Sweetened Beverage and Metabolic Syndrome Components in Adolescents.

ORIGIN	ALITY REPORT		
	3% 19% INTERNET SOURCES	9% PUBLICATIONS	7% STUDENT PAPERS
PRIMAR	RY SOURCES		
1	download.atlantis-press.c	om	9%
2	ejournal.undip.ac.id		3%
3	Submitted to Universitas	Negeri Semara	ng 2%
4	repozitorij.svkst.unist.hr		1%
5	www.ncbi.nlm.nih.gov		1%
6	Submitted to Quest Intern Perak Student Paper	ational Univers	ity 1 %
7	www.researchgate.net		1%
8	publikasi.polije.ac.id		1%

9

<1%

10	Hanieh-Sadat Ejtahed, Zahra Bahadoran, Parvin Mirmiran, Fereidoun Azizi. "Sugar-Sweetened Beverage Consumption Is Associated with Metabolic Syndrome in Iranian Adults: Tehran Lipid and Glucose Study", Endocrinology and Metabolism, 2015 Publication	<1%
11	www.rsmricher.com	<1%
12	d.lib.msu.edu Internet Source	<1%
13	Kang, Y.H "The association of serum gamma glutamyltransferase with components of the metabolic syndrome in the Korean adults", Diabetes Research and Clinical Practice, 200708 Publication	<1%
14	www.nutritionandmetabolism.com	<1%
15	pubs.sciepub.com Internet Source	<1%
16	pakar.unnes.ac.id	<1%

17	bmcresnotes.biomedcentral.com	<1%
18	www.degruyter.com	<1%
19	www.endocrineconnections.com	<1%
20	circ.ahajournals.org	<1%
21	Nabila Bouatia-Naji, Vincent Vatin, Cécile Lecoeur, Barbara Heude et al. "Secretory granule neuroendocrine protein 1 (SGNE1) genetic variation and glucose intolerance in severe childhood and adult obesity", BMC Medical Genetics, 2007 Publication	<1%
22	Bray, G. A "Energy and Fructose From Beverages Sweetened With Sugar or High- Fructose Corn Syrup Pose a Health Risk for Some People", Advances in Nutrition An International Review Journal, 2013. Publication	<1%
23	Fructose High Fructose Corn Syrup Sucrose and Health, 2014. Publication	<1%
24	Betty R. Vohr, Charlotte M. Boney. "Gestational	<1%

diabetes: The forerunner for the development of maternal and childhood obesity and metabolic syndrome?", The Journal of Maternal-Fetal & Neonatal Medicine, 2009 Publication

25

27

"Metabolic Syndrome", Springer Science and	<1 %
Business Media LLC, 2016	
Publication	

<1%

<1%

- Gaurang Deshpande, Rudo F Mapanga, M 26 Faadiel Essop. "Frequent sugar-sweetened beverage consumption and the onset of cardiometabolic diseases: cause for concern?", Journal of the Endocrine Society, 2017 Publication
 - Noora Mattsson, Tapani Rönnemaa, Markus Juonala, Jorma S. A. Viikari, Olli T. Raitakari. "Childhood predictors of the metabolic syndrome in adulthood. The Cardiovascular Risk in Young Finns Study", Annals of Medicine, 2009 Publication
- Trevor S Ferguson, Marshall K Tulloch-Reid, <1% 28 Novie OM Younger, Jennifer M Knight-Madden et al. "Prevalence of the metabolic syndrome and its components in relation to socioeconomic status among Jamaican young adults: a crosssectional study", BMC Public Health, 2010 Publication

Exclude quotes	
----------------	--

Exclude bibliography On

On

Exclude matches

Off