

Low zinc serum levels and high blood lead levels among school-age children in coastal area

by Fronthea Swastawati

Submission date: 07-Jan-2020 04:34PM (UTC+0700)

Submission ID: 1239738749

File name: C14-IOP Conference Series Earth and Environmental Science, Vol 55, Conference 1.pdf (1,002.34K)

Word count: 4836

Character count: 25360

PAPER • OPEN ACCESS

Low zinc serum levels and high blood lead levels among school-age children in coastal area

² To cite this article: Adriyan Pramono *et al* 2017 *IOP Conf. Ser.: Earth Environ. Sci.* **55** 012058

² View the [article online](#) for updates and enhancements.

Related content

- [Living in Prone Flooding Area: in Coastal Areas of Semarang](#)
W P Tyas
- [Correlation between iron deficiency anemia and intestinal parasitic infection in school-age children in Medan](#)
D M Darlan, F R Ananda, M I Sari et al.
- [Serum iron parameters in liver cirrhosis](#)
G A Siregar and W Maail

Low zinc serum levels and high blood lead levels among school-age children in coastal area

Adriyan Pramono¹, Binar Panunggal^{1*}, M.Zen Rahfiludin², Fronthea Swastawati³

¹Department of Nutrition/Center of Nutrition Research (CENURE), Faculty of Medicine, Diponegoro University, Jl.Dr.Sutomo No.18, Semarang 50231, Indonesia, adriyanpramono@fk.undip.ac.id

² Faculty of Public Health, Diponegoro University, Jl. Prof.Sudarto SH, Semarang 50277, Indonesia, rahfiludin@yahoo.com

³ Faculty of Marine and Fisheries, Diponegoro University, Jl. Prof.Sudarto SH, Semarang 50277, Indonesia, fronthea_thp@yahoo.co.id

*Email: panunggalbinar@gmail.com

Abstract. The coverage of environmental lead toxicant was quiet wide. Lead exposure recently has been expected to be associated with zinc deficiency and blood indices disturbance. Emphasizing on children, which could absorb more than 50 % of lead that enters the body. Lead became the issue on the coastal area due to it has polluted the environment and waters as the source of fisheries products. This was a cross sectional study to determined nutritional status, blood lead levels, zinc serum levels, blood indices levels, fish intake among school children in coastal region of Semarang. This study was carried out on the school children aged between 8 and 12 years old in coastal region of Semarang. Nutritional status was figured out using anthropometry measurement. Blood lead and zinc serum levels were analyzed using the Atomic Absorbent Spectrophotometry (AAS) at a wavelength of 213.9 nm for zinc serum and 283.3 nm for blood lead. Blood indices was measured using auto blood hematology analyzer. Fish intake was assessed using 3-nonconsecutive days 24-hours food recall. The children had high lead levels (median 34.86 µg/dl, range 11.46 - 58.86 µg/dl) compared to WHO cut off. Zinc serum levels was low (median 18.10 µg/dl, range 10.25 - 41.39 µg/dl) compared to the Joint WHO/UNICEF/IAEA/IZiNCG cut off. Approximately 26.4% of children were anemic. This study concluded that all school children had high blood lead levels, low zinc serum, and presented microcytic hypochromic anemia. This phenomenon should be considered as public health concern.

Keywords: blood lead, zinc serum, coastal region, microcytic hypochromic anemia, school children

1. Introduction 26

Zinc (Zn) is an essential trace element plays role as a cofactor of more than 100 metalloenzyme, plays an important part in cell regeneration, metabolism, growth and immune function [1]. Zinc deficiency is associated with suboptimal growth, diarrhea, and decreased of immunity [2]. *World Health*



2 Content from this work may be used under the terms of the [Creative Commons Attribution 3.0 licence](https://creativecommons.org/licenses/by/3.0/). Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Organization (WHO) estimates the prevalence of zinc deficiency in the world's population is ranged between 4-11% [11]. Additionally, about 5 to 30% zinc deficiency has occurred in children and adolescents both in developed and developing countries [3]. Especially in the developing countries, harmful environmental exposure recently has been associated with the deficiency of Zn. Nutritional zinc deficiency can be caused from lead (Pb) exposure through dangerous cycle that increases lead absorption and increases Zn secretion consecutively [4]. Previous studies in animal experiments summarized that the conditions of marginal Zn deficiency, could increase blood lead levels in the presence of Pb exposure [5,6].

Among school children, chronic Pb exposure is correlated with nutritional deficiencies, anemia, impairments to physical growth, cognitive deficient and learning disorders [7 - 10]. Anemia is a major public health problem especially in developing countries. Additionally, WHO reported that of 25.4% anemia has been found in school children [11]. Anemia is a state of decreasing of erythrocytes and hemoglobin, whereas those condition impaired its function to carry oxygen through body tissues sufficiently [12,13]. Hemoglobin is composed of four globin chain polypeptide whereas in each chain contains a heme molecule containing Iron [14]. As well as iron, zinc also plays an important role in the formation of hemoglobin. In heme biosynthesis, δ -ALA dehydratase enzyme that catalyzes charge 2 δ -ALA into molecules that is highly dependent on zinc [15]. A study conducted in Atlanta gives us information that there is a significant correlation between serum zinc concentrations and hemoglobin levels since children who suffered iron deficiency anemia tend to be deficiency of zinc serum [16]. Normal zinc level has a consequence on erythrocytes life period, due to zinc contributes in protecting erythrocytes from oxidative stress and cell damage [17]. Increasing of blood lead levels could also interfered with erythropoiesis by inhibiting the synthesis of protoporphyrin and the absorption of iron (Fe) thus increasing the risk anemia. Furthermore, lead also affects the morphology of erythrocytes [18].

Nowadays, rapid growth of industrialization and its waste became environmental issues in coastal region of Indonesia. The industrial waste that was dumped in the coastal region induced the increasing level of lead pollutant in this area [19]. The coverage of environmental lead toxicant was quiet wide, including the exposure of fisheries products, public water source, and air pollution [20]. Lead exposure among children is an increasing problem globally, adversely affecting the quality of life among large numbers of children. The impact of lead exposure can affect all ages, whereas the children could absorbed more than 50 % of lead that enters the body, compared with adults who absorbed only 15-35 % [21]. The major vehicle of its exposure in the human body consecutively through the food consumption (65 %), the water consumption (20 %) or through the air (15 %) which were contaminated by lead [22]. In area with lead exposure, anemia and/or zinc deficiency might be occurred with or without clinical manifestation [20].

Lead exposure can be seen as only a trivial environment and health problem. A study depicted even though given iron and zinc supplementation on lead-exposed children did not reduce blood lead levels [23]. Recently in Indonesia study about lead exposure among children mostly conducted in urban area, but not in coastal region. Especially among school children in Semarang city coastal region, the linking between blood lead levels with serum zinc level, anemia, and red blood cell indices has been not fully investigated.

2. Materials and Methods

2.1. Study population and site

A cross sectional study was conducted on an elementary school in Tambak lorok coastal region, Semarang, Central Java Indonesia. Tambak lorok is located less than 3 Kilometers from coastal java sea and near to harbor industrial region (Figure 1). About 80 subjects in this study were recruited using consecutive sampling method and met the inclusion criteria. During data collection, 8 subjects were dropped out due to did not agree to collect the blood. There were 72 children have completed data collection.

Inclusion criteria in this study: age between 8 to 12 years, and did not suffer from infectious illness (diarrhea, pneumonia, tuberculosis, and parasites) which were evidenced by looking at health records in primary health care. Demographic characteristics focused on parental educational level, parental occupation, source of water consumption, consumption of seafood products, age, and sex. Body weight was measured using calibrated digital body weight scale with accuracy 0.1 Kilogram. Body height was obtained using board stadiometer with accuracy 0.1 Centimeters. Median z score for weight for age, height for age, and BMI for age were performed to describe children nutritional status.

2.2. Biochemical collection and analysis

Biochemical data in this study consists of serum zinc level, red blood cells indices (Hb, Ht, mcv, mch, mchc), and blood lead level. About 10 mL venous blood has taken in the morning day before subjects have a breakfast using disposable plastic syringe and immediately transferred into three sterile-vacutainer. A potassium anticoagulant - EDTA (Ethylene Diamine Tetra Acetate) vacutainer was injected with 2 mL venous blood. This vacutainer was used to measure red blood cells (RBCs, in $\times 10^6$ /mL), hemoglobin (Hb, in g/dL), hematocrit (Ht, in g/dL) mean corpuscle volume (MCV), mean corpuscle hemoglobin (MCH), and mean corpuscle hemoglobin concentration (MCHC). Blood indices were analyzed using automated Nihon Cohden Celltac E MEK-7222K hematology analyzer. Ferritin serum was analyzed using ELISA method.

The other two non-additives vacutainer were injected with 3 mL for serum zinc measurement and 5 mL for blood lead measurement. Approximately 3 mL whole blood in non-additive vacutainer was centrifuged, then the blood serum was prepared to zinc serum analysis. Substantially 5 mL whole blood was centrifuged and incorporated with APDC (Ammonium Pyrrolidine Dithiocarbamate) 2%. The incorporated solution was then re-centrifuged, allowed to stand for 5 minutes and mixed with 2 mL of butyl acetate. That mixed solution was centrifuged again until separation occurs. The reading of serum zinc level and blood lead level were operated using the Atomic Absorbent Spectrophotometry (AAS) at a wavelength of 213.9 nm for zinc serum and 283.3 nm for blood lead (Shimadzu AA6401F, Japan). Biochemical measurement was organized in the micronutrient and IDD Center laboratory, Diponegoro University, Indonesia.

Dietary assessment was conducted to determine fish intake among this population. The three-non consecutive days 24-hour food recall method was carried out to describe fish intake. In this study, local fish products have also been collected from this coastal area. Environmental assessment has been conducted on drinking water samples were collected using standard methods. Lead contents from fish products was measured using Spectrophotometry. Lead analysis of local fish products and drinking water samples were carried out in Center of Prevention on Industrial Pollution Laboratory, Ministry of Industrial, Republic Indonesia.

2.3. Statistical analysis

The SPSS for IBM version 19 and Microsoft excel software were performed for statistical analysis. All the study variables were tested for normality by the Kolmogorov-Smirnov test. The Mann Whitney test was operated to compare the blood lead levels and zinc serum levels between subjects with anemia and non anemia. The Correlation between blood lead level and serum z, hb, ht, RBC, mcv, mch, and ferritin were performed using Spearman rank's correlation analysis. Statistical significance was considered when p value < 0.05.

2.4. Ethics

The study protocol was approved by the Board of Medical Ethics on Faculty of Medicine Diponegoro University / Dr. Kariadi Hospital with no. 534/EC/FK-RSDK/2015. The participants, who were recruited in this study, clearly informed about the purpose of investigations and expected outcomes. Informed consent was obtained and signed by the parents of the subjects before the study began.

3. Results

3.1. Demographic characteristics of respondents

The coastal region in this study was taken place at Tambak Lorok area, the northern area of Semarang, Central Java. This area borders along with java sea. The main harbor of Central Java province has been located in this coastal area. Industrial area that was built in this region may contributed to environmental and health problems issue. Figure 1 depicts environmental situation in this coastal region.




Figure 1. Environmental situation of Semarang coastal region (authors documented).

3.2. Biochemical data analysis

Table 1 shows values of haemoglobin, hematocrit, red blood indices, zinc serum levels, blood lead levels among all children. All subjects had blood lead levels ≥ 10 ($\mu\text{g}/\text{dl}$), low zinc serum levels (< 65 $\mu\text{g}/\text{dl}$), and median of ferritin describes < 15 ng/ml .

Table 1. Mean, SD, median, minimum and maximum values of blood indices, zinc serum, blood lead levels, and ferritin among subjects (N=72)

Biochemical parameters	Median	Range (min – max)
Haemoglobin (g/dl)	12.85	10,00 – 15,6
Hematocrit (g/dl)	35.40	28,70 – 44,70
RBC ($\times 10^6$ mm^3)	4.82	4.21 – 6.00
MCV (μ^3)	74.45	52.10 – 81.05
MCH	27.00	18.10 – 29.70
MCHC	36.30	34.10 – 38.30
Ferritin (ng/ml)	13.56	2.28 – 44.68
Zinc serum ($\mu\text{g}/\text{dl}$)	18.10	10,25 – 41,39
Blood Lead Level ($\mu\text{g}/\text{dl}$)	34.86	11.46 – 58.86

24

This study was carried out to 72 children with age range from 9 to 13 years with a mean value of 10.16 \pm 1.02 years. Approximately 66.7% of children's age were categorized between 9 and 10 years old.

Surprisingly, about 90.3% of children were stunting, consists of mild stunting 34.7%, moderate stunting 45.8%, and severe stunting 9.7%. We can see from table 1 about 26.4% children were anemic. Father's occupation of subjects largely as fisherman (59.7%). Both father and mother of children had literate, mostly graduated from Senior High School (father 51.4%, mother 45.8%) (Table 2).

Table 2. Characteristics of school age children (N=72)

Characteristics	N (%)
Sex	
Male	39 (54.2)
Female	33 (45.8)
Age	
9 – 10	48 (66.7)
11 – 12	24 (33.3)
Height for Age Nutritional Status	
Normal	7 (9.7)
Mild Stunting	25 (34.7)
Moderate Stunting	33 (45.8)
Severe Stunting	7 (9.7)
Anemia Status	
Anemic	19 (26.4)
Non Anemic	53 (73.6)
Father literacy	
Graduated elementary school	13 (18.1)
Graduated junior high school	22 (30.6)
Graduated senior high school	37 (51.4)
Mother literacy	
Graduated elementary school	16 (22.2)
Graduated junior high school	23 (31.9)
Graduated senior high school	33 (45.8)
Father's occupation	
Salesman	5 (6.9)
Fisherman	43 (59.7)
Labor	24 (33.3)
Mother's occupation	
Household	36 (50.0)
Salesman	2 (2.8)
Labor	34 (47.2)

Comparison between mean values of different hematological parameters, blood lead levels, zinc serum levels and serum ferritin in anemic and non anemic were figured out in this study (Table 3). Regarding the hematological parameters, nearly all values were significantly lower among the anemic than the non-anemic group except for blood lead levels, which showed a highly significant elevation among the anemic group. However, according to World Health Organization (WHO) cut off for blood lead levels, all children in this study had a blood lead ≥ 10 $\mu\text{g}/\text{dl}$ (high blood lead level group {HBLLs}). As for the RBC count, ferritin, and zinc serum, no statistically significant difference was detected between the groups.

Table 1 Comparison between mean values of different hematological parameters, zinc serum, blood lead levels, and serum level of ferritin in anemic and non-anemic groups.

Biochemical parameters	Anemic Group (N=19)	Non Anemic Group (N=53)	Test of Significance
RBC ($\times 10^6 \text{ mm}^3$)	4.82 \pm 0.49	4.88 \pm 0.33	t = - 0.56 p = 0.58
Hb (g/dl)	11.65 \pm 0.62	13.26 \pm 0.77*	t = - 9.03 p = 0.000
Hct	32.66 \pm 1.44	36.47 \pm 1.88*	t = - 9.05 p = 0.000
MCV(μ^3)	68.43 \pm 7.50	74.89 \pm 3.74*	t = - 4.84 p = 0.000
MCH	24.45 \pm 3.05	27.26 \pm 1.65**	t = - 5.01 p = 0.000
MCHC	35.66 \pm 0.92	36.39 \pm 0.75*	t = - 3.09 p = 0.005
Ferritin (ng/ml)	15.59 \pm 8.39	15.36 \pm 8.58	t = 0.10 p = 0.92
Zinc serum ($\mu\text{g/dl}$)	20.13 \pm 9.64	20.44 \pm 8.31	t = - 0.12 p = 0.22
Blood lead level ($\mu\text{g/dl}$)	41.95 \pm 10.61	32.03 \pm 10.33**	t = 3.52 p = 0.001

*independent t-test were performed (significant if $p < 0.05$)

**Mann Whitney test were performed (significant if $p < 0.05$)

Table 4 reveals the correlation between the different red blood indices, serum zinc levels and the blood levels of lead. Blood lead levels had a significant negative correlation with haemoglobin, hematocrit, and zinc serum ($r = -0.330$, $r = -0.328$, respectively $p < 0.005$ and $r = -0.265$, $p < 0.05$).

Table 4. Correlation of different hematological parameters, serum iron and ferritin levels in relation to blood lead, copper and zinc.

Hematological parameters	Blood lead levels r-value (p-value)	Zinc serum levels r-value (p-value)
RBC ($\times 10^6 \text{ mm}^3$)	-0.066 (0.585)	-0.128 (0.285)
Hb (g/dl)	-0.330 (0.005)	0.119 (0.319)
Hct	-0.328 (0.005)	0.101 (0.396)
MCV(μ^3)	-0.124 (0.298)	0.130 (0.276)
MCH	-0.109 (0.363)	0.148 (0.214)
MCHC	-0.106 (0.375)	0.144 (0.227)
Ferritin (ng/ml)	0.178 (0.135)	0.075 (0.529)
Zinc serum levels	-0.265 (0.025)	-
Blood lead levels	-	-0.265 (0.025)

3.3. Environmental assessment

Environmental assessment has been conducted on drinking water samples and dietary fish intake. Local fish products were collected based on the information from dietary fish intake interview. Lead content of drinking water and local fish products can be seen in Table 5.

Table 5. Fish intake (g per day) and lead levels of drink water and local fish products

Parameters	Mean	Median	Range (min – max)
Fish intake (g/day)		40	10.0 – 128
Lead from drinking water (µg/dl)		0	0
Lead from some type of fish (mg/Kg)			
- Green mussels (Pena viridis)	1.13		
- Scallop shell (Anadara granosa)	0.76		
- Kipper fish (Scatophagidae)	0.18		
- Snapper fish (Lates calalifer)	0.13		
- Mullet fish (Mugil cephalus)	0.18		

4. Discussion

All school children in this study (100%) had BLL ≥ 10 µg/dl and zinc serum levels < 65 µg/dl, similar to a study done by Hegazy et al [7], who also reported about more than a half subjects (63.3%) had BLL ≥ 10 µg/dl. However the current study is similar to the results obtained for children population India [24]. The cut off value of 10 µg/dl defined by the WHO guideline for Childhood Lead Poisoning as a limit for an elevated blood lead level primarily is based on neurological toxicity and has wide range of toxicity including neurobehavioral impairment [21]. Recently, even though blood lead levels (BLL) less than 10 µg/dl is considered safe, a study confirmed that BLL < 10 µg/dl has associated with cognitive deficits [25]. Thus our data showed 45 children had low zinc serum levels (< 65 µg/dl). The Joint WHO/UNICEF/IAEA/IZINCG asserted zinc serum levels < 65 µg/dl has been recognized as serious public health problems [26].

Schwartz et al [28] reported that children living near lead smelters in the US of Idaho, had blood lead levels approximately 25 µg/dl and were correlated with anaemia. In addition, Jain et al [29], reported that children with BLL > 10 µg/dl had 1.7 times risk to be moderate anaemia, in contrast with our finding showed that 73.6% of children had normal haemoglobin. From et al [10], suggested that haemoglobin level did not correlate well with BLLs. However, our finding described that Hb, Hct, MCV, MCH and MCHC values of children with anaemia lower in comparison to anemic group. Moreover high BLLs has negatively associated with lower zinc serum and some blood indices levels. Lead absorption occurs predominantly in the duodenum and jejunum. The process of lead entry into the body may carry on through food, drink or by air. Approximately 90 % of lead absorbed by the blood binds to red blood cells [27]. The children absorbed up to 50% of lead from food and/or drink through the gastrointestinal tract and will be included in the metabolic processes of the body. Those lower red blood indices is similar with hypochromic microcytic anaemia which can be caused by blood lead toxicity [7]. Lead uses anaemia by impairing heme synthesis and increasing the rate of red blood cell destruction [31]. Although a causal pathway cannot be determined, yet the study findings clearly demonstrate the differences of BLLs between anaemia status.

In the present study zinc serum level of the anemic group is insignificantly different than non-anemic group. Lead (Pb) has the same valiancy number (Pb²⁺) as zinc (Zn²⁺) and iron (Fe²⁺), whereas on the cellular transport by Dimetal Transporter-1 (DMT-1) competition may be occurred [32]. Lead (Pb²⁺) will always be dominant to carry on by DMT-1 because of its density higher than Zn²⁺ and Fe²⁺

[27]. Lead (Pb) inhibits δ -ALA dehydratase enzyme that catalyzes δ -ALA into molecules, which is highly dependent to zinc on haemoglobin synthesis [15].

In the current study high BLLs among school children may be due to air pollution, sewage contamination, and probably from food consumption which were contaminated with Lead (Pb). Based on dietary interview, children consume fish regularly (range between 10 g/day and 128 g/day). Thus based on lead measurement content on some fish products showed variety lead content (mg/Kg fish products), green mussels (*Perna viridis*) had the highest content of lead (1.13mg/Kg). Green mussels (*Perna viridis*) has been vended on street food vendor near by school building. However a causal pathway between some fish products intake and blood lead levels cannot be firmed only from this data. A large epidemiology study on general northern coastal communities should remain a concern because of the nature of accumulation.

In developing countries such as Indonesia, control of lead contamination is much slower and the negative health effects getting more sporadic. The present work revealed that lead contamination should be considered as public health concern in northern coastal population of Semarang, Central Java, Indonesia.

5. Conclusion

In the present study, all subjects has high blood lead levels (BLLs) and low zinc serum. Blood lead levels (BLLs) were negatively associated with the hematological parameters and zinc serum levels. Lead (Pb) was not presented in drinking water, but Pb was discovered in some fish products that may be regularly consume by the children. Lead exposure could be controlled and strides should be taken to reduce zinc deficiency and anaemia among children population at coastal region.

6. References

- [1]. Osredkar J, Sustar N. 2011. Copper and zinc, biological role and significance of copper/zinc imbalance. *J Clin Toxicol Suppl* 3, 1 – 18.
- [2]. Prasad AS. 2003. Zinc deficiency: Has been known of for 40 years but ignored by global health organizations. *BMJ*, 326(7386), 409 - 410.
- [3]. Wessells KR, Brown KH. 2012. Estimating the global prevalence of zinc deficiency: results based on zinc availability in national food supplies and the prevalence of stunting. *PLoS One*, 7(11), e50568.
- [4]. Mason LH, Harp JP, Han DY. 2014. Pb neurotoxicity: neuropsychological effects of lead toxicity. *BioMed Res Int*, 2014, 1 – 8.
- [5]. Bushnell PJ, Levin ED. 1982. Effects of zinc deficiency on lead toxicity in rats. *Neurobeh. Toxicol. Ter*, 5(3), 283-288.
- [6]. Ashraf MH, Fosmire GJ. 1985. Effects of marginal zinc deficiency on subclinical lead toxicity in the rat neonate. *J Nutr*, 115(3), 334-346.
- [7]. Hegazy AA, Zaher MM, Abd el-hafez MA, Morsy AA, Saleh RA. 2010. Relation between anemia and blood levels of lead, copper, zinc and iron among children. *BMC Res Notes*, 3(1), 133.
- [8]. Costa de Almeida GR, de Freitas Tavares CF, de Souza AM, et al. 2010. Whole blood, serum, and saliva lead concentrations in 6- to 8-year-old children. *Sci. Total Environ*, 408, 1551–1556.
- [9]. Kordas K, Canfield RL, Lopez P, et.al. 2006. Deficits in cognitive function and achievement in Mexican first-graders with low blood lead concentrations. *Environ. Res*, 100(3), 371-386.
- [10]. Gardner RM, Kippler M, Tofail F, et al. 2013. Environmental exposure to metals and children's growth to age 5 years: a prospective cohort study. *Am. J. Epidemiol*, 177(12), 1356-1367.
- [11]. World Health Organization. Worldwide prevalence of anemia 1993-2005: WHO global database on anemia. 2008. Download from: http://apps.who.int/iris/bitstream/10665/43894/1/9789241596657_eng.pdf
- [12]. Ferreira MU, da Silva-Nunes M, Bertolino CN, et al. 2007. Anemia and iron deficiency in school children, adolescents, and adults: a community-based study in rural Amazonia. *Am J Public Health* 97(2): 237-239.

- [13]. World Health Organization. Haemoglobin concentrations for the diagnosis of anemia and assessment of severity. 2011. Download from: <http://www.who.int/vmnis/indicators/haemoglobin.pdf>
- [14]. Thomas C, Lumb AB. 2012. Physiology of haemoglobin. *CEACCP*, 12(5), 251-256.
- [15]. Scinicariello F, Murray HE, Moffett DB, et al. 2007. Lead and δ -aminolevulinic acid dehydratase polymorphism: where does it lead? A meta-analysis. *Environ Health Persp*, 115 (1), 35-41.
- [16]. Cole CR, Grant FK, Swaby-Ellis ED, et al. 2010. Zinc and iron deficiency and their interrelations in low-income African American and Hispanic children in Atlanta. *Am J Clin Nutr*, 91(4), 1027 - 1034.
- [17]. Ryu MS, Guthrie GJ, Maki AB, et al. 2012. Proteomic analysis shows the up regulation of erythrocyte dematin in zinc-restricted human subjects. *Am J Clin Nutr*, 95(5), 1096 - 1102.
- [18]. Shah F, Kazi TG, Afridi HI, et al. 2010. Environmental exposure of lead and iron deficit anemia in children age ranged 1–5years: a cross sectional study. *Sci Total Environ*, 408(22), 5325 - 5330.
- [19]. Anetor JI. 2013. Rising environmental cadmium levels in developing countries: threat to genome stability and health. *Niger J Physiol Sci*, 27(2), 103-115.
- [20]. Kordas K, Lonnerdal B, Stoltzfus RJ. 2007. Interactions between Nutrition and Environmental Exposures: Effects on Health Outcomes in Women and Children. *J Nutr*, 137(12), 2794 – 7.
- [21]. World Health Organization. Childhood lead poisoning. 2010. Download from: <http://www.who.int/ceh/publications/leadguidance.pdf>
- [22]. Liu J, McCauley L, Compber C, et al. 2011. Regular breakfast and blood lead levels among preschool children. *Environ Health*, 10(1), 28.
- [23]. Rosado JL, Lopez P, Kordas K, et al. 2006. Iron and/or zinc supplementation did not reduce blood lead concentrations in children in a randomized, placebo-controlled trial. *J Nutr*, 136(9), 2378-2383.
- [24]. Patel AB, Williams SV, Frumkin H. 2001. Blood lead in children and its determinants in Nagpur, India. *Int J Occup Environ Health*, 7: 119-26.
- [25]. Lanphear BP, Dietrich K, Auinger P, Cox C. 2000. Cognitive deficits associated with blood lead concentrations < 10 microg/dL in US children and adolescents. *Public health reports*, 115(6), 521.
- [26]. De Benoist B, Darnton-Hill I, Davidsson L, Fontaine O, Hotz C. 2007. Conclusions of the joint WHO/UNICEF/IAEA/IZiNCG interagency meeting on zinc status indicators. *Food and nutrition bulletin*, 28(3 suppl3), S480-S484.
- [27]. Kasperczyk A, Prokopowicz A, Dobrakowski M, Pawlas N, Kasperczyk S. 2012. The effect of occupational lead exposure on blood levels of zinc, iron, copper, selenium and related proteins. *Biological trace element research*, 150(1-3), 49-55.
- [28]. Schwaetz J, Landrigan PJ, Baker EL. Lead induced anemia: dose response relation and evidence for a threshold. *Am J Public Health*. 1990, 80: 165-8.
- [29]. Jain NB, Laden F, Guller U, Shankar A, Kazani S, Garshick E. 2005. Relation between blood lead levels and childhood anemia in India. *American Journal of Epidemiology*, 161(10), 968-973.
- [30]. Fromm P, Kristal-Bonch E, Benbassat J. 1999. Lead exposure in battery-factors workers is not associated with anemia. *J Occup Environ Med*, 41: 120-3.
- [31]. Bradman A, Eskenazi B, Sutton P, Goldman LR. 2001. Iron deficiency associated with higher blood lead level in children living in contaminated environments. *Environ Health Perspect*, 109: 1079-84.
- [32]. Kordas K, Lonnerdal B, Stoltzfus RJ. 2007. Interactions between nutrition and environmental exposures: effects on health outcomes in women and children. *J Nutr*, 137, 2794.

Low zinc serum levels and high blood lead levels among school-age children in coastal area

ORIGINALITY REPORT

19%

SIMILARITY INDEX

16%

INTERNET SOURCES

17%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

1

bmcrenotes.biomedcentral.com

Internet Source

9%

2

repository.ubaya.ac.id

Internet Source

2%

3

Maria Mexitalia, Yesi Oktavia Dewi, Adriyan Pramono, Mohammad Syarofil Anam. "Effect of tuberculosis treatment on leptin levels, weight gain, and percentage body fat in Indonesian children", Korean Journal of Pediatrics, 2017

Publication

1%

4

Amal A Hegazy, Manal M Zaher, Manal A Abd el-hafez, Amal A Morsy, Raya A Saleh. "Relation between anemia and blood levels of lead, copper, zinc and iron among children", BMC Research Notes, 2010

Publication

1%

5

Ivan S. Ferraz, Júlio C. Daneluzzi, Hélio Vannucchi, Alceu A. Jordão Jr. et al. "Nível sérico de zinco e sua associação com

1%

deficiência de vitamina A em crianças pré-escolares", *Jornal de Pediatria*, 2007

Publication

6	www.omicsonline.org Internet Source	<1%
7	www.epa.gov Internet Source	<1%
8	ajpregu.physiology.org Internet Source	<1%
9	www.lifesciencesite.com Internet Source	<1%
10	jn.nutrition.org Internet Source	<1%
11	rd.springer.com Internet Source	<1%
12	Bemnet Amare, Beyene Moges, Bereket Fantahun, Ketema Tafess et al. "Micronutrient levels and nutritional status of school children living in Northwest Ethiopia", <i>Nutrition Journal</i> , 2012 Publication	<1%
13	www.jil.go.jp Internet Source	<1%
14	Tasneem G. Kazi, Faheem Shah, Haffeezur Rehman Shaikh, Hassan Imran Afridi, Afzal	<1%

Shah, Naeemullah, Sadaf Sadia Arain.
"Exposure of lead to mothers and their new born
infants, residents of industrial and domestic
areas of Pakistan", Environmental Science and
Pollution Research, 2013

Publication

15

article.sciencepublishinggroup.com

Internet Source

<1%

16

Nils Milman. "Anemia—still a major health
problem in many parts of the world!", Annals of
Hematology, 2011

Publication

<1%

17

Pascal Haefliger. "Mass Lead Intoxication from
Informal Used Lead-Acid Battery Recycling in
Dakar, Senegal", Environmental Health
Perspectives, 05/14/2009

Publication

<1%

18

jissn.biomedcentral.com

Internet Source

<1%

19

Adnan M. Massadeh, Wail A. Hayajneh,
Abdullah Shorman, Mohammad Y. Gharaibeh,
Moh'd Mohanad A. Al-Dabet. "Correlation
between lead and iron in children's blood in
Jordan", Toxicological & Environmental
Chemistry, 2013

Publication

<1%

edlawcenter.org

20

Internet Source

<1%

21

academic.oup.com

Internet Source

<1%

22

www.friendsofbonobos.org

Internet Source

<1%

23

www.operaevening.com

Internet Source

<1%

24

G. A. Mostafa. "Blood lead levels in Egyptian children from high and low lead-polluted areas: impact on cognitive function", *Acta Neurologica Scandinavica*, 07/2009

Publication

<1%

25

www.sdiarticle2.org

Internet Source

<1%

26

ister.ir

Internet Source

<1%

27

Mathee, A.. "Reductions in blood lead levels among school children following the introduction of unleaded petrol in South Africa", *Environmental Research*, 200603

Publication

<1%

28

Veena Kalra, Jitendra Kumar Sahu, Puneet Bedi, R. M. Pandey. "Blood Lead Levels among School Children after Phasing-Out of Leaded

<1%

Petrol in Delhi, India", The Indian Journal of Pediatrics, 2013

Publication

29

Amal A Hegazy. "Relation between anemia and blood levels of lead, copper, zinc and iron among children", BMC Research Notes, 2010

Publication

<1%

30

Jorge L. Rosado, Patricia López, Katarzyna Kordas, Gonzalo García-Vargas et al. "Iron and/or Zinc Supplementation Did Not Reduce Blood Lead Concentrations in Children in a Randomized, Placebo-Controlled Trial", The Journal of Nutrition, 2006

Publication

<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On