

ISSN-0976-0245 (Print) • ISSN-0976-5506 (Electronic)

Volume 9

Number 9

September 2018



Indian Journal of Public Health Research & Development

An International Journal

SCOPUS IJPHRD CITATION SCORE

Indian Journal of Public Health Research and Development
Scopus coverage years: from 2010 to 2017 Publisher:
R.K. Sharma, Institute of Medico-Legal Publications
ISSN:0976-0245E-ISSN: 0976-5506 Subject area: Medicine:
Public Health, Environmental and Occupational Health
CiteScore 2015- 0.02
SJR 2015- 0.105
SNIP 2015- 0.034



Website:

www.ijphrd.com

Indian Journal of Public Health Research & Development

EXECUTIVE EDITOR

Prof Vidya Surwade

Prof Dept of Community Medicine SIMS, Hapur

INTERNATIONAL EDITORIAL ADVISORY BOARD

1. **Dr. Abdul Rashid Khan B. Md Jagar Din**, (*Associate Professor*)
Department of Public Health Medicine, Penang Medical College, Penang, Malaysia
2. **Dr. V Kumar** (*Consulting Physician*)
Mount View Hospital, Las Vegas, USA
3. **Basheer A. Al-Sum**,
Botany and Microbiology Deptt, College of Science, King Saud University,
Riyadh, Saudi Arabia
4. **Dr. Ch Vijay Kumar** (*Associate Professor*)
Public Health and Community Medicine, University of Buraimi, Oman
5. **Dr. VMC Ramaswamy** (*Senior Lecturer*)
Department of Pathology, International Medical University, Bukit Jalil, Kuala Lumpur
6. **Kartavya J. Vyas** (*Clinical Researcher*)
Department of Deployment Health Research,
Naval Health Research Center, San Diego, CA (USA)
7. **Prof. PK Pokharel** (*Community Medicine*)
BP Koirala Institute of Health Sciences, Nepal

NATIONAL SCIENTIFIC COMMITTEE

1. **Dr. Anju Ade** (*Associate Professor*)
Navodaya Medical College, Raichur, Karnataka
2. **Dr. E. Venkata Rao** (*Associate Professor*) Community Medicine,
Institute of Medical Sciences & SUM Hospital, Bhubaneswar, Orissa.
3. **Dr. Amit K. Singh** (*Associate Professor*) Community Medicine,
VCSG Govt. Medical College, Srinagar – Garhwal, Uttarakhand
4. **Dr. R G Viveki** (*Associate Professor*) Community Medicine,
Belgaum Institute of Medical Sciences, Belgaum, Karnataka
5. **Dr. Santosh Kumar Mulage** (*Assistant Professor*)
Anatomy, Raichur Institute of Medical Sciences Raichur(RIMS), Karnataka
6. **Dr. Gouri Ku. Padhy** (*Associate Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Dr. Ritu Goyal** (*Associate Professor*)
Anaesthesia, Sarswathi Institute of Medical Sciences, Panchsheel Nagar
8. **Dr. Anand Kalaskar** (*Associate Professor*)
Microbiology, Prathima Institute of Medical Sciences, AP
9. **Dr. Md. Amirul Hassan** (*Associate Professor*)
Community Medicine, Government Medical College, Ambedkar Nagar, UP
10. **Dr. N. Girish** (*Associate Professor*) Microbiology, VIMS&RC, Bangalore
11. **Dr. BR Hungund** (*Associate Professor*) Pathology, JNMC, Belgaum.
12. **Dr. Sartaj Ahmad** (Assistant Professor),
Medical Sociology, Department of Community Medicine, Swami Vivekananda Subharti
University, Meerut, Uttar Pradesh, India
13. **Dr Sumeeta Soni** (Associate Professor)
Microbiology Department, B.J. Medical College, Ahmedabad, Gujarat, India

NATIONAL EDITORIAL ADVISORY BOARD

1. **Prof. Sushanta Kumar Mishra** (Community Medicine)
GSL Medical College – Rajahmundry, Karnataka
2. **Prof. D.K. Srivastava** (*Medical Biochemistry*)
Jamia Hamdard Medical College, New Delhi
3. **Prof. M Sriharibabu** (*General Medicine*) GSL Medical College, Rajahmundry,
Andhra Pradesh
4. **Prof. Pankaj Datta** (*Principal & Prosthodontist*)
Indraprastha Dental College, Ghaziabad

NATIONAL EDITORIAL ADVISORY BOARD

5. **Prof. Samarendra Mahapatro** (*Pediatrician*)
Hi-Tech Medical College, Bhubaneswar, Orissa
6. **Dr. Abhiruchi Galhotra** (*Additional Professor*) Community and Family
Medicine, All India Institute of Medical Sciences, Raipur
7. **Prof. Deepti Pruthvi** (*Pathologist*) SS Institute of Medical Sciences &
Research Center, Davangere, Karnataka
8. **Prof. G S Meena** (*Director Professor*)
Maulana Azad Medical College, New Delhi
9. **Prof. Pradeep Khanna** (*Community Medicine*)
Post Graduate Institute of Medical Sciences, Rohtak, Haryana
10. **Dr. Sunil Mehra** (*Paediatrician & Executive Director*)
MAMTA Health Institute of Mother & Child, New Delhi
11. **Dr Shailendra Handu**, *Associate Professor*, Phrma, DM (Pharma, PGI
Chandigarh)
12. **Dr. A.C. Dhariwal**: *Directorate* of National Vector Borne Disease
Control Programme, Dte. DGHS, Ministry of Health Services, Govt. of
India, Delhi

Print-ISSN: 0976-0245-Electronic-ISSN: 0976-5506, Frequency: Monthly

Indian Journal of Public Health Research & Development is a double blind peer reviewed international journal. It deals with all aspects of Public Health including Community Medicine, Public Health, Epidemiology, Occupational Health, Environmental Hazards, Clinical Research, and Public Health Laws and covers all medical specialties concerned with research and development for the masses. The journal strongly encourages reports of research carried out within Indian continent and South East Asia.

The journal has been assigned International Standards Serial Number (ISSN) and is indexed with Index Copernicus (Poland). It is also brought to notice that the journal is being covered by many international databases. The journal is covered by EBSCO (USA), Embase, EMCare & Scopus database. The journal is now part of DST, CSIR, and UGC consortia.

Website : www.ijphrd.com

©All right reserved. The views and opinions expressed are of the authors and not of the Indian Journal of Public Health Research & Development. The journal does not guarantee directly or indirectly the quality or efficacy of any product or service featured in the advertisement in the journal, which are purely commercial.

Editor

Dr. R.K. Sharma
Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

Printed, published and owned by

Dr. R.K. Sharma
Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019

Published at

Institute of Medico-legal Publications
501, Manisha Building, 75-76, Nehru Place,
New Delhi-110019



Indian Journal of Public Health Research & Development

www.ijphrd.com

Contents

Volume 9, Number 9

September 2018

1. The Role of Chest Physiotherapy on Pulmonary Outcome in Patients with Renal Dysfunction 1
A. Sankarganesh, T. Senthilkumar, N. Venketesh, P. Soundararajan
2. A Prospective Study Comparing Combined Viscoanalostomy–Trabeculectomy to Trabeculectomy for the Management of Primary Open Angle Glaucoma 6
Anand Verma M. S., Prerana Agarwal, Pratibha Kaushal
3. Comparison between Empirical and Variational Mode Decomposition Based on Percentage Variation in Entropy Feature from Glaucoma Image 10
Bhupendra Singh Kirar, Dheeraj Kumar Agrawal
4. Isolation, Identification of Bacteria Associated with Mobile Phones and Their Antibiotic Susceptibility . 16
Ch. M. Kumari Chitturi, P. Jeevana Lakshmi
5. Effectiveness of Lumbar Flexors and Extensors Muscle Strengthening Exercises on Waist-Hip Ratio in Overweight Adolescents 21
D. Malarvizhi, A. M. Dinesh
6. Influence of Good Governance Implementation on Healthcare Performance in Three Provinces in Indonesia 27
Dumilah Ayuningtyas, RR Mega Utami, Ni Nyoman Dwi Sutrisnawati, Misnaniarti
7. Pattern of Skin Diseases in Children Attending Anganwadis a Cross Sectional Study 32
Hemalatha Umashankar, Mangala Subramanian
8. Clinical Spectrum of Hiv Infection in Children 37
H S Rajani, D Narayanappa
9. Compatibility of Sodium Fluoride Patch as an Innovation Model of Transferring Fluoride in Dental Care: A Quantitative Study Using in Vitro & in Vivo Rabbit Skin 42
Diyah Fatmasari, Endah Aryati Eko Ningtyas, Tri Wiyatini, Arwani, Ismi Rajiani
10. Spatial Distribution Characteristics And Differences Larva *Habitat An. Barbirostris* and *An. Subpictus* in the District Bulukumba 47
Iwan Suryadi, Seviana Rinawati, Siti Rachmawati
11. Prevalence and Pattern of Anxiety and Depressive Disorders in Pregnant Women Attending Antenatal Clinic 52
Pallabi Sahu, Jagadish Hansa, Debi Prasad Mohanty, Suvendu Narayan Mishra
12. Dental Treatment Demands, Needs and Utilization Among Geriatric Patients: A Tertiary Health Care-Based Retrospective Study 59
Mahima Jain, Manuel S. Thomas, Ramya Shenoy

13. Nurses' Knowledge and Practice Toward Oral Care for Intubated Patients 65
Mohammed Baqer Abbas Al-Jubouri, Sabah Abdullah Jaafar
14. Association of Food Patterns, Central Obesity Measures and Metabolic Risk Factors for Coronary Heart Disease in Adult Men 71
Monika Jain, Chetna Singh, Priya Agarwal
15. Screening of *Chlamydia trachomatis* Infection among Childbearing Age Group Women in a Tertiary Center in South India 77
Swathi. S, Naveen kumar. C, Abarna. V, Jayapradha. S, Srikumar. R
16. Problem Focus Coping Model to Face Working Environment Stressors Prevents Unsafe Action among Workers in a Steel Construction Plant 82
Noeroel Widajati
17. Knowledge and Attitude about Human Immunodeficiency Virus/Acquired Immune Deficiency Syndrome among Married Men in Kangrali, India: A Cross-Sectional Study 89
Sagar G. Khandekar MPH, Padmaja R Walvekar
18. A Study of Environmental Factors Affecting Nutritional Status of Under 5 Children in Rural Area of North India 94
Parul Jain, Amrit Virk, Zahid Ali Khan, Anshu Mittal, Harshpreet Singh, Muzamil Nazir
19. Epidemiological Pattern of Hand Injuries and Impact of Machine-Cut Hand Injuries in a Tertiary Care Hospital in South India 100
Akriti Gupta, Rajesh Kamath, Anil Bhat, Varalakshmi Chandra Sekaran, Prajwal Salins, Brayal D'Souza, Sagarika Kamath
20. X Rays Induced Oxidative Stress in Cerebral Tissue of Albino Wistar Rats 106
Sudha K, Reshma K, Gaya Pr, Vinitha Dsouza, Charu Yadav
20. Study of Some Risk Factors Causing Infertility in Females at Tertiary Care Centre 110
Salphale Shweta N., Mundada Vinod D., Lokhande G. S., Kuril B. M., Doibale M. K.
21. Learning Styles and Academic Outcome among Nursing Students-Systematic Review 114
Shalini G Nayak, Prima J J D'Souza
22. Goiter and Hypothyroidism among Elementary School Children in Lowland Agricultural Area, Brebes District Indonesia 120
Apoina Kartini, Suhartono, Dina R. Pangestuti, M. Sakundarno Adi, Suratman, Rasipin
23. Comparison between Stretching and Stabilization Exercises on Upper Trapezius Muscle Fatigue in Mobile Phone Users 126
Suresh Jothi, M.Lakshmi Prasanna, Sivakumar V. P.R.
24. Correlation of Hip Muscle Strength and Patellofemoral Pain Syndrome in Men 132
Suresh Jothi, M. Keerthiga, Sivakumar V. P. R.
25. Relationship between Depression and Oral Health Related Quality of Life among Institutionalized Elderly Population of Mysore City, India—A Cross Sectional Study 137
Nanditha Kumar M., Thippeswamy H M, Raghavendra Swamy K N
26. Canine Index: A Tool for Determination of Sex 143
V. Anu, Abinaya Vijayakumar, Jude Ritheesh Roy, V. Ravi Pavan, R. Ram Kiran, R. S. Besen Jas, S. Abinaya

Goiter and Hypothyroidism among Elementary School Children in Lowland Agricultural Area, Brebes District Indonesia

Apoina Kartini¹, Suhartono², Dina R. Pangestuti¹, M. Sakundarno Adi³, Suratman⁴, Rasipin⁵

¹Department of Public Health Nutrition, ²Department of Environmental Health, ³Department of Epidemiology, Faculty of Public Health, Diponegoro University, Semarang, Indonesia; ⁴School of Public Health, Faculty of Health Sciences, Jenderal Soedirman University, Purwokerto, Indonesia; ⁵Brebes District Health Office, Brebes, Indonesia

ABSTRACT

The most common cause of goiter is the lack of iodine intake and usually occurs in many communities in upland areas. Pesticides are widely used in the agricultural area, and it was probably the cause of goiter. This research is aimed to identify goiter and hypothyroidism among elementary school children in lowland agricultural areas. Cross-sectional study recruited sample of 100 children aged 9-12 years old from three elementary schools in Brebes District, Indonesia. Goiter existence determined by palpation of a trained nutritionist; level of thyroid stimulating hormone (TSH) measured with mini vidas test kits; and Urinary Iodine Concentration (UIC) measured with acid digestion method. Univariate and bivariate data analysis (Chi-square test, risk estimate) were applied. Proportion of goiter and hypothyroidism was 53.0% and 17.4% respectively. Median UIC was 346 µg/dL. Proportion of goiter in the children whose fathers were farmer and non-farmer was 80,8% and 43,2% respectively ($p=0.002$; Prevalence Ratio=1.9; 95% CI=1.3-2.6). The proportion of hypothyroidism in the children whose fathers were farmers (29.4%) tends to be higher than non-farmers (13.5%). However, there was no significance difference proportion of hypothyroidism in both group ($p=0.255$; Prevalence Ratio=2,2; 95% CI=0.8-6.0).

Keywords: Goiter, Hypothyroidism, Children, Agricultural Area, Indonesia

INTRODUCTION

Goiter is an enlargement of thyroid gland located in the neck caused by malfunctioning or changing gland structures or its morphology. The enlargement of the thyroid gland can influence the positions of organs. The negative effects of goiter can be a cosmetic problem and difficulties in swallowing and breathing¹. Thyroid gland has a function to produce thyroid hormones, thyroxine (T4) and triiodothyronine (T3). Thyroid hormone plays an important role in a process of body growth, brain development, nervous system, and teeth and skeletal

development². The lack of thyroid hormone will increase a level of Thyroid Stimulating Hormone (TSH), a type of hormone that increases a synthesis of thyroid hormone and stimulates an enlargement of thyroid gland³. Goiter occurred on children is responsible for growth and development disruption such as stunting, low Intelligence Quotient, and mental disorders⁴.

Goiter in an endemic area is mainly caused by the lack of iodine intakes as the critical raw material in the process of thyroid hormone synthesis. Endemic goiter, well-known as Iodine Deficiency Disorder (IDD)⁵, is generally found in highland areas due to low iodine levels in soil, water, and agricultural products. Goiter rate in lowland areas is also quite high even though the content of iodine levels in soil, water, and agricultural commodities is sufficient⁶. Related to this phenomenon, some theories stated that thyroid dysfunctions occurred due to exposures of heavy metals in the environment such as lead, mercury, cadmium, polychlorinated biphenyl (PCB), and pesticide^{7,8}.

Corresponding Author:

Suhartono

Jalan Prof. Soedarto, SH Tembalang,

Semarang 1269, Indonesia

Phone: +62 81 127 5410

Email: suhartono_damas@yahoo.com

Bulakamba, one of the Subdistricts in Brebes District, is the area with high-intensity pesticide use and depends on agricultural products (paddy, shallots, corn, green bean, and chili). Meanwhile, Kluwut Public Health Center located in the Bulakamba Subdistrict was the highest total goiter rate (TGR) among elementary school students in Brebes District (38.5%). The result of the measurement of UIC demonstrated that median UIC ranged from 286-293µg/L (Brebes District Health Offices, unpublished data 2010). These values were not categorized as iodine deficiency⁹.

MATERIAL AND METHOD

A cross-sectional study design was conducted in Brebes district, Central Java Indonesia in 2012. Minimum sample size were 97 students calculated using the formula for estimating the population proportion¹⁰. Notwithstanding, as many as 100 students aged 9-12 years old randomly selected from three elementary schools.

Variables of characteristics were collected using a structured questionnaire by a trained interviewer. Thyroid gland palpation undertaken by trained nutritionists worked at Brebes District Health Office determined the occurrence of goiter. The thyroid size was graded according to the joint criteria of WHO, UNICEF and ICCIDD (non-palpable goiter = grade 0, palpable but not visible goiter = grade 1 and palpable and visible goiter = grade 2)⁹.

Thyroid stimulating hormone (TSH) levels collected from sub-samples of 69 subjects were measured with a mini VIDAS® (bioMérieux S.A.), a compact automated immunoassay system based on the Enzyme Linked Fluorescent Assay (ELFA) principles at an accredited clinic laboratory. Non-fasting peripheral venous blood samples were obtained in the morning from 09.00 to 11.00. Subjects were categorized suffering from hypothyroidism when the TSH levels were ≥ 4.5 µIU/L². Meanwhile, UIC levels were measured using a method of acid digestion with persulfate ammonium. Spot urine samples were obtained from sub-samples of 66 subjects.

Univariate analysis was used to describe frequencies for categorical data, mean±standard deviation, and range for continuous data. Chi-square test was performed to analyze the association between two variables.

Ethics approval was obtained from the health ethics committee of Faculty of Public Health, Diponegoro

University, Semarang, Indonesia. Written permission to conduct this study was obtained from the head of Education Office Brebes District and the head teachers and chiefs of the schools involved. Parents also signed an informed consent form.

FINDINGS

A proportion of female was higher than a proportion of male in three elementary schools. Fathers and mothers educational status were mostly middle or less, respectively 95% and 97%. Meanwhile, as many as 26% of fathers worked as a farmworker and mothers as a farmworker as many as 21%. (Table 1).

Table 1: Characteristics of Subjects

Characteristics	Frequency (%)
Age (years):	
9	6 (6.0)
10	17 (17.0)
11	52 (52.0)
12	25 (25.0)
Sex:	
Male	43 (43.0)
Female	57 (57.0)
Fathers educational status:	
Middle or less	95 (95.0)
High school	5 (5.0)
Fathers occupation:	
Farmworker	26 (26.0)
Non-farmworker	74 (74.0)
Mothers educational status:	
Middle or less	97 (97.0)
High school	3 (3.0)
Mother's occupation:	
Unemployment	33 (33.0)
Farmworker	21 (21.0)
Non-farmworker	46 (46.0)

The overall prevalence of goiter was found to be 53.0%. Prevalence of Grade 1 goiter was 27.0% and that of grade 2 was 26.0%. The results of TSH level measurement on 69 samples demonstrated that as many as 17% of them were categorized suffering from hypothyroidism (TSH > 4.5 µIU/L). Meanwhile, the results of UIC level measurement showed that there was no subject suffering from iodine deficiency (> 100 (mg/L) (Table 2).

Table 2: The occurrence of goiter, hypothyroidism, and median UIC

Characteristics	Frequency (%)
The occurrence of goiter (n = 100):	
Yes	53 (53.0)
No	47 (47.0)
Grade:	
Grade 0	47 (47.0)
Grade 1	27 (27.0)
Grade 2	26 (26.0)

Conted...

Hypothyroidism (n = 69):	
Yes	12 (17.4)
No	57 (82.6)
TSH (mean±SD, range)	3,1±1,87 (0,4-11,4)
UIC (median, range) (mg/L) (n=66)	346 (192-349)

A proportion of goiter among subjects who had fathers working as a farmworker was equal to 80.8%, two times higher than that of subjects who had fathers working as non-farmworker (43.2%) (Table 3).

Table 3: The occurrence of goiter based on fathers' occupation

Fathers' occupation	The occurrence of goiter		P	Prevalence Ratio (95% CI)
	Yes	No		
Farmworker	21 (80.8)	5 (19.2)	0.002*	1.9 (1.3-2.6)
Non-farmworker	32 (43.2)	42 (56.8)		

Significant <0.05

A proportion of hypothyroidism among subjects who had fathers working as a farmworker was equal to 29.4%, two times higher than that of subjects who had fathers working as non-farmworker (13.5%) (Table 4).

Table 4: The occurrence of hypothyroidism based on fathers' occupation

Fathers' occupation	The occurrence of hypothyroidism		P	Prevalence Ratio (95% CI)
	Yes	No		
Farmworker	5 (29.4)	12 (70.6)	0.255	2.2 (0.8-6.0)
Non-farmworker	7 (13.5)	45 (86.5)		

DISCUSSION

This study found as many as 53% of elementary school students located in lowland areas suffered from goiter. Brebes is an agricultural area producer of shallot in Central Java province, Indonesia. The farmers usually spray pesticides 2-3 times per week, even every day in the rainy season. The yield rate of shallot in Brebes in 2014 as much as 121.46 quintals/hectare¹¹.

Goiter, a manifestation of thyroid hyperplasia and increased thyroid vascularity, results from a compensatory increase in the release of thyroid-stimulating hormone (TSH) as a result of lower triiodothyronine (T₄) levels and is traditionally detected and evaluated using inspection and palpation¹². We used palpation in our study because ultrasonography is cumbersome and costly to carry out¹³, and palpation is regarded as an acceptable and

simple alternative. Palpation of the thyroid is important in assessing goiter prevalence, which is relatively easy to conduct, and training of personnel to do it¹⁴.

UIC levels greater than 100 µg/L (Table 2) mean that there was no case of iodine deficiency on all research participants. Iodine deficiency early in life impairs cognition and growth, but iodine status is also a key determinant of thyroid disorders in adults. Dietary iodine intake is required for the production of thyroid hormone. Consequences of iodine deficiency include goiter, intellectual impairments, growth retardation, neonatal hypothyroidism, and increased pregnancy loss and infant mortality^{15,16}. A main cause of goiter commonly is due to lack of iodine intakes. Iodine is the critical raw material for a process of thyroid hormone biosynthesis. Iodine plays an important role in growth and development, and brain function¹⁷.

There are several accepted methods for the monitoring of population iodine status¹⁸. Because 90% of ingested iodine is renal excreted, median spot urinary iodine concentrations (UIC) serve as a biomarker for recent dietary iodine intake. Population iodine sufficiency is defined by median UIC of 100–299 µg/L in school-aged children⁹. A source of iodine in the environment depends on a location/geographical factor. Generally, sufferers of goiter are commonly found in highland areas (mountainous regions) because there is a lack of iodine levels in water and soil, where iodine has been washed away by glaciation and flooding¹⁵. On the other hand, oceans are the world's main repositories of iodine and very little of earth iodine is actually found in the soil. The deposition of iodine in the soil occurs due to volatilization from ocean water, a process aided by ultraviolet radiation. The coastal regions of the world are much richer in iodine content than the soils further inland¹⁹.

IDD has a strong relationship with a geographical factor because it is generally found in highland areas among school age children. This age group is very susceptible to iodine intakes obtained from the environment. School-age children are considered as an appropriate target group for determining iodine deficiency due to their susceptibility to iodine deficiency, easily accessibility as a study group and representativeness of their community society as a whole²⁰. Some studies demonstrated that the occurrence of goiter was due to a lack of iodine intake and generally found in highland areas. A study conducted at Kayseri, Turkey, where the goiter is endemic, goiter prevalence was 54.8% and median urinary iodine level was 9.54 µg/dl indicating mild iodine deficiency²¹. The prevalence of goiter among children 6-12 years old in Lay Armachiho Northwest Ethiopia was found to be 37.6%, whereas 70.3% of the subjects had inadequate iodine content (<15 ppm)²⁰. A study in high altitude areas of Saudi Arabia revealed the goiter prevalence was 7.4% and about 71% of the participants had UIC less than 100µg/L²². Other studies demonstrated that goiter cases were found in areas with sufficient iodine intakes (non-endemic areas) such as coastal or lowland areas. Study at adult more than 18 years old in an iodine-sufficient area in Chengdu China revealed prevalences of goiter was 8.8%²³. A study by Suhartono in District of Brebes found pesticide exposure as a risk factor for hypothyroidism among women at childbearing age living in an agricultural area, Brebes District, Indonesia²⁴.

The proportion of goiter and hypothyroidism was higher among farmworkers' children than among non-farmworker's children (Table 4). However, Iodine intakes in all research participants were sufficient. Pesticides might be a risk factor for goiter. A pesticide is a chemical widely used to increase agricultural products and to decrease food-borne or vector-borne diseases²⁵. The use of a synthetic pesticide can contaminate soil, water, grass, and other vegetation. In addition to killing the insects or weeds, pesticides can be toxic to a number of other organisms, including birds, fish, beneficial insects, and non-target plants. Insecticides are generally the most acute toxic class of pesticides, herbicides but can also pose a risk to non-target organisms²⁶. Pesticide residues leftover from agriculture not only contaminated crops but also the environment, such as ambient air, surface water, and soil. These findings reinforce the concern about pollution by organophosphates in areas surrounding agriculture areas of pesticide use²⁷. Ideally, toxic effects of pesticide are on target organisms (pests) but in fact, the toxicity of most of the active ingredients of pesticide is not specific. Therefore, it is very harmful to human health²⁸.

Pesticides are categorized as endocrine disrupting chemicals (EDCs)²⁹. Exposure by EDCs can disrupt thyroid function that is well-known as thyroid disrupting chemicals (TDCs)³⁰. Thyroid dysfunction has an impact on growth and development of children². EDCs can bind and activate a variety of hormone receptors and then mimics the action of natural hormones (agonist action). EDCs also can bind to the receptors without activating them. Antagonist action would block and inhibit the action of the receptors. In addition, EDCs also can interfere with the synthesis, transport, metabolism and elimination of hormones, thus reducing the concentrations of natural hormones³¹.

CONCLUSIONS

The proportion of goiter and hypothyroidism among children was two times higher in subjects whose fathers as farmworkers than non-farmworkers although the iodine intake was adequate.

Conflict of Interest: The authors declare that they have no conflict of interest.

Source of Funding: This study was funded by Faculty of Public Health, Diponegoro University, Semarang, Indonesia.

REFERENCES

1. Guyton AC and Hall JE. *Textbook of Medical Physiology*. 12th ed. Philadelphia: Saunders, Elsevier; 2011.
2. Stone MB and Wallace RB. *Medicare Coverage of Routine Screening for Thyroid Dysfunction*. Washington DC: The National Academic Press; 2003. <http://www.nap.edu/catalog/10682.html>.
3. American Thyroid Association. www.thyroid.org. http://www.thyroid.org/wp-content/uploads/patients/brochures/Goiter_brochure.pdf. Published 2016. Accessed March 14, 2017.
4. Dunn JT. Iodine should be routinely added to complementary foods. *J Nutr*. 2003;133:3008S-3010S.
5. Eastman C. Iodine Deficiency Disorders Continue to be a Problem in the Asia Pacific Region. *J ASEAN Fed Endocr Soc*. 2012;27(2):170-175. doi:10.15605/jafes.027.02.06.
6. Sulchan M. Goiter in the coastal areas (case study in Pati Regency): an ecological nutrition problem? *Indones J IDD*. 2007;6(1):17-22.
7. Wade M, Parent S, Finnson K, et al. Thyroid toxicity due to subchronic exposure to a complex mixture of 16 organochlorines, lead, and cadmium. *Toxicol Sci*. 2002;67(2):207-218. doi:10.1093/toxsci/67.2.207.
8. Henson M, Chedrese P. Endocrine disruption by cadmium, a common environmental toxicant with paradoxical effects on reproduction. *Exp Biol Med*. 2004;229(5):383-392.
9. World Health Organisation. Assessment of the iodine deficiency disorders and monitoring their elimination. *WHO, Geneva*. 2007:1-107. doi:ISBN 978 92 4 159582 7.
10. Lemeshow S, Hosmer Jr D, Klar J, Lwanga S. *Adequacy of Sample Size in Health Studies*. New York: John Wiley & Sons Ltd.; 1990.
11. Central Statistics Office. *Brebes Regency in Figures*. Brebes; 2016.
12. WHO/NMH/NHD/EPG/14.5. Goitre as a determinant of the prevalence and severity of iodine deficiency disorders in populations. 2014:14-19.
13. Peterson S, Sanga a, Eklöf H, et al. Classification of thyroid size by palpation and ultrasonography in field surveys. *Lancet*. 2000;355(9198):106-110. doi:10.1016/S0140-6736(99)07221-9.
14. World Health Organization. Indicators for assessing iodine deficiency disorders and their control through salt iodization. 1994:1-55.
15. Pearce EN, Andersson M, Zimmermann MB. Global Iodine Nutrition: Where Do We Stand in 2013? *Thyroid*. 2013;23(5):523-528. doi:10.1089/thy.2013.0128.
16. Zimmermann M, Boelaert K. Iodine deficiency and thyroid disorders. *Lancet Diabetes Endocrinol*. 2015;3(4):286-295. doi:10.1016/S2213-8587(14)70225-6.
17. Zimmermann M. The role of iodine in human growth and development. *Semin Cell Dev Biol*. 2011;22(6):645-652. doi:10.1016/j.semcdb.2011.07.009.
18. Zimmermann M, Andersson M. Assessment of iodine nutrition in populations: Past, present, and future. *Nutr Rev*. 2012;70(10):553-570. doi:10.1111/j.1753-4887.2012.00528.x.
19. Patrick L. Iodine: deficiency and therapeutic considerations. *Altern Med Rev*. 2008;13(2):116-127 12p. <http://search.ebscohost.com/login.aspx?direct=true&db=c8h&AN=105798795&%5Cnlang=ja&site=ehost-live>.
20. Mesele M, Degu G, Gebrehiwot H. Prevalence and associated factors of goiter among rural children aged 6-12 years old in Northwest Ethiopia, cross-sectional study. *BMC Public Health*. 2014;14(1):130. doi:10.1186/1471-2458-14-130.
21. Bayram F, Borazan B, Torun E, et al. The Prevalence of Endemic Goiter and Iodine Deficiency and Evaluation of Thyroid Functions in an Area of Central Anatolia. *Turkish J Endocrinol Metab*. 2003;1:37-43.
22. Omar MS, Desouky DES. Environmental, urinary iodine status and prevalence of goitre among schoolchildren in a high altitude area of Saudi

- Arabia. *Pakistan J Med Sci.* 2015;31(2):414-419. doi:10.12669/pjms.312.6637.
23. Liu Y, Huang H, Zeng J, Sun C. Thyroid volume, goiter prevalence, and selenium levels in an iodine-sufficient area: a cross-sectional study. *BMC Public Health.* 2013;13(37):1153. doi:10.1186/1471-2458-13-1153.
 24. Suhartono, Djokomoeljanto R, Hadisaputro S, Subagio H, Kartini A, Suratman. Paparan pestisida sebagai faktor risiko hipotiroidisme pada wanita usia subur di daerah pertanian. *Media Med Indones.* 2012;46(2):91-99.
 25. Weiss B, Amler S, Amler R. Pesticides. *Pediatrics.* 2004;113(4):1030-1036.
 26. Aktar MW, Sengupta D, Chowdhury A. Impact of pesticides use in agriculture: their benefits and hazards. *Interdiscip Toxicol.* 2009;2(1):1-12. doi:10.2478/v10102-009-0001-7.
 27. Harnpicharnchai K, Chaiear N, Charerntanyarak L. Residues of organophosphate pesticides used in vegetable cultivation in ambient air, surface water and soil in Bueng Niam subdistrict, Khon Kaen, Thailand. *Southeast Asian J Trop Med Public Health.* 2013;44(6):1088-1097.
 28. Costa L. Toxic effects of pesticides. In: *Toxicology: The Basic Science of Poisons.* ; 2008:883-930.
 29. Diamanti-Kandarakis E, Bourguignon J, Giudice L, et al. Endocrine-Disrupting Chemicals: An Endocrine Society Scientific Statement. *Endocr Rev.* 2009;30(4):293-342. doi:10.1210/er.2009-0002.
 30. Crofton KM. Thyroid disrupting chemicals: Mechanisms and mixtures. *Int J Androl.* 2008;31(2):209-222. doi:10.1111/j.1365-2605.2007.00857.x.
 31. Mnif W, Hassine A, Bouaziz A, Bartegi A, Thomas O, Roig B. Effect of endocrine disruptor pesticides: A review. *Int J Environ Res Public Health.* 2011;8(6):2265-2303. doi:10.3390/ijerph8062265.