LEMBAR HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW KARYA ILMIAH : PROSIDING

Judul Karya Ilmiah (Prosiding)	:	Zeolite/Magnetite Composites as	Cata	lyst on the Synthesis of Methyl Ester (MES) from
		Cooking Oil		
Nama/ Jumlah Penulis	:	Sriatun, Adi Darmawan, Sriyanti,	Wu	ri Cahyani, Hendri Widyandari
Status Pengusul	:	Penulis pertama		
Identitas Prosiding	:	a. Judul Prosiding	:	IOP Conference Series: Journal of Physics
		b. ISBN/ISSN	:	1742-6588, E-ISSN:1742-6596
		c. Thn Terbit, Tempat Pelaks.	:	2018/1025/012135
		d. Penerbit/Organiser	:	IOP Publishing
		e. Alamat Repository/Web	:	https://iopscience.iop.org/journal/1742-6596
		Alamat Artikel	:	https://iopscience.iop.org/article/10.1088/1742-
				<u>6596/1025/1/012135/pdf</u>
				Url Turnitin (17%)

Kategori Publikasi Makalah (beri √pada kategori yang tepat) f. Terindeks di (jika ada)

:

Prosiding Forum Ilmiah Internasional
 Prosiding Forum Ilmiah Nasional

https://doc-

:

SCOPUS (SJR 0,221)

pak.undip.ac.id/2874/27/Turnitin%20C27.pdf

Hasil Penilaian Peer Review :

	Nilai Mak	Nilai Maksimal Prosiding				
Komponen Yang Dinilai	Internasional	Nasional	Yang Diperoleh			
a. Kelengkapan unsur isi prosiding (10%)	3		3			
b. Ruang lingkup dan kedalaman pembahasan (30%)	9		9			
c. Kecukupan dan kemutahiran data/informasi dan metodologi (30%)	9		8,6			
d. Kelengkapan unsur dan kualitas terbitan /prosiding (30%)	9		8,6			
Total = (100%)	30,00		29,2			
Penulis Utam	a: 0,6 x 29,2 = 17,5	2	·			

Catatan Penilaian Paper oleh Reviewer :

 Kesesuaian dan kelengkapan unsur isi prosiding: Paper ini telah berisi unsur-unsur penting dalam karya ilmiah, hubungan antara judul, abstak, pendahuluan sampai pembahasan telah sesuai. Nilai 3
 Ruang lingkup dan kedalaman pembahasan:

2. Ruang lingkup dan kedalaman pembanasan: Ruang lingkup tentang komposit zeolit/magnetit sebagai katalis dalam sintesis MES dari minyak goreng bekas. Pembahasan dibahas dengan baik dan detil. Nilai 9

- 3. Kecukupan dan kemutakhiran data/informasi dan metodologi: Data disampaikan dengan baik dan lengkap dan cukup mutakhir, metodologi cukup detil. Nilai 8,6
- 4. Kelengkapan unsur dan kualitas terbitan/prosiding:
- Unsur terbitan lengkap, kualitas terbitan prosiding cukup baik. Nilai 8,6

Semarang, 3 April 2020 Reviewer 1

Dr. Bambang Cahyono, MS NIP. 196303161988101001 Unit Kerja : Departemen Kimia FSM UNDIP

LEMBAR HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW KARYA ILMIAH : PROSIDING

Judul Karya Ilmiah (Prosiding)	: Zeo Coo	lite/Magnetite Composites as obtain Oil	Cata	alyst on the Synthesis of Methyl Ester`(MES) from
Nama/ Jumlah Penulis	: Sria	atun. Adi Darmawan. Srivanti.	Wu	ri Cahvani. Hendri Widvandari
Status Pengusul	: Pen	ulis pertama		
Identitas Prosiding	: a.	Judul Prosiding	:	IOP Conference Series: Journal of Physics
	b.	ISBN/ISSN	:	1742-6588, E-ISSN:1742-6596
	с.	Thn Terbit, Tempat Pelaks.	:	2018/1025/012135
	d.	Penerbit/Organiser	:	IOP Publishing
	e.	Alamat Repository/Web	:	https://iopscience.iop.org/journal/1742-6596
		Alamat Artikel	:	https://iopscience.iop.org/article/10.1088/1742-
				6596/1025/1/012135/pdf
				Url Turnitin (9%)
				http://eprints.undip.ac.id/80669/26/C26 Zeolitem
				agnetite_composites_as_catalysts_on_the_Synthe
				sis_of_Methyl_Esters_(MES)_from_cooking_oil.
				<u>pdf</u>
	f.	Terindeks di (jika ada)	:	SCOPUS (SJR 0,221)
Kategori Publikasi Makalah (beri √pada kategori yang tepat)	:	Prosiding Forum Ilmia	ah Ir ah N	nternasional Jasional

(beri ✓ pada kategori yang tepat)

Hasil Penilaian Peer Review :

	Nilai Mak	simal Prosiding	Nilai Akhir
Komponen Yang Dinilai	Internasional	Nasional	Yang Diperoleh
a. Kelengkapan unsur isi prosiding (10%)	3		3
b. Ruang lingkup dan kedalaman pembahasan (30%)	9		9
 Kecukupan dan kemutahiran data/informasi dan metodologi (30%) 	9		8
d. Kelengkapan unsur dan kualitas terbitan /prosiding (30%)	9		8
Total = (100%)	30,00		28
Penulis Utam	a: 0,6 x 28= 16,8		

Catatan Penilaian Paper oleh Reviewer :

1. Kesesuaian dan kelengkapan unsur isi prosiding: Unsur isi prosiding lengkap dan sesuai dengan standart IOP Publishing. Nilai 3 2. Ruang lingkup dan kedalaman pembahasan:

Ruang lingkup artikel ini adalah katalis komposit zeolit/magnetit untuk sintesis ME dari minyak goreng bekas dengan keterbaruan yang tinggi. Pembahasan diulas dengan baik (dalam) disertai literartur pendukung. Nilai 9

3. Kecukupan dan kemutakhiran data/informasi dan metodologi: Data yang disajikan cukup memadai sebagai artikel dan informasi yang disampaikan ada yang tahun kemautakhirannya kurang, metodologi diulas dengan detil sehingga peneliti lain bisa mengulang dengan baik. Nilai 8

4. Kelengkapan unsur dan kualitas terbitan/ prosiding: Unsur terbitan lengkap, kualitas terbitan ada yang kurang terbaca (gambar) da salah penulisan (typo). Nilai 8

Semarang, 14 Maret 2020 Reviewer 2

Drs. Gunawan, M.Si, Ph.D NIP.196408251991031001 Unit Kerja : Departemen Kimia FSM UNDIP

LEMBAR HASIL PENILAIAN SEJAWAT SEBIDANG ATAU *PEER REVIEW* KARYA ILMIAH : PROSIDING

Judul Karya Ilmiah (Prosiding) Nama/ Jumlah Penulis Status Pengusul Identitas Prosiding	: Ze Est : Sri : Per : a. b. c. d. e.	olite/Magnetite Composites ter`(MES) from Cooking Oi atun, Adi Darmawan, Sriyan nulis pertama Judul Prosiding ISBN/ISSN Thn Terbit, Tempat Pelaks. Penerbit/Organiser Alamat Repository/Web Alamat Artikel	as (1 nti, : : :	Catalyst on the Synthesis of Methyl Wuri Cahyani, Hendri Widyandari IOP Conference Series: Journal of Physics 1742-6588, E-ISSN:1742-6596 2018/1025/012135 IOP Publishing https://iopscience.iop.org/journal/1742-6596 https://iopscience.iop.org/article/10.1088/174 2-6596/1025/1/012135/pdf Url Turnitin (9%) http://eprints.undip.ac.id/80669/26/C26 Zeolitem agnetite_composites_as_catalysts_on_the_Synthe sis_of_Methyl_Esters_(MES)_from_cooking_oil. pdf
	f.	Terindeks di (jika ada)	:	SCOPUS (SJR 0,221)

Kategori Publikasi Makalah (beri ✓ pada kategori yang tepat) : V Prosiding Forum Ilmiah Internasional Prosiding Forum Ilmiah Nasional

Hasil Penilaian Peer Review :

	Nilai l		
Komponen Yang Dinilai	Reviewer I	Reviewer II	Nilai Rata- rata
a. Kelengkapan unsur isi prosiding (10%)	3	3	3
b. Ruang lingkup dan kedalaman pembahasan (30%)	9	9	9
c. Kecukupan dan kemutahiran data/informasi dan metodologi (30%)	8,6	8	8,3
d. Kelengkapan unsur dan kualitas terbitan/prosiding(30%)	8,6	8	8,3
Total = (100%)	29,2	28	28,6
Penulis Utama(rata-rata): 0,6	6 x 28,6 = 17,16		

Semarang, 3 April 2020

Reviewer 1

Dr. Bambang Cahyono, MS NIP. 196303161988101001 Unit Kerja : Departemen Kimia FSM UNDIP

Reviewer 2

Drs. Gunawan, M.Si, Ph.D NIP. 196408251991031001 Unit Kerja : Departemen Kimia FSM UNDIP

THE MINISTRY OF RESEARCH, TECHNOLOGY AND HIGHER EDUCATION THE REPUBLIC OF INDONESIA DIPONEGORO UNIVERSITY

FACULTY OF SCIENCE AND MATHEMATICS

CERTIFICATE

Decree of Dean Number : 1440/UN7.5.8/HK/2017

This is to certify that

Sriatun

as

PRESENTER

In the 7th International Seminar on New Paradian, and Innovation of Natural Science and Its Application (ISNPINSA-7) held on 17 October 2014 at Grand Candi Hotel Semarang Indonesia

with paper entrancies follows:

Zeolite / magnetite composites as catalysts on the synthesis of methyl esters (MES) from cooking oil

Prof. Dr. Widowati, S.Si M.Si. NIP 195902141994032002 7 ISNPINSA COMMITTEE, CHAIRMAN

Dr. Bull Warsito, S.Si, M.Si. NIP. 197508241999031003

PAPER • OPEN ACCESS

The 7th International Seminar on New Paradigm and Innovation on Natural Science and Its Application

To cite this article: 2018 J. Phys.: Conf. Ser. 1025 011001

View the article online for updates and enhancements.

Related content

- Workshop and International Seminar on Science of Complex Natural Systems
- Once a physicist: Jon Palfreman
- International Seminar on Mathematics and Physics in Sciences and Technology 2017 (ISMAP 2017)



IOP ebooks[™]

Bringing together innovative digital publishing with leading authors from the global scientific community.

Start exploring the collection-download the first chapter of every title for free.

PREFACE

The 7th International Seminar on New Paradigm and Innovation on Natural Sciences and Its Application (ISNPINSA-7) is annual conferences organized by Faculty of Sciences and Mathematics (FSM) Diponegoro University and has been successfully conducted since 2011. The aims of ISNPINSA are to facilitate brain storming and state of the art information in field of sciences and mathematics; to increase innovation of technology that can be applied in industries; to contribute in formulating strategy to increase the role of science for community; and to stimulate collaboration between industries, researchers and government to increase community welfare. The theme of 7th ISNPINSA in 2017 is "Science and Data Science for Sustainable Development Goals".

The scope of the field of participants comes from various fields including biology, physics, chemistry, statistics, mathematics, informatics, environment, public health, and relevant fields that contribute to sustainable development. The conference was held in Semarang, Indonesia on October, 17th, 2017. There were three keynote speakers and three invited speaker who came from Japan, Italy, Malaysia, Philipines and Indonesia. The number of participants of this seminar were more than 200 consist of researchers, lecturers, postgraduate and undergraduate students from various universities and after the selection process there are 132 articles selected to be published in the present conference proceeding.

The Editors

Dr. Budi Warsito Sapto Purnomo Putro, Ph.D. Ali Khumaeni, Ph.D.

LIST OF REFEREES

- 1. Prof. Widowati
- 2. Prof. Mustafid
- 3. Prof. Wahyu Setia Budi
- 4. Sapto P Putro, PhD
- 5. Dr. Munifatul Izzati
- 6. Dr. Budi Warsito
- 7. Dr. Eng. Ali Khumaeni
- 8. Dr. Di Asih I Maruddani
- 9. Dr. Muhammad Nur, DEA
- 10. Hendri Widyandari, PhD
- 11. Dr. Kusworo Adi
- 12. Dr. Heri Sutanto
- 13. Dr. Endang Kusdiyantini, DEA
- 14. Dr. Jafron W. Hidayat
- 15. Rully Rahadian, PhD
- 16. Anto Budiharjo, PhD
- 17. Dr. Tri Retnaningsih Soeprobowati
- 18. Ismiyarto, PhD
- 19. Dr. Retno Ariadi Lusiana
- 20. Dr. Tarno, MSi
- 21. Adi Wibowo, PhD

LIST OF SPEAKERS

Keynote Speaker:

Prof. Dr. Masahiko Tani	University of Fukui, Japan
Prof. Dr. Norsarahaida Saidina Amin	Universiti Teknologi Malaysia
Mario Rosario Guarracino, PhD.	Instituto di Calcolo e Reti ad Alte Prestazioni-
	National Research Council (ICAR-CNR), Italy
Dr. dr. Budi Wiweko, Sp.OG-KFER.	University of Indonesia, Jakarta
Invited Speaker:	
Prof. Elmer S. Estacio, PhD.	National Institute of Phyics, University of the
Ismiyarto, S.Si., M.Si., PhD. Dr. Eng. Adi Wibowo, S.Si., M.Kom.	Philippines, Manila, Diponegoro University, Semarang, Indonesia Diponegoro University, Semarang, Indonesia

LIST OF COMMITTEES

Steering Committee

Prof. Muhammad Zainuri, Diponegoro University, Indonesia Prof. Widowati, S.Si., M.Si., Diponegoro University, Indonesia Prof. Heru Susanto, Diponegoro University, Indonesia Sapto Purnomo P, Ph.D, Diponegoro University, Indonesia Drs. Bayu Surarso, M.Sc., Ph.D., Diponegoro University, Indonesia

Scientific Committee

Prof. Norsarahida Saidina Amin, Universiti Teknologi Malaysia
Prof. Dr. Masahiko Tani, University of Fukui, Japan
Mario Rosario Guarracino, Ph.D, ICAR-CNR
Prof. Elmer S. Estacio, University of the Philippines Diliman
Dr. dr. Budi Wiweko, Sp.OG-KFER, University of Indonesia
Prof. Mustafid, Ph.D., Diponegoro University, Indonesia
Prof. Wahyu Setia Budi, Diponegoro University, Indonesia
Dr. Munifatul Izzati, Diponegoro University, Indonesia
Dr. Muhammad Nur, DEA, Diponegoro University, Indonesia
Ismiyarto, Ph.D., Diponegoro University, Indonesia
Adi Wibowo, Ph.D., Diponegoro University, Indonesia

Organizing Committee

Chairman: Dr. Budi Warsito, M.Si. Vice Chairman: Dr. Eng. Ali Khumaeni Secretary: Dr. Di Asih I Maruddani, M.Si.

Members:

Yayuk Astuti, Ph.D Nurdin Bahtiar, S.Si., M.Kom. Rismiyati, B.Eng, M.Cs. Pandji Triadyaksa, SSi, M.Sc. Nur Azizah, SE Joko Rustianto, S.IP. Iys Syabilla Rusda, SIP Fajar Budi Handoyo

Secretariat:

Alik Maulidiyah, M.Sc.

Treasurer:

Titik Eryanti, SE Choiriyah, SE

Sections

Program:

Drs. Agus Setyo Utomo, S.Sos, MM & Team

Food: Dra. Sri Harumaningsih, S.IP. & Team

Papers, Presentations and Proceeding:

Rahmawan Bagus Trianto, S.Kom. Nur Rohmat, Amd

Transportation and Accomodation: Deby Yuniarto & Team

Public Relation and Documentation: Joko Rustianto, S.IP. & Team

This site uses cookies. By continuing to use this site you agree to our use of cookies. To find out more, see our Privacy and Cookies policy. ፡

□ NOTICE: Ensuring subscriber access to content on IOPscience throughout the coronavirus outbreak - see our remote access guidelines.

Table of contents

Volume 1025 2018

Next issue > • Previous issue

The 7th International Seminar on New Paradigm and Innovation on Natural Science and Its Application 17 October 2017, Semarang, Indonesia

Accepted papers received: 04 May 2018 Published online: 30 May 2018

View all abstracts

OPEN ACCESS			01100
The 7th Internation	onal Seminar on No	ew Paradigm and Innovation on Natural Science and Its	
	View article	PDF	
OPEN ACCESS			011002
Peer review state	ment		
	View article	🔁 PDF	
Papers			
Material Physiscs	;		
OPEN ACCESS			012001
Dye-sensitized so	olar cell simulation	performance using MATLAB	
Alvin Muhammad I	Habieb, Muhammad I	rwanto, Ilham Alkian, Fitri Khalimatus Sya'diyah, Hendri Widiyan	dari and

	View article	🔁 PDF
--	--------------	-------

OPEN ACCESS Lipid production from tapioca wasters simultaneous BOD_COD and pitroe	ewater by culture of <i>Scenedesmus sp</i> . with	012075
Romaidi Muhammad Hasanudin Khusn	ul Kholifah, Alik Maulidiyah, Santo P. Putro, Akira Kikuchi at	nd
Toshifumi Sakaguchi		
+ Open abstract View article	🔁 PDF	
OPEN ACCESS		012076
The bacterial diversity associated with Fab.) from Pemalang Coast, Indones	ith bacterial diseases on Mud Crab (Scylla serrata sia	
Sarjito, Desrina, AHC Haditomo and S. F	Budi Prayitno	
← Open abstract	🔁 PDF	
OPEN ACCESS		012077
The effect of organic quail egg supp (<i>Rattus Norvegicus</i> L.) during the la	lementation on the blood lipid profile of white mice ctation period	
Sri lestari purba, Tyas Rini Saraswati and	Sri Isdadiyanto	
+ Open abstract	🔁 PDF	
Applications of bloactive material fi learning-memory capability and more caused oxidative stress in rats	toric activity: a case study of physiological aging and a	ging-
+ Open abstract View article	PDF	
OPEN ACCESS	of Dongue Heemorphagie favor in Semerang City	012079
Impact of Chinate of the Incidence of	asih. Subartono and Hansari	
+ Open abstract View article	PDF	
OPEN ACCESS Effect of blood estrogen and proget	corona an soverity of minor PAS	012080
S Literri T W D Dehendie A Deried T T	A hundring on d E L Assertioni	
+ Open abstract ■ View article	PDF	
OPEN ACCESS Correlation between macrobenthic s (abiotic) adjacent aquaculture areas	tructure (biotic) and water-sediment characteristics at Tembelas Island, indonesia	012081
Jeanny Sharani Jafron W Hidayat and S	anto P. Putro	

	SOC) wi-fi microco	ontroller for multistation measurement of water surface	012029
Survono Survono S	Sonic Sensor	Widowati and Satrivo Adhy	
View abstract	View article	PDF	
OPEN ACCESS			012030
Analysis of open Film	ings and wide of lea	af on multileaf Colimators Using Gafchromic RTQA2	
Evi Setiawati, Assy	vifa Lailla Rachma and	l M. Hidayatullah	
	View article	PDF	
Ecology and Biod	liversity		
OPEN ACCESS Potential Re-utili Improvement	ization of Composte	ed Mangrove Litters for Pond Environment Quality	012031
Endah Dwi Hastuti	, Rini Budi Hastuti and	d Riche Hariyati	
	View article	PDF	
OPEN ACCESS The abundance a and its implicatic Erry Wiryani, Murr	nd importance valu on to the spring ningsih and Jumari	e of tree in "Sendang Kalimah Toyyibah" surrounding	012032
OPEN ACCESS The abundance a and its implication Erry Wiryani, Murre + View abstract	nd importance valu on to the spring ningsih and Jumari View article	e of tree in "Sendang Kalimah Toyyibah" surrounding	012032
 OPEN ACCESS The abundance a and its implication Erry Wiryani, Murret View abstract OPEN ACCESS The invasive aliee Ontoloe Island, F 	nd importance valu on to the spring ningsih and Jumari Twiew article on plants threatened Flores-Indonesia	e of tree in "Sendang Kalimah Toyyibah" surrounding PDF the balance of ecosystem in conservative area in	012032
OPEN ACCESS The abundance a and its implication Erry Wiryani, Murre + View abstract OPEN ACCESS The invasive aliee Ontoloe Island, F Ignatius Antonius M	nd importance valu on to the spring ningsih and Jumari To View article on plants threatened Flores-Indonesia Mboka Segu Wake, Tri	e of tree in "Sendang Kalimah Toyyibah" surrounding PDF the balance of ecosystem in conservative area in i Retnaningsih Soeprobowati and Jumari	012032
 OPEN ACCESS The abundance a and its implication Erry Wiryani, Murret View abstract OPEN ACCESS The invasive aliee Ontoloe Island, F Ignatius Antonius M View abstract 	and importance valu on to the spring ningsih and Jumari Toiew article on plants threatened Flores-Indonesia Mboka Segu Wake, Tri	e of tree in "Sendang Kalimah Toyyibah" surrounding PDF the balance of ecosystem in conservative area in i Retnaningsih Soeprobowati and Jumari PDF	012032
 OPEN ACCESS The abundance a and its implication Erry Wiryani, Murn View abstract OPEN ACCESS The invasive aliee Ontoloe Island, F Ignatius Antonius N View abstract OPEN ACCESS Extended benefitt forest ecosystem 	Ind importance value on to the spring ningsih and Jumari I View article on plants threatened Flores-Indonesia Mboka Segu Wake, Tri I View article View article	e of tree in "Sendang Kalimah Toyyibah" surrounding PDF the balance of ecosystem in conservative area in i Retnaningsih Soeprobowati and Jumari PDF instrument of economic valuated in Petungkriyono	012032
 OPEN ACCESS The abundance a and its implication Erry Wiryani, Murn View abstract OPEN ACCESS The invasive aliee Ontoloe Island, H Ignatius Antonius M View abstract OPEN ACCESS Extended benefitt forest ecosystem Irma Damayanti, A 	Ind importance value on to the spring ningsih and Jumari I View article on plants threatened Flores-Indonesia Mboka Segu Wake, Tri I View article Cost analysis as an services zis Nur Bambang and	e of tree in "Sendang Kalimah Toyyibah" surrounding PDF the balance of ecosystem in conservative area in i Retnaningsih Soeprobowati and Jumari PDF instrument of economic valuated in Petungkriyono Tri Retnaningsih Soeprobowati	012032
 OPEN ACCESS The abundance a and its implication Erry Wiryani, Murret View abstract OPEN ACCESS The invasive aliee Ontoloe Island, H Ignatius Antonius N View abstract OPEN ACCESS Extended benefitt forest ecosystem Irma Damayanti, A View abstract 	Ind importance value on to the spring ningsih and Jumari I View article on plants threatened Flores-Indonesia Mboka Segu Wake, Tri I View article t cost analysis as an services zis Nur Bambang and I View article	e of tree in "Sendang Kalimah Toyyibah" surrounding PDF the balance of ecosystem in conservative area in i Retnaningsih Soeprobowati and Jumari PDF instrument of economic valuated in Petungkriyono Tri Retnaningsih Soeprobowati PDF	012032

020	Journ	al of Physics: Conference Series, Volume 1025, 2018 - IOPscience	
Application of In Growth (Zea Mag	organic Fertilizer V ys L.)	With NanoChisil and Nanosilica on Black Corn Plant	012128
Lolita Dewi Pertam	niningsih, Erma Prihas	tanti, Sarjana Parman, Agus Subagio and Ngadiwiyana	
+ View abstract	View article	🔁 PDF	
OPEN ACCESS			012129
Synthesis of Silic	con Carbide Fiber a	s semiconductor substrate for Betavoltaic cell	
Masripah, Agustino	• Zulys and Jan Setiaw	van	
	View article	🔁 PDF	
OPEN ACCESS			012130
Total flavonoid c	ontent and formula	tion antioxidant cream stem of <i>jatropha multifida</i> l.	
Yuvianti Dwi Frany	voto, Lia Kusmita, Mu	atmainah and Riega Demma Angrena	
+ View abstract	View article	🔁 PDF	
OPEN ACCESS			012131
Ammonium Rem Using <i>Ecotru</i> Inc	noval Efficiency in a social second	Fixed Bed Reactor and Suspended Growth Reactor	
Rusyda Syahidah, S	Sudarno and Badrus Z	aman	
+ View abstract	View article	PDF	
OPEN ACCESS			012132
Antioxidant activ	vity from encapsula	ted Cinnamaldehyde-Chitosan	
Bonita Ariestiani, P	Purbowatingrum, Ngao	liwiyana, Ismiyarto, Enny Fachriyah and Khikmah Nurani	
	Tiew article	🔁 PDF	
OPEN ACCESS			012133
Sulfonated polys	tyrene and its chara	cterization as a material of electrolyte polymer	
Ngadiwiyana, Ismi	yarto, Gunawan, RS P	urbowatiningrum, N.B.A. Prasetya, T.D. Kusworo and H. Susanto	
	View article	🔁 PDF	
OPEN ACCESS			012135
Zeolite/magnetite	e composites as cata	alysts on the Synthesis of Methyl Esters (MES) from	
Sriatun, Adi Darma	wan, Sriyanti, Wuri C	Cahyani and Hendri Widyandari	
	View article	🔁 PDF	
JOURNAL LINK	XS		

Journal home

4/18/2020

IOP Conference Series: Materials Science and Engineering - IOPscience

N V Danilina and S V Privezentseva 2020 IOP Conf. Ser.: Mater. Sci. Eng. 775 012008			
Specific features of diaphragm wall construction			
D A Mironov et al 2020 IOP Conf. Ser.: Mater. Sci. Eng. 775 012051			
+ View abstract 🔄 View article 🄁 PDF			
Media technologies in shaping urban identity			
T V Karakova et al 2020 IOP Conf. Ser.: Mater. Sci. Eng. 775 012038			
+ View abstract View article PDF			

JOURNAL LINKS

Journal home

Information for organizers

Information for authors

Search for published proceedings

Contact us

Reprint services from Curran Associates

2009-present IOP Conference Series: Materials Science and Engineering doi:10.1088/issn.1757-899X Online ISSN: 1757-899X Print ISSN: 1757-8981

Source details

Journal of Physics: Conference Series	CiteScore 2018	(i)
Scopus coverage years: from 2005 to Present	0.51	Ŭ
Publisher: Institute of Physics Publishing		
ISSN: 1742-6588 E-ISSN: 1742-6596	SJR 2018	ŝ
Subject area: (Physics and Astronomy: General Physics and Astronomy)	0.221	0
View all documents > Set document alert Save to source list Journal Homepage	SNIP 2018	Û
	0.454	U

CiteScore	CiteScore rank & trend Ci	teScore presets Scopus content coverage		
CiteScore	e 2018 ~	Calculated using data from 30 April, 2019	CiteScore rai	nk 🛈
0 5 1	Citation Count 2018	11,243 Citations >	Category	Rank Percentile
0.51	= Documents 2015 - 2017*	= 21,896 Documents >	Physics and Astronomy	#167/216 21st
*CiteScore incl	udes all available document types	View CiteScore methodology > CiteScore FAQ >	General Physics and Astronomy	
CiteScore	eTracker 2019 🛈	Last updated on <i>09 April, 2020</i> Updated monthly	View CiteScore tree	nds > our site &
0 50	Citation Count 2019	18,254 Citations to date >	,	0
0.37	= Documents 2016 - 2018	3 31,135 Documents to date>		

Metrics displaying this icon are compiled according to Snowball Metrics a, a collaboration between industry and academia.

About Scopus	Language	Customer Service
What is Scopus	日本語に切り替える	Help
Content coverage	切换到简体中文	Contact us
Scopus blog	切換到繁體中文	
Scopus API	Русский язык	
Privacy matters		

ELSEVIER

Terms and conditions a Privacy policy a



view

Author search Sources

⑦ 🟦 Create account

Int Sign in

Sriatun,

Universitas Diponegoro, Semarang, Indonesia
 Connect to ORCID
 View Mendeley profile

🤌 Edit profile 🛛 🗘 Set alert 🔗 Potential author matches 🕞 Export to SciVal

Metrics overview





Most contributed Topics 2015-2021 ()

Nanofiltration Membranes; Pervaporation; Water Flux 2 documents Transesterification; Cooking Fats and Oils; Rubber Seed Oil 1 document Coal Fly Ash; Zeolite; Kaolin 1 document

View all Topics

	8 Documents	Cited by 26 Documents	14 Co-Authors	Topics		
	Note: Scopus Preview users	s can only view an author's last 10	documents, while mos	t other features are	e	
	disabled. Do you hav documents and featu	re access through your institution ures.	Check your institution	's access to view al	I	
	Export all Add all t	o list		2	Sort by Date (newest)	~
> View list in search results format	Conference Paper • O Characteristic O	^{pen Access} f ZSM-5 catalyst supported	by nickel and mol	ybdenum		0
> View 198 references	Sriatun,, Susanto, H	., Widayat,, Darmawan, A.				Cited by
riangle Set document alert	IOP Conference Series: N	Materials Science and Engineering, 2019,	509(1), 012138			
	View abstract \vee R	elated documents				
	Conference Paper • O	pen Access				
	Zeolite/magneti	te composites as catalysts	on the Synthesis o	f Methyl Esters)	0
	(MES) from cool	king oil				Cited by
	Sriatun,, Darmawan	, A., Sriyanti,, Cahyani, W., Widyaı	ndari, H.			

Journal of Physics: Conference Series, 2018, 1025(1), 012135

Journal of Physics: Conference Series

also developed by scimago:

Viz Tools

SCIMAGO INSTITUTIONS RANKINGS

SJR

Scimago Journal & Country Rank

Journal Rankings

Home

Country Rankings

Help About Us

Enter Journal Title, ISSN or Publisher Name

Journal of Physics: Conference Series 8

Country	United Kingdom - IIII SIR Ranking of United Kingdom	65			
Subject Area and Category	Physics and Astronomy Physics and Astronomy (miscellaneous)	UJ			
Publisher	Institute of Physics	H Index			
Publication type	Journals				
ISSN	426588, 17426596				
Coverage)05-ongoing				
Scope	he open access Journal of Physics: Conference Series (JPCS) provides a fast, versatile and cost-effective proceedings aublication service.				
0	Homepage				
	How to publish in this journal				
	Contact				
	igsirphi Join the conversation about this journal				





4/18/2020

Journal of Physics: Conference Series



R Rose 2 months ago

Dear SCImago Team, It is stated in the JPCS homepage that the journal is a proceeding publication

Artikel Luar Negeri 1 (Japan)

Lipid production from tapioca wastewater by culture of Scenedesmus sp. with simultaneous BOD, COD and nitrogen removal

Romaidi^{1*}, Muhammad Hasanudin¹, Khusnul Kholifah¹, Alik Maulidiyah², Sapto P. Putro², Akira Kikuchi^{3,4}, Toshifumi Sakaguchi⁵

¹Department of Biology, Faculty of Science and Technology, Universitas Islam Negeri Maulana Malik Ibrahim Malang, Indonesia

²Faculty of Science and Mathematics, Diponegoro University, Semarang, Indonesia ³Faculty of Agriculture, University of Brawijaya, Malang, Indonesia

⁴Institute of Environmental and Water Resource Management, Universiti Teknologi Malaysia (UTM)

⁵Faculty of Environmental and Life Science, Prefectural University of Hiroshima, Japan

*Corresponding's author: romaidi@bio.uin-malang.ac.id

Abstract. The use of microalgae to produce biodiesel or possibly remove nutrients from industrial wastewater has gained important attention during recent years due to their photosynthetic rate and its versatile nature to grow in various wastewater systems. In this study, a microalgae, Scenedesmus sp., was cultured to enhance the lipid production and nutrients removal from tapioca wastewater sample. To assess lipid production, Scenedesmus sp. was cultured in different concentration of tapioca wastewater sample (from 0 to 100 %), and nutrient removal including BOD, COD, NH4, NO2, NO3 level by Scenedesmus sp. was assessed in 100% of tapioca wastewater culture. After 8 days of culture, it was found out that 50% of tapioca wastewater sample resulted in highest concentration of lipid content than that of the other concentrations. The level of environment indicator as nutrient removal such as BOD, COD, NH₄, NO₂, NO₃ were also decreased up to 74%, 72%, 95%, 91%, and 91%, respectively. The pH condition changed from initial condition acidic (pH: 4) to neutral or basic condition (pH: 7-8) as recommended in wastewater treatment system. This research provided a novel approach and achieved efficient simultaneous lipid production and nutrients removal from tapioca wastewater sample by Scenedesmus's culture system.

Keyword: Scenedesmus sp., tapioca wastewater, lipid production

1. Introduction

The energy crisis is one of the most important problems faced by all people over the world in the 21st century. The highest consumption of fossil fuels has result in greenhouse effect and causes global climate change [1,2]. One of a type of renewable energy is microalgae biomass-based biofuel, which is considered as one of the most potent substitutes for fossil fuel [2]. However, to increase the production of microalgae biomass, several strategies should be developed, such as modification of culture medium and environmental factors. Hence, one of promising strategies is using wastewater sample as

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd

Zeolite/magnetite composites as catalysts on the Synthesis of Methyl Esters (MES) from cooking oil

Sriatun¹, Adi Darmawan¹, Sriyanti¹, Wuri Cahyani¹, Hendri Widyandari²

¹) Department of Chemistry, Faculty of Science and Mathematics, Diponegoro University

²) Department of Physics Faculty of Science and Mathematics, Diponegoro University Jl. Prof. Soedharto, SH, Tembalang, Semarang 50275, Indonesia

E-mail: sriatun@live.undip.ac.id

Abstract. The using of zeolite/magnetite composite as a catalyst for the synthesis of methyl esters (MES) of cooking oil has been performed. In this study the natural magnetite was extracted from the iron sand of Semarang marina beach and milled by high energy Milling (HEM) with ball: magnetite ratio: 1:1. The composites prepared from natural zeolite and natural magnetite with zeolite: magnetite ratio 1:1; 2:1; 3:1 and 4:1. Preparation of methyl ester was catalyzed by composite of zeolite/magnetite through transeserification reaction, it was studied on variation of catalyst concentration (w/v) 1%, 3%, 5% and 10% to feed volume. The reaction product are mixture of methyl Oleic (MES), methyl Palmitic (MES) and methyl Stearic (MES). Character product of this research include density, viscosity, acid number and iodine number has fulfilled to SNI standard 7182: 2015.

Keywords: composite, zeolite/magnetite, catalyst, MES

1. Introduction

Increased transportation demand leads to an increase in energy demand globally. Transportation sector is a sector that consumes the most energy, so it can increase environmental emission. Transportation using fuel oil is mainly processed petroleum products, while the world's oil reserves and Indonesia began to thin out. In an effort to find solutions to these problems, various studies reveal the existence of a variety of raw materials from renewable resources capable of being processed into a more environmentally friendly fuel as it reduces CO₂ emissions.

Methyl esters are a major component of biodiesel which is the most appropriate solution to replace the fuel fossils as the world's major source of energy transport, because biodiesel is a material fuel that can replace petrol diesel on a machine. Biodiesel is consisting of a mixture of mono-alkyl esters of long chain fatty acids, which are used as an alternative to fuel from diesel engines. Biodiesel or methyl esters are obtained from the transesterification of animal fats, vegetable oils, and even cooking oil waste, with catalytic process [1].

Actually Vegetable oil has a calorific value that is almost the same with conventional fuel, but usage directly as fuel is still encountered obstacles. When compared to biodiesel, vegetable oils have a much higher viscosity value at temperature 40°C. This may cause problems in the pumping and spraying process when they used as fuel [2]. The incandescent point (flash point) also much larger than diesel oil, this is it inhibit the injection process and result incomplete combustion. The use of oil

Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

vegetable as a fuel also leaves residue carbon in the engine injector. Efforts to reduce viscosity and flash point vegetable oils, among others, by dilution of oil with solvents, emulsification, pyrolysis, and transesterification. The transesterification process is the most common way done because it does not require energy and temperature tall one. This reaction will produce methyl or ethyl esters, depending on the type of alcohol reacted. Methanol is used more as a source of alcohol because the chain is shorter, more polar and the price cheaper than other alcohols [3].

Both homogeneous and heterogeneous catalysts can be used in transeterification reactions. Basic homogeneous catalysts such as KOH as well as sodium and potassium alkoxide [3], NaOH [4], while acidic ones such as sulfuric acid, sulfonic acid and hydrochloric acid are usually used as catalysts in transesterification reactions. The using of homogeneous catalysts has a disadvantage that is difficulty in the separation of methyl ester products. The lack of homogeneous catalysts can be overcome by using a heterogeneous catalyst that is a solid catalyst, becaused many advantages are obtained when transesterification reactions apply the use of heterogeneous catalysts such as catalysts easily separated from the product, not corrosive and safer and more environmentally friendly [5-7]. Researches on heterogeneous/solid catalysts has been performed such as alkaline or alkaline earth oxides, alkali metals, zeolites, hydrotalcites [8-12], CaO [13], natural bentonite-supported CaO [14].

Zeolite is a porous material widely used as a catalyst based on its properties that have Bronsted acid sites and Lewis acid sites contained in pore zeolite. However, zeolites have less optimum activity and selectivity to convert reactants into desired products. Therefore we need an active phase that can minimize the less expected conditions. The widely used active phases are generally metals or metal oxides. The utilization of natural zeolite as catalyst support for transesterification of palm oil has reported. In the research, KOH was supported into natural zeolit [15]. In this work used magnetite (Fe₃O₄). Magnetite is in great demand as a catalyst because of its biocompatibility, stability, large surface area, and super paramagnetic properties [16]. Magnetite (Fe₃O₄) and γ -Fe₂O₃ which had spiral structure are used to improve biodiesel production [17]. However the magnetite embedded in zeolite as a catalyst in the transesterification reaction of used cooking oil has not been reported. As we know that in the Central Java region has great natural resource potential that is natural zeolite that found in Bayat Klaten area and iron sand/magnetite which is contained in sand of Marina Semarang beach. Therefore, in this research the utilization of both materials as raw material to preparation composite zeolite/magnetite. Furthermore the composite was applied as a catalyst for the transesterification reaction of used cooking oil to produce methyl esters.

2. Experimental Method

2.1. Materials

Sand from Marina beach Semarang, natural zeolite from Bayat Klaten. Both areas are located in Central Java, used cooking oil. Fluoric acid p.a, methanol absolute, polyethylenglycol (PEG) 4000 p.a reagents were supplied by Merck.

2.2. Instruments

X-ray diffraction (XRD) Rigaku Multiplex with Cu K α radiation ($\lambda = 1.54184$ A°) at generator voltage 40 kV and current 40 mA. Scanning Electron Microscopy (SEM) JEOL, JED 2300. GCMS-TQ8030 Shimadzu Serial O207051.

2.3. Preparation zeolite/magnetite composite catalyst

Zeolite as catalyst support was prepared by activation of natural zeolite from Bayat Klaten using HF 1% solution. The activated zeolite washing until it reaches a neutral pH, then dried to a constant weight. Characterization using XRD is used to determine the constituent components and the structure of material. Magnetite was prepared by extraction methodof sand from marina beach. Treatment by using HEM-3D (High energy milling) method with planetary ball milling is used to purify and reduce the size of magnetite particles. The mass ratio of magnetite:ball was 1:1, speed 400 rpm for 5 h. The milled magnetite characterized by X-ray Diffraction (XRD) and Scanning Electron Microscopy

ISNPINSA-7

IOP Publishing IOP Conf. Series: Journal of Physics: Conf. Series 1025 (2018) 012135 doi:10.1088/1742-6596/1025/1/012135

(SEM). The catalyst of zeolite/magnetite composite was prepared in a way activated natural zeolite was mixed with milled magnetite (Fe_3O_4). The various of zeolite:magnetiteratio mass were 1:1, 2:1, 3:1 and 4:1. The next step, addition of aquadest to the mixture and stirred in a magnetic stirrer at room temperature at 1200 rpm. An amount of PEG 4000 solution as binder was slowly added suspension. The product was dried and characterized by X-ray Diffraction (XRD.

2.4. Application zeolite/magnetite composite as transesterification catalyst

Catalytic activity tests of catalyst to transesterification reaction with used cooking oil as feed. Previously step the used cooking oil are filtered to eliminate impurities and heated to remove the water. Transesterification of used cooking oil in excess methanol (v/v) (1:12) by composite catalyst was carried out at 60°C for 4 hours. The concentration variations of catalyst were 1%, 3%, 5% and 10% (w/v). The resulting product is separated in a separating funnel for 8 hours to form 2 layers. The top layer was methyl ester and the bottom layer was glycerol. The top layer is separated and evaporated to remove the excess methanol which not participate in reaction. Furthermore, the components of the product transesterification were determined by gas chromatography mass spectrometry (GC-MS). Besides that the physical characters such as density, viscosity, acid number, iodine number and content of water also determined. Product methyl ester is weighed.

3. Results and Discussions

3.1. Characterization of zeolite/magnetite composite catalvst

Zeolite activation is done by using fluoric acid. HF solution is selected for zeolite activation because HF only slightly lowers the radius of the zeolite pores, so the radius of zeolite is still large enough and easy to do further modification. The aim of activation by fluoric acid to form decationation that influenced the zeolite surface area to increase due to the decrease of impurity covering the pores of zeolite. The increase of surface area are needed, therefore the catalytic performance of zeolite as catalyst support can be more optimal. The activation process causing the skeleton of zeolite easy to substitute with metal.

The X-ray diffraction of activated natural zeolite shown in table 1. The highest intensity peak is at $2\theta = 25.56$ °. Identification of peaks mordenite minerals in activated natural zeolites was matched with JCPDS 6-239 data while clinoptilonite minerals were matched with JCPDS data 39-1383 as shown in the Table 1.

Component	2θ Standart (deg)	2θ observed
Component	(405)	(405)
	13.2	13.36
Mordenite	19.7	19.52
JCPDS 6-239	25.7	25.56
	27.7	27.62
Clinoptilolite	22.1	22.18
JCPDS 39-1383	26.6	26.54
	29.9	29.98

Table 1.	XRD	data	of	activated	natural
----------	-----	------	----	-----------	---------

The effect of the separation using permanent magnet can be seen in figure 1. It is clearly difference in color and size of the iron sand.



Sand before extraction

Extracted material (magnetite)



Figure 1. The difference of magnetite appearance

Extracted materials from sand is magnetite/iron sand has more dark color. This is due to the reduction of impurities from the iron sand. After milling treatment using HEM-3D, it appears that magnetite has a finer size than before the HEM-3D process. The diminished magnetic particle size results in increased surface area, where it is necessary to enlarge sites that react with feeds. The evidence of magnetite in milled material can be seen in Table 2.

Component	20 Standart (deg)	2θ observed (deg)
	30.083	30.093
Magnetite	35.434	35.462
JCPDS 89-4319	43.064	43.099
	56.949	56.981
	62.536	62.592

Table 2. The 2O XRD data of milled material

The catalytic performance of the magnetite will be affected by the high magnetite surface area. However, due to the small particle size, agglomeration often occurs if the catalyst applied at high temperature. Therefore, magnetite particles supported in the zeolite matrix to prepare composite zeolite/magnetite as a heterogeneous catalyst. This step is expected can be improve both catalytic properties at the transesterification stage in the preparation of methyl esters from used cooking oil. The presence of magnetite in composite that the magnetic properties is expected to facilitate the separation so as to prevent the formation of hydrogen bonds so as to reduce the purity of glycerol. The evidence showing the composite has been formed on Table 3.

	20 Standart	2θ observed
Component	(deg)	(deg)
	13.2	-
Mordenite	19.7	-
JCPDS 6-239	25.7	25.6
	27.7	27.54
Clinoptilolite	22.1	22.14
JCPDS 39-1383	26.6	26.58

Table 3. The 2O XRD data of composite

	29.9	28.22
Magnetite	30.083	30.1
JCPDS 89-4319	35.434	35.46
	43.064	43.12
	56.949	56.92
	62.536	62.56

3.2. Structure morphology of magnetite

A slightly difference was observed in surface structure of the appearance of the magnetite and after treatment by HEM-3D. The magnetite extracted from the sand shows the area of crystals or magnetite particles appear less clear and regular. In addition, the images in Figure 2 also reveal that the material still contains impurities. While on the magnetite of treatment with HEM-3D has a crystal or particles with the edge specified. From the figure 2 also investigated that the particle size of milled magnetit seems more smooth and homogeneous. It indicates that the milling treatment using a planetary ball mill (high energy milling/HEM-3D) causing a very strong collision between the magnetite particles and the spheres/balls on the planetary ball mill, therefore reducing on magnetite particles size.



Magnetite before HEM-3D treatment Milled magnetite with HEM-3D Figure 2. SEM images of magnetite

3.3. Zeolite/magnetite composite as catalyst in transesterification reaction

In this transesterification reaction, triglycerides in used cooking oil react with methanol to produce methyl esters and glycerol. This reaction will more quickly achieve equilibrium with the addition of a catalyst. In biodiesel (methyl ester) synthesis, methanol is mostly used as alcohol because of its low cost, availability and production of alkyl esters with lower viscosity [18]. In this research, transesterification reaction of used cooking oil with methanol was catalyzed by zeolite/magnetite composite as heterogeneous catalysts. This composite were formed from zeolite superseded by magnetite. The use of this zeolite/magnetite composite as heterogenous catalysts. This composite as heterogenous catalyst will be more effectively used in transesterification reactions because it has two active sites of zeolite and magnetite that work simultaneously and for faster triglyceride splitting. Furthermore, this is because the zeolite/magnetite catalyst has two catalytically active sites namely the active site of zeolite in the form of Bronsted acid and the active site of the magnetite in the form of oxide. The Bronsted acid in this zeolite initiates triglycerides to convert the C = O bond to a C-OH bond because the Brønsted acid is able to attack the C = O bond to the C-OH bond. Whereas in this magnetite the iron oxide FeO will bind to methanol.

Based on GC-MS analysis, its known that in this transesterification reaction produce methyl ester compounds.



Figure 3. Chromatogram GC of transesterification product

Figure 3 shows the chromatogram GC of transesterification product. Moreover, fragmentation using mass spectroscopy (MS) indicates the presence of methyl esters in the reaction product. Methyl ester of transesterification product without catalyst produces methyl oleic/9-octadecanoic acid (m/z = 296) compound which is the highest peak fragmentation compound followed by methyl palmitic/hexadecanoic acid (m/z = 270) and Methyl stearic/octadecanoic acid (m/z = 270), meanwhile the using of concentration 1% ZM (zeolite/magnetite composite) catalyst resulted the methyl Palmitat (m/z = 270) as the highest peak fragmentation compound followed by methyl Oleic (m/z = 296). For concentration 10% ZM catalyst produces the methyl oleic (m/z = 296) compound which is the highest peak fragmentation followed by Methyl palmitic (m/z = 270).

The influence of the catalyst concentration ranging 1-10 % (w/v) on the transesterification of the used cooking oil is presented in Figure 4. The Fig. 4 shows transesterification reaction efficiency to yielded methyl ester strongly depends upon the catalyst amount, therefore increasing of the catalyst amount would increases the yield percentage. However, utilization of zeolite/magnetite composite catalyst in large amount less effective precisely because the content of water in methyl ester product is

too great. It beyond which no considerable change in oil conversion is observed due to increasing the viscosity of the mixture. Mansir [19] reveals that catalytic conversion of triglycerides to biodiesel/methyl ester involves the use of catalyst which lowers the activation energy and speeds up the reaction. In this process, the amount of catalyst utilized affects the biodiesel content. Usually biodiesel production reaction involves the use of catalyst. Lower catalyst loading is applied in the initial stage of the reaction and the conversion yield of biodiesel is measured. The subsequent catalyst loading will be increased periodically at constant reaction parameters until optimum biodiesel content from a particular feedstock is observed.



Figure 4. Correlation between Yield of methyl ester and catalyst concentration

3.4. The properties of methyl ester as transesterification product

Investigation of the product of transesterification addressed to its physical properties. The physical properties include density, viscosity, acid number, iodine number and water content. The properties should be determined because reflecting the quality of the methyl ester. Hence it can be decided whether the methyl ester is feasible as a fuel substitute or not.

3.4.1. Density. Density methyl ester as a product from transesterification reaction using various catalyst presents in Figure 5.



Figure 5. Density of methyl ester using various of catalyst

In Fig. 5 can observe the effect of zeolite/magnetite catalyst concentration to the density of methyl ester. The using of zeolite only, magnetite or composite zeolite/magnetite catalyst causes the density decreased sharply. At various concentrations of catalysts the use of composite catalysts showed less

ISNPINSA-7 IOP Publishing IOP Conf. Series: Journal of Physics: Conf. Series **1025** (2018) 012135 doi:10.1088/1742-6596/1025/1/012135

density than zeolite or magnetite only as catalyst. This suggests that composite catalysts are more effective because of the role of multiple active sites derived in magnetite (Fe₃O₄) as acid sites that initiate reactions with methanol and acid-active sites in zeolite. In all variations of concentration catalyst the range 1-10%, composite which was synthesized by zeolite/magnetite ratio 2:1 shows the lowest density.

3.4.2. Viscosity. The viscosity of methyl ester often called kinematic viscosity is the value of methyl ester treacly as fuel at 40°C. Viscosity of methyl ester as a product from transesterification using various catalyst presents in Figure 6.



Figure 6. Viscosity of methyl ester using various of catalyst

Viscosity is one of the important properties that must be known if the methyl esters are used instead of fuels. This is because these properties correlate with their performance on the machine. When overly high viscosity will make the fuel become larger droplets that will result in a deposit on the machine, however if the viscosity is too low it will produce a spray that is too smooth to form a rich zone that causes the formation of soot [20]. The role of the catalyst in influencing the viscosity properties has a tendency similar to that of the methyl ester density. Hence in all variations of concentration catalyst the range 1-10%, composite which was synthesized by zeolite/magnetite ratio 2:1 shows the lowest viscosity.

3.4.3. Acid number. The acid number is milligram amount of KOH needed for neutralized 1 gram of sample. The acid number is calculated from the percentage value of fatty acids free. The acid number of methyl ester as a product from transesterification using various catalyst presents in Figure 7.



Figure 7. Viscosity of methyl ester using various of catalyst

Large acid numbers show large free fatty acids derived from oil hydrolysis or due to poor processing. The higher the acid number the lower the quality. In this work, the catalyst plays a very significant role in decreasing the acid number. In all variations of concentration catalyst the range 1-10%, composite which was synthesized by zeolite/magnetite ratio 2:1 shows the lowest acid number. When the reaction using zeolite/magnetite ratio 2:1 composite as catalyst cause the acid number increasing slightly. This data reveals that the need for magnetite in assisting the process of initiation of reactions between feeds and methanol.

3.4.4. Iodine number. The iodine number called iodine absorption number is the mass of iodine in grams absorbed at 100 grams of a chemical under the test conditions used. Iodine numbers are often used to determine the amount of unsaturation in fatty acids. The unsaturation is in the form of a double bond, which will react with the iodine compound. The higher the iodine number, the more the C = C bonds present in the fat [21]. Iodine number of methyl ester as a product from transesterification using various catalyst presents in Figure 8.



Figure 8. Iodine number of methyl ester using various of catalyst

The higher the iodine value, the higher the number of double-bonded fatty acids contained in the methyl ester. The more double bonds held in the sample have greater potential to experience the polymerization process due to lower stability. The influence of the rise of methyl ester unsaturation can cause CO_2 gas to increase the degree of unsaturation associated with the iodine number. The longer the carbon chain the lower the CO_2 emissions and the higher the iodine number the lower the CO_2 emissions produced [20]. If the methyl ester would be use as biofuel so the number of iodine must

be low. Therefore based on the data in Fig. 8, the low iodine number is belonged by methyl esters of transesterified results catalyzed by zeolites. The explanation argument for this fact is that iodine is most likely absorbed by the pores of the zeolite, where it cannot occur in magnetite because magnetite is not a porous material. This is also cause the highest iodine number on magnetite catalyst. The ability adsorption of zeolite is reduced when zeolite compose with magnetite.

3.4.5. Water content. Water content is the difference of weight methyl ester before and after evaporating. In this work data of water content in methyl ester presents in Figure 9. In this data there is a tendency on catalyst has similarity of role with the viscosity data. This means that the lowest water content in the use of catalysts only zeolite, and the highest water content in magnetite. The role of the pores present in the zeolite structure absorb the water, therefore the catalyst having relatively high porosity as in the zeolite/magnetite 3: 1 composite also shows a low water content.



Figure 9. The water content of methyl ester using various of catalyst

4. Conclusion

Preparation of methyl ester was catalyzed by composite of zeolite/magnetite through transeserification reaction using variation of catalyst concentration 1%, 3%, 5% and 10% (w/v) produce a mixture of methyl Oleic (MES) Palmitic Acid (MES) and Stearic Acid (MES). Character product such as density, viscosity, acid number and iodine number has fulfilled to SNI standard 7182: 2015.

Acknowlegment

This research financial support by funding of Besides APBN DPA SUKPA LPPM Diponegoro University, Fiscal Year 2017. We convey gratefully to Chemistry Departement for the facilities to carry out this research.

References

- [1] Srivastava, A., Prasad, R., 2000, *Renewable and Sustainable Energy Reviews*, **4**(2), 111-133.
- [2] Sinha, S., Agarwal, A.K., Garg, S., 2008, *Energy Conversion and Management*, **49** (5), 1248-1257.
- [3] Ma, F., Hanna, M.A., 1999, *Bioresource Technology*, **70** (1), 1-15.
- [4] Keera, S.T., El Sabagh, S.M., Taman, A.R., 2011, *Fuel*, **90** (1), 42-47.
- [5] Kumar, M., Sharma, M.P., 2016, *Renewable and Sustainable Energy Reviews*, **56**, 1129-1138.
- [6] Wang, J.X., Chen, K.T., Wu, J.S., Wang, P.H., Huang, S.T., Chen, C.C., 2012, *Fuel Processing Technology*, **104**, 167-173.
- [7] Dai, Y.M., Chen, K.T., Wang, P.H., Chen, C.C., 2016, Advanced Powder Technology, 27 (6), 2432-2438.
- [8] Lu, Y., Zhang, Z., Xu, Y., Liu, Q., Qian, G., 2015, *Bioresource Technology*, **190**, 438-441.

- [9] Amani, H., Ahmad, Z., Hameed, B.H., 2014, Applied Catalysis A, 487, 16-25.
- [10] Paterson, G., Issariyakul, T., Baroi, C., Bassi, A., Dalai, A., 2013, Catalysis Today, 212, 157-163.
- [11] Dai, Y.M., Kao, I.H., Chen, C.C., 2017, *Journal of Taiwan Institute of Chemical Engineers*, **70**, 260-266.
- [12] Habaki, H., Hayashi, T., Egashira, R., 2014, *Journal of Environmental Chemical Engineering*, 2 (3),1543-1550.
- [13] Kouzu, M., Hidaka, J., 2012, Fuel, 93, 1-2.
- [14] Shan, R., Zhao, C., Yuan, H., Wang, S.X., 2017, *Energy Conversion and Management*, **138**, 547-555.
- [15] IndraKusuma, R., Hadinoto, J.P., Ayucitra, A., Soetaredjo, F.E., Ismadji, S., 2013, *Applied Clay Science*, **74**, 121-126.
- [16] Lu, A.H., Salabas, E.E.L., Schüth, F., 2007, Angewandte Chemie International Edition 46 (8), 1222-1244.
- [17] Teja A.S. and Koh, P.Y., 2009, *Progress in crystal growth and characterization of materials* **55** (1), 22-45.
- [18] Wittcoff, H.A., Reuben, B.G., 1996, *Industrial organic chemicals*, John Wiley & Sons, New York.
- [19] Mansir, N., Taufiq-Yap, Y.H., Rashid, U., Lokman, I.M., 2017, Energy Conversion and Management, 141, 171-182.
- [20] Prihandana R. and Hendroko, R., 2008, Energi Hijau: Pilihan Bijak Menuju Negeri Mandiri Energi, Niaga Swadaya.
- [21] Thomas, A., 2002, *Fats and Fatty Oils*, Ullmann's Encyclopedia of Industrial Chemistry, Weinheim: Wiley-VCH.

Zeolitemagnetite composites as catalysts on the Synthesis of Methyl Esters (MES) from cooking oil by Sriatun Sriatun

Submission date: 25-May-2019 08:10AM (UTC+0700) Submission ID: 1135637183 File name: lysts_on_the_Synthesis_of_Methyl_Esters_MES_from_cooking_oil.pdf (776.43K) Word count: 4468 Character count: 23535 Journal of Physics: Conference Series

PAPER · OPEN ACCESS

Zeolite/magnetite composites as catalysts on the Synthesis of Methyl Esters (MES) from cooking oil

1 To cite this article: Sriatun *et al* 2018 *J. Phys.: Conf. Ser.* **1025** 012135

16 View the <u>article online</u> for updates and enhancements.



IOP ebooks[™]

Bringing you innovative digital publishing with leading voices to create your essential collection of books in STEM research.

Start exploring the collection - download the first chapter of every title for free.

This content was downloaded from IP address 180.246.99.173 on 19/10/2018 at 03:41

ISNPINSA-7

IOP Publishing

IOP Conf. Series: Journal of Physics: Conf. Series 1025 (2018) 012135 doi:10.1088/1742-6596/1025/1/012135

Zeolite/magnetite composites as catalysts on the Synthesis of Methyl Esters (MES) from cooking oil

Sriatun¹, Adi Darmawan¹, Sriyanti¹, Wuri Cahyani¹, Hendri Widyandari²

¹) Department of Chemistry, Faculty of Science and Mathematics, Diponegoro Un 5versity

²) Department of Physics Faculty of Science and Mathematics, Diponegoro University Jl. Prof. Soedharto, SH, Tembalang, Semarang 50275, Indonesia

E-mail: sriatun@live.undip.ac.id

Abstract. The using of zeolite/magnetite composite as a catalyst for the synthesis of methyl esters (MES) of cooking oil has been performed. In this study the natural magnetite was extracted from the iron sand of Semarang marina beach and milled by high energy Milling (HEM) with ball: magnetite ratio: 1:1. The composites prepared from natural zeolite and natural magnetite with zeolite: magnetite ratio 1:1; 2:1; 3:1 and 4:1. Preparation of methyl ester was catalyzed by composite of zeolite/magnetite through transeserification reaction, it was studied on variation of catalyst concentration (w/v) 1%, 3%, 5% and 10% to feed volume. The reaction product are mixture of methyl Oleic (MES), methyl Palmitic (MES) and methyl Stearic (MES). Character product of this research include density, viscosity, acid number and iodine number has fulfilled to SNI standard 7182: 2015.

Keywords: composite, zeolite/magnetite, catalyst, MES

1. Introduction

Increased transportation demand leads to an increase in energy demand globally. Transportation sector is a sector that consumes the most energy, so it can increase environmental emission. Transportation using fuel oil is mainly processed petroleum products, while the world's oil reserves and Indonesia began to thin out. In an effort to find solutions to these problems, various studies reveal the existence of a variety of raw materials from renewable resources capable of being processed into a more environmentally friendly fuel as it reduces CO₂ emissions.

Methyl esters are a major component of biodiesel which is the most appropriate solution to replace the fuel fossils as the world's major source of energy tizhsport, because biodiesel is a material fuel that can replace petrol diesel on a machine. Biodiesel is consisting of a mixture of mono-alkyl esters of long chain fatty acids, which are used as an alternative to fuel from diesel engines. Biodiesel or methyl esters are obtained from the transesterification of animal fats, vegetable oils, and even cooking oil waste, with catalytic process [1].

Actually Vegetable oil has a calorific value that is almost the same with conventional fuel, but usage directly as fuel is still encountered obstacles. When compared to biodiesel, vegetable oils have a much higher viscosity value at temperature 40°C. This may cause problems in the pumping and spraying process when they used as fuel [2]. The incandescent point (flash point) also much larger than diesel oil, this is it inhibit the injection process and result incomplete combustion. The use of oil



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI. Published under licence by IOP Publishing Ltd 1

ISNPINSA-7	IOP Publishing
IOP Conf. Series: Journal of Physics: Conf. Series 1025 (2018) 012135	doi:10.1088/1742-6596/1025/1/012135

vegetable as a fuel also leaves residue carbon in the engine injector. Efforts to reduce viscosity and flash point vegetable oils, among others, by dilution of oil with solvents, emulsification, pyrolysis, and transesterification. The transesterification process is the most common way done because it does not require energy and temperature tall one. This reaction will produce methyl or ethyl esters, depending on the type of alcohol reacted. Methanol is used more as a source of alcohol because the chain is shorter, more polar and the price cheaper than other alcohols [3].

Both homogeneous and heterogeneous catalysts can be used in transeterification reactions. Basic homogeneous cataly 18 such as KOH as well as sodium and potassium alkoxide [3], NaOH [4], while acidic ones such as sulfuric acid, sulfonic acid and hydrochloric acid are usually used as catalysts in transesterification reactions. The using of homogeneous catalysts has a disadvantage that is difficulty in the separation of methyl ester products. The lack of homogeneous catalysts can be overcome by using a heterogeneous catalyst that is a solid catalyst, becaused many advantages are obtained when transesterification reactions apply the use of heterogeneous catalysts such as catalysts easily separated from the product, not corrosive and safer and more environmentally friendly [5-7]. Researches on heterogeneous/solid catalysts has been performed such as alkaline or alkaline earth oxides, alkali metals, zeolites, hydrotalcites [8-12], CaO [13], natural bentonite-supported CaO [14].

Zeolite is a porous material widely used as a catalyst based on its properties that have Bronsted acid sites and Lewis acid sites contained in pore zeolite. However, zeolites have less optimum activity and selectivity to convert reactants into desired products. Therefore we need an active phase that can minimize the less expected conditions. The videly used active phases are generally metals or metal oxides. The utilization of natural zeolite as catalyst support for transesterification of palm oil has reported. In the research, KOH was supported into r20 ral zeolit [15]. In this work used magnetite (Fe₃O₄). Magnetite is in great demand as a catalyst because of its biocompatibility, stability, large surface area, and super paramagnetic properties [16]. Magnetite (Fe₃O₄) and γ -Fe₂O₃ which had spiral **37** actuate are used to improve biodiesel production [17]. However the magnetite embedded in zeolite as a catalyst in the transesterification reaction of used cooking oil has not been reported. As we know that in the Central Java region has great natural resource potential that is natural zeolite that found in Bayat Klaten area and iron sand/magnetite which is contained in sand of Marina Semarang beach. Therefore, in this research the utilization of both materials **a** a catalyst for the transesterification reaction of used cooking oil to produce methyl esters.

2. Experimental Method

2.1. Materials

Sand from Marina beach Semarang, natural zeolite from Bayat Klaten. Both areas are located in Central Java, used cooking oil. Fluoric acid p.a, methanol absolute, polyethylenglycol (PEG) 4000 p.a reagents were supplied by Merck.

2.2. Instruments

23

X-ray diffraction (XRD) Rigaku Multiplex with Cu K α radiation ($\lambda = 1.54184$ Ű) at generator voltage 40 kV and current 40 mA. Scanning Electron Microscopy (SEM) JEOL, JED 2300. GCMS-TQ8030 Shimadzu Serial O207051.

2.3. Preparation zeolite/magnetite composite catalyst

Zeolite as catalyst support was prepared by activation of natural zeolite from Bayat Klaten using HF 1% solution. The activated zeolite washing until it reaches a neutral pH, then dried to a constant weight. Characterization using XRD is used to determine the constituent components and the structure of material. Magnetite was prepared by extraction methodof sand from marina beach. Treatment by using HEM-3D (High energy milling) method with planetary ball milling is used to purify and reduce the size of magnetic particles. The mass ratio of magnetite:ball was 1:1, speed 400 rpm for 5 h. The milled magnetite characterized by X-ray Diffraction (XRD) and Scanning Electron Microscopy

ISNPINSA-7

IOP Publishing

IOP Conf. Series: Journal of Physics: Conf. Series 1025 (2018) 012135 doi:10.1088/1742-6596/1025/1/012135

(SEM). The catalyst of zeolite/magnetite composite was prepared in a way activated natural zeolite was mixed with milled magnetite (Fe_3O_4). The various of zeolite 15 gnetiteratio mass were 1:1, 2:1, 3:1 and 4:1. The next step, addition of aquadest to the mixture and stirred in a magnetic stirrer at room temperature at 1200 rpm. An amount of PEG 4000 solution as binder was slowly added suspension. The product was dried and characterized by X-ray Diffraction (XRD.

2.4. Application zeolite/magnetite composite as transesterification catalyst

Catalytic activity tests of catalyst to transesterification reaction with used cooking oil as feed. Previously step the used cooking oil are filtered to eliminate impurities and heated to remove the water. Transesterification of used cooking oil in excess methanol (v/v) (1:12) by composite catalyst was carried out at 60°C for 4 hours. The continuation variations of catalyst were 1%, 3%, 5% and 10% (w/v). The resulting product is separated in a separating funnel for 8 hours to form 2 layers. The **9** layer was methyl ester and the bottom layer was glycerol. The top layer is separated and evaporated to remove the excess methanol which not participate in reaction. Furthermore, the components of the product transesterification were determined by gas chromatography mass spectrometry (GC-MS). Besides that the physical characters such as density, viscosity, acid number, iodine number and content of water also determined. Product methyl ester is weighed.

3. Results and Discussions

3.1. Characterization of zeolite/magnetite composite catalyst

Zeolite activation is done by using fluoric acid. HF solution is selected for zeolite activation because HF only slightly lowers the radius of the zeolite pores, so the radius of zeolite is still large enough and easy to do further modification. The aim of activation by fluoric acid to form decationation that influenced the zeolite surface area to increase due to the decrease of impurity covering the pores of zeolite. The increase of surface area are needed, therefore the catalytic performance of zeolite as catalyst support can be more optimal. The activation process causing the skeleton of zeolite easy to substitute with metal.

The X-ray diffraction of activated natural zeolite shown in table 1. The highest intensity peak is at $2\theta = 25.56^{\circ}$. Identification of peaks mordenite minerals in activated natural zeolites was matched with JCPDS 6-239 data while clinoptilonite minerals were matched with JCPDS data 39-1383 as shown in the Table 1.

Component	20 Standart (deg)	2θ observed (deg)
	13.2	13.36
Mordenite	19.7	19.52
JCPDS 6-239	25.7	25.56
	27.7	27.62
Clinoptilolite	22.1	22.18
JCPDS 39-1383	26.6	26.54
	29.9	29.98

Table 1. XRD data of activated natural

The effect of the separation using permanent magnet can be seen in figure 1. It is clearly difference in color and size of the iron sand.

ISNPINSA-7 IOP Publishing IOP Conf. Series: Journal of Physics: Conf. Series **1025** (2018) 012135 doi:10.1088/1742-6596/1025/1/012135



Sand before extraction





Milled material (milled magnetite)

Figure 1. The difference of magnetite appearance

Extracted material (magnetite)

Extracted materials from sand is magnetite/iron sand has more dark color. This is due to the reduction of impurities from the iron sand. After milling treatment using HEM-3D, it appears that magnetite has a finer size than before the HEM-3D process. The diminished magnetic particle size results in increased surface area, where the is necessary to enlarge sites that react with feeds. The evidence of magnetite in milled material can be seen in Table 2.

Component	20 Standart (deg)	2θ observed (deg)
	30.083	30.093
Magnetite	35.434	35.462
JCPDS 89-4319	43.064	43.099
	56.949	56.981
	62.536	62.592

 Table 2. The 2O XRD data of milled material

The catalytic performance of the magnetite will be affected by the high magnetite surface area. However, due to the small particle size, agglomeration often occurs if the catalyst applied at high temperature. Therefore, magnetite particles supported in the zeolite matrix to prepare composite zeolite/magnetite as a heterogeneous catalyst. This step is expected can be improve both catalytic properties at the transesterification stage in the preparation of methyl esters from used cooking oil. The presence of magnetite in composite that the magnetic properties is expected to facilitate the separation so as to prevent the formation of hydrogen bonds so as to reduce the purity of glycerol. The evidence showing the composite has been formed on Table 3.

Table 3. The 20 XR	D data of composite
--------------------	---------------------

Component	2θ Standart (deg)	2θ observed (deg)
	13.2	-
Mordenite	19.7	-
JCPDS 6-239	25.7	25.6
	27.7	27.54
Clinoptilolite	22.1	22.14
JCPDS 39-1383	26.6	26.58

ISNPINSA-7

IOP Publishing

IOP Conf. Series: Journal of Physics: Conf. Series 1025 (2018) 012135 doi:10.1088/1742-6596/1025/1/012135

	29.9	28.22
Magnetite	30.083	30.1
JCPDS 89-4319	35.434	35.46
	43.064	43.12
	56.949	56.92
	62.536	62.56

3.2. Structure morphology of magnetite

A slightly difference was observed in surface structure of the appearance of the magnetite and after treatment by HEM-3D. The magnetite extracted from the sand shows the area of crystals or magnetite particles appear less clear and regular. In addition, the images in Figure 2 also reveal that the material still contains impurities. While on the magnetite of treatment with HEM-3D has a crystal or particles with the edge specified. From the figure 2 also investigated that the particle size of milled magnetit seems more smooth and homogeneous. It indicates that the milling treatment using a planetary ball mill (high energy milling/HEM-3D) causing a very strong collision between the magnetite particles and the spheres/balls on the planetary ball mill, therefore reducing on magnetite particles size.



Magnetite before HEM-3D treatment Figure 2. SEM images of magnetite

3.3. Zeolite/magnetite composite as catalyst in transesterification reaction 31 In this transesterification reaction, triglycerides in used cooking oil react with methanol to produce methyl esters and glycerol. This reaction will more quickly achieve equilibrium with the addition of a catalyst. In biodiesel (methyl ester) synthesis, methanol is mostly used as alcohol because of its low 36 t, availability and production of alkyl esters with lower viscosity [18]. In this research, transesterification reaction of used cooking oil with methanol was catalyzed by zeolite/magnetite composite as heterogeneous catalysts. This composite were formed from zeolite superseded by magnetite. The use of this zeolite/magnetite composite as heterogenous catalyst will be more effectively used in transesterification reactions because it has two active sites of zeolite and magnetite that work simultaneously and for faster triglyceride splitting. Furthermore, this is because the zeolite/magnetite catalyst has two catalytically active sites namely the active site of zeolite in the form of Bronsted acid and the act 25 site of the magnetite in the form of oxide. The Bronsted acid in this zeolite initiates trigging to convert the C = O bond to a C-OH bond because the Brønsted acid is able to attack the C = O bond to the C-OH bond. Whereas in this magnetite the iron oxide FeO will bind to methanol.

Based on GC-MS analysis, its known that in this transesterification reaction produce methyl ester compounds.

ISNPINSA-7 IOP Publishing IOP Conf. Series: Journal of Physics: Conf. Series **1025** (2018) 012135 doi:10.1088/1742-6596/1025/1/012135



Figure 3. Chromatogram GC of transesterification product

Figure 3 shows the chromatogram GC of transesterification product. Moreover, fragmentation using mass spectroscopy (MS) indicates the presence of methyl esters in the reaction product. Methyl ester of transesterification product without catalyst produces methyl oleic/9-octadecanoic acid (m/z = 296) compour 30 which is the highest peak fragmentation compound followed by methyl palmitic/hexadecanoic acid (m/z = 270) and Methyl stearic/octadecanoic acid (m/z = 270), meanwhile the using of concentration 1% ZM (zeolite/magnetite composite) catalyst resulted the methyl Palmitat (m/z = 270) as the highest peak fragmentation compound followed by methyl Oleic (m/z = 296). For concentration 10% ZM catalyst produces the methyl oleic (m/z = 296) compound which is the highest peak fragmentation followed by Methyl palmitic (m/z = 270).

The influence of the catalyst concentration ranging 1-10 % (w/v) on the transesterification of the used cooking oil is posented in Figure 4. The Fig. 4 shows transesterification reaction efficiency to yielded methyl ester strongly depends upon the catalyst amount, therefore increasing of the catalyst amount would increases the yield percentage. However, utilization of zeolite/magnetite composite catalyst in large amount less effective precisely because the content of water in methyl ester product is

ISNPINSA-7 IOP Publishing IOP Conf. Series: Journal of Physics: Conf. Series **1025** (2018) 012135 doi:10.1088/1742-6596/1025/1/012135

too great. It beyond which no considerable change in oil coversion is observed due to increasing the viscosity of the mixture. Mansir [19] reveals that catalytic conversion of triglycerides to biodiesel/methyl ester involves the use of catalyst which lowers the activation energy and speeds up the reaction. In this process, the amount of catalyst utilized affects the biodiesel content. Usually biodiesel production reaction involves the use of catalyst. Lower catalyst loading is applied in the initial stage of the reaction and the conversion yield of biodiesel is measured. The subsequent catalyst loading will be increased periodically at constant reaction parameters until optimum biodiesel content from a particular feedstock is observed.



Figure 4. Correlation between Yield of methyl ester and catalyst concentration

3.4. The properties of methyl ester as transesterification product

Investigation of the product of transesterification addressed to its physical properties. The physical properties include density, viscosity, acid number, iodine number and water content. The properties should be determined because reflecting the quality of the methyl ester. Hence it can be decided whether the methyl ester is feasible as a fuel substitute or not.

3.4.1. Density. Density methyl ester as a product from transesterification reaction using various catalyst presents in Figure 5.



Figure 5. Density of methyl ester using various of catalyst

In Fig. 5 can observe the effect of zeolite/magnetite catalyst concentration to the density of methyl ester. The using of zeolite only, magnetite or composite zeolite/magnetite catalyst causes the density decreased sharply. At various concentrations of catalysts the use of composite catalysts showed less

ISNPINSA-7	IOP Publishing
IOP Conf. Series: Journal of Physics: Conf. Series 1025 (2018) 012135	doi:10.1088/1742-6596/1025/1/012135

density than zeolite or magnetite only as catalyst. This suggests that composite catalysts are more effective because of the role of multiple active sites derived in magnetite (Fe_3O_4) as acid sites that initiate reactions with methanol and acid-active sites in zeolite. In all variations of concentration catalyst the range 1-10%, composite which was synthesized by zeolite/magnetite ratio 2:1 shows the lowest density.

3.4.2. Viscosity. The viscosity of methyl ester often called kinematic viscosity is the value of methyl ester treacly as fuel at 40° C. Viscosity of methyl ester as a product from transesterification using various catalyst presents in Figure 6.



Figure 6. Viscosity of methyl ester using various of catalyst

Viscosity is one of the important properties that must be known if the methyl esters are used instead of fuels. This is because these properties correlate with their performance on the machine. When overly high viscosity will make the fuel become larger droplets that will result in a deposit on the machine, however if the viscosity is too low it will produce a spray that is too smooth to form a rich zone that causes the formation of soot [20]. The role of the catalyst in influencing the viscosity properties has a tendency similar to that of the methyl ester density. Hence in all variations of concentration catalyst the range 1-10%, composite which was synthesized by zeolite/magnetite ratio 2:1 shows the lowest viscosity.

3.4.3. Acid number. The acid number is milligram amount of KOH needed for neutralized 1 gram of sample. The acid number is calculated from the percentage value of fatty acids free. The acid number of methyl ester as a product from transesterification using various catalyst presents in Figure 7.

8

ISNPINSA-7

IOP Publishing

IOP Conf. Series: Journal of Physics: Conf. Series 1025 (2018) 012135 doi:10.1088/1742-6596/1025/1/012135



Figure 7. Viscosity of methyl ester using various of catalyst

Large acid numbers show large free fatty acids derived from oil hydrolysis or due to poor processing. The higher the acid number the lower the quality. In this work, the catalyst plays a very significant role in decreasing the acid number. In all variations of concentration catalyst the range 1-10%, composite which was synthesized by zeolite/magnetite ratio 2:1 shows the lowest acid number. When the reaction using zeolite/magnetite ratio 2:1 composite as catalyst cause the acid number increasing slightly. This data reveals that the need for magnetite in assisting the process of initiation of reactions between feeds and methanol.

3.4.4. Iodine number. The iodine number called iodine absorption number is the mass of iodine in grams absorbed at 100 grams of a chemical under the test conditions used. Iodine numbers are often used to determine the amount of unsaturation in fatty acids. The unsaturation is in the form of a double bond, which will react with the iodine compound. The higher the iodine number, the more the C = C bonds present in the fat [21]. Iodine number of methyl ester as a product from transesterification using various catalyst presents in Figure 8.



Figure 8. Iodine number of methyl ester using various of catalyst

The higher the iodine value, the higher the number of double-bonded fatty acids contained in the methyl ester. The more double bonds held in the sample have greater potential to experience the polymerization process due to lower stability. The influence of the rise of methyl ester unsaturation can cause CO_2 gas to increase the degree of unsaturation associated with the iodine number. The longer the carbon chain the lower the CO_2 emissions and the higher the iodine number the lower the CO_2 emissions produced [20]. If the methyl ester would be use as biofuel so the number of iodine must

ISNPINSA-7	IOP Publishing
IOP Conf. Series: Journal of Physics: Conf. Series 1025 (2018) 012135	doi:10.1088/1742-6596/1025/1/012135

be low. Therefore based on the data in Fig. 8, the low iodine number is belonged by methyl esters of transesterified results catalyzed by zeolites. The explanation argument for this fact is that iodine is most likely absorbed by the pores of the zeolite, where it cannot occur in magnetite because magnetite is not a porous material. This is also cause the highest iodine number on magnetite catalyst. The ability adsorption of zeolite is reduced when zeolite compose with magnetite.

3.4.5. Water content. Water content is the difference of weight methyl ester before and after evaporating. In this work data of water content in methyl ester presents in Figure 9. In this data there is a tendency on catalyst has similarity of role with the viscosity data. This means that the lowest water content in the use of catalysts only zeolite, and the highest water content in magnetite. The role of the pores present in the zeolite structure absorb the water, therefore the catalyst having relatively high porosity as in the zeolite/magnetite 3: 1 composite also shows a low water content.



Figure 9. The water content of methyl ester using various of catalyst

4. Conclusion

Preparation of methyl ester was catalyzed by composite of zeolite/magnetite through transeserification reaction using variation of catalyst concentration 1%, 3%, 5% and 10% (w/v) produce a mixture of methyl Oleic (MES) Palmitic Acid (MES) and Stearic Acid (MES). Character product such as density, viscosity, acid number and iodine number has fulfilled to SNI standard 7182: 2015.

Acknowlegment

This research financial support by funding of Besides APBN DPA SUKPA LPPM Diponegoro University, Fiscal Year 2017. We convey gratefully to Chemistry Departement for the facilities to carry out this research.

Refer 61ces

- 13 vastava, A., Prasad, R., 2000, Renewable and Sustainable Energy Reviews, 4(2), 111-133. [1]
- [2] Sinha, S., Agarwal, A.K., Garg, S., 2008, Energy Conversion and Management, 49 (5), 1248-6 1257.
 6 a, F., Hanna, M.A., 1999, *Bioresource Technology*, 70 (1), 1-15.
- [3]
- Keera, S.T., El Sabagh, S.M., Tar331, A.R., 2011, Fuel, 90 (1), 42-47. [4]
- Kolmar, M., Sharma, M.P., 2016, Renewable and Sustainable Energy Reviews, 56, 1129-1138. [5]
- Wang, J.X., Chen, K.T., Wu, J.S., Wang, P.H., Huang, S.T., Chen, C.C., 2012, Fuel Processing [6] 9 Technology, 104, 167-173.
- Dai, Y.M., Chen, K.T., Wang, P.H., Chen, C.C., 2016, Advanced Powder Technology, 27 (6), [7] 28 2432-2438.
- [8] Lu, Y., Zhang, Z., Xu, Y., Liu, Q., Qian, G., 2015, Bioresource Technology, 190, 438-441.

ISNP	NSA-7

IOP Publishing

IOP Conf. Series: Journal of Physics: Conf. Series 1025 (2018) 012135 doi:10.1088/1742-6596/1025/1/012135

- [9] 24 nani, H., Ahmad, Z., Hameed, B.H., 2014, Applied Catalysis A, 487, 16-25.
- [10] Paterson, G., Issariyakul, T., Baroi, C., Bassi, A., Dalai, A., 2013, Catalysis Today, 212, 157-27 163.
- [11] Dai, Y.M., Kao, I.H., Chen, C.C., 2017, Journal of Taiwan Institute of Chemical Engineers, 70, 260-266.
- [12] Habaki, H., Hayashi, T., Egashira, R., 2014, *Journal of Environmental Chemical Engineering*, 2 (3),1543-1550.
- [13] Kouzu, M., Hidaka, J., 2012, Fuel, 93, 1-2.
- [14] Shan, R., Zhao, C., Yuan, H., Wang, S.X., 2017, Energy Conversion and Management, 138, 547-555.
- [15] IndraKusuma, R., Hadinoto, J.P., Ayucitra, A., Soetaredjo, F.E., Ismadji, S., 2013, Applied Clay 11 Science, 74, 121-126.
- [16] Lu, A.H., Salabas, E.E.L., Schüth, F., 2007, Angewandte Chemie International Edition 46 (8), 8 1222-1244.
 [17] Teja A.S. and Koh, P.Y., 2009, *Progress in crystal growth and characterization of materials* 55
- (1), 22-45.
- (1), 22-45.
 [18] Wittcoff, H.A., Reuben, B.G., 1996, *Industrial organic chemicals*, John Wiley & Sons, New 21 York.
- [19] Mansir, N., Taufiq-Yap, Y.H., Rashid, U., Lokman, I.M., 2017, Energy Conversion and 19 Management, 141, 171-182.
- [20] Prihandana R. and Hendroko, R., 2008, Energi Hijau: Pilihan Bijak Menuju Negeri Mandiri 12 Energi, Niaga Swadaya.
- [21] Thomas, A., 2002, Fats and Fatty Oils, Ullmann's Encyclopedia of Industrial Chemistry, Weinheim: Wiley-VCH.

Zeolitemagnetite composites as catalysts on the Synthesis of Methyl Esters (MES) from cooking oil

ORIGINALITY REPORT



E. Rashtizadeh, F. Farzaneh, M. Ghandi. "A 3 comparative study of KOH loaded on double aluminosilicate layers, microporous and mesoporous materials as catalyst for biodiesel production via transesterification of soybean oil", Fuel, 2010

1%

4	Konga, Ajay Kumar, Annapurna Subhash Muchandi, and Gomathi Priya Ponnaiah. "Soxhlet extraction of Spirogyra sp. algae: an alternative fuel", Biofuels, 2016. Publication	1%
5	jcc.undip.ac.id Internet Source	1%
6	Submitted to University of Bath	1%
7	Ary Budi Mulyono, Bambang Sugiarto, Muchammad Taufiq Suryantoro, Hari Setiapraja et al. "Effect of hydrotreating in biodiesel on the growth of deposits in the combustion chamber as a solution for the deposits reduction in the usage of biodiesel", E3S Web of Conferences, 2018 Publication	1%
8	Submitted to Panjab University Student Paper	1%
9	www.thaiscience.info Internet Source	1%
10	Submitted to University of Hong Kong Student Paper	<1%
11	scholarbank.nus.edu.sg	<1%

12	WWW.eijo.in Internet Source	<1%
13	www.i-scholar.in	< 1 %
14	Submitted to Macquarie University Student Paper	< 1 %
15	Istadi, I., Didi D. Anggoro, Luqman Buchori, Dyah A. Rahmawati, and Dinnia Intaningrum. "Active Acid Catalyst of Sulphated Zinc Oxide for Transesterification of Soybean Oil with Methanol to Biodiesel", Procedia Environmental Sciences, 2015. Publication	<1%
16	Lale Budi Hutami Rahayu, Ika Oktavia Wulandari, Djoko Herry Santjojo, Akhmad Sabarudin. " Synthesis and Characterization of Fe O Nanoparticles using Polyvinyl Alcohol (PVA) as Capping Agent and Glutaraldehyde (GA) as Crosslinker ", IOP Conference Series: Materials Science and Engineering, 2018	<1%

Publication

17

www.esciencecentral.org

18 S.T. Keera, S.M. El Sabagh, A.R. Taman. "Transesterification of vegetable oil to biodiesel fuel using alkaline catalyst", Fuel, 2011 <1% <1%

- A Supriyanto, I Lestari, N Citrasari, S S Putri. " The bioconversion of vegetable waste extract from Osowilangun Central Market Surabaya into bioethanol using ", IOP Conference Series: Earth and Environmental Science, 2019 Publication
- Li, Yufei, Xiaoyu Du, Chao Wu, Xueying Liu, Xia Wang, and Ping Xu. "An efficient magnetically modified microbial cell biocomposite for carbazole biodegradation", Nanoscale Research Letters, 2013.

<1%

Publication

21

Hashem Sharghi, Pezhman Shiri, Mahdi Aberi. "An overview on recent advances in the synthesis of sulfonated organic materials, sulfonated silica materials, and sulfonated carbon materials and their catalytic applications in chemical processes", Beilstein Journal of Organic Chemistry, 2018 Publication



25	O'Hara, P "Fatty acid synthesis in developing leaves of Brassica napus in relation to leaf growth and changes in activity of 3-oxoacyl-ACP reductase", FEBS Letters, 20010112 Publication	< 1 %
26	Yujie Wang, Minghui Zhang, Xuefeng Ding. "Biodiesel production from soybean oil using modified calcium loaded on rice husk activated carbon as a low-cost basic catalyst", Separation Science and Technology, 2017 Publication	<1%
27	Submitted to Fujairah College Student Paper	<1%
28	www.ncbi.nlm.nih.gov Internet Source	< 1 %
29	Submitted to Institute of Graduate Studies, UiTM	-1
	Student Paper	< %
30	Student Paper Caroline Calloni, Rafaela Dall Agnol, Luana Soares Martínez, Fábio de Siqueira Marcon, Sidnei Moura, Mirian Salvador. "Jaboticaba (Plinia trunciflora (O. Berg) Kausel) fruit reduces oxidative stress in human fibroblasts cells (MRC-5)", Food Research International, 2015 Publication	<1% <1%

Student Paper

< **|**%

Submitted to Universidade de Sao Paulo

Student Paper

32

<1%

33	Submitted to Higher Education Commission Pakistan Student Paper	<1%
34	Sinha, S "Biodiesel development from rice bran oil: Transesterification process optimization and fuel characterization", Energy Conversion and Management, 200805 Publication	<1%
35	Guan, X.H "Competitive adsorption of organic matter with phosphate on aluminum hydroxide", Journal of Colloid And Interface Science, 20060401 Publication	<1%
36	Ricky Indra Kusuma, Johan Prabowo Hadinoto, Aning Ayucitra, Felycia Edi Soetaredjo, Suryadi Ismadji. "Natural zeolite from Pacitan Indonesia, as catalyst support for transesterification of palm oil", Applied Clay Science, 2013 Publication	<1%
37	www.locus.ufv.br Internet Source	<1%
38	www.degruyter.com	<1%
	Submitted to UWC ChangShu China	



Exclude quotes	Off	Exclude matches	Off
Exclude bibliography	Off		

Zeolitemagnetite composites as catalysts on the Synthesis of Methyl Esters (MES) from cooking oil

GRADEMARK REPORT

FINAL GRADE	GENERAL COMMENTS
/100	Instructor

PAGE 1	
PAGE 2	
PAGE 3	
PAGE 4	
PAGE 5	
PAGE 6	
PAGE 7	
PAGE 8	
PAGE 9	
PAGE 10	
PAGE 11	
PAGE 12	