

Daftar Lampiran C1

1. Hasil Penilaian Sejawat Sebidang atau peer review Karya Ilmiah
2. Surat Pernyataan dan Klarifikasi Artikel Bukan Bagian dari Desertasi
3. Tabel Of Conten Jurnal “Indonesian Journal of Chemistry” Vol 18, No 1 (2018)
4. Bukri Tim Editor dari Lebih 4 Negara
5. Abstrak Artikel dari 2 Negara ke 1
6. Abstrak Artikel dari 2 Negara ke 2
7. Abstrak Arikel yang bersangkutan
8. Bukti Jurnal terindeks SCOPUS
9. Bukti Abstrak artikel yang bersangkutan di SCOPUS
10. Bukti Impac Faktor (SJR) Scimago Journal & Country Rank
11. Kronologi Proses Submisi Artikel Pada Jurnal Hingga Terbit
12. Surat Keterangan Penyesuaian Judul Artikel dari Editor
13. Bukti korespondensi
14. Artikel lengkap
15. Hasil Turnitin

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel) : Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent

Jumlah Penulis : 3 orang

Status Pengusul : Penulis Utama

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : Indonesian Journal of Chemistry
- b. Nomor ISSN : 1411-9420 (print); 2460-1578 (on-line)
- c. Vol, No., Bln Thn : Vol 18, No 1 (2018), Hal. 159 -165
- d. Penerbit : Gadjah Mada University
- e. DOI artikel (jika ada) : <https://doi.org/10.22146/ijc.22197>
- f. Alamat web jurnal : <https://jurnal.ugm.ac.id/ijc/>
- g. Terindex : <https://jurnal.ugm.ac.id/ijc/article/view/22197/18879>
url Turnitin: (18%)
<https://doc-pak.undip.ac.id/2950/1/Turnitin%20C1.pdf>
Jurnal Internasional bereputasi (Q3) terindeks SCOPUS (SJR=0,215)
<https://www.scopus.com/sourceid/21100223536?origin=resultlist>

Kategori Publikasi Jurnal Ilmiah : ☒ Jurnal Ilmiah Internasional
(beri ✓ pada kategori yang tepat) ☐ Jurnal Ilmiah Nasional Terakreditasi
☐ Jurnal Ilmiah Nasional Tidak Terakreditasi


Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah			Nilai Akhir Yang Diperoleh
	Internasional	Nasional Terakreditasi	Nasional Tidak Terakreditasi	
	☒	☐	☐	
a. Kelengkapan unsur isi jurnal (10%)	4,00			4
b. Ruang lingkup dan kedalaman pembahasan (30%)	12,00			12
c. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12,00			12
d. Kelengkapan unsur dan kualitas terbitan/jurnal (30%)	12,00			11,6
Total = (100%)	40,00			39,6
Penulis Utama : $0,6 \times 39,6 = 23,76$				

Catatan Penilaian artikel oleh Reviewer :

- Kesesuaian dan kelengkapan unsur isi jurnal:**
Isi artikel ini sudah memenuhi kelengkapan sebuah jurnal ilmiah, unsur-unsur jurnal juga sudah sesuai dan sinkron. Judul abstrak, pendahuluan sampai kesimpulan lengkap. Nilai 4
- Ruang lingkup dan kedalaman pembahasan:**
Pembahasan pada makalah ini sudah baik. Metode penelitian dan hasil disampaikan dengan baik dan didukung oleh referensi yang memadai. Nilai 12
- Kecukupan dan kemutakhiran data/informasi dan metodologi:**
Metode yang digunakan dalam artikel ini cukup mutakhir dan referensi yang digunakan sudah mencukupi dan mutakhir (< 10 th), data baik dan dibahas dengan baik. Nilai 12
- Kelengkapan unsur dan kualitas terbitan:**
Terlihat jurnal ini berkuwalitas dan unsur-unsurnya lengkap mencakup cover, dewan redaksi dan daftar isi semua memenuhi syarat jurnal ilmiah. Nilai 11,6

Semarang, 30 Maret 2020
Reviewer 1



Dr. Bambang Cahyono
NIP. 196303161988101001
Unit Kerja : Departemen Kimia FSM UNDIP

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel) : Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent

Jumlah Penulis : 3 orang

Status Pengusul : Penulis Utama

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : Indonesian Journal of Chemistry
- b. Nomor ISSN : 1411-9420 (print); 2460-1578 (on-line)
- c. Vol, No., Bln Thn : Vol 18, No 1 (2018), Hal. 159 -165
- d. Penerbit : Gadjah Mada University
- e. DOI artikel (jika ada) : <https://doi.org/10.22146/ijc.22197>
- f. Alamat web jurnal : <https://jurnal.ugm.ac.id/ijc/>
- Alamat Artikel : <https://jurnal.ugm.ac.id/ijc/article/view/22197/18879>
- url Turnitin: (18%)
<https://doc-pak.undip.ac.id/2950/1/Turnitin%20C1.pdf>
- g. Terindex : Jurnal Internasional bereputasi (Q3) terindeks SCOPUS (SJR=0,215)
<https://www.scopus.com/sourceid/21100223536?origin=resultslist>

Kategori Publikasi Jurnal Ilmiah : ☒ Jurnal Ilmiah Internasional
 (beri ✓ pada kategori yang tepat) ☐ Jurnal Ilmiah Nasional Terakreditasi
☐ Jurnal Ilmiah Nasional Tidak Terakreditasi

Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Maksimal Jurnal Ilmiah			Nilai Akhir Yang Diperoleh
	Internasional ☒	Nasional Terakreditasi ☐	Nasional Tidak Terakreditasi ☐	
e. Kelengkapan unsur isi jurnal (10%)	4,00			4
f. Ruang lingkup dan kedalaman pembahasan (30%)	12,00			12
g. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12,00			12
h. Kelengkapan unsur dan kualitas terbitan/jurnal (30%)	12,00			11
Total = (100%)	40,00			39
Penulis Utama: 0,6 x 39 = 23,4				

Catatan Penilaian artikel oleh Reviewer :

- Kesesuaian dan kelengkapan unsur isi jurnal:**
 Lengkap dan sesuai dengan yang dipersyaratkan jurnal internasional.
 Artikel yang dimuat, dibahas dan disusun secara lengkap. Nilai 4
- Ruang lingkup dan kedalaman pembahasan:**
 Data yang disajikan dibahas dengan detil dan lengkap, disertai referensi pendukung. Nilai 12
- Kecukupan dan kemutakhiran data/informasi dan metodologi:**
 Kualitas dan kuantitas pustaka baik, sebagian besar tahun terbit kurang dari 10 th (90%).
 Metodologi disajikan detil sehingga bisa diulang peneliti lain. Nilai 12
- Kelengkapan unsur dan kualitas terbitan:**
 Kualitas terbitan baik dan masuk dalam Q3, hanya ada beberapa gambar yang kurang jelas. Nilai 11

Semarang, 23 Maret 2020
 Reviewer 2

Drs. Gunawan, M.Si, Ph.D
 NIP.196408251991031001
 Unit Kerja : Departemen Kimia FSM UNDIP

LEMBAR
HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH : JURNAL ILMIAH

Judul Jurnal Ilmiah (Artikel) : Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent

Jumlah Penulis : 3 orang

Status Pengusul : Penulis Utama

Identitas Jurnal Ilmiah :

- a. Nama Jurnal : Indonesian Journal of Chemistry
- b. Nomor ISSN : 1411-9420 (print); 2460-1578 (on-line)
- c. Vol, No., Bln Thn : Vol 18, No 1 (2018), Hal. 159 -165
- d. Penerbit : Gadjah Mada University
- e. DOI artikel (jika ada) : <https://doi.org/10.22146/ijc.22197>
- f. Alamat web jurnal : <https://jurnal.ugm.ac.id/ijc/>
- Alamat Artikel : <https://jurnal.ugm.ac.id/ijc/article/view/22197/18879>
- url Turnitin: (18%)
<https://doc-pak.undip.ac.id/2950/1/Turnitin%20C1.pdf>
- g. Terindex : Jurnal Internasional bereputasi (Q3) terindeks SCOPUS (SJR=0,215)
<https://www.scopus.com/sourceid/21100223536?origin=resultslist>

Kategori Publikasi Jurnal Ilmiah : ☒ Jurnal Ilmiah Internasional
 (beri ✓ pada kategori yang tepat) ☐ Jurnal Ilmiah Nasional Terakreditasi
☐ Jurnal Ilmiah Nasional Tidak Terakreditasi

Hasil Penilaian *Peer Review* :

Komponen Yang Dinilai	Nilai Reviewer		Nilai Rata-rata
	Reviewer I	Reviewer II	
a. Kelengkapan unsur isi jurnal (10%)	4	4	4
b. Ruang lingkup dan kedalaman pembahasan (30%)	12	12	12
c. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12	12	12
d. Kelengkapan unsur dan kualitas terbitan/jurnal (30%)	11,6	11	11,3
Total = (100%)	39,6	39	39,3
Penulis Utama (rata-rata): $0,6 \times 39,3 = 23,58$			

Semarang, 30 Maret 2020

Reviewer 2

Drs. Gunawan, M.Si, Ph.D
 NIP.196408251991031001
 Unit Kerja : Departemen Kimia FSM UNDIP

Reviewer 1

Dr. Bambang Cahyono
 NIP. 196303161988101000
 Unit Kerja : Departemen Kimia FSM UNDIP

SURAT PERNYATAAN DAN KLARIFIKASI ARTIKEL BUKAN BAGIAN DARI DESERTASI

Yang bertanda tangan dibawah ini saya:

Nama : Sriatun, S.Si, M.Si
NIP : 197103151997022001
Departemen : Kimia
Fakultas : Sains dan Matematika
Jenis : artikel penelitian
Judul : Synthesis of Zeolite from Sugarcane Bagasse Ash Using
Cetyltrimethylammonium Bromide as Structure Directing Agent

Menyatakan dan mengklarifikasi bahwa artikel tersebut bukan merupakan bagian dari desertasi S3 saya. Adapun sebagai penguat hal tersebut adalah:

1. Artikel tersebut merupakan luaran penelitian yang berjudul **Pengembangan Katalis Heterogen Berbasis Silika Alumina dari Ampas tebu dan Aplikasinya pada Pembuatan Biodiesel dari Minyak Goreng Sisa Pakai** dari skim hibah bersaing tahun pelaksanaan 2014, dengan perjanjian pelaksanaan penelitian No. 84-68/UN7.5.1/PG/2014. Dalam Surat Pernyataan Validasi Karya Ilmiah, asal usul karya ilmiah dan sumber pendanaan telah disebutkan bahwa artikel tersebut berasal dari Program Hibah Bersaing tahun 2014 dengan nomer kontrak seperti tersebut di atas.
2. Pada bagian Acknowledgment dari artikel tersebut sudah tertulis sebagai berikut:
Acknowledgment
Sriatun, Taslimah, and Suyati, L., gratefully acknowledge financial support from The Directorate of Research and Community Service, Ministry of Research, Technology and High Education via Grants Compete Award (Number: 84-68/UN7.5.1/PG/2014) and Chemistry Department of Diponegoro University for the facilities to carry out this research.
3. Artikel tersebut saya submit pada jurnal Indonesian Journal of Chemistry UGM pada tanggal **18 Februari 2017**.

[Home](#) > [User](#) > [Author](#) > [Submissions](#) > #22197 > [Summary](#)

#22197 Summary

[SUMMARY](#) [REVIEW](#) [EDITING](#)

Submission


Authors	Sriatun Sriatun, Taslimah Taslimah, Linda Suyati
Title	Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent
Original file	22197-41689-1-SM.DOCX 2017-02-18
Supp. files	None
Submitter	Mrs Sriatun Sriatun ☐
Date submitted	February 18, 2017 - 09:27 PM
Section	Articles
Editor	Nuryono Nuryono ☐
Abstract Views	1540

Saya terdaftar sebagai mahasiswa S3 pada Departemen Teknik Kimia Fakultas Teknik UNDIP pada bulan September 2016, dimana pada masa submit artikel saya belum ada penentuan judul disertasi serta kegiatan laboratorium terkait studi S3 saya.

4. Promotor studi S3 : Prof. Dr.rer.nat Heru Susanto, ST, MT, MM
Co-promotor 1 : Prof. Dr. Ir. Widayat, ST, MT
Co-promotor 2 : Adi Darmawan, S.Si, M.Si, Ph.D
Pada artikel tersebut tidak terdapat nama promotor maupun co-promotor pada daftar author (penulis).

Demikian surat klarifikasi ini saya buat dengan sebenar-benarnya.


Mengetahui
Promotor/Co-promotor



Prof. Dr. Ir. Widayat, ST, MT
NIP. 197206091998031001

Semarang, 16 Oktober 2020

membuat



Sriatun, S.Si, M.Si
NIP. 197103151997022001



Menu

[Home](#) [About](#) [Login](#) [Register](#) [Search](#) [Current](#) [Archives](#) [Announcements](#) [Statistics](#) [Indexing & Abstracting](#) [Journal History](#) [Contact](#)
[Home](#) > [Archives](#) > **Vol 18, No 1 (2018)**

Vol 18, No 1 (2018)

Accredited by DIKTI No : 58/DIKTI/Kep/2013

Date : 22 August 2013



Table of Contents

Articles

- | | |
|---|-------|
| <p>The Modification of PVDF Membrane via Crosslinking with Chitosan and Glutaraldehyde as the Crosslinking Agent</p> <p><i>Romaya Sitha Silitonga, Nurul Widiastuti, Juhana Jaafar, Ahmad Fauzi Ismail, Muhammad Nidzhom Zainol Abidin, Ihsan Wan Azelee, Mahesan Naidu</i></p> <p> 10.22146/ijc.25127 Abstract views : 2339 PDF views : 2949</p> | 1-6 |
| <p>Physical Mixture Interaction of Acetaminophenol with Naringenin</p> <p><i>Normyzatul Akmal Abd Malek, Hamizah Mohd Zaki, Mohammad Noor Jalil</i></p> <p> 10.22146/ijc.26732 Abstract views : 1166 PDF views : 1227</p> | 18-25 |
| <p>An Ellagic Acid Derivative and Its Antioxidant Activity of Stem Bark Extracts of <i>Syzygium polycepalum</i> Miq. (Myrtaceae)</p> <p><i>Tukiran Tukiran, Andika Pramudya Wardana, Nurul Hidayati, Kuniyoshi Shimizu</i></p> <p> 10.22146/ijc.25467 Abstract views : 2291 PDF views : 2643</p> | 26-34 |
| <p>One-Pot Eco-Safe Saccharin-Catalyzed Procedure for Expedient and Convenient Synthesis of Dihydropyrano[2,3-c]pyrazole, Tetrahydrobenzo[b]pyran and Pyrano[2,3-d]pyrimidinone Scaffolds as a Green and Versatile Catalyst</p> <p><i>Farzaneh Mohamadpour, Mojtaba Lashkari, Nourallah Hazeri</i></p> <p> 10.22146/ijc.28450 Abstract views : 1750 PDF views : 1674</p> | 7-17 |
| <p>Liquid Chromatography Mass Spectrometry (LC-MS) Fingerprint Combined with Chemometrics for Identification of Metabolites Content and Biological Activities of <i>Curcuma aeruginosa</i></p> <p><i>Dewi Anggraini Septaningsih, Latifah Kosim Darusman, Farit Mochamad Afendi, Rudi Heryanto</i></p> <p> 10.22146/ijc.25456 Abstract views : 3240 PDF views : 4581</p> | 43-52 |
| <p>Cytotoxic Triterpenoids from the Stembark of <i>Aglaia argentea</i> (Meliaceae)</p> <p><i>Ace Tatang Hidayat, Kindi Farabi, Desi Harneti, Nurlelasari Nurlelasari, Rani Maharani, Ida Nurfarida,</i></p> | 35-42 |

Subscribing on:



ARTICLE IN PRESS

List of the accepted articles for future issues.

FUTURE ISSUES

Vol. 20, no. 6 (December 2020).

[Focus & Scope](#)[Author Guidelines](#)[Author Fees](#)[Online Submission](#)[Publication Ethics](#)[Plagiarism Policy](#)[Editorial Board](#)[Peer Reviewers](#)[Order Journal](#)[Visitor Statistics](#)

USER

Username

Password

☐ Remember me

JOURNAL CONTENT

Search

Search Scope

Browse

- [▶ By Issue](#)
- [▶ By Author](#)
- [▶ By Title](#)
- [▶ Other Journals](#)

INFORMATION

- [▶ For Readers](#)
- [▶ For Authors](#)
- [▶ For Librarians](#)

Unang Supratman, Yoshihito Shiono

 10.22146/ijc.25052  Abstract views : 1552 |  views : 1642

A Green Synthesis of a Novel Calix[4]Resorcinarene from 7-Hydroxycitronellal Using Microwave Irradiation

53-59

Ratnaningsih Eko Sardjono, Iqbal Musthapa, Iis Rosliana, Fitri Khoerunnisa, Galuh Yuliani

 10.22146/ijc.25466  Abstract views : 1601 |  views : 1683
Structure-Activity Relationship Study on the Ethyl *p*-Methoxycinnamate as an Anti-Inflammatory Agent

60-65

Ismiarni Komala, Supandi Supandi, Nurhasni Nurhasni, Ofa Suzanti Betha, Eka Putri, Syarifatul Mufidah, Muhammad Fikry Awaludin, Mida Fahmi, Muhammad Reza, Nurkhayati Putri Indriyani

 10.22146/ijc.26162  Abstract views : 1884 |  views : 1964
Antidiabetic Activities and Identification of Chemical Compound from Samama (*Anthocephalus macrophyllus* (Roxb) Havil)

66-74

Laela Nur Anisah, Wasrin Syafii, Gustan Pari, Rita Kartika Sari

 10.22146/ijc.25492  Abstract views : 1745 |  views : 2187
The Effect of *Pseudomonas aeruginosa* Addition on 1,1,1-Trichloro-2,2-bis(4-chlorophenyl)ethane (DDT) Biodegradation by Brown-rot Fungus *Fomitopsis pinicola*

75-81

Atmira Sariwati, Adi Setyo Purnomo

 10.22146/ijc.25158  Abstract views : 1455 |  views : 1442

The Physico-Mechanical Properties and Release Kinetics of Eugenol in Chitosan-Alginate Polyelectrolyte Complex Films as Active Food Packaging

82-91

Baiq Amelia Riyandari, Suherman Suherman, Dwi Siswanta

 10.22146/ijc.26525  Abstract views : 1625 |  views : 1679

Synthesis and Characterization of Chitosan Linked by Methylene Bridge and Schiff Base of 4,4'-Diaminodiphenyl Ether-Vanillin

92-101

Ahmad Fatoni, Poedji Loekitowati Hariani, Hermansyah Hermansyah, Aldes Lesbani

 10.22146/ijc.25866  Abstract views : 2607 |  views : 3322
Synthesis of Chalcone Derivatives and Their *in vitro* Anticancer Test Against Breast (T47D) and Colon (WiDr) Cancer Cell Line

102-107

Chairil Anwar, Yogo Dwi Prasetyo, Sabirin Matsjeh, Winarto Haryadi, Eti Nurwening Sholikhah, Nendrowati Nendrowati

 10.22146/ijc.26864  Abstract views : 1875 |  views : 2502

Synthesis and Kinetic Study of the Urea Controlled Release Composite Material: Sodium Lignosulfonate from Isolation of Wood Sawdust-Sodium Alginate-Tapioca

108-115

Arif Cahyo Imawan, Yehezkiel Steven Kurniawan, Muhammad Fernadi Lukman, Jumina Jumina, Triyono Triyono, Dwi Siswanta

 10.22146/ijc.26597  Abstract views : 1981 |  views : 1212 |  views : 1235

A New Thiourea Compound as Potential Ionophore for Metal Ion Sensor

116-120

Fatimatul Akma Awang Ngah, Emma Izzati Zakariah, Imran Fakhar, Nurul Izzaty Hassan, Lee Yook Heng, Bohari Yamin, Siti Aishah Hasbullah

 10.22146/ijc.27078  Abstract views : 1482 |  views : 1583

Synthesis of A Novel Carrier Compound Thiazoethyl Methyl Eugenoxycetate from Eugenol and Its Use in the Bulk Liquid Membrane Technique

121-126

Muhammad Cholid Djunaidi, Pratama Jujur Wibawa, Ratna Hari Murti

 10.22146/ijc.25075  Abstract views : 1614 |  views : 1585
Improvement of Cycling Performance of $\text{Na}_{2/3}\text{Co}_{2/3}\text{Mn}_{1/3}\text{O}_2$ Cathode by PEDOT/PSS Surface Coating for Na Ion Batteries

127-136

Yatim Lailun Ni'mah, Ju Hsiang Cheng, Ming Yao Cheng, Wei Nien Su, Bing Joe Hwang

 10.22146/ijc.24893  Abstract views : 1760 |  views : 1727

Spatial Distribution of Heavy Metals in the Surface Sediments of the Southern Coast of Pacitan, Indonesia

137-144

Defri Yona, Mochamad Arif Zainul Fuad, Nurin Hidayati

 10.22146/ijc.22400  Abstract views : 1244 |  views : 1548

Activated Carbon from Spent Brewery Barley Husks for Cadmium Ion Adsorption from Aqueous Solution

145-152

Ilesanmi Osasona, Kayode Aiyedatiwa, Jonathan Johnson, Oluwabamise Lekan Faboya

 10.22146/ijc.22422  Abstract views : 1439 |  views : 1765

KEYWORDS

HPLC QSAR TiO₂

adsorption

antioxidant catalyst

characterization chitosan

eugenol extraction fly ash

immobilization kinetics

methylene blue molecular

docking photocatalyst silica

sol-gel synthesis

transesterification zeolite


Indones. J. Chem.
indexed by:

Scopus®

Indonesian Journal of
ChemistryQ3 Chemistry
(miscellaneous)
best quartileSJIR 2019
0.24

powered by scimagojr.com

CURRENT ISSUE



Artikel 2 negara, abstrak terlampir

Artikel 2 negara, abstrak terlampir

- Synthesis, Radiochemical Purity Control and Stability of Scandium-46-1,4,7-triazacyclononane-1,4,7-triacetic acid (^{46}Sc -NOTA) 153-158
Duyeh Setiawan, Iwan Hastiawan, Asri Nurul Bashiroh
 10.22146/ijc.24467  Abstract views : 1835 |  views : 1542
- Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent 159-165
Sriatun Sriatun, Taslimah Taslimah, Linda Suyati
 10.22146/ijc.22197  Abstract views : 1538 |  views : 2163
- Hydrocracking of α -Cellulose Using Co, Ni, and Pd Supported on Mordenite Catalysts 166-172
Wega Trisunaryanti, Triyono Triyono, Ria Armunanto, Lathifah Puji Hastuti, Desinta Dwi Ristiana, Resi Vita Ginting
 10.22146/ijc.26491  Abstract views : 2133 |  views : 2124
- Density Functional Theory (DFT) and Natural Bond Orbital (NBO) Analysis of Intermolecular Hydrogen Bond Interaction in "Phosphorylated Nata De Coco - Water" 173-178
Sitti Rahmawati, Cynthia Linaya Radiman, Muhamad Abdulkadir Martoprawiro
 10.22146/ijc.25170  Abstract views : 1645 |  views : 2077
- Molecular Docking Analysis of Ginger Active Compound on Transient Receptor Potential Cation Channel Subfamily V Member 1 (TRPV1) 179-185
Fifteen Aprila Fajrin, Agung Endro Nugroho, Rina Susilowati, Arief Nurrochmad
 10.22146/ijc.28172  Abstract views : 1448 |  views : 1737
- The Role of E27-K31 and E56-K10 Salt-Bridge Pairs in the Unfolding Mechanism of the B1 Domain of Protein G 186-193
Tony Ibnu Sumaryada, Kania Nur Sawitri, Setyanto Tri Wahyudi
 10.22146/ijc.23934  Abstract views : 1391 |  views : 1582



Menu

[Home](#) [About](#) [Login](#) [Register](#) [Search](#) [Current](#) [Archives](#) [Announcements](#) [Statistics](#) [Indexing & Abstracting](#) [Journal History](#) [Contact](#)
[Home](#) > [About the Journal](#) > **Editorial Team**

Editorial Team (Terdiri dari lebih 4 negara sebagai editor)

Editor-in-Chief

Nuryono Nuryono, Laboratory of Inorganic Chemistry, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Managing Editor

Dwi Siswanta, Laboratory of Analytical Chemistry, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Editorial Board

Akhmad Syoufian, Laboratory of Physical Chemistry, Department of Chemistry, Universitas Gadjah Mada

Hideaki Hisamoto, Department of Applied Chemistry, Osaka Prefecture University, Japan

Dr. Hendrik Oktendy Lintang, Indonesia

Iqmal Tahir, Laboratory of Physical Chemistry, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Indriana Kartini, Laboratory of Inorganic Chemistry, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Prof. Joe da Costa, Department of Chemical Engineering, University of Queensland, Australia

Dr. Maurizio Barbieri, Department of Earth Science, Sapienza University, Italy

Ming Cai, Tongji University, China

Muhammad Idham Darussalam Mardjan, Laboratory of Organic Chemistry, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Pornthep Sompornpisut, Department of Chemistry, Chulalongkorn University, Thailand

Praveen Kumar Sharma, Department of Chemistry, Lovely Professional University, Punjab, India

Assoc. Prof. Dr. Roswanira Abdul Wahab, Department of Chemistry, Universiti Teknologi Malaysia, Malaysia

Dr. Saprizal Hadisaputra, Chemistry Education Division, Faculty of Science and Education, University of Mataram, Indonesia

Prof. Dr. Taghreed Hashim Al-Noor, Ibn-Al-Haitham Education College /University of Baghdad, Iraq

Tutik Dwi Wahyuningsih, Laboratory of Organic Chemistry, Department of Chemistry, Universitas Gadjah Mada

Satya Candra Wibawa Sakti, Laboratory of Inorganic Chemistry, Department of Chemistry, Universitas Airlangga, Indonesia

Tri Joko Raharjo, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Winifred Uduak Anake, Department of Chemistry, College of Science and Technology, Covenant University, Nigeria

Administration Support

Djoko Prihandono, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Aulia Sukma Hutama, Laboratory of Physical Chemistry, Department of Chemistry, Universitas Gadjah Mada, Indonesia

Aulia Ratri Hapsari, Department of Chemistry, Faculty of Mathematics and Natural Sciences, Universitas Gadjah Mada, Indonesia

Indonesian Journal of Chemistry (ISSN 1411-9420 / 2460-1578) - Chemistry Department, Universitas Gadjah Mada, Indonesia.

01457551 View The Statistics of Indones. J. Chem.

Subscribing on:



ARTICLE IN PRESS

List of the accepted articles for future issues.

FUTURE ISSUES

Vol 20 no 6 (December 2020).

[Focus & Scope](#)
[Author Guidelines](#)
[Author Fees](#)
[Online Submission](#)
[Publication Ethics](#)
[Plagiarism Policy](#)
[Editorial Board](#)
[Peer Reviewers](#)
[Order Journal](#)
[Visitor Statistics](#)

USER

Username

Password

☐ Remember me

JOURNAL CONTENT

Search

Search Scope

All

Browse

- ▶ [By Issue](#)
- ▶ [By Author](#)
- ▶ [By Title](#)
- ▶ [Other Journals](#)

INFORMATION

- ▶ [For Readers](#)
- ▶ [For Authors](#)
- ▶ [For Librarians](#)



Abstrak artikel 2 negara ke 1

Activated Carbon from Spent Brewery Barley Husks for Cadmium Ion Adsorption from Aqueous Solution

<https://doi.org/10.22146/ijc.22422>

Ilesanmi Osasona^(1*), Kayode Aiyedatiwa⁽²⁾, Jonathan Johnson⁽³⁾, Oluwabamise Lekan Faboya⁽⁴⁾

(1) Afe Babalola University, Ado - Ekiti

(2) Afe Babalola University, Ado - Ekiti

(3) Afe Babalola University, Ado - Ekiti

(4) Afe Babalola University, Ado - Ekiti

(*) Corresponding Author

Abstract

This study investigated the feasibility of using acid activated carbon prepared from brewery spent barley husks for the adsorption of cadmium from aqueous solution. The effects of operation parameters such as pH, contact time, adsorbent dosage, concentration and temperature were verified. The amount of cadmium adsorbed increased with increase in solution pH, initial solution concentration and with the amount of adsorbent dosed. A time of 5 minutes was required for attainment of equilibrium. The equilibrium data obtained were analysed using both Langmuir and Freundlich isotherm models and the data were better described by Langmuir model with correlation coefficient of 0.9183. The thermodynamic parameters revealed that the removal of cadmium by the activated carbon was exothermic and spontaneous. Thus, activated carbon obtained from brewery spent barley husk can be employed as an economically viable low-cost adsorbent for removing cadmium from aqueous solution.

Keywords

Cadmium; adsorbent; thermodynamic; equilibrium

Full Text:

References

- [1] Volesky, B., and Holan, Z.R., 1995, Biosorption of heavy metals, *Biotechnol. Prog.*, 11 (3), 235–250.
- [2] Pagana, A.E., Sklar, S.D., Kikkinides, E.S., and Zaspalis, V.T., 2008, Microporous ceramic membrane technology for the removal of arsenic and chromium ions from contaminated water, *Microporous Mesoporous Mater.*, 110 (1), 150–156.
- [3] Donia, A.M., Atia, A.A., and Mabrouk, D.H., 2011, Fast kinetic and efficient removal of As(V) from aqueous solution using anion exchange resins, *J. Hazard. Mater.*, 191 (1-3), 1–7.
- [4] Sud, D., Mahajan, G., and Kaur, M.P., 2008, Agricultural waste material as potential adsorbent for sequestering heavy metal ions from aqueous solutions - A review, *Bioresour. Technol.*, 99 (14), 6017–6027.
- [5] Khezami, L., and Capart, R., 2005, Removal of chromium(VI) from aqueous solution by activated carbons: Kinetic and equilibrium studies, *J. Hazard. Mater.*, 123 (1-3), 223–231.
- [6] Osasona, I., Adebayo, A.O., and Ajayi, O.O., 2013, Adsorptive removal of chromium(VI) from aqueous solution using cow hooves, *J. Sci. Res. Rep.*, 2 (1), 288–303.
- [7] Nasuha, N., Hameed, B.H., and Mohd Din, A.T., 2010, Rejected tea as a potential low-cost adsorbent for the removal of methylene blue, *J. Hazard. Mater.*, 175 (1-3), 126–132.
- [8] Musatto, S.I., 2014, Brewer's spent grain: A valuable feedstock for industrial applications, *J. Sci. Food Agric.*, 94 (7), 1264–1275.

Subscribing on:



ARTICLE IN PRESS

List of the accepted articles for future issues.

FUTURE ISSUES

Vol 20 no 6 (December 2020).

[Focus & Scope](#)

[Author Guidelines](#)

[Author Fees](#)

[Online Submission](#)

[Publication Ethics](#)

[Plagiarism Policy](#)

[Editorial Board](#)

[Peer Reviewers](#)

[Order Journal](#)

[Visitor Statistics](#)

USER

Username

Password

☐ Remember me

ABOUT THE AUTHORS

Ilesanmi Osasona

✉ *Corresponding Author

Afe Babalola University, Ado - Ekiti
Nigeria
Department of Chemical Sciences

Kayode Aiyedatiwa

Afe Babalola University, Ado - Ekiti
Nigeria
Department of Chemical Sciences

Jonathan Johnson

Afe Babalola University, Ado - Ekiti
Nigeria
Department of Chemical Sciences

Oluwabamise Lekan Faboya

Afe Babalola University, Ado - Ekiti



Abstrak artikel 2 negara ke 2

Improvement of Cycling Performance of Na_{2/3}Co_{2/3}Mn_{1/3}O₂ Cathode by PEDOT/PSS Surface Coating for Na Ion Batteries

<https://doi.org/10.22146/ijc.24893>

Yatim Lailun Ni'mah^(1*), Ju Hsiang Cheng⁽²⁾, Ming Yao Cheng⁽³⁾, Wei Nien Su⁽⁴⁾, Bing Joe Hwang⁽⁵⁾

(1) Department of Chemistry, Institut Teknologi Sepuluh Nopember

(2) Department of Chemical Engineering, National Taiwan University of Science and Technology

(3) Department of Chemical Engineering, National Taiwan University of Science and Technology

(4) Department of Chemical Engineering, National Taiwan University of Science and Technology

(5) Department of Chemical Engineering, National Taiwan University of Science and Technology National Synchrotron Radiation Research Center

(*) Corresponding Author

Abstract

The surface-modified Na_{2/3}Co_{2/3}Mn_{1/3}O₂ is coated with a conductive Poly (3,4-Ethylene dioxy thiophene)-poly (styrene sulfonate) (PEDOT/PSS) polymer, and their resulting electrochemical properties were investigated as Na-ion battery cathode. The surface-modified Na_{2/3}Co_{2/3}Mn_{1/3}O₂ cathode material exhibits a high discharge capacity and good rate capability due to enhanced electron transport by surface PEDOT/PSS. The presence of PEDOT/PSS surface layer suppresses the growth of a resistive layer, while the dissolution of transition metals of the active cathode materials is inhibited as well. The resulting surface-modified Na_{2/3}Co_{2/3}Mn_{1/3}O₂ shows superior cycling performance, which is much stable than the pristine one as being the Na-ion battery cathode.

Keywords

Sodium ion battery; PEDOT/PSS; cathode; surface coating

Full Text:

References

- [1] Armand, M., and Tarascon, J.M., 2008, Building better batteries, *Nature*, 451 (7179), 652–657.
- [2] Zu, C.X., and Li, H., 2011, Thermodynamic analysis on energy densities of batteries, *Energy Environ. Sci.*, 4 (8), 2614–2624.
- [3] Yang, Z., Zhang, J., Kintner-Meyer, M.C.W., Lu, X., Choi, D., Lemmon, J.P., and Liu, J., 2011, Electrochemical energy storage for green grid, *Chem. Rev.*, 111 (5), 3577–3613.
- [4] Palomares, V., Serras, P., Villaluenga, I., Hueso, K.B., Carretero-González, J., and Rojo, T., 2012, Na-ion batteries, recent advances and present challenges to become low cost energy storage systems, *Energy Environ. Sci.*, 5, 5884–5901.
- [5] Wang, L., Lu, Y., Liu, J., Xu, M., Cheng, J., Zhang, D., and Goodenough, J.B., 2013, A superior low-cost cathode for a Na-ion battery, *Angew. Chem, Int. Ed.*, 52, 1964–1967.
- [6] Berthelot, R., Carlier, D., and Delmas, C., 2011, Electrochemical investigation of P2-Na_xCoO₂ phase diagram, *Nat. Mater.*, 10 (1), 74–80.
- [7] Cao, Y., Xiao, L., Wang, W., Choi, D., Nie, Z., Yu, J., Saraf, L.V., Yang, Z., and Liu, J., 2011, Reversible sodium ion insertion in single crystalline manganese oxide nanowires with long cycle life, *Adv. Mater.*, 23 (28), 3155–3160.
- [8] Yamada, Y., Doi, T., Tanaka, I., Okada, S., and Yamaki, J., 2011, Liquid-phase synthesis of highly dispersed NaFeF₃ particles and their electrochemical properties for sodium-ion batteries, *J. Power Sources*, 196 (10), 4837–4841.
- [9] Lee, K.T., Ramesh, T.N., Nan, F., Botton, G., and Nazar, L.F., 2011, Topochemical synthesis of sodium metal phosphate olivines for sodium-

Subscribing on:



ARTICLE IN PRESS

List of the accepted articles for future issues.

FUTURE ISSUES

Vol 20 no 6 (December 2020).

[Focus & Scope](#)

[Author Guidelines](#)

[Author Fees](#)

[Online Submission](#)

[Publication Ethics](#)

[Plagiarism Policy](#)

[Editorial Board](#)

[Peer Reviewers](#)

[Order Journal](#)

[Visitor Statistics](#)

USER

Username

Password

☐ Remember me

ABOUT THE AUTHORS

Yatim Lailun Ni'mah

*Corresponding Author
Department of Chemistry,
Institut Teknologi Sepuluh
Nopember

Ju Hsiang Cheng

Department of Chemical
Engineering, National Taiwan
University of Science and
Technology
Taiwan, Province of China

Ming Yao Cheng

Department of Chemical
Engineering, National Taiwan
University of Science and
Technology
Taiwan, Province of China

Wei Nien Su

Department of Chemical
Engineering, National Taiwan
University of Science and
Technology



Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent

<https://doi.org/10.22146/ijc.22197>

Sriatun Sriatun^(1*), Taslimah Taslimah⁽²⁾, Linda Suyati⁽³⁾

(1) Diponegoro University

(2) Diponegoro University

(3) Diponegoro University

(*) Corresponding Author

Abstract

The purpose of this research is to synthesize zeolite from sugarcane bagasse ash using cetyltrimethylammonium bromide as structure directing agent. This research used cetyltrimethylammonium bromide surfactant to invent the high porosity, surface area, acidity and thermal stability of synthesized zeolite. The Silica was extracted by alkaline fusion using sodium hydroxide solution. The synthesis was conducted by hydrothermal process at 100 °C for 7 days, ageing process for 24 h and calcination at 500 °C for 5 h. The ratio of Si/Al (v/v) was 1, 15 and 25, meanwhile the concentration of cetyltrimethylammonium bromide was 5×10^{-4} M, 1×10^{-3} M and 1×10^{-2} M. The result showed all of product have strong absorbance at $950\text{--}1050\text{ cm}^{-1}$ and $620\text{--}690\text{ cm}^{-1}$, $420\text{--}460\text{ cm}^{-1}$, double ring at $520\text{--}570\text{ cm}^{-1}$, pore opening at $300\text{--}370\text{ cm}^{-1}$. Vibration of -OH as silanol group or water was indicated by broad absorbance at $3400\text{--}3450\text{ cm}^{-1}$. The diffractograms XRD showed that the product had high crystallinity. The composition of product on ratio Si/Al 1 with concentration of cetyltrimethylammonium 10^{-2} M is sodalite, the ratio Si/Al 15 and 25 are NaP1 and SiO₂ quartz and contain 12.23% and 12.19% of Si, 4.17% and 13.18% of Al, respectively. Observation on SEM revealed that the crystal produced using cetyltrimethylammonium bromide were homogenous and regular in shape.

Keywords

surfactant; cetyltrimethylammonium bromide; synthesis; zeolite; sugarcane bagasse

Full Text:

References

- [1] Srinivasan, R., and Sathiya, K., 2010, Experimental study on bagasse ash in concrete, *Int. J. Serv. Learn. Eng.*, 5 (2), 60–66.
- [2] *Indonesian Statistics*, 2012, <http://www.bps.go.id>.
- [3] Hanafi, A., and Nandang, A., 2010, Studi pengaruh bentuk silika dari ampas tebu terhadap kekuatan produk keramik, *Jurnal Kimia Indonesia*, 5 (1), 35–38.
- [4] Norsuraya, S., Fazlena, H., and Norhasyimi, R., 2016, Sugarcane bagasse as a renewable Source of silica to synthesize Santa Barbara Amorphous-15 (SBA-15), *Procedia Eng.*, 148, 839–846.
- [5] Cardoso, A.M., Paprocki, A., Ferret, L.S., Azevedo, C.M.N., and Pires, M., 2015, Synthesis of zeolite Na-P1 under mild conditions using Brazilian coal fly ash and its application in wastewater treatment, *Fuel*, 139, 59–67.
- [6] Moisés, M.P., da Silva, C.T.P., Meneguín, J.G., Girotto, E.M., and Radovanovic, E., 2013, Synthesis of zeolite NaA from sugarcane bagasse ash, *Mater. Lett.*, 108, 243–246.
- [7] Maia, A.Á.B., Neves, R.F., Angélica, R.S., and Pöhlmann, H., 2015, Synthesis, optimisation and characterisation of the zeolite NaA using kaolin waste from the Amazon Region. Production of zeolites KA, MgA and CaA, *Appl. Clay Sci.*, 108, 55–60.

Subscribing on:



ARTICLE IN PRESS

List of the accepted articles for future issues.

FUTURE ISSUES

Vol 20 no 6 (December 2020).

[Focus & Scope](#)

[Author Guidelines](#)

[Author Fees](#)

[Online Submission](#)

[Publication Ethics](#)

[Plagiarism Policy](#)

[Editorial Board](#)

[Peer Reviewers](#)

[Order Journal](#)

[Visitor Statistics](#)

USER

Username

Password

☐ Remember me

ABOUT THE AUTHORS

Sriatun Sriatun

✉ *Corresponding Author

Diponegoro University
Indonesia
Chemistry Department

Taslimah Taslimah

Diponegoro University
Indonesia
Chemistry Department

Linda Suyati

Diponegoro University
Indonesia
Chemistry Department

ARTICLE TOOLS

[Print this article](#)

[Indexing metadata](#)

[How to cite item](#)

[Email this article \(Login required\)](#)



Source details

Indonesian Journal of Chemistry

Open Access ⓘ

Scopus coverage years: 2010, from 2012 to Present

Publisher: Gadjah Mada University

ISSN: 1411-9420

Subject area: Chemistry: General Chemistry

[View all documents >](#)

[Set document alert](#)

[📁 Save to source list](#) [Journal Homepage](#)

CiteScore 2019
1.2

ⓘ

SJR 2019
0.242

ⓘ

SNIP 2019
0.832

ⓘ

[CiteScore](#) [CiteScore rank & trend](#) [Scopus content coverage](#)

Improved CiteScore methodology

CiteScore 2019 counts the citations received in 2016-2019 to articles, reviews, conference papers, book chapters and data papers published in 2016-2019, and divides this by the number of publications published in 2016-2019. [Learn more >](#)

CiteScore 2019 ▾

$$1.2 = \frac{399 \text{ Citations 2016 - 2019}}{344 \text{ Documents 2016 - 2019}}$$

Calculated on 06 May, 2020

CiteScoreTracker 2020 ⓘ

$$1.3 = \frac{455 \text{ Citations to date}}{360 \text{ Documents to date}}$$

Last updated on 07 September, 2020 • Updated monthly

CiteScore rank 2019 ⓘ

Category	Rank	Percentile
Chemistry		
General Chemistry	#254/398	36th

[View CiteScore methodology >](#) [CiteScore FAQ >](#) [Add CiteScore to your site 🔗](#)

About Scopus

[What is Scopus](#)
[Content coverage](#)
[Scopus blog](#)
[Scopus API](#)
[Privacy matters](#)

Language

[日本語に切り替える](#)
[切换到简体中文](#)
[切换到繁體中文](#)
[Русский язык](#)

Customer Service

[Help](#)
[Contact us](#)

ELSEVIER

[Terms and conditions ↗](#) [Privacy policy ↗](#)

Copyright © Elsevier B.V. All rights reserved. Scopus® is a registered trademark of Elsevier B.V.

We use cookies to help provide and enhance our service and tailor content. By continuing, you agree to the use of cookies.

RELX



< Back to results | 1 of 1

[↗ Export](#)
[⬇ Download](#)
[🖨 Print](#)
[✉ E-mail](#)
[📄 Save to PDF](#)
[★ Add to List](#)
[More... >](#)

View at Publisher

 Indonesian Journal of Chemistry [Open Access](#)
 Volume 18, Issue 1, 2018, Pages 159-165

Synthesis of zeolite from sugarcane bagasse ash using cetyltrimethylammonium bromide as structure directing agent [\(Article\)](#)

[\(Open Access\)](#)

 Sriatun [✉](#) Taslimah, Suyati, L. [👤](#)

Department of Chemistry, Faculty of Science and Mathematics, Diponegoro University, Jl. Prof. Soedarto, SH, Tembalang, Semarang, 50275, Indonesia

Abstract

[View references \(22\)](#)

The purpose of this research is to synthesize zeolite from sugarcane bagasse ash using cetyltrimethylammonium bromide as structure directing agent. This research used cetyltrimethylammonium bromide surfactant to invent the high porosity, surface area, acidity and thermal stability of synthesized zeolite. The Silica was extracted by alkaline fusion using sodium hydroxide solution. The synthesis was conducted by hydrothermal process at 100 °C for 7 days, ageing process for 24 h and calcination at 500 °C for 5 h. The ratio of Si/Al (v/v) was 1, 15 and 25, meanwhile the concentration of cetyltrimethylammonium bromide was 5×10^{-4} M, 1×10^{-3} M and 1×10^{-2} M. The result showed all of product have strong absorbance at $950\text{--}1050\text{ cm}^{-1}$ and $620\text{--}690\text{ cm}^{-1}$, $420\text{--}460\text{ cm}^{-1}$, double ring at $520\text{--}570\text{ cm}^{-1}$, pore opening at $300\text{--}370\text{ cm}^{-1}$. Vibration of --OH as silanol group or water was indicated by broad absorbance at $3400\text{--}3450\text{ cm}^{-1}$. The diffractograms XRD showed that the product had high crystallinity. The composition of product on ratio Si/Al 1 with concentration of cetyltrimethylammonium 10^{-2} M is sodalite, the ratio Si/Al 15 and 25 are NaP1 and SiO_2 quartz and contain 12.23% and 12.19% of Si, 4.17% and 13.18% of Al, respectively. Observation on SEM revealed that the crystal produced using cetyltrimethylammonium bromide was more homogenous and regular in shape. © 2018, Gadjah Mada University. All rights reserved.


SciVal Topic Prominence ⓘ

Topic: Coal Fly Ash | Zeolite | Kaolin

Prominence percentile: 96.330 ⓘ

Chemistry database information ⓘ

Substances



Author keywords

Cetyltrimethylammonium bromide Sugarcane bagasse Surfactant Synthesis Zeolite

Funding details

Funding sponsor	Funding number	Acronym
Universitas Diponegoro		UNDIP

Metrics ⓘ [View all metrics >](#)

1 Citation in Scopus
33rd percentile

0.11 Field-Weighted
Citation Impact



PlumX Metrics

 Usage, Captures, Mentions,
Social Media and Citations
beyond Scopus.

Cited by 1 document

Synthesis of silica-rich zeolite using quaternary ammonium-based templates

 Sriatun, S. , Susanto, H. , Widayat, W.
(2020) *Journal of Physics: Conference Series*
[View details of this citation](#)

Inform me when this document is cited in Scopus:

[Set citation alert >](#)
[Set citation feed >](#)

Related documents

A Brief Review on the Synthesis of Zeolites from Hazardous Wastes

 Mallapur, V.P. , Oubagaradin, J.U.K.
(2017) *Transactions of the Indian Ceramic Society*

Selective synthesis of zeolites A and X from two industrial wastes: Crushed stone powder and aluminum ash

 Kuroki, S. , Hashishin, T. , Morikawa, T.
(2019) *Journal of Environmental Management*

Zero-waste process for the transformation of a hazardous aluminum waste into a raw material to obtain zeolites

López-Delgado, A. , Robla, J.I. , Padilla, I.

Funding sponsor	Funding number	Acronym	(2020) Journal of Cleaner Production
	84-68/UN7.5.1/PG/2014		View all related documents based on references
Funding text Sriatun, Taslimah, and Suyati, L., gratefully acknowledge financial support from The Directorate of Research and Community Service, Ministry of Research, Technology and High Education via Grants Compete Award (Number: 84-68/UN7.5.1/PG/2014) and Chemistry Department of Diponegoro University for the facilities to carry out this research.			
			Find more related documents in Scopus based on:
			Authors > Keywords >

ISSN: 14119420
Source Type: Journal
Original language: English

DOI: 10.22146/ijc.22197
Document Type: Article
Publisher: Gadjah Mada University

References (22)

View in search results format >

☐ All ☐ Export ☐ Print ☐ E-mail ☐ Save to PDF ☐ Create bibliography

- ☐ 1 Srinivasan, R., Sathiya, K.
Experimental study on bagasse ash in concrete
(2010) *Int. J. Serv. Learn. Eng.*, 5 (2), pp. 60-66. Cited 62 times.
- ☐ 2 (2012)
<http://www.bps.go.id>
- ☐ 3 Hanafi, A., Nandang, A.
Studi pengaruh bentuk silika dari ampas tebu terhadap kekuatan produk keramik
(2010) *Jurnal Kimia Indonesia*, 5 (1), pp. 35-38. Cited 3 times.
- ☐ 4 Norsuraya, S., Fazlena, H., Norhasyimi, R.
Sugarcane Bagasse as a Renewable Source of Silica to Synthesize Santa Barbara Amorphous-15 (SBA-15) ([Open Access](#))

(2016) *Procedia Engineering*, 148, pp. 839-846. Cited 39 times.
<http://sciencedirect.proxy.undip.ac.id:2048/science/journal/18777058>
doi: 10.1016/j.proeng.2016.06.627

[View at Publisher](#)
- ☐ 5 Cardoso, A.M., Paprocki, A., Ferret, L.S., Azevedo, C.M.N., Pires, M.
Synthesis of zeolite Na-P1 under mild conditions using Brazilian coal fly ash and its application in wastewater treatment

(2015) *Fuel*, 139, pp. 59-67. Cited 86 times.
<http://www.journals.elsevier.com.proxy.undip.ac.id:2048/fuel/>
doi: 10.1016/j.fuel.2014.08.016

[View at Publisher](#)
- ☐ 6 Moisés, M.P., Da Silva, C.T.P., Meneguín, J.G., Giroto, E.M., Radovanovic, E.
Synthesis of zeolite NaA from sugarcane bagasse ash

(2013) *Materials Letters*, 108, pp. 243-246. Cited 46 times.
doi: 10.1016/j.matlet.2013.06.086

[View at Publisher](#)



SJR

Scimago Journal & Country Rank

Enter Journal Title, ISSN or Publisher Name

[Home](#)[Journal Rankings](#)[Country Rankings](#)[Viz Tools](#)[Help](#)[About Us](#)[?](#) [x](#)

Global Chemical Conference

Providing a Platform for The Industry to Collectively Innovate, Adapt & Overcome Problems

reutersevents.com

OPEN

Indonesian Journal of Chemistry

12

H Index

Country	Indonesia - IIII SIR Ranking of Indonesia
Subject Area and Category	Chemistry Chemistry (miscellaneous)
Publisher	Gadjah Mada University - IIII SCImago Institutions Ranking
Publication type	Journals
ISSN	14119420
Coverage	2010, 2012-2020
Scope	Indonesian Journal of Chemistry is a peer-reviewed, open access journal that publishes original research articles, review articles, as well as short communication in all areas of chemistry, including educational chemistry, applied chemistry, and chemical engineering.
	Homepage
	How to publish in this journal
	Contact
	Join the conversation about this journal

Free English Writing Tool

Makes sure everything you type is easy to read and mistake-free. Try now

Grammarly

DOWNLOAD

Global Chemical Conference


Providing a Platform for The Industry to Collectively Innovate, Adapt & Overcome Problems

reutersevents.com


OPEN

Quartiles


Chemistry (miscellaneous)




SJR




Citations per document




Total Cites Self-Cites



External Cites per Doc Cites per Doc



% International Collaboration



Citable documents Non-citable documents

Cited documents Uncited documents

400

400

**Indonesian Journal of
Chemistry**

Q3

Chemistry
(miscellaneous)

best quartile

SJIR 2019

0.24

powered by scimagojr.com

← Show this widget in
your own websiteJust copy the code below
and paste within your html
code:`<a href="https://www.scimaç`

Free English Writing Tool

Makes sure everything you type is easy to read and mistake-free. Try now

Grammarly

DOWNLOAD

U **Ulin Sahami** 8 months ago

Excellent

reply

M **Melanie Cornelia** 2 years ago

selamat pagi, saya dosen di Teknologi Pangan UPH....saya mau kirim article penelitian ke journal ini, rata-rata berapa lama review nya? terimakasih infonya

reply

M **Manton Rais** 2 years ago

nice

reply

Leave a comment

Name

Email

(will not be published)

KRONOLOGI PROSES SUBMISI ARTIKEL PADA JURNAL HINGGA TERBIT

Judul saat awal submit "***The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash***". Judul saat submit menjadi identitas judul ketika korespondensi maupun perbaikan artikel. Setelah direview oleh tim reviewer jurnal, penambahan data dan perbaikan narasi pada pembahasan serta kesimpulan oleh penulis (author), maka agar judul disesuaikan karena kata effect pada judul menjadi kurang tepat. Sehingga judul diganti menjadi "***Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent***"

Indonesian Journal of Chemistry

ISSN: 1411-9420 (print); 2460-1578 (online)

Published by Chemistry Department Universitas Gadjah Mada



Home | About | User Home | Search | Current | Archives | Announcements | Statistics | Indexing & Abstracting | Journal History | Contact

Home > User > Author > Archive

Archive

[ACTIVE](#) [ARCHIVE](#)

ID	MM-DD SUBMIT	SEC	AUTHORS	TITLE	VIEWS	STATUS
22197	02-18	ART	Sriatun, Taslimah, Suyati	SYNTHESIS OF ZEOLITE FROM SUGARCANE BAGASSE ASH USING...	2159	Vol 18, No 1 (2018)

1 - 1 of 1 Items

Start a New Submission

[CLICK HERE](#) to go to step one of the five-step submission process.

Subscribing on:



ARTICLE IN PRESS

List of the accepted articles for future issues.

FUTURE ISSUES

Vol. 20, no. 6 (December 2020).

Focus & Scope

Author Guidelines

Author Fees

Online Submission

Publication Ethics

Plagiarism Policy

Editorial Board

Indonesian Journal of Chemistry (ISSN 1411-9420 / 2460-1578) - Chemistry Department, Universitas Gadjah Mada, Indonesia.

[Home](#) > [User](#) > [Author](#) > [Submissions](#) > [#22197](#) > [Summary](#)

#22197 Summary

SUMMARY REVIEW EDITING

Submission

Authors	Sriatun Sriatun, Taslimah Taslimah, Linda Suyati
Title	Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent
Original file	22197-41689-1-SM.DOCX 2017-02-18
Supp. files	None
Submitter	Mrs Sriatun Sriatun
Date submitted	February 18, 2017 7:09:27 PM
Section	Articles
Editor	Nuryono Nuryono
Abstract Views	1535

Author Fees

Subscribing on:



ARTICLE IN PRESS

List of the accepted articles
for future issues.

FUTURE ISSUES

Vol. 20, no. 6 (December 2020).

Focus & Scope

Author Guidelines

Author Fees

Online Submission

Publication Ethics

Plagiarism Policy

Editorial Board

Peer Reviewers


Judul awal submission

Judul yang muncul di sini adalah judul setelah perubahan terakhir dan yang terpublikasi

JUDUL AWAL SUBMISSION

The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash.

Judul yang masuk ketika submit merupakan identitas artikel dalam korespondensi.



PROSES REVIEW

REVIEW ROUND 1

Indonesian Journal of Chemistry

ISSN: 1411-9420 (print); 2460-1578 (online)

Published by Chemistry Department Universitas Gadjah Mada

[Home](#) | [About](#) | [User Home](#) | [Search](#) | [Current](#) | [Archives](#) | [Announcements](#) | [Statistics](#) | [Indexing & Abstracting](#) | [Journal History](#) | [Contact](#)

Home > User > Author > Submissions > #22197 > Review

#22197 Review

SUMMARY
REVIEW
EDITING

Submission

Authors	Sriatun Sriatun, Taslimah Taslimah, Linda Suyati
Title	Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent
Section	Articles
Editor	Nuryono Nuryono

Peer Review

Round 1

Review Version	22197-41693-2-RV.PDF 2017-02-22
Initiated	2017-02-22
Last modified	2017-05-14

Subscribing on:

ARTICLE IN PRESS
List of the accepted articles for future issues.

FUTURE ISSUES
Vol. 20, no. 6 (December 2020).

- Focus & Scope
- Author Guidelines
- Author Fees
- Online Submission
- Publication Ethics
- Plagiarism Policy
- Editorial Board
- Peer Reviewers

Editor/Author Correspondence

Editor
2017-06-28 06:32 AM

Subject: [IJC] Editor Decision

DELETE

Mrs sriatun sriatun:

We have reached a decision regarding your submission to Indonesian Journal of Chemistry, "The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash".

Our decision is: Revisions Required

The comments of reviewers can be read below or/and in attachment

The Revised article has to be submitted within three weeks after receiving this email.

Best regards,

Nuryono Nuryono
Laboratory of Inorganic Chemistry,
Department of Chemistry,
Universitas Gadjah Mada, Yogyakarta
Phone +628156800908
Fax +62274545188
nuryono_mipa@ugm.ac.id
Editor in Chief
Indonesian Journal of Chemistry
Indexed in SCOPUS since 2012

Reviewer A:

Additional Comment::

The present article reports the Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash. The results are great and worth to be published in this journal. However, there are some issues need to be addressed:

1. The novelty of this research has not been described in detail, what advantage this study from previous similar studies, is there any important breakthroughs resulting from this research?
2. Several additional data are necessarily required to support author's arguments within the manuscript


Reviewer B:

Additional Comment::

Novelty aspect of this research is not explicitly described in this article. In addition, the data presented in this article is also relatively less significant to support the conclusion. Characterization data needs to be enriched by other characterization data fsuch as DTA and BET data.

Indonesian Journal of Chemistry
<https://jurnal.ugm.ac.id/ijc>
Indexed by SCOPUS since 2012

RECEIVED 28 JUN 2017 06:32 AM; ACCEPTED 28 JUN 2017 06:32 AM; PUBLISHED 28 JUN 2017 06:32 AM



IMPROVEMENT OF ARTICLE

Title: The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash

REVIEWER	COMMENT	DESCRIPTION OF IMPROVEMENT
Reviewer A	The results are great and worth to be published in this journal	
	There are some issues need to be addressed: 1. The novelty of this research has not been described in detail, what advantage this study from previous similar studies, is there any important breakthroughs resulting from this research?	Description of the virtues and advantage of this study has been added at the last paragraph on page 3. Based on previously studies information, there has never been conducted researches using CTAB surfactants as directing agent in synthesis of zeolite from sugarcane bagasse ash. Therefore, in this study, we report the influence of cetyltrimethylammonium bromide (CTAB) as structure directing agent in synthesis zeolite using sugarcane bagasse ash as source of silica. It would examine how the effect of the CTAB as directing agent on shape of crystalline and crystallinity the product. As a comparison, zeolite from sugarcane bagasse ash in the absence of structure directing agent was also synthesized.
	Is there any important breakthroughs resulting from this research?	Yes The important breakthroughs resulting from this research that this research is uplifting a low cost preparation of NaP1 zeolite from low cost materials. The results of this study are expected to increase the economic value of waste bagasse into material into high-value material, because this products had many potential applications such as an adsorbent, catalyst and fertilizers. In addition this study also to contributing to the field of material synthesis with renewable resources. Finally by using bagasse as raw material to make zeolite then the accumulation of waste in the environment can be reduced.
	2. Several additional data are necessarily required to support author's arguments within the manuscript (e.g. XRD and BET).	XRD datas has been presented in Figure 3, but not all synthesis products are analyzed by XRD. Only products synthesized without CTAB and with CTAB concentrations of 0.01 M. This concentration is a highest concentration, hat is 10 times to CMC so it is expected to provide information quite different from before. The explanation about the data is given at pages 9-10.

IMPROVEMENT LIST IJC SRIATUN.doc [Compatibility Mode] - Word		
References Mailings Review View Help Nitro Pro 10 Tell me what you want to do		
Font Paragraph Styles		
1 2 3 4 5 6 7 8		
		addition this study also to contributing to the field of material synthesis with renewable resources. Finally by using bagasse as raw material to make zeolite then the accumulation of waste in the environment can be reduced.
	2. Several additional data are necessarily required to support author's arguments within the manuscript (e.g. XRD and BET).	XRD datas has been presented in Figure 3, but not all synthesis products are analyzed by XRD. Only products synthesized without CTAB and with CTAB concentrations of 0.01 M. This concentration is a highest concentration, hat is 10 times to CMC so it is expected to provide information quite different from before. The explanation about the data is given at pages 9-10.
		For BET data was added in page 11. The description is given at pages 10-11.

Reviewer B	Novelty aspect of this research is not explicitly described in this article. In addition, the data presented in this article is also, relatively less significant to support the conclusion.	The novelty aspect was described more explicit. In this article has been added several data such as SEM for other samples, TGA and BET to support in discussion and conclusion.
	Characterization data needs to be enriched by other characterization data such as DTA and BET data.	Several analyzes (TGA and BET) to characterize the product have been added to support the discussion and conclusions.

ROUND 2

Editor
2017-07-25 03:43 PM

Subject: [IJC] Editor Decision

[DELETE](#)

Dear Mrs sriatun sriatun:

We have reached a decision regarding your submission to Indonesian Journal of Chemistry, "The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash".

Our decision is: Revisions Required
The comments of reviewers can be read below or/and in attachment

The Revised article has to be submitted within one week after receiving this email.

Best regards,

Nuryono Nuryono
Laboratory of Inorganic Chemistry,
Department of Chemistry,
Universitas Gadjah Mada, Yogyakarta
Phone +628156800908
Fax +62274545188
nuryono_mipa@ugm.ac.id
Editor in Chief
Indonesian Journal of Chemistry
Indexed in SCOPUS since 2012

Indonesian Journal of Chemistry
<https://jurnal.ugm.ac.id/ijc>
Indexed by SCOPUS since 2012

[SUMMARY](#) [REVIEW](#) [EDITING](#)

Submission

Authors	Sriatun Sriatun, Taslimah Taslimah, Linda Suyati 
Title	Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent
Section	Articles
Editor	Nuryono Nuryono 

Peer Review

Round 1

Review Version	22197-41693-2-RV.PDF 2017-02-22
Initiated	2017-02-22
Last modified	2017-05-14
Uploaded file	None

Editor Decision

Decision	Accept Submission 2017-08-05
Notify Editor	 Editor/Author Email Record  2017-08-05
Editor Version	22197-41987-1-ED.PDF 2017-02-22 22197-41987-2-ED.DOCX 2017-07-25
Author Version	22197-56851-1-ED.DOCX 2017-07-25 DELETE 22197-56851-2-ED.DOCX 2017-08-05 DELETE
Upload Author Version	Pilih File Tidak ada file yang dipilih Upload

for future issues.

FUTURE ISSUES

Vol. 20, no. 6 (December 2020).

Focus & Scope

Author Guidelines

Author Fees

Online Submission

Publication Ethics

Plagiarism Policy

Editorial Board

Peer Reviewers

Order Journal

Visitor Statistics

USER

You are logged in as...
sriatun

[My Journals](#)

[My Profile](#)

[Log Out](#)

ACCEPTED

Editor
2017-08-05 11:11 AM

Subject: [IJC] Editor Decision

DELETE

Mrs sriatun sriatun:

We have reached a decision regarding your submission to Indonesian Journal of Chemistry, "The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash".

Our decision is to: Accept Submission

Thank you for your contribution to Indonesian Journal of Chemistry.

Best regards,

Nuryono Nuryono
Laboratory of Inorganic Chemistry,
Department of Chemistry,
Universitas Gadjah Mada, Yogyakarta
Phone +628156800908
Fax +62274545188
nuryono_mipa@ugm.ac.id


Editor in Chief
Indonesian Journal of Chemistry
Indexed in SCOPUS since 2012

Indonesian Journal of Chemistry
<https://jurnalugm.ac.id/ijc>
Indexed by SCOPUS since 2012

PROOF ARTIKEL DAN PERUBAHAN JUDUL

File untuk VIEW PROOF pada OJS IJC kami simpan sebagai file complete proof 2219767454-1-PB.doc tertanggal 1 November 2017 judul masih The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash.

☐	Name	Date modified	Type	Size
	1047-surat-Biaya Administrasi Penerbi...	11/13/2017 12:36 PM	PDF Document	317 KB
	22197-67454-1-PB (3).doc	10/31/2017 11:56 AM	DOC File	472 KB
	22197-67454-1-PB-Sriatun - sdh cek (...)	2/15/2018 9:12 AM	PDF Document	3,710 KB
	24467-62754-1-PB.doc	11/5/2017 10:57 PM	DOC File	171 KB
	complete proof 22197-67454-1-PB.doc	11/1/2017 10:48 PM	DOC File	472 KB
	complete proof Nov 22197-67454-1-P...	11/5/2017 10:40 PM	DOC File	474 KB
	Doc1.docx	11/2/2017 2:48 PM	Microsoft Word D...	534 KB
	IJC_17_3_2017.pdf	11/23/2017 7:32 PM	PDF Document	430 KB
	Perbaikan-22197-67454-1-PB-Sriatun ...	2/15/2018 9:42 AM	DOC File	479 KB




Dari tambahan data dan perbaikan pembahasan dan kesimpulan yang disarankan oleh reviewer (review Round 1) maka judul "The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash" dirasa kurang tepat, sehingga judul diganti menjadi Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent.

SETELAH PERUBAHAN JUDUL

Perubahan judul kami simpan pada file complete proff Nov 22197-67454-1-P.doc tertanggal 5 November 2020. Adapun tampilannya seperti dibagian bawah ini.

☐ Name	Date modified	Type	Size
1047-surat-Biaya Administrasi Penerbi...	11/13/2017 12:36 PM	PDF Document	317 KB
22197-67454-1-PB (3).doc	10/31/2017 11:56 AM	DOC File	472 KB
22197-67454-1-PB-Sriatun - sdh cek (...)	2/15/2018 9:12 AM	PDF Document	3,710 KB
24467-62754-1-PB.doc	11/5/2017 10:57 PM	DOC File	171 KB
complete proof 22197-67454-1-PB.doc	11/1/2017 10:48 PM	DOC File	472 KB
complete proof Nov 22197-67454-1-P...	11/5/2017 10:40 PM	DOC File	474 KB
Doc1.docx	11/2/2017 2:48 PM	Microsoft Word D...	534 KB
IUC_17_3_2017.pdf	11/23/2017 7:32 PM	PDF Document	430 KB
Perbaikan-22197-67454-1-PB-Sriatun ...	2/15/2018 9:42 AM	DOC File	479 KB



Perubahan judul juga sudah terdapat pada file terakhir yaitu file perbaikan -22197-67454-1-PB-Sriatun tertanggal 15 Februari 2018 (sebelum terbit).

☐ Name	Date modified	Type	Size
1047-surat-Biaya Administrasi Penerbita...	11/13/2017 12:36 PM	PDF Document	317 KB
22197-67454-1-PB (3).doc	10/31/2017 11:56 AM	DOC File	472 KB
22197-67454-1-PB-Sriatun - sdh cek (ST...	2/15/2018 9:12 AM	PDF Document	3,710 KB
24467-62754-1-PB.doc	11/5/2017 10:57 PM	DOC File	171 KB
complete proof 22197-67454-1-PB.doc	11/1/2017 10:48 PM	DOC File	472 KB
complete proof Nov 22197-67454-1-PB...	11/5/2017 10:40 PM	DOC File	474 KB
Doc1.docx	11/2/2017 2:48 PM	Microsoft Word D...	534 KB
IJC_17_3_2017.pdf	11/23/2017 7:32 PM	PDF Document	430 KB
☒ Perbaikan-22197-67454-1-PB-Sriatun - s...	2/15/2018 9:42 AM	DOC File	479 KB

Layout References Mailings Review View Help Nitro Pro 10 Tell me what you want to do

Font Paragraph Styles

12 A⁺ A⁻ Aa A² A³ A⁴ A⁵ A⁶ A⁷ A⁸ A⁹ A¹⁰ A¹¹ A¹² A¹³ A¹⁴ A¹⁵ A¹⁶ A¹⁷ A¹⁸ A¹⁹ A²⁰ A²¹ A²² A²³ A²⁴ A²⁵ A²⁶ A²⁷ A²⁸ A²⁹ A³⁰ A³¹ A³² A³³ A³⁴ A³⁵ A³⁶ A³⁷ A³⁸ A³⁹ A⁴⁰ A⁴¹ A⁴² A⁴³ A⁴⁴ A⁴⁵ A⁴⁶ A⁴⁷ A⁴⁸ A⁴⁹ A⁵⁰ A⁵¹ A⁵² A⁵³ A⁵⁴ A⁵⁵ A⁵⁶ A⁵⁷ A⁵⁸ A⁵⁹ A⁶⁰ A⁶¹ A⁶² A⁶³ A⁶⁴ A⁶⁵ A⁶⁶ A⁶⁷ A⁶⁸ A⁶⁹ A⁷⁰ A⁷¹ A⁷² A⁷³ A⁷⁴ A⁷⁵ A⁷⁶ A⁷⁷ A⁷⁸ A⁷⁹ A⁸⁰ A⁸¹ A⁸² A⁸³ A⁸⁴ A⁸⁵ A⁸⁶ A⁸⁷ A⁸⁸ A⁸⁹ A⁹⁰ A⁹¹ A⁹² A⁹³ A⁹⁴ A⁹⁵ A⁹⁶ A⁹⁷ A⁹⁸ A⁹⁹ A¹⁰⁰ A¹⁰¹ A¹⁰² A¹⁰³ A¹⁰⁴ A¹⁰⁵ A¹⁰⁶ A¹⁰⁷ A¹⁰⁸ A¹⁰⁹ A¹¹⁰ A¹¹¹ A¹¹² A¹¹³ A¹¹⁴ A¹¹⁵ A¹¹⁶ A¹¹⁷ A¹¹⁸ A¹¹⁹ A¹²⁰ A¹²¹ A¹²² A¹²³ A¹²⁴ A¹²⁵ A¹²⁶ A¹²⁷ A¹²⁸ A¹²⁹ A¹³⁰ A¹³¹ A¹³² A¹³³ A¹³⁴ A¹³⁵ A¹³⁶ A¹³⁷ A¹³⁸ A¹³⁹ A¹⁴⁰ A¹⁴¹ A¹⁴² A¹⁴³ A¹⁴⁴ A¹⁴⁵ A¹⁴⁶ A¹⁴⁷ A¹⁴⁸ A¹⁴⁹ A¹⁵⁰ A¹⁵¹ A¹⁵² A¹⁵³ A¹⁵⁴ A¹⁵⁵ A¹⁵⁶ A¹⁵⁷ A¹⁵⁸ A¹⁵⁹ A¹⁶⁰ A¹⁶¹ A¹⁶² A¹⁶³ A¹⁶⁴ A¹⁶⁵ A¹⁶⁶ A¹⁶⁷ A¹⁶⁸ A¹⁶⁹ A¹⁷⁰ A¹⁷¹ A¹⁷² A¹⁷³ A¹⁷⁴ A¹⁷⁵ A¹⁷⁶ A¹⁷⁷ A¹⁷⁸ A¹⁷⁹ A¹⁸⁰ A¹⁸¹ A¹⁸² A¹⁸³ A¹⁸⁴ A¹⁸⁵ A¹⁸⁶ A¹⁸⁷ A¹⁸⁸ A¹⁸⁹ A¹⁹⁰ A¹⁹¹ A¹⁹² A¹⁹³ A¹⁹⁴ A¹⁹⁵ A¹⁹⁶ A¹⁹⁷ A¹⁹⁸ A¹⁹⁹ A²⁰⁰ A²⁰¹ A²⁰² A²⁰³ A²⁰⁴ A²⁰⁵ A²⁰⁶ A²⁰⁷ A²⁰⁸ A²⁰⁹ A²¹⁰ A²¹¹ A²¹² A²¹³ A²¹⁴ A²¹⁵ A²¹⁶ A²¹⁷ A²¹⁸ A²¹⁹ A²²⁰ A²²¹ A²²² A²²³ A²²⁴ A²²⁵ A²²⁶ A²²⁷ A²²⁸ A²²⁹ A²³⁰ A²³¹ A²³² A²³³ A²³⁴ A²³⁵ A²³⁶ A²³⁷ A²³⁸ A²³⁹ A²⁴⁰ A²⁴¹ A²⁴² A²⁴³ A²⁴⁴ A²⁴⁵ A²⁴⁶ A²⁴⁷ A²⁴⁸ A²⁴⁹ A²⁵⁰ A²⁵¹ A²⁵² A²⁵³ A²⁵⁴ A²⁵⁵ A²⁵⁶ A²⁵⁷ A²⁵⁸ A²⁵⁹ A²⁶⁰ A²⁶¹ A²⁶² A²⁶³ A²⁶⁴ A²⁶⁵ A²⁶⁶ A²⁶⁷ A²⁶⁸ A²⁶⁹ A²⁷⁰ A²⁷¹ A²⁷² A²⁷³ A²⁷⁴ A²⁷⁵ A²⁷⁶ A²⁷⁷ A²⁷⁸ A²⁷⁹ A²⁸⁰ A²⁸¹ A²⁸² A²⁸³ A²⁸⁴ A²⁸⁵ A²⁸⁶ A²⁸⁷ A²⁸⁸ A²⁸⁹ A²⁹⁰ A²⁹¹ A²⁹² A²⁹³ A²⁹⁴ A²⁹⁵ A²⁹⁶ A²⁹⁷ A²⁹⁸ A²⁹⁹ A³⁰⁰ A³⁰¹ A³⁰² A³⁰³ A³⁰⁴ A³⁰⁵ A³⁰⁶ A³⁰⁷ A³⁰⁸ A³⁰⁹ A³¹⁰ A³¹¹ A³¹² A³¹³ A³¹⁴ A³¹⁵ A³¹⁶ A³¹⁷ A³¹⁸ A³¹⁹ A³²⁰ A³²¹ A³²² A³²³ A³²⁴ A³²⁵ A³²⁶ A³²⁷ A³²⁸ A³²⁹ A³³⁰ A³³¹ A³³² A³³³ A³³⁴ A³³⁵ A³³⁶ A³³⁷ A³³⁸ A³³⁹ A³⁴⁰ A³⁴¹ A³⁴² A³⁴³ A³⁴⁴ A³⁴⁵ A³⁴⁶ A³⁴⁷ A³⁴⁸ A³⁴⁹ A³⁵⁰ A³⁵¹ A³⁵² A³⁵³ A³⁵⁴ A³⁵⁵ A³⁵⁶ A³⁵⁷ A³⁵⁸ A³⁵⁹ A³⁶⁰ A³⁶¹ A³⁶² A³⁶³ A³⁶⁴ A³⁶⁵ A³⁶⁶ A³⁶⁷ A³⁶⁸ A³⁶⁹ A³⁷⁰ A³⁷¹ A³⁷² A³⁷³ A³⁷⁴ A³⁷⁵ A³⁷⁶ A³⁷⁷ A³⁷⁸ A³⁷⁹ A³⁸⁰ A³⁸¹ A³⁸² A³⁸³ A³⁸⁴ A³⁸⁵ A³⁸⁶ A³⁸⁷ A³⁸⁸ A³⁸⁹ A³⁹⁰ A³⁹¹ A³⁹² A³⁹³ A³⁹⁴ A³⁹⁵ A³⁹⁶ A³⁹⁷ A³⁹⁸ A³⁹⁹ A⁴⁰⁰ A⁴⁰¹ A⁴⁰² A⁴⁰³ A⁴⁰⁴ A⁴⁰⁵ A⁴⁰⁶ A⁴⁰⁷ A⁴⁰⁸ A⁴⁰⁹ A⁴¹⁰ A⁴¹¹ A⁴¹² A⁴¹³ A⁴¹⁴ A⁴¹⁵ A⁴¹⁶ A⁴¹⁷ A⁴¹⁸ A⁴¹⁹ A⁴²⁰ A⁴²¹ A⁴²² A⁴²³ A⁴²⁴ A⁴²⁵ A⁴²⁶ A⁴²⁷ A⁴²⁸ A⁴²⁹ A⁴³⁰ A⁴³¹ A⁴³² A⁴³³ A⁴³⁴ A⁴³⁵ A⁴³⁶ A⁴³⁷ A⁴³⁸ A⁴³⁹ A⁴⁴⁰ A⁴⁴¹ A⁴⁴² A⁴⁴³ A⁴⁴⁴ A⁴⁴⁵ A⁴⁴⁶ A⁴⁴⁷ A⁴⁴⁸ A⁴⁴⁹ A⁴⁵⁰ A⁴⁵¹ A⁴⁵² A⁴⁵³ A⁴⁵⁴ A⁴⁵⁵ A⁴⁵⁶ A⁴⁵⁷ A⁴⁵⁸ A⁴⁵⁹ A⁴⁶⁰ A⁴⁶¹ A⁴⁶² A⁴⁶³ A⁴⁶⁴ A⁴⁶⁵ A⁴⁶⁶ A⁴⁶⁷ A⁴⁶⁸ A⁴⁶⁹ A⁴⁷⁰ A⁴⁷¹ A⁴⁷² A⁴⁷³ A⁴⁷⁴ A⁴⁷⁵ A⁴⁷⁶ A⁴⁷⁷ A⁴⁷⁸ A⁴⁷⁹ A⁴⁸⁰ A⁴⁸¹ A⁴⁸² A⁴⁸³ A⁴⁸⁴ A⁴⁸⁵ A⁴⁸⁶ A⁴⁸⁷ A⁴⁸⁸ A⁴⁸⁹ A⁴⁹⁰ A⁴⁹¹ A⁴⁹² A⁴⁹³ A⁴⁹⁴ A⁴⁹⁵ A⁴⁹⁶ A⁴⁹⁷ A⁴⁹⁸ A⁴⁹⁹ A⁵⁰⁰ A⁵⁰¹ A⁵⁰² A⁵⁰³ A⁵⁰⁴ A⁵⁰⁵ A⁵⁰⁶ A⁵⁰⁷ A⁵⁰⁸ A⁵⁰⁹ A⁵¹⁰ A⁵¹¹ A⁵¹² A⁵¹³ A⁵¹⁴ A⁵¹⁵ A⁵¹⁶ A⁵¹⁷ A⁵¹⁸ A⁵¹⁹ A⁵²⁰ A⁵²¹ A⁵²² A⁵²³ A⁵²⁴ A⁵²⁵ A⁵²⁶ A⁵²⁷ A⁵²⁸ A⁵²⁹ A⁵³⁰ A⁵³¹ A⁵³² A⁵³³ A⁵³⁴ A⁵³⁵ A⁵³⁶ A⁵³⁷ A⁵³⁸ A⁵³⁹ A⁵⁴⁰ A⁵⁴¹ A⁵⁴² A⁵⁴³ A⁵⁴⁴ A⁵⁴⁵ A⁵⁴⁶ A⁵⁴⁷ A⁵⁴⁸ A⁵⁴⁹ A⁵⁵⁰ A⁵⁵¹ A⁵⁵² A⁵⁵³ A⁵⁵⁴ A⁵⁵⁵ A⁵⁵⁶ A⁵⁵⁷ A⁵⁵⁸ A⁵⁵⁹ A⁵⁶⁰ A⁵⁶¹ A⁵⁶² A⁵⁶³ A⁵⁶⁴ A⁵⁶⁵ A⁵⁶⁶ A⁵⁶⁷ A⁵⁶⁸ A⁵⁶⁹ A⁵⁷⁰ A⁵⁷¹ A⁵⁷² A⁵⁷³ A⁵⁷⁴ A⁵⁷⁵ A⁵⁷⁶ A⁵⁷⁷ A⁵⁷⁸ A⁵⁷⁹ A⁵⁸⁰ A⁵⁸¹ A⁵⁸² A⁵⁸³ A⁵⁸⁴ A⁵⁸⁵ A⁵⁸⁶ A⁵⁸⁷ A⁵⁸⁸ A⁵⁸⁹ A⁵⁹⁰ A⁵⁹¹ A⁵⁹² A⁵⁹³ A⁵⁹⁴ A⁵⁹⁵ A⁵⁹⁶ A⁵⁹⁷ A⁵⁹⁸ A⁵⁹⁹ A⁶⁰⁰ A⁶⁰¹ A⁶⁰² A⁶⁰³ A⁶⁰⁴ A⁶⁰⁵ A⁶⁰⁶ A⁶⁰⁷ A⁶⁰⁸ A⁶⁰⁹ A⁶¹⁰ A⁶¹¹ A⁶¹² A⁶¹³ A⁶¹⁴ A⁶¹⁵ A⁶¹⁶ A⁶¹⁷ A⁶¹⁸ A⁶¹⁹ A⁶²⁰ A⁶²¹ A⁶²² A⁶²³ A⁶²⁴ A⁶²⁵ A⁶²⁶ A⁶²⁷ A⁶²⁸ A⁶²⁹ A⁶³⁰ A⁶³¹ A⁶³² A⁶³³ A⁶³⁴ A⁶³⁵ A⁶³⁶ A⁶³⁷ A⁶³⁸ A⁶³⁹ A⁶⁴⁰ A⁶⁴¹ A⁶⁴² A⁶⁴³ A⁶⁴⁴ A⁶⁴⁵ A⁶⁴⁶ A⁶⁴⁷ A⁶⁴⁸ A⁶⁴⁹ A⁶⁵⁰ A⁶⁵¹ A⁶⁵² A⁶⁵³ A⁶⁵⁴ A⁶⁵⁵ A⁶⁵⁶ A⁶⁵⁷ A⁶⁵⁸ A⁶⁵⁹ A⁶⁶⁰ A⁶⁶¹ A⁶⁶² A⁶⁶³ A⁶⁶⁴ A⁶⁶⁵ A⁶⁶⁶ A⁶⁶⁷ A⁶⁶⁸ A⁶⁶⁹ A⁶⁷⁰ A⁶⁷¹ A⁶⁷² A⁶⁷³ A⁶⁷⁴ A⁶⁷⁵ A⁶⁷⁶ A⁶⁷⁷ A⁶⁷⁸ A⁶⁷⁹ A⁶⁸⁰ A⁶⁸¹ A⁶⁸² A⁶⁸³ A⁶⁸⁴ A⁶⁸⁵ A⁶⁸⁶ A⁶⁸⁷ A⁶⁸⁸ A⁶⁸⁹ A⁶⁹⁰ A⁶⁹¹ A⁶⁹² A⁶⁹³ A⁶⁹⁴ A⁶⁹⁵ A⁶⁹⁶ A⁶⁹⁷ A⁶⁹⁸ A⁶⁹⁹ A⁷⁰⁰ A⁷⁰¹ A⁷⁰² A⁷⁰³ A⁷⁰⁴ A⁷⁰⁵ A⁷⁰⁶ A⁷⁰⁷ A⁷⁰⁸ A⁷⁰⁹ A⁷¹⁰ A⁷¹¹ A⁷¹² A⁷¹³ A⁷¹⁴ A⁷¹⁵ A⁷¹⁶ A⁷¹⁷ A⁷¹⁸ A⁷¹⁹ A⁷²⁰ A⁷²¹ A⁷²² A⁷²³ A⁷²⁴ A⁷²⁵ A⁷²⁶ A⁷²⁷ A⁷²⁸ A⁷²⁹ A⁷³⁰ A⁷³¹ A⁷³² A⁷³³ A⁷³⁴ A⁷³⁵ A⁷³⁶ A⁷³⁷ A⁷³⁸ A⁷³⁹ A⁷⁴⁰ A⁷⁴¹ A⁷⁴² A⁷⁴³ A⁷⁴⁴ A⁷⁴⁵ A⁷⁴⁶ A⁷⁴⁷ A⁷⁴⁸ A⁷⁴⁹ A⁷⁵⁰ A⁷⁵¹ A⁷⁵² A⁷⁵³ A⁷⁵⁴ A⁷⁵⁵ A⁷⁵⁶ A⁷⁵⁷ A⁷⁵⁸ A⁷⁵⁹ A⁷⁶⁰ A⁷⁶¹ A⁷⁶² A⁷⁶³ A⁷⁶⁴ A⁷⁶⁵ A⁷⁶⁶ A⁷⁶⁷ A⁷⁶⁸ A⁷⁶⁹ A⁷⁷⁰ A⁷⁷¹ A⁷⁷² A⁷⁷³ A⁷⁷⁴ A⁷⁷⁵ A⁷⁷⁶ A⁷⁷⁷ A⁷⁷⁸ A⁷⁷⁹ A⁷⁸⁰ A⁷⁸¹ A⁷⁸² A⁷⁸³ A⁷⁸⁴ A⁷⁸⁵ A⁷⁸⁶ A⁷⁸⁷ A⁷⁸⁸ A⁷⁸⁹ A⁷⁹⁰ A⁷⁹¹ A⁷⁹² A⁷⁹³ A⁷⁹⁴ A⁷⁹⁵ A⁷⁹⁶ A⁷⁹⁷ A⁷⁹⁸ A⁷⁹⁹ A⁸⁰⁰ A⁸⁰¹ A⁸⁰² A⁸⁰³ A⁸⁰⁴ A⁸⁰⁵ A⁸⁰⁶ A⁸⁰⁷ A⁸⁰⁸ A⁸⁰⁹ A⁸¹⁰ A⁸¹¹ A⁸¹² A⁸¹³ A⁸¹⁴ A⁸¹⁵ A⁸¹⁶ A⁸¹⁷ A⁸¹⁸ A⁸¹⁹ A⁸²⁰ A⁸²¹ A⁸²² A⁸²³ A⁸²⁴ A⁸²⁵ A⁸²⁶ A⁸²⁷ A⁸²⁸ A⁸²⁹ A⁸³⁰ A⁸³¹ A⁸³² A⁸³³ A⁸³⁴ A⁸³⁵ A⁸³⁶ A⁸³⁷ A⁸³⁸ A⁸³⁹ A⁸⁴⁰ A⁸⁴¹ A⁸⁴² A⁸⁴³ A⁸⁴⁴ A⁸⁴⁵ A⁸⁴⁶ A⁸⁴⁷ A⁸⁴⁸ A⁸⁴⁹ A⁸⁵⁰ A⁸⁵¹ A⁸⁵² A⁸⁵³ A⁸⁵⁴ A⁸⁵⁵ A⁸⁵⁶ A⁸⁵⁷ A⁸⁵⁸ A⁸⁵⁹ A⁸⁶⁰ A⁸⁶¹ A⁸⁶² A⁸⁶³ A⁸⁶⁴ A⁸⁶⁵ A⁸⁶⁶ A⁸⁶⁷ A⁸⁶⁸ A⁸⁶⁹ A⁸⁷⁰ A⁸⁷¹ A⁸⁷² A⁸⁷³ A⁸⁷⁴ A⁸⁷⁵ A⁸⁷⁶ A⁸⁷⁷ A⁸⁷⁸ A⁸⁷⁹ A⁸⁸⁰ A⁸⁸¹ A⁸⁸² A⁸⁸³ A⁸⁸⁴ A⁸⁸⁵ A⁸⁸⁶ A⁸⁸⁷ A⁸⁸⁸ A⁸⁸⁹ A⁸⁹⁰ A⁸⁹¹ A⁸⁹² A⁸⁹³ A⁸⁹⁴ A⁸⁹⁵ A⁸⁹⁶ A⁸⁹⁷ A⁸⁹⁸ A⁸⁹⁹ A⁹⁰⁰ A⁹⁰¹ A⁹⁰² A⁹⁰³ A⁹⁰⁴ A⁹⁰⁵ A⁹⁰⁶ A⁹⁰⁷ A⁹⁰⁸ A⁹⁰⁹ A⁹¹⁰ A⁹¹¹ A⁹¹² A⁹¹³ A⁹¹⁴ A⁹¹⁵ A⁹¹⁶ A⁹¹⁷ A⁹¹⁸ A⁹¹⁹ A⁹²⁰ A⁹²¹ A⁹²² A⁹²³ A⁹²⁴ A⁹²⁵ A⁹²⁶ A⁹²⁷ A⁹²⁸ A⁹²⁹ A⁹³⁰ A⁹³¹ A⁹³² A⁹³³ A⁹³⁴ A⁹³⁵ A⁹³⁶ A⁹³⁷ A⁹³⁸ A⁹³⁹ A⁹⁴⁰ A⁹⁴¹ A⁹⁴² A⁹⁴³ A⁹⁴⁴ A⁹⁴⁵ A⁹⁴⁶ A⁹⁴⁷ A⁹⁴⁸ A⁹⁴⁹ A⁹⁵⁰ A⁹⁵¹ A⁹⁵² A⁹⁵³ A⁹⁵⁴ A⁹⁵⁵ A⁹⁵⁶ A⁹⁵⁷ A⁹⁵⁸ A⁹⁵⁹ A⁹⁶⁰ A⁹⁶¹ A⁹⁶² A⁹⁶³ A⁹⁶⁴ A⁹⁶⁵ A⁹⁶⁶ A⁹⁶⁷ A⁹⁶⁸ A⁹⁶⁹ A⁹⁷⁰ A⁹⁷¹ A⁹⁷² A⁹⁷³ A⁹⁷⁴ A⁹⁷⁵ A⁹⁷⁶ A⁹⁷⁷ A⁹⁷⁸ A⁹⁷⁹ A⁹⁸⁰ A⁹⁸¹ A⁹⁸² A⁹⁸³ A⁹⁸⁴ A⁹⁸⁵ A⁹⁸⁶ A⁹⁸⁷ A⁹⁸⁸ A⁹⁸⁹ A⁹⁹⁰ A⁹⁹¹ A⁹⁹² A⁹⁹³ A⁹⁹⁴ A⁹⁹⁵ A⁹⁹⁶ A⁹⁹⁷ A⁹⁹⁸ A⁹⁹⁹ A¹⁰⁰⁰ A¹⁰⁰¹ A¹⁰⁰² A¹⁰⁰³ A¹⁰⁰⁴ A¹⁰⁰⁵ A¹⁰⁰⁶ A¹⁰⁰⁷ A¹⁰⁰⁸ A¹⁰⁰⁹ A¹⁰¹⁰ A¹⁰¹¹ A¹⁰¹² A¹⁰¹³ A¹⁰¹⁴ A¹⁰¹⁵ A¹⁰¹⁶ A¹⁰¹⁷ A¹⁰¹⁸ A¹⁰¹⁹ A¹⁰²⁰ A¹⁰²¹ A¹⁰²² A¹⁰²³ A¹⁰²⁴ A¹⁰²⁵ A¹⁰²⁶ A¹⁰²⁷ A¹⁰²⁸ A¹⁰²⁹ A¹⁰³⁰ A¹⁰³¹ A¹⁰³² A¹⁰³³ A¹⁰³⁴ A¹⁰³⁵ A¹⁰³⁶ A¹⁰³⁷ A¹⁰³⁸ A¹⁰³⁹ A¹⁰⁴⁰ A¹⁰⁴¹ A¹⁰⁴² A¹⁰⁴³ A¹⁰⁴⁴ A¹⁰⁴⁵ A¹⁰⁴⁶ A¹⁰⁴⁷ A¹⁰⁴⁸ A¹⁰⁴⁹ A¹⁰⁵⁰ A¹⁰⁵¹ A¹⁰⁵² A¹⁰⁵³ A¹⁰⁵⁴ A¹⁰⁵⁵ A¹⁰⁵⁶ A¹⁰⁵⁷ A¹⁰⁵⁸ A¹⁰⁵⁹ A¹⁰⁶⁰ A¹⁰⁶¹ A¹⁰⁶² A¹⁰⁶³ A¹⁰⁶⁴ A¹⁰⁶⁵ A¹⁰⁶⁶ A¹⁰⁶⁷ A¹⁰⁶⁸ A¹⁰⁶⁹ A¹⁰⁷⁰ A¹⁰⁷¹ A¹⁰⁷² A¹⁰⁷³ A¹⁰⁷⁴ A¹⁰⁷⁵ A¹⁰⁷⁶ A¹⁰⁷⁷ A¹⁰⁷⁸ A¹⁰⁷⁹ A¹⁰⁸⁰ A¹⁰⁸¹ A¹⁰⁸² A¹⁰⁸³ A¹⁰⁸⁴ A¹⁰⁸⁵ A¹⁰⁸⁶ A¹⁰⁸⁷ A¹⁰⁸⁸ A¹⁰⁸⁹ A¹⁰⁹⁰ A¹⁰⁹¹ A¹⁰⁹² A¹⁰⁹³ A¹⁰⁹⁴ A¹⁰⁹⁵ A¹⁰⁹⁶ A¹⁰⁹⁷ A¹⁰⁹⁸ A¹⁰⁹⁹ A¹¹⁰⁰ A¹¹⁰¹ A¹¹⁰² A¹¹⁰³ A¹¹⁰⁴ A¹¹⁰⁵ A¹¹⁰⁶ A¹¹⁰⁷ A¹¹⁰⁸ A¹¹⁰⁹ A¹¹¹⁰ A¹¹¹¹ A¹¹¹² A¹¹¹³ A¹¹¹⁴ A¹¹¹⁵ A¹¹¹⁶ A¹¹¹⁷ A¹¹¹⁸ A¹¹¹⁹ A¹¹²⁰ A¹¹²¹ A¹¹²² A¹¹²³ A¹¹²⁴ A¹¹²⁵ A¹¹²⁶ A¹¹²⁷ A¹¹²⁸ A¹¹²⁹ A¹¹³⁰ A¹¹³¹ A¹¹³² A¹¹³³ A¹¹³⁴ A¹¹³⁵ A¹¹³⁶ A¹¹³⁷ A¹¹³⁸ A¹¹³⁹ A¹¹⁴⁰ A¹¹⁴¹ A¹¹⁴² A¹¹⁴³ A¹¹⁴⁴ A¹¹⁴⁵ A¹¹⁴⁶ A¹¹⁴⁷ A¹¹⁴⁸ A¹¹⁴⁹ A¹¹⁵⁰ A¹¹⁵¹ A¹¹⁵² A¹¹⁵³ A¹¹⁵⁴ A¹¹⁵⁵ A¹¹⁵⁶ A¹¹⁵⁷ A¹¹⁵⁸ A¹¹⁵⁹ A¹¹⁶⁰ A¹¹⁶¹ A¹¹⁶² A¹¹⁶³ A¹¹⁶⁴ A¹¹⁶⁵ A¹¹⁶⁶ A¹¹⁶⁷ A¹¹⁶⁸ A¹¹⁶⁹ A¹¹⁷⁰ A¹¹⁷¹ A¹¹⁷² A¹¹⁷³ A¹¹⁷⁴ A¹¹⁷⁵ A¹¹⁷⁶ A¹¹⁷⁷ A¹¹⁷⁸ A¹¹⁷⁹ A¹¹⁸⁰ A¹¹⁸¹ A¹¹⁸² A¹¹⁸³ A¹¹⁸⁴ A¹¹⁸⁵ A¹¹⁸⁶ A¹¹⁸⁷ A¹¹⁸⁸ A¹¹⁸⁹ A¹¹⁹⁰ A¹¹⁹¹ A¹¹⁹² A¹¹⁹³ A¹¹⁹⁴ A¹¹⁹⁵ A¹¹⁹⁶ A¹¹⁹⁷ A¹¹⁹⁸ A¹¹⁹⁹ A¹²⁰⁰ A¹²⁰¹ A¹²⁰² A¹²⁰³ A¹²⁰⁴ A¹²⁰⁵ A¹²⁰⁶ A¹²⁰⁷ A¹²⁰⁸ A<

Berikut adalah permintaan VIEWPROOF sebelum terbit.

Search mail

328 of 338

[IJC] Copyediting Review Request Inbox x

Dwi Siswanta
Dear Mrs sriatun sriatun: Your submission "The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash" for Indon

Sep 15, 2017, 1:56 PM

Sriatun Sriatun
Dear Mr. Dwi Siswanta Indonesian Journal of Chemistry My submission " The Effect of Cetyltrimethylammonium Bromide Surfactant in Synthesis of Zeolite From Suga

Sep 19, 2017, 11:29 AM

Dwi Siswanta <dsiswanta@ugm.ac.id>
to me

Sep 20, 2017, 2:42 PM

Subject: The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash

EDIT DELETE

Please check the copyedit comment from the editor and respond accordingly

1. No space between number and "%"
2. use symbol "x" for scientific notation e.g. 1×10^{-3}
3. Fig. 2a -> Fig 2; Fig 2b -> Fig. 3; Fig 2c -> Fig. 4.
4. Improve the quality of Fig 2a, 2b and 2c; the axis line must be black 1 pt; Arial font 9 pt; without outside border
5. Fig. 3; without outside border line; black axes line (1 pt)
6. Fig. 5; without outside border line; black axes line (1 pt); use "." for decimal; dimension 8 cm (width) x 8 cm (height)
7. Improve Fig. 6. Please consult good journal article on how to show the TGA result.
8. Please follow the IJC style for referencing.

Best regards

Search mail

325 of 338

[IJC] Proofreading Request (Author) Inbox x

Dwi Siswanta <dsiswanta@ugm.ac.id>
to me

Mon, Oct 30, 2017, 12:38 PM

Mrs Sriatun Sriatun:

Your submission "The Effect of Cetyltrimethylammonium Bromide Surfactant in Synthesis of Zeolite from Sugarcane Bagasse Ash" to Indonesian Journal of Chemistry now needs to be proofread by following these steps.

1. Click on the Submission URL below.
2. Log into the journal and view PROOFING INSTRUCTIONS
3. Click on VIEW PROOF in Layout and proof the galley in the one or more formats used.
4. Enter corrections (typographical and format) in Proofreading Corrections.
5. Save and email corrections to Layout Editor and Proofreader.
6. Send the COMPLETE email to the editor.

Submission URL: <https://journal.ugm.ac.id/ijc/author/submissionEditing/22197>
Username: sriatun

Dwi Siswanta
Laboratory of Analytical Chemistry,
Department of Chemistry,
Universitas Gadjah Mada
Phone +628157951198
Fax +62545188
dsiswanta@ugm.ac.id

TRACKING PADA OJS IJC

Copyediting

[COPYEDIT INSTRUCTIONS](#)

Copyeditor None

REVIEW METADATA

	REQUEST	UNDERWAY	COMPLETE
1. Initial Copyedit	—	—	2017-02-22
File: 22197-58428-2-CE.DOCX 2017-08-05			
2. Author Copyedit	2017-09-15	2017-09-18	2017-10-13
File: 22197-59026-3-CE.DOCX 2017-09-19			
Pilih File Tidak ada file yang dipilih Upload			
3. Final Copyedit	2017-10-13	—	2017-02-22
File: 22197-58428-4-CE.DOCX 2017-10-24			

Copyedit Comments 2017-09-15

Layout

Layout Editor Djoko Prihandono

	REQUEST	UNDERWAY	COMPLETE	VIEWS
22197-66497-1-LE.DOCX 2017-10-24	2017-10-24	2017-10-25	2017-10-30	

Galley Format

FILE		
1. Full Text PDF VIEW PROOF	22197-72634-3-PB.PDF 2018-02-28	2159

Supplementary Files

FILE	
None	

Layout Comments No Comments

VIEW PROOF (GALLEY FORMAT)

JUDUL “SYNTHESIS OF ZEOLIT FROM SUGARCANE BAGASSE ASH USING CETYLTRIMETHYLAMMONIUM AS STRUCTURE DIRECTING AGENT” pada file galley format dan terbit pada tanggal 28 Februari 2018 sebagai artikel pada Jurnal IJC Vol 18 No. 1 halaman 159-165.

[Back to Submission Editing](#)

Microsoft Word - 18-1-21-159-165-Sriatun - sdh cek

1 / 7

Indones. J. Chem., 2018, 18 (1), 159 - 165

159

Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent

Sriatun^{*}, Taslimah, and Linda Suyati

Department of Chemistry, Faculty of Science and Mathematics, Diponegoro University,
Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275, Indonesia

Received February 18, 2017; Accepted October 24, 2017

ABSTRACT

The purpose of this research is to synthesize zeolite from sugarcane bagasse ash using cetyltrimethylammonium bromide as structure directing agent. This research used cetyltrimethylammonium bromide surfactant to invent the high porosity, surface area, acidity and thermal stability of synthesized zeolite. The Silica was extracted by alkaline fusion using sodium hydroxide solution. The synthesis was conducted by hydrothermal process at 100 °C for 7 days, ageing process for 24 h and calcination at 500 °C for 5 h. The ratio of Si/Al (v/v) was 1, 15 and 25, meanwhile the concentration of cetyltrimethylammonium bromide was 5x10⁻⁴ M, 1x10⁻³ M and 1x10⁻² M. The result showed all of product have strong absorbance at 950–1050 cm⁻¹ and 620–690 cm⁻¹, 420–460 cm⁻¹, double ring at 520–570 cm⁻¹, pore opening at 300–370 cm⁻¹. Vibration of –OH as silanol group or water was indicated by broad absorbance at 3400–3450 cm⁻¹. The diffractograms XRD showed that the product had high crystallinity. The composition of product on ratio Si/Al 1 with concentration of cetyltrimethylammonium 10⁻² M is sodalite, the ratio Si/Al 15 and 25 are NaP1 and SiO₂ quartz and contain 12.23% and 12.19% of Si, 4.17% and 13.18% of Al, respectively. Observation on SEM revealed that the crystal produced using cetyltrimethylammonium bromide was more homogenous and regular in shape.

Keywords: surfactant; cetyltrimethylammonium bromide; synthesis; zeolite; sugarcane bagasse

ABSTRAK

Tujuan penelitian ini adalah membuat zeolit dari ampas tebu menggunakan setiltrimetilamonium bromida

SURAT KETERANGAN

Yang bertanda tangan di bawah ini:

Nama : Prof. Dr.rer.nat. Nuryono, M.S
Instansi : Departemen Kimia FMIPA Universitas Gajah Mada
Kedudukan : Editor in Chief Jurnal Indonesian Journal of Chemistry (IJC) UGM
Alamat : Jl. Sekip Utara, BLS 21, 55281, Senolowo, Sinduadi, Kec. Mlati,
Kabupaten Sleman, Daerah Istimewa Yogyakarta 55281
E-mail : nuryono_mipa@ugm.ac.id

Menerangkan bahwa artikel yang berjudul **Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent** yang terbit pada Jurnal Indonesian Journal of Chemistry (IJC) Vol. 18 No.1 benar-benar merupakan paper final yang dihasilkan dari review manuskrip yang disubmit dengan judul mula-mula **The Effect of Cetyltrimethylammonium Bromide Surfactant in Synthesis of Zeolite from Sugarcane Bagasse Ash**. Perubahan judul ini merupakan rekomendasi setelah proofreading akhir paper layout, sebelum terbit, untuk meningkatkan relevansi antara judul dan isi penelitian yang dilaporkan.

Demikian surat keterangan ini kami buat dengan sesungguhnya, agar dapat dipergunakan sebagaimana mestinya.

Yogyakarta, 17 Oktober 2020

Yang menerangkan,



Prof. Dr. rer. nat. Nuryono, M.S
Editor in Chief Indonesian Journal of Chemistry



Q ijc

Compose

Inbox

191

Starred

Snoozed

Sent

More

Meet

Start a meeting

Join a meeting

Chat



Sriatun



[IJC] Submission Acknowledgement Inbox x

**Prof. Dr.rer.nat. Nuryono, MS** <nuryono_mipa@ugm.ac.id>

to me

Mrs sriatun sriatun:

Thank you for submitting the manuscript, "The Effect of cetyltrimethylammonium bromide surfactant in synthesis of zeolite from sugarcane bagasse ash" to Indonesian Journal of Chemistry. With the journal management system that we are using, you will be able to track progress through the editorial process by logging in to the journal web site:

Manuscript URL: <https://jurnal.ugm.ac.id/ijc/author/submission/22197>

Username: sriatun

If you have any questions, please contact me. Thank you for considering journal as a venue for your work.

Prof. Dr.rer.nat. Nuryono, MS
Indonesian Journal of Chemistry

Indonesian Journal of Chemistry
<https://jurnal.ugm.ac.id/ijc>



Q ijc

Compose

Inbox

191

Starred

Snoozed

Sent

More

Meet

Start a meeting

Join a meeting

Chat



Sriatun

[IJC] Editor Decision Inbox x**Nuryono Nuryono** <nuryono_mipa@ugm.ac.id>

to me

Mrs sriatun sriatun:

We have reached a decision regarding your submission to Indonesian J of Chemistry, "The Effect of cetyltrimethylammonium bromide surfactant synthesis of zeolite from sugarcane bagasse ash".

Our decision is: Revisions Required

The comments of reviewers can be read below or/and in attachment

The Revised article has to be submitted within three weeks after receiving this email.

Best regards,

Nuryono Nuryono
Laboratory of Inorganic Chemistry,
Department of Chemistry,
Universitas Gadjah Mada, Yogyakarta
Phone +628156800908
Fax +62274545188



Q ijc

Compose

Inbox

191

Starred

Snooked

Sent

More

Meet

Start a meeting

Join a meeting

Chat



Sriatun

[IJC] Copyediting Review Request Inbox x**Nuryono Nuryono** <nuryono_mipa@ugm.ac.id>

to me, ijc

Dear Mrs sriatun sriatun:

Your submission "The Effect of cetyltrimethylammonium bromide surfact synthesis of zeolite from sugarcane bagasse ash" for Indonesian Journe Chemistry has been through the first step of copyediting, and is available for you to review by following these steps.

1. Click on the Submission URL below.
2. Log into the journal and click on the File that appears in Step 1.
3. Open the downloaded submission.
4. Review the text, including copyediting proposals and Author Queries.
5. Make any copyediting changes that would further improve the text.
6. When completed, upload the file in Step 2.
7. Click on METADATA to check indexing information for completeness ; accuracy.
8. Send the COMPLETE email to the editor and copyeditor.

Submission URL: <https://jurnal.ugm.ac.id/ijc/author/submissionEditing/2/>

Username: sriatun



Q ijc

Compose

Inbox

191

Starred

Snoozed

Sent

More

Meet

Start a meeting

Join a meeting

Chat



Sriatun



revision article

**Sriatun Sriatun** <sriatun@live.undip.ac.id>

to MS

To. Mr. Nuryono
Editor in Chief
Indonesian Journal of Chemistry

Thank you Mr. Nuryono, for the second opportunity improv
synthesis of zeolite from sugarcane bagasse ash".

I re-submitted the article that I have revised along accordi
my article can be published in Indonesian Journal of Cherr

Best regards

Sriatun

Laboratory of Inorganic Chemistry,
Dept of Chemistry FSM Undip

--

2 Attachments

[Compose](#)**Inbox**

191

Starred

Snoozed

Sent

More

Meet[Start a meeting](#)[Join a meeting](#)**Chat**

Sriatun



[IJC] Editor Decision Inbox x

**Nuryono Nuryono** <nuryono_mipa@ugm.ac.id>
to me

Mrs sriatun sriatun:

We have reached a decision regarding your submission to Indonesian J of Chemistry, "The Effect of cetyltrimethylammonium bromide surfactant synthesis of zeolite from sugarcane bagasse ash".

Our decision is to: Accept Submission

Thank you for your contribution to Indonesian Journal of Chemistry.

Best regards,

Nuryono Nuryono
Laboratory of Inorganic Chemistry,
Department of Chemistry,
Universitas Gadjah Mada, Yogyakarta
Phone +628156800908
Fax +62274545188
nuryono_mipa@ugm.ac.id

Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent

Sriatun*, Taslimah, and Linda Suyati

Department of Chemistry, Faculty of Science and Mathematics, Diponegoro University,
Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275, Indonesia

Received February 18, 2017; Accepted October 24, 2017

ABSTRACT

The purpose of this research is to synthesize zeolite from sugarcane bagasse ash using cetyltrimethylammonium bromide as structure directing agent. This research used cetyltrimethylammonium bromide surfactant to invent the high porosity, surface area, acidity and thermal stability of synthesized zeolite. The Silica was extracted by alkaline fusion using sodium hydroxide solution. The synthesis was conducted by hydrothermal process at 100 °C for 7 days, ageing process for 24 h and calcination at 500 °C for 5 h. The ratio of Si/Al (v/v) was 1, 15 and 25, meanwhile the concentration of cetyltrimethylammonium bromide was 5×10^{-4} M, 1×10^{-3} M and 1×10^{-2} M. The result showed all of product have strong absorbance at $950\text{--}1050\text{ cm}^{-1}$ and $620\text{--}690\text{ cm}^{-1}$, $420\text{--}460\text{ cm}^{-1}$, double ring at $520\text{--}570\text{ cm}^{-1}$, pore opening at $300\text{--}370\text{ cm}^{-1}$. Vibration of --OH as silanol group or water was indicated by broad absorbance at $3400\text{--}3450\text{ cm}^{-1}$. The diffractograms XRD showed that the product had high crystallinity. The composition of product on ratio Si/Al 1 with concentration of cetyltrimethylammonium 10^{-2} M is sodalite, the ratio Si/Al 15 and 25 are NaP1 and SiO_2 quartz and contain 12.23% and 12.19% of Si, 4.17% and 13.18% of Al, respectively. Observation on SEM revealed that the crystal produced using cetyltrimethylammonium bromide was more homogenous and regular in shape.

Keywords: surfactant; cetyltrimethylammonium bromide; synthesis; zeolite; sugarcane bagasse

ABSTRAK

Tujuan penelitian ini adalah membuat zeolit dari ampas tebu menggunakan setiltrimetilamonium bromida sebagai agen pengarah struktur. Dalam penelitian ini surfaktan setiltrimetilamonium bromida digunakan untuk meningkatkan porositas, luas permukaan, keasaman dan stabilitas termal. Silika diekstrak dari abu ampas tebu melalui penambahan alkali yaitu NaOH 6 M. Sintesis dilakukan secara hidrotermal pada 100 °C selama 7 hari, ageing selama 24 jam dan kalsinasi pada 500 °C selama 5 jam. Variasi rasio silikat/aluminat (v/v) adalah 1, 15 and 25, sedangkan variasi konsentrasi surfaktan 5×10^{-4} M, 1×10^{-3} M dan 1×10^{-2} M. Hasil penelitian menunjukkan bahwa keseluruhan produk menghasilkan serapan yang kuat pada daerah bilangan gelombang $950\text{--}1050\text{ cm}^{-1}$, $620\text{--}690\text{ cm}^{-1}$, $420\text{--}460\text{ cm}^{-1}$. Vibrasi double ring pada $520\text{--}570\text{ cm}^{-1}$, pore opening $300\text{--}370\text{ cm}^{-1}$. Vibrasi gugus --OH yang menunjukkan keberadaan silanol dan air ditunjukkan oleh serapan pada $3400\text{--}3450\text{ cm}^{-1}$. Data XRD menunjukkan bahwa produk sintesis mempunyai kristalinitas tinggi. Pada rasio Silikat/Aluminat 1 menghasilkan zeolite tipe sodalit sedangkan rasio 15 dan 25 menghasilkan NaP1 dan SiO_2 kuarsa. Pada rasio Si/Al 1 dan 25 berturut-turut mempunyai Si sebesar 12,23% dan 12,19% serta Al sebesar 4,17% and 13,18%. Hasil SEM menunjukkan bahwa produk sintesis yang menggunakan setiltrimetilamonium bromida mempunyai bentuk yang lebih homogen dan teratur.

Kata Kunci: surfaktan; setiltrimetilamonium bromida; sintesis; zeolit; ampas tebu

INTRODUCTION

Currently, there has been an attempt to utilize the large amount of bagasse ash, such as the residue from an in-line sugar industry and the bagasse-biomass fuel in electric generation industry [1]. Sugarcane bagasse is a solid waste produced in large amount from sugar mills. Sugarcane milling industry produced 35–40% of bagasse and Indonesia has large potential to produce

bagasse. Production of sugarcane in Indonesia was 53,612,133 tons [2]. The compositions of sugarcane product are 52.9% of liquid waste, 32% of bagasse, 4.5% of molasses, 7.05% of sugar and 0.1% ash. Generally sugarcane bagasse contained of 52.67% water, 55.89% organic carbon, 0.25% total nitrogen, 0.16% P_2O_5 and 0.38% K_2O . When this sugarcane bagasse is burned under controlled conditions, it also gives ash having amorphous silica. After sugarcane

* Corresponding author.
Email address : sriatun@live.undip.ac.id

bagasse converted to ashes, the content of silica (SiO_2) to be 64.65% [3]. According to Norsurayaa et al. [4] the SiO_2 content in sugarcane bagasse ash from sugarcane juice industries in Shah Alam, Malaysia much higher, that was 88.19%. The consideration of a large number of sugarcane bagasse and the content of silica caused the development of new procedures for its productive reuse such as using as a source of silica. In the recent decades, many efforts are being carried out to explore other potential applications such as an adsorbent, fertilizers and cement mixture. So, the accumulation of waste and environment pollution can be avoided.

There are many methods applicated to increase the economic values of sugarcane bagasse ash. One of these methods was extraction by alkaline fusion. This extraction produced silica as sodium silicate and a small amount of aluminate. If the sodium silicate solution was mixed with sodium aluminate solution would be occurred condensation polymerization under hydrothermal condition [5]. Most researchers have investigated some materials as low cost silicon and aluminum sources for synthesis of various zeolites. Synthesis of NaA zeolite from sugarcane bagasse ash [6], from kaolin waste [7], NaP1 zeolite from high silica fly ash [8-10], from illite-smectite [11], from aluminum solid waste [12]. NaX and NaA [13] NaY was synthesized from rice husk silica [14] and bentonite [15]. Synthesis of zeolite as micro-mesoporous materials has been developed continuously. As micro-mesoporous materials zeolite was expected possess ordered meso-structure, porous, high surface area, acidity and thermal stability. There are some variables which should be controlled in synthesis processes such as the ratio of Si/Al, temperature and time in hydrothermal treatment and the presence of structure directing agent (SDA). The most versatile variable can be used to influence the product of synthesis was structure directing agent, because they influenced on the nucleation and crystallization processes during the formation the framework of zeolite. Organic molecules as structure directing agent must have high chemical stability in order to resist in the hydrothermal condition of the crystallization process and must be soluble in synthesis medium [16].

Selection the properties of directing agent such as size, shape, flexibility and hydrophilicity can lead to the framework of zeolites have channel with different dimensions. Several investigations have carried out using directing has controlled the morphology of MTT and MFI using TMPD (N^1, N^1, N^3, N^3 -tetramethylpropane-1,3-diamine) and HMPD (N^1, N^1, N^3, N^3 , 2,2-hexamethylpropane-1,3-diamine) as linier and branch molecule [17]. Directing agent also used to control the growth of crystal [18].

Based on previously studies information, there have never been conducted researches using

cetyltrimethylammonium bromide (CTAB) surfactants as directing agent in synthesis of zeolite from sugarcane bagasse ash. Therefore, in this study, we report the using of cetyltrimethylammonium bromide (CTAB) as structure directing agent in synthesis of zeolite from sugarcane bagasse ash as source of silica. It would examine how the effect of the CTAB as directing agent on shape of crystalline and crystallinity of the product. As a comparison, zeolite from sugarcane bagasse ash in the absence of structure directing agent was also synthesized.

The important breakthroughs resulting from this research that this research is uplifting a low cost preparation of NaP1 zeolite from low cost materials. The results of this study are expected to increase the economic value of bagasse waste into high-value material, because these products had many potential applications such as an adsorbent, catalyst and fertilizers. In addition, this study also to contributing to the field of material synthesis with renewable resources. Finally by using bagasse as raw material to make zeolite then the accumulation of waste in the environment can be reduced.

EXPERIMENTAL SECTION

Materials

In this research sugarcane bagasse collected from sugarcane industry in the region of Klaten, Central Java. First step, Sugarcane bagasse was burned then ashed in furnace at 700 °C during 4 h. Furthermore, extraction by alkaline fusion step, 96 g of sugarcane bagasse ash was reacted with about 300 mL 6 M NaOH solution. The mixture was stirred at room temperature for 24 h and filtered. The supernatant was sodium silicate solution. Determination of the content of SiO_2 , Al_2O_3 and others oxide was conducted by AAS. NaOH pellets, $\text{Al}(\text{OH})_3$ powder, cetyltrimethylammonium bromide powder, HF and HCl solution were purchase from Merck.

Instrumentation

The sample functional group is determined by FTIR spectrometer Shimadzu. The diffraction patterns of the samples were measured by X-ray diffractometer (XRD) Rigaku Multiplex with Cu K α radiation ($\lambda = 1.54184 \text{ \AA}$) at generator voltage 40 kV and current 40 mA. The simultaneous scanning electron microscope (SEM) and Energy-Dispersive X-ray (EDX) were performed using JEOL JSM 6510/LV/A/LA. The BET surface areas, pore volume and average pore size of samples were determined from N_2 adsorption isotherms at liquid nitrogen temperature (-195.7 °C)

using a Quantachrome NovaWin2, Quantachrome Instrument version 2.2. The thermal gravimetric analysis (TGA) analysis of samples was determined using LINSEIS STA Platinum Series, Platinum evaluation V1.0.138.

Procedure

An early step was preparation of sodium aluminate solution from dilution of 20 g sodium hydroxide in 100 mL aquadest. The solution was boiled, then 8.5 g aluminium hydroxide was added into solution until homogenous solution achieved. For preparation of zeolite, a number of sodium silicate supernatant was mixed with sodium aluminate solution and cetyltrimethylammonium bromide surfactant (cationic surfactant) solution. The compositions of precursor and surfactant are provided in Table 1. The mixture was stirred at specific duration until gel formed, the process was continued by hydrothermal treatment in Teflon container at 100 °C during 7 days (168 h), then aging process at room temperature for 24 h. The final steps, the precipitates were filtered, washed, dried, and calcined at 500 °C for 5 h. The product was characterized by FTIR and XRD. The provision code in this research was given in Table 1.

RESULT AND DISCUSSION

In this research, content of silica and alumina in sugarcane bagasse ash is Si 70.82% and Al 1.04% respectively, this shows that the method of extraction with alkali NaOH 6 M is effective to extract Si and Al. Thus, sugarcane bagasse ash is a potential source of silica.

The success of zeolite synthesis is demonstrated by the fingerprint absorption of FTIR from the synthesized material. It was investigated synthesis of zeolite on various of composition of sodium silicate and sodium aluminate 1, 15 and 25 (v/v). The comparison between the properties of synthesized product before and after modified with surfactant as structure directing agent is needed to find out of the influence. The FTIR spectra of synthesis product were given in Fig. 1, 2, and 3.

It has been known that every type of zeolite has two kind of specific infrared vibration, they are internal and external linkage of tetrahedral (TO_4). In all products, there can be observed the internal linkage, there is asymmetric stretching vibration of TO_4 ($\text{T} = \text{Si}$ or Al) and appears at 950–1250 cm^{-1} wavenumber, whereas the symmetric stretching vibration at 650–720 and 500 cm^{-1} . Furthermore, the wavenumber at 1050–1150 cm^{-1} was asymmetric stretching vibration, 750–820 cm^{-1} is symmetric stretching vibration in the external linkage. All

Table 1. The composition of precursor and surfactant

Sample Code	Ratio of Precursor (v/v)		Surfactant CTAB (M)
	Sodium silicate	Sodium aluminate	
A	1	1	0
A1	1	1	5×10^{-4} (1/2 cmc)
A2	1	1	1×10^{-3} (cmc)
A3	1	1	1×10^{-2} (10 cmc)
B	15	1	0
B1	15	1	5×10^{-4} (1/2 cmc)
B2	15	1	1×10^{-3} (cmc)
B3	15	1	1×10^{-2} (10 cmc)
C	25	1	0
C1	25	1	5×10^{-4} (1/2 cmc)
C2	25	1	1×10^{-3} (cmc)
C3	25	1	1×10^{-2} (10 cmc)




Fig 1. FTIR spectra of synthesized zeolite with ratio Si/Al (v/v) 1 (A) without CTAB surfactant (A1) with CTAB 5×10^{-4} M (A2) with CTAB 1×10^{-3} M (A3) with CTAB 1×10^{-2} M




Fig 2. FTIR spectra of synthesized zeolite with ratio Si/Al (v/v) 15 (B) without CTAB surfactant (B1) with CTAB 5×10^{-4} M (B2) with CTAB 1×10^{-3} M (B3) with CTAB 1×10^{-2} M




Fig 3. FTIR spectra of synthesized zeolite with ratio Si/Al (v/v) 25 (C) without CTAB surfactant (C1) with CTAB 5×10^{-4} M (C2) with CTAB 1×10^{-3} M (C3) with CTAB 1×10^{-2} M

of spectra were indicating that all the product of synthesis had Si-O-Si or Si-O-Al groups as a specific fingerprint of zeolite type. The absorption at $500\text{--}650\text{ cm}^{-1}$ is double ring [20]. Double ring is specific character of zeolite type, because it is an external linkage between polyhedral. From the FTIR data's, the zeolite was formed as synthesis product for all.

When the cetyltrimethylammonium (CTAB) as cationic surfactant is used as a directing molecule in the synthesis process of zeolite, so it would be interacted electrostatically with the silicate/ $[\text{SiO}_4]^{4-}$ and aluminate/ $[\text{AlO}_4]^{5-}$ anions primary building units. This interaction depends on surfactant concentration. The effect of surfactant presence can be observed on spectra A1, A2, and A3 in Fig. 1, B1, B2, and B3 in Fig. 2, C1, C2, and C3 in Fig. 3. It could be clearly observed that the presence of cetyltrimethylammonium (CTAB) surfactants has significantly affected the intensity and patterns of absorption at the wavenumber $950\text{--}1250$ and $3400\text{--}3500\text{ cm}^{-1}$. The lower absorption intensity of wave number at $3400\text{--}3500\text{ cm}^{-1}$ in the CTAB modified zeolite showed the low levels of silanol groups in the sample, where this corresponds to the formation of framework of zeolite. Besides that, the stronger bonding of Si-O-Si and Si-O-Al in zeolite framework characterized by increasing of intensity absorption bands at wave numbers $950\text{--}1250\text{ cm}^{-1}$ after the addition of CTAB.

In the FTIR spectra of zeolite modified by CTAB surfactant, no new absorption was observed. The spectra have similarity to the absorption region of zeolite which synthesized without surfactant. It was showed that the calcination treatment at 600°C to the modified zeolite by CTAB has removed and decomposed the CTAB as structure directing and all undesired organic compounds.




Fig 4. The pattern of diffractogram synthesized zeolite (A) Ratio Si/Al 1 without CTAB (A3) Ratio Si/Al 1 with CTAB 1×10^{-2} M (B) Ratio Si/Al 15 without CTAB (B3) Ratio Si/Al 15 with CTAB 1×10^{-2} M (C) Ratio Si/Al 25 without CTAB (C3) Ratio Si/Al 25 with CTAB 1×10^{-2} M

Thus the shape and pore size of modified zeolite was expected homogenously.

Fig. 4 shows the pattern of diffractogram XRD of synthesis product. Inorganic material has different shape and structure crystal, due to the diffractogram pattern was characteristic.

The intensity of diffractogram indicated the crystal quality of material. It could investigate the kind of synthesis product by CTAB different with synthesis product without CTAB. In addition, the composition of precursor also affected the kind of product. The synthesis product by ratio of Si/Al (v/v) 1 without CTAB was sodalite, because the XRD pattern has similarity to the sodalite standard in Treacy [21]. The sodalite is plain type of zeolite. The same product was obtained on ratio Si/Al (v/v) 1 used CTAB as structure directing. Although the product is the same but it is observed in XRD that the using of CTAB makes the peak of the diffractogram sharper and the product more pure.

If the ratio of Si/Al (v/v) increased to 15 or 25, the synthesis product was NaP1 zeolite. The presence of CTAB surfactant as structure directing was not affected on the type zeolite but on the main peak of diffractogram. The shifting of 2θ and the changing intensity of peaks at sodalite and NaP1 were known from the Fig. 4. All of the peaks on NaP1 diffractogram NaP1 as synthesis product modified by CTAB shifted to higher angle. Overall the intensity of peaks increased, thus it was concluded that the product modified by CTAB more crystalline.

Fig. 5 is SEM photograph corresponding to the synthesis product. From the XRD data, the products by ratio Si/Al (v/v) 1 is sodalite and from the SEM it appears




Fig 5. The SEM image of synthesized zeolite (A) Ratio Si/Al 1 without CTAB (A3) Ratio Si/Al 1 with CTAB 1×10^{-2} M (B) Ratio Si/Al 15 without CTAB (B3) Ratio Si/Al 15 with CTAB 1×10^{-2} M (C) Ratio Si/Al 25 without CTAB (C3) Ratio Si/Al 25 with CTAB 1×10^{-2} M

Table 2. The chemical composition of synthesized zeolite

Element	Wt (%)					
	A1	A3	B1	B3	C1	C3
O	48.03	45.78	53.14	53.09	53.20	42.59
Na	5.25	17.96	12.65	14.05	13.80	18.49
Mg		00.25				0.34
Al	3.72	18.97	10.81	11.47	11.52	17.25
Si	3.26	17.04	11.18	11.69	11.90	16.6
S		-	1.68	1.74	1.63	3.57
K		-	1.67	0.44	0.55	1.16
Ca				0.16		

Description: (A) Ratio Si/Al 1 without CTAB (A3) Ratio Si/Al 1 with CTAB 1×10^{-2} M (B) Ratio Si/Al 15 without CTAB (B3) Ratio Si/Al 15 with CTAB 1×10^{-2} M (C) Ratio Si/Al 25 without CTAB (C3) Ratio Si/Al 25 with CTAB 1×10^{-2} M

that the materials had regular spherical crystal and homogenous. At ratio Si/Al (v/v) 15 and 25 the materials had the same type, rectangular shape with thin platy crystal, however at ratio Si/Al (v/v) 15 has smaller in size. It showed different shape to the ratio Si/Al (v/v) 1. It could be investigated that the shape and size of granular/crystal depends on the composition of precursor, it means the ratio of Si/Al. The use of surfactant CTAB as a directing agent affects the size and homogeneity of the product particles. In SEM it appears that the crystalline product of synthesis with

CTAB has a more porous, regular and firm form crystal. The data's from SEM consistent with the result to the XRD, that the using of surfactant CTAB produce a more crystalline materials.

Meanwhile the chemical composition of products synthesis in Table 2. Sample A1 and A3 are sodalite as the product of synthesis. For sample B1, B3, C1, and C3 are NaP1. The entire product showed that ratio of Si/Al almost 1, these are consistent with the BET result that this product is micropore material. The use of surfactant CTAB as a pore directing agent increases the




Fig 6. The isotherms adsorption of the synthesized product ratio Si/Al (v/v) 1




Fig 7. TGA of the synthesized product ratio Si/Al (v/v) 1

number of ions silicate and aluminate that interacted to form the product, so the using of surfactant is needed in this process. This is appropriate to the Wang [22] that the surfactants such as PEG (polyethylene glycol) and CTAB play an important role during the formation of nano crystal IM-5 zeolite.

BET (Brunauer–Emmett–Teller) method is used for multi molecular adsorption on solid surface, so the surface area of a material can be known with a surface area gauge that uses the BET method principle. The measurement of surface area with BET model used nitrogen as adsorbate. This measurement is based on isothermal adsorption data nitrogen at a temperature of 77.35 K. Isothermal adsorption with BET principle is a type physical isotherms. BET analysis is given in Fig. 6. The isotherm could be classified as type I of Brunauer's

classification showing closed hysteresis loops. The calculated surface area of the prepared zeolite is relatively low 22.062 m²/g. This reveals the existence only a small number of active sorption sites. The total pore volume, V_p was found to be 8.56×10^{-2} cm³/g and average pore radius was determined as 7.737 Å. It means the NaP1 of this research has a micropore structure.

The TGA (thermogravimetric analysis) graph for the synthesized zeolite from sugarcane bagasse ash is given in Fig. 7. From the figure, it can be seen that the sample is lost the weight at temperature 120 °C, it means the loosely bond or free water molecules from product zeolite cage. Then the reducing weight also occurred at 270- which is due to the decomposition of the CTAB surfactant residues on the product zeolite surface.

CONCLUSION

In this study zeolite were successfully obtained from sugarcane bagasse ash by hydrothermal method at 100 °C for 168 h. The type of zeolite was determined by the ratio of Si/Al in term the SiO₄⁴⁻ and AlO₄⁵⁻ which formed the zeolite framework. The ratio Si/Al 1 (v/v) produced sodalite, while the ratio Si/Al 15 and 25 produced NaP1. The presence of CTAB surfactant micelles in synthesis zeolite from sugarcane bagasse ash could improve the homogeneity and crystallinity of the product zeolite. It means the product has homogeneous and orderly structure. But the introduction of CTAB micelles in the synthesis failed in terms of generating the mesopores structure because both of sodalite and NaP1 still had micropores structure.

ACKNOWLEDGEMENT

Sriatun, Taslimah, and Suyati, L., gratefully acknowledge financial support from The Directorate of Research and Community Service, Ministry of Research, Technology and High Education via Grants Compete Award (Number: 84-68/UN7.5.1/PG/2014) and Chemistry Department of Diponegoro University for the facilities to carry out this research.

REFERENCES

- [1] Srinivasan, R., and Sathiya, K., 2010, Experimental study on bagasse ash in concrete, *Int. J. Serv. Learn. Eng.*, 5 (2), 60–66.
- [2] *Indonesian Statistics*, 2012, <http://www.bps.go.id>.
- [3] Hanafi, A., and Nandang, A., 2010, Studi pengaruh bentuk silika dari ampas tebu terhadap kekuatan produk keramik, *Jurnal Kimia Indonesia*, 5 (1), 35–38.
- [4] Norsuraya, S., Fazlena, H., and Norhasyimi, R., 2016, Sugarcane bagasse as a renewable Source of silica to synthesize Santa Barbara Amorphous-15 (SBA-15), *Procedia Eng.*, 148, 839–846.
- [5] Cardoso, A.M., Paprocki, A., Ferret, L.S., Azevedo, C.M.N., and Pires, M., 2015, Synthesis of zeolite Na-P1 under mild conditions using Brazilian coal fly ash and its application in wastewater treatment, *Fuel*, 139, 59–67.
- [6] Moisés, M.P., da Silva, C.T.P., Meneguín, J.G., Giroto, E.M., and Radovanovic, E., 2013, Synthesis of zeolite NaA from sugarcane bagasse ash, *Mater. Lett.*, 108, 243–246.
- [7] Maia, A.Á.B., Neves, R.F., Angélica, R.S., and Pöllmann, H., 2015, Synthesis, optimisation and characterisation of the zeolite NaA using kaolin waste from the Amazon Region. Production of zeolites KA, MgA and CaA, *Appl. Clay Sci.*, 108, 55–60.
- [8] Kazemian, H., Naghdali, Z., Kashani, T.G., and Farhadi, F., 2010, Conversion of high silicon fly ash to Na-P1 zeolite: Alkaline fusion followed by hydrothermal crystallization, *Adv. Powder Technol.*, 21 (3), 279–283.
- [9] Babajide, O., Musyoka, N., Petrik, L., and Ameer, F., 2012, Novel zeolite Na-X synthesized from fly ash as a heterogeneous catalyst in biodiesel production, *Catal. Today*, 190 (1), 54–60.
- [10] Musyoka, N.M., Petrik, L.F., Fatoba, O.O., and Hums, E., 2013, Synthesis of zeolites from coal fly ash using mine waters, *Miner. Eng.*, 53, 9–15.
- [11] Baccouche, A., Srasra, E., and El Maaoui, M., 1998, Preparation of Na-P1 and sodalite octahydrate zeolites from interstratified illite–smectite, *Appl. Clay Sci.*, 13 (4), 255–273.
- [12] Sánchez-Hernández, R., López-Delgado, A., Padilla, I., Galindo, R., and López-Andrés, S., 2016, One-step synthesis of NaP1, SOD and ANA from a hazardous aluminum solid waste, *Microporous Mesoporous Mater.*, 226, 267–277.
- [13] Purnomo, C.W., Salim, C., and Hinode, H., 2012, Synthesis of pure Na–X and Na–A zeolite from bagasse fly ash, *Microporous Mesoporous Mater.*, 162, 6–13.
- [14] Wittayakun, J., Khemthong, P., and Prayoonpokarach, S., 2008, Synthesis and characterization of zeolite NaY from rice husk silica, *Korean J. Chem. Eng.*, 25 (4), 861–864.
- [15] Faghihian, H., and Godazandeha, N., 2009, Synthesis of nano crystalline zeolite Y from bentonite, *J. Porous Mater.*, 16 (3), 331–335.
- [16] Almeida, R.K.S., Gómez-Hortigüela, L., Pinar, A.B., and Pérez-Pariente, J., 2016, Synthesis of ferrierite by a new combination of co-structure-directing agents: 1,6-bis(N-methylpyrrolidinium)hexane and tetramethylammonium, *Microporous Mesoporous Mater.*, 232, 218–226.
- [17] Lee, H.J., Kim, S.H., Kim, J.H., Park, S.J., and Cho, S.J., 2014, Synthesis and characterization of zeolites MTT and MFI, with controlled morphologies using mixed structure directing agents, *Microporous Mesoporous Mater.*, 195, 205–215.
- [18] Hasan, F., Singh, R., Li, G., Zhao, D., and Webley, P.A., 2012, Direct synthesis of hierarchical LTA zeolite via a low crystallization and growth rate technique in presence of cetyltrimethylammonium bromide, *J. Colloid Interface Sci.*, 382 (1), 1–12.
- [19] Su, S., Ma, H., and Chuan, X., 2016, Hydrothermal synthesis of zeolite A from K-feldspar and its crystallization mechanism, *Adv. Powder Technol.*, 27 (1), 139–144.
- [20] Weitkamp, J., and Puppe, L., 1999, *Catalysis and Zeolites: Fundamentals and Applications*, Springer-Verlag Berlin Heidelberg.
- [21] Treacy, M.M.J., and Higgins, J.B., 2001, *Collection of Simulated XRD Powder Pattern for Zeolite*, 4th rev. ed., Elsevier Science.
- [22] Wang, L., Yang, W., Xin, C., Ling, F., Sun, W., Fang, X., and Yang, R., 2012, Synthesis of nano-zeolite IM-5 by hydrothermal method with the aid of PEG and CTAB, *Mater. Lett.*, 69, 16–19.

Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent

by Sriatun Sriatun

Submission date: 25-May-2019 08:33AM (UTC+0700)

Submission ID: 1135642929

File name: Cetyltrimethylammonium_Bromide_as_Structure_Directing_Agent.pdf (429.2K)

Word count: 4250

Character count: 21744

Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent

Sriatun*, Taslimah, and Linda Suyati

Department of Chemistry, Faculty of Science and Mathematics, Diponegoro University,
Jl. Prof. Soedarto, SH, Tembalang, Semarang 50275, Indonesia

Received February 18, 2017; Accepted October 24, 2017

ABSTRACT

The purpose of this research is to synthesize zeolite from sugarcane bagasse ash using cetyltrimethylammonium bromide as structure directing agent. This research used cetyltrimethylammonium bromide surfactant to invent the high porosity, surface area, acidity and thermal stability of synthesized zeolite. The Silica was extracted by alkaline fusion using sodium hydroxide solution. The synthesis was conducted by hydrothermal process at 100 °C for 7 days, ageing process for 24 h and calcination at 500 °C for 5 h. Ti_{23} ratio of Si/Al (v/v) was 1, 15 and 25, meanwhile the concentration of cetyltrimethylammonium bromide was 5×10^{-4} M, 1×10^{-3} M and 1×10^{-2} M. The result showed all of product have strong absorbance at 950–1050 cm^{-1} and 620–690 cm^{-1} , 420–460 cm^{-1} , double ring at 520–570 cm^{-1} , pore opening at 300–370 cm^{-1} . Vibration of –OH as silanol group or water was indicated by broad absorbance at 3400–3450 cm^{-1} . The diffractograms XRD showed that the product had high crystallinity. The composition of product on ratio Si/Al 1 with concentration of cetyltrimethylammonium 10^{-2} M is sodalite, the ratio Si/Al 15 and 25 are NaP1 and SiO₂ quartz and contain 12.23% and 12.19% of Si, 4.17% and 13.18% of Al, respectively. Observation on SEM revealed that the crystal produced using cetyltrimethylammonium bromide was more homogenous and regular in shape.

Keywords: surfactant; cetyltrimethylammonium bromide; synthesis; zeolite; sugarcane bagasse

ABSTRAK

Tujuan penelitian ini adalah membuat zeolit dari ampas tebu menggunakan setiltrimetilamonium bromida sebagai agen pengarah struktur. Dalam penelitian ini surfaktan setiltrimetilamonium bromida digunakan untuk meningkatkan porositas, luas permukaan, keasaman dan stabilitas termal. Silika diekstrak dari abu ampas tebu melalui penambahan alkali yaitu NaOH 6 M. Sintesis dilakukan secara hidrotermal pada 100 °C selama 7 hari, ageing selama 24 jam dan kalsinasi pada 500 °C selama 5 jam. Variasi rasio silikat/aluminat (v/v) adalah 1, 15 and 25, sedangkan variasi konsentrasi surfaktan 5×10^{-4} M, 1×10^{-3} M dan 1×10^{-2} M. Hasil penelitian menunjukkan bahwa keseluruhan produk menghasilkan serapan yang kuat pada daerah bilangan gelombang 950–1050 cm^{-1} , 620–690 cm^{-1} , 420–460 cm^{-1} . Vibrasi double ring pada 520–570 cm^{-1} , pore opening 300–370 cm^{-1} . Vibrasi gugus –OH yang menunjukkan keberadaan silanol dan air ditunjukkan oleh serapan pada 3400–3450 cm^{-1} . Data XRD menunjukkan bahwa produk sintesis mempunyai kristalinitas tinggi. Pada rasio Silikat/Aluminat 1 menghasilkan zeolite tipe sodalit sedangkan rasio 15 dan 25 menghasilkan NaP1 dan SiO₂ kuarsa. Pada rasio Si/Al 1 dan 25 berturut-turut mempunyai Si sebesar 12,23% dan 12,19% serta Al sebesar 4,17% and 13,18%. Hasil SEM menunjukkan bahwa produk sintesis yang menggunakan setiltrimetilamonium bromida mempunyai bentuk yang lebih homogen dan teratur.

Kata Kunci: surfaktan; setiltrimetilamonium bromida; sintesis; zeolit; ampas tebu

INTRODUCTION

Currently, there has been an attempt to utilize the large amount of bagasse ash, such as the residue from an in-line sugar industry and the bagasse-bioass fuel in electric generation industry [1]. Sugarcane bagasse is a solid waste produced in large amount from sugar mills. Sugarcane milling industry produced 35–40% of bagasse and Indonesia has large potential to produce

bagasse. Production of sugarcane in Indonesia was 53,612,133 tons [2]. The compositions of sugarcane product are 52.9% of liquid waste, 32% of bagasse, 4.5% of molasses, 7.05% of sugar and 0.1% ash. Generally sugarcane bagasse contained of 52.67% water, 55.89% organic carbon, 0.25% total nitrogen, 0.16% P₂O₅ and 0.38% K₂O. When this sugarcane bagasse is burned under controlled conditions, it also gives ash having amorphous silica. After sugarcane

* Corresponding author.
Email address : sriatun@live.undip.ac.id

bagasse converted to ashes, the content of silica (SiO_2) to be 64.65% [3]. According to Norsuray^[27] et al. [4] the SiO_2 content in sugarcane bagasse ash from sugarcane juice industries in Shah Alam, Malaysia much higher, that was 88.19%. The consideration of a large number of sugarcane bagasse and the content of silica caused the development of new procedures for its productive reuse such as using as a source of silica. In the recent decades, many efforts are being carried out to explore other potential applications such as an adsorbent, fertilizers and cement mixture. So, the accumulation of waste and environment pollution can be avoided.

There are many methods applicated to increase the economic values of sugarcane bagasse ash. One of these methods was extraction by alkaline fusion. This extraction produced silica as sodium silicate and a small amount of aluminate. If the sodium silicate solution was mixed with sodium aluminate solution would be occurred condensation polymerization under hydrothermal condition [5]. Most res^[22]chers have investigated some materials as low cost silicon and aluminum sources for synthesis of various zeolites. Synthesis of NaA zeolite from sugarcane bagasse ash [6], from kaolin waste [7], NaP1 zeolite from high silica fly ash [8-10], from illite-smectite [11], from aluminum solid waste [12]. NaX and NaA [13] NaY was synthesized from rice husk silica [14] and bentonite [15]. Synthesis of zeolite as micro-mesoporous materials has been developed continuously. As micro-mesoporous materials zeolite was expected possess ordered meso-structure, porous, high surface area, acidity and thermal stability. There are some variables which should be controlled in synthesis processes such as the ratio of Si/Al, temperature and time in hydrothermal treatment and the presence of structure directing agent (SDA). The most versatile variable can be used to influence the product of synthesis was structure directing agent, because they influenced on the nucleation and crystallization processes during the formation the framework of zeolite. Organic molecules as structure directing agent must have high chemical stability in order to resist in the hydrothermal condition of the crystallization process and must be soluble in synthesis medium [16].

Selection the properties of directing agent such as size, shape, flexibility and hydrophilicity can lead to the framework of zeolites have channel with different dimensions. Several investigations have carried out using directing has cor^[20]lled the morphology of MTT and MFI using TMPD (N^1, N^1, N^3, N^3 -tetramethylpropane-1,3-diamine) and HMPD ($N^1, N^1, N^3, N^3, 2, 2$ -hexamethylpropane-1,3-diamine) as linier and branch molecule [17]. Directing agent also used to control the growth of crystal [18].

Based on previously studies information, there have never been conducted researches using

cetyltrimethylammonium bromide (CTAB) surfactants as directing agent in synthesis of zeolite from sugarcane bagasse ash. Therefore, in this study, we report the using of cetyltrimethylammon^[1]ium bromide (CTAB) as structure directing agent in synthesis of zeolite from sugarcane bagasse ash as source of silica. It would examine how the effect of the CTAB as directing agent on shape of crystalline and crystallinity of the product. As a comparison, zeolite from sugarcane bagasse ash in the absence of structure directing agent was also synthesized.

The important breakthroughs resulting from this research that this research is uplifting a low cost preparation of NaP1 zeolite from low cost materials. The results of this study are expected to increase the economic value of bagasse waste into high-value material, because these products had many potential applications such as an adsorbent, catalyst and fertilizers. In addition, this study also to contributing to the field of material synthesis with renewable resources. Finally by using bagasse as raw material to make zeolite then the accumulation of waste in the environment can be reduced.

EXPERIMENTAL SECTION

Materials

In this research sugarcane bagasse collected from sugarcane industry in the region of Klaten, Central Java. First step, Sugarcane bagasse was burned then ashed in furnace at 700 °C during 4 h. Furthermore, extraction by alkaline fusion step, 96 g of sugarcane bagasse ash was rea^[26]d with about 300 mL 6 M NaOH solution. The mixture was stirred at room temperature for 24 h and filtered. The supernatant was sodium silicate solution. Determination of the content of SiO_2 , Al_2O_3 and others oxide was conducted by AAS. NaOH pellets, $\text{Al}(\text{OH})_3$ powder, cetyltrimethylammonium bromide powder, HF and HCl solution were purchase from Merck.

Instrumentation

The sample functional group is determined by FTIR ^[6]ectrometer Shimadzu. The diffraction patterns of the samples were measured by X-ray diffractometer (XRD) Rigaku Multiplex with Cu $K\alpha$ radiation ($\lambda = 1.54184 \text{ \AA}$) at generator ^[8]oltage 40 kV and current 40 mA. The simultaneous scanning electron microscope (SEM) and Energy-Dispersive X-ray (EDX) were ^[30]formed using JEOL JSM 6510/LV/A/LA. The BET surface areas, pore volume and average pore size of samples were determined from N_2 adsorption isotherms at liquid nitrogen temperature (-195.7 °C)

using a Quantachrome NovaWin2, Quantachrome Instrument version 2.2. The thermal gravimetric analysis (TGA) analysis of samples was determined using LINSEIS STA Platinum Series, Platinum evaluation V1.0.138.

Procedure

An early step was preparation of sodium aluminate solution from dilution of 20 g sodium hydroxide in 100 mL aquadest. The solution was boiled, then 8.5 g aluminium hydroxide was added into solution until homogenous solution achieved. For preparation of zeolite, a number of sodium silicate supernatant was mixed with sodium aluminate solution and cetyltrimethylammonium bromide surfactant (cationic surfactant) solution. The compositions of precursor and surfactant are provided in Table 1. The mixture was stirred at specific duration until gel formed, the process was continued by hydrothermal treatment in Teflon container $24 \pm 100^\circ\text{C}$ during 7 days (168 h), then aging process at room temperature for 24 h. The final steps, the precipitates were filtered, washed, dried, and calcined at 500°C for 5 h. The product was characterized by FTIR and XRD. The provision code in this research was given in Table 1.

RESULT AND DISCUSSION

In this research, content of silica and alumina in sugarcane bagasse ash is Si 70.82% and Al 1.04% respectively, this shows that the method of extraction with alkali NaOH 6 M is effective to extract Si and Al. Thus, sugarcane bagasse ash is a potential source of silica.

The success of zeolite synthesis is demonstrated by the fingerprint absorption of FTIR from the synthesized material. It was investigated synthesis of zeolite on various of composition of sodium silicate and sodium aluminate 1, 15 and 25 (v/v). The comparison between the properties of synthesized product before and after modified with surfactant as structure directing agent is needed to find out of the influence. The FTIR spectra of synthesis product were given in Fig. 1, 2, and 3.

It has been known that every type of zeolite has two kind of specific infrared vibration, they are internal and external linkage of tetrahedral (TO_4). In all products, there can be observed the internal linkage, there is asymmetric stretching vibration of TO_4 ($T = \text{Si}$ or Al) and appears at $950\text{--}1250\text{ cm}^{-1}$ wavenumber, whereas the symmetric stretching vibration at $650\text{--}720$ and 500 cm^{-1} . Furthermore, the wavenumber at $1050\text{--}1150\text{ cm}^{-1}$ was asymmetric stretching vibration, $750\text{--}820\text{ cm}^{-1}$ is symmetric stretching vibration in the external linkage. All

Table 1. The composition of precursor and surfactant

Sample Code	Ratio of Precursor (v/v)		Surfactant CTAB (M)
	Sodium silicate	Sodium aluminate	
A	1	1	0
A1	1	1	5×10^{-4} (1/2 cmc)
A2	1	1	1×10^{-3} (cmc)
A3	1	1	1×10^{-2} (10 cmc)
B	15	1	0
B1	15	1	5×10^{-4} (1/2 cmc)
B2	15	1	1×10^{-3} (cmc)
B3	15	1	1×10^{-2} (10 cmc)
C	25	1	0
C1	25	1	5×10^{-4} (1/2 cmc)
C2	25	1	1×10^{-3} (cmc)
C3	25	1	1×10^{-2} (10 cmc)




Fig 1. FTIR spectra of synthesized zeolite with ratio Si/Al (v/v) 1 (A) without CTAB surfactant (A1) with CTAB 5×10^{-4} M (A2) with CTAB 1×10^{-3} M (A3) with CTAB 1×10^{-2} M




Fig 2. FTIR spectra of synthesized zeolite with ratio Si/Al (v/v) 15 (B) without CTAB surfactant (B1) with CTAB 5×10^{-4} M (B2) with CTAB 1×10^{-3} M (B3) with CTAB 1×10^{-2} M




Fig 3. FTIR spectra of synthesized zeolite with ratio Si/Al (v/v) 25 (C) without CTAB surfactant (C1) with CTAB 5×10^{-4} M (C2) with CTAB 1×10^{-3} M (C3) with CTAB 1×10^{-2} M

of spectra were indicating that all the product of synthesis had Si-O-Si or Si-O-Al groups as a specific fingerprint of zeolite type. The absorption at $500\text{--}650\text{ cm}^{-1}$ is double ring [20]. Double ring is specific character of zeolite type, because it is an external linkage between polyhedral. From the FTIR data's, the zeolite was formed as synthesis product for all.

When the cetyltrimethylammonium (CTAB) as cationic surfactant is used as a directing molecule in the synthesis process of zeolite, so it would be interacted electrostatically with the silicate/ $[\text{SiO}_4]^{4-}$ and aluminate/ $[\text{AlO}_4]^{5-}$ anions primary building units. This interaction depends on surfactant concentration. The effect of surfactant presence can be observed on spectra A1, A2, and A3 in Fig. 1, B1, B2, and B3 in Fig. 2, C1, C2, and C3 in Fig. 3. It could be clearly observed that the presence of cetyltrimethylammonium (CTAB) surfactants has significantly affected the intensity and patterns of absorption at the wavenumber $950\text{--}1250$ and $3400\text{--}3500\text{ cm}^{-1}$. The lower absorption intensity of wave number at $3400\text{--}3500\text{ cm}^{-1}$ in the CTAB modified zeolite showed the low levels of silanol groups in the sample, where this corresponds to the formation of framework of zeolite. Besides that, the stronger bonding of Si-O-Si and Si-O-Al in zeolite framework characterized by increasing of intensity absorption bands at wave numbers $950\text{--}1250\text{ cm}^{-1}$ after the addition of CTAB.

In the FTIR spectra of zeolite modified by CTAB surfactant, no new absorption was observed. The spectra have similarity to the absorption region of zeolite which synthesized without surfactant. It was showed that the calcination treatment at 600°C to the modified zeolite by CTAB has removed and decomposed the CTAB as structure directing and all undesired organic compounds.




Fig 4. The pattern of diffractogram synthesized zeolite (A) Ratio Si/Al 1 without CTAB (A3) Ratio Si/Al 1 with CTAB 1×10^{-2} M (B) Ratio Si/Al 15 without CTAB (B3) Ratio Si/Al 15 with CTAB 1×10^{-2} M (C) Ratio Si/Al 25 without CTAB (C3) Ratio Si/Al 25 with CTAB 1×10^{-2} M

Thus the shape and pore size of modified zeolite was expected homogenously.

Fig. 4 shows the pattern of diffractogram XRD of synthesis product. Inorganic material has different shape and structure crystal, due to the diffractogram pattern was characteristic.

The intensity of diffractogram indicated the crystal quality of material. It could investigate the kind of synthesis product by CTAB different with synthesis product without CTAB. In addition, the composition of precursor also affected the kind of product. The synthesis product by ratio of Si/Al (v/v) 1 without CTAB was sodalite, because the XRD pattern has similarity to the sodalite standard in Treacy [21]. The sodalite is plain type of zeolite. The same product was obtained on ratio Si/Al (v/v) 1 used CTAB as structure directing. Although the product is the same but it is observed in XRD that the using of CTAB makes the peak of the diffractogram sharper and the product more pure.

If the ratio of Si/Al (v/v) increased to 15 or 25, the synthesis product was NaP1 zeolite. The presence of CTAB surfactant as structure directing was not affected on the type zeolite but on the main peak of diffractogram. The shifting of 2θ and the changing intensity of peaks at sodalite and NaP1 were known from the Fig. 4. All of the peaks on NaP1 diffractogram NaP1 as synthesis product modified by CTAB shifted to higher angle. Overall the intensity of peaks increased, thus it was concluded that the product modified by CTAB more crystalline.

Fig. 5 is SEM photograph corresponding to the synthesis product. From the XRD data, the products by ratio Si/Al (v/v) 1 is sodalite and from the SEM it appears




Fig 5. The SEM image of synthesized zeolite (A) Ratio Si/Al 1 without CTAB (A3) Ratio Si/Al 1 with CTAB 1×10^{-2} M (B) Ratio Si/Al 15 without CTAB (B3) Ratio Si/Al 15 with CTAB 1×10^{-2} M (C) Ratio Si/Al 25 without CTAB (C3) Ratio Si/Al 25 with CTAB 1×10^{-2} M

Table 2. The chemical composition of synthesized zeolite

Element	Wt (%)					
	A1	A3	B1	B3	C1	C3
O	48.03	45.78	53.14	53.09	53.20	42.59
Na	5.25	17.96	12.65	14.05	13.80	18.49
Mg		00.25				0.34
Al	3.72	18.97	10.81	11.47	11.52	17.25
Si	3.26	17.04	11.18	11.69	11.90	16.6
S		-	1.68	1.74	1.63	3.57
K		-	1.67	0.44	0.55	1.16
Ca				0.16		

Description: (A) Ratio Si/Al 1 without CTAB (A3) Ratio Si/Al 1 with CTAB 1×10^{-2} M (B) Ratio Si/Al 15 without CTAB (B3) Ratio Si/Al 15 with CTAB 1×10^{-2} M (C) Ratio Si/Al 25 without CTAB (C3) Ratio Si/Al 25 with CTAB 1×10^{-2} M

that the materials had regular spherical crystal and homogenous. At ratio Si/Al (v/v) 15 and 25 the materials had the same type, rectangular shape with thin platy crystal, however at ratio Si/Al (v/v) 15 has smaller in size. It showed different shape to the ratio Si/Al (v/v) 1. It could be investigated that the shape and size of granular/crystal depends on the composition of precursor, it means the ratio of Si/Al. The use of surfactant CTAB as a directing agent affects the size and homogeneity of the product particles. In SEM it appears that the crystalline product of synthesis with

CTAB has a more porous, regular and firm form crystal. The data's from SEM consistent with the result to the XRD, that the using of surfactant CTAB produce a more crystalline materials.

Meanwhile the chemical composition of products synthesis in Table 2. Sample A1 and A3 are sodalite as the product of synthesis. For sample B1, B3, C1, and C3 are NaP1. The entire product showed that ratio of Si/Al almost 1, these are consistent with the BET result that this product is micropore material. The use of surfactant CTAB as a pore directing agent increases the




Fig 6. The isotherms adsorption of the synthesized product ratio Si/Al (v/v) 1




Fig 7. TGA of the synthesized product ratio Si/Al (v/v) 1

number of ions silicate and aluminate that interacted to form the product, so the using of surfactant is needed in this process. This is appropriate to the Wang [22] 3 at the surfactants such as PEG (polyethylene glycol) and CTAB play an important role during the formation of nano crystal IM-5 zeolite.

BET (Brunauer–Emmett–Teller) method is used for multi molecular adsorption on solid surface, so the surface area of a material can be known with a surface area gauge that uses the BET method principle. The measurement of surface area with BET model used nitrogen as adsorbate. This measurement is based on isothermal adsorption data nitrogen at a temperature of 77.35 K. Isothermal adsorption with BET principle is a type physical isotherms. BET analysis is given in Fig. 6. The isotherm could be classified as type I of Brunauer's

classification showing closed hysteresis loops. The calculated surface area of the prepared zeolite is relatively low 22.062 m²/g. This reveals the existence only a small number of active sorption sites. The total pore volume, V_p was found to be 8.56×10^{-2} cm³/g and average pore radius was determined as 7.737 Å. It means the NaP1 of this research has a micropore structure.

The TGA (thermogravimetric analysis) graph for the synthesized zeolite from sugarcane bagasse ash is given in Fig. 7. From the figure, it can be seen that the sample is lost the weight at temperature 120 °C, it means the loosely bond or free water molecules from product zeolite cage. Then the reducing weight also occurred at 270- which is due to the decomposition of the CTAB surfactant residues on the product zeolite surface.

CONCLUSION

In this study zeolite were successfully obtained from sugarcane bagasse ash by hydrothermal method at 100 °C for 168 h. The type of zeolite was determined by the ratio of Si/Al in term the SiO₄⁴⁻ and AlO₄⁵⁻ which formed the zeolite framework. The ratio Si/Al 1 (v/v) produced sodalite, while the ratio Si/Al 15 and 25 produced NaP1. The presence of CTAB surfactant micelles in synthesis zeolite from sugarcane bagasse ash could improve the homogeneity and crystallinity of the product zeolite. It means the product has homogeneous and orderly structure. But the introduction of CTAB micelles in the synthesis failed in terms of generating the mesopores structure because both of sodalite and NaP1 still had micropores structure.

ACKNOWLEDGEMENT

Sriatun, Taslimah, and Suyati, L., gratefully acknowledge financial support from The Directorate of Research and Community Service, Ministry of Research, Technology and High Education via Grants Compete Award (Number: 84-68/UN7.5.1/PG/2014) and Chemistry Department of Diponegoro University for the facilities to carry out this research.

REFERENCES

- [1] Srinivasan, R., and Sathiya, K., 2010, Experimental study on bagasse ash in concrete, *Int. J. Serv. Learn. Eng.*, 5 (2), 60–66.
- [2] Indonesian Statistics, 2012, <http://www.bps.go.id>.
- [3] Hanafi, A., and Nandang, A., 2010, Studi pengaruh bentuk silika dari ampas tebu terhadap kekuatan produk keramik, *Jurnal Kimia Indonesia*, 5 (1), 35–38.
- [4] Norsuraya, S., Fazlena, H., and Norhasyimi, R., 2016, Sugarcane bagasse as a renewable Source of silica to synthesize Santa Barbara Amorphous-15 (SBA-15), *Procedia Eng.*, 148, 839–846.
- [5] Cardoso, A.M., Paprocki, A., Ferret, L.S., Azevedo, C.M.N., and Pires, M., 2015, Synthesis of zeolite Na-P1 under mild conditions using Brazilian coal fly ash and its application in wastewater treatment, *Water*, 7, 139, 59–67.
- [6] Moisés, M.P., da Silva, C.T.P., Meneguín, J.G., Giroto, E.M., and Radovanovic, E., 2013, Synthesis of zeolite NaA from sugarcane bagasse ash, *Mater. Res. Lett.*, 108, 243–246.
- [7] Maia, A.Á.B., Neves, R.F., Angélica, R.S., and Pöhlmann, H., 2015, Synthesis, optimisation and characterisation of the zeolite NaA using kaolin waste from the Amazon Region. Production of zeolites KA, MgA and CaA, *Appl. Clay Sci.*, 108, 41–60.
- [8] Kazemian, H., Naghdali, Z., Kashani, T.G., and Farhadi, F., 2010, Conversion of high silicon fly ash to Na-P1 zeolite: Alkaline fusion followed by hydrothermal crystallization, *Adv. Powder Technol.*, 21 (3), 279–283.
- [9] Babajide, O., Musyoka, N., Petrik, L., and Ameer, F., 2012, Novel zeolite Na-X synthesized from fly ash as a heterogeneous catalyst in biodiesel production, *Catal. Today*, 190 (1), 54–60.
- [10] Musyoka, N.M., Petrik, L.F., Fatoba, O.O., and Hums, E., 2013, Synthesis of zeolites from coal fly ash using mine waters, *Miner. Eng.*, 53, 9–15.
- [11] Baccouche, A., Srasra, E., and El Maaoui, M., 1998, Preparation of Na-P1 and sodalite octahydrate zeolites from interstratified illite–montmorillonite, *Appl. Clay Sci.*, 13 (4), 255–273.
- [12] Sánchez-Hernández, R., López-Delgado, A., Padilla, I., Galindo, R., and López-Andrés, S., 2016, One-step synthesis of NaP1, SOD and ANA from a hazardous aluminum solid waste, *Microporous Mesoporous Mater.*, 226, 267–277.
- [13] Purnomo, C.W., Salim, C., and Hinode, H., 2012, Synthesis of pure Na-X and Na-A zeolite from bagasse fly ash, *Microporous Mesoporous Mater.*, 152, 6–13.
- [14] Wittayakun, J., Khemthong, P., and Prayoonpokarach, S., 2008, Synthesis and characterization of zeolite NaY from rice husk silica, *Korean J. Chem. Eng.*, 25 (4), 861–864.
- [15] Faghihian, H., and Godazandeha, N., 2009, Synthesis of nano crystalline zeolite Y from bentonite, *J. Porous Mater.*, 16 (3), 331–335.
- [16] Almeida, R.K.S., Gómez-Hortigüela, L., Pinar, A.B., and Pérez-Pariente, J., 2016, Synthesis of ferrierite by a new combination of co-structure-directing agents: 1,6-bis(N-methylpyrrolidinium)hexane and tetramethylammonium, *Microporous Mesoporous Mater.*, 232, 218–226.
- [17] Lee, H.J., Kim, J., Kim, J.H., Park, S.J., and Cho, S.J., 2014, Synthesis and characterization of zeolites MTT and MFI, with controlled morphologies using mixed structure directing agents, *Microporous Mesoporous Mater.*, 195, 205–215.
- [18] Hasan, F., Singh, R., Li, G., Zhao, D., and Webley, P.A., 2012, Direct synthesis of hierarchical LTA zeolite via a low crystallization and growth rate technique in presence of cetyltrimethylammonium bromide, *J. Colloid Interface Sci.*, 382 (1), 1–12.
- [19] Su, S., Ma, H., and Chuan, X., 2016, Hydrothermal synthesis of zeolite A from K-feldspar and its crystallization mechanism, *Adv. Powder Technol.*, 27 (1), 139–144.
- [20] Weitkamp, J., and Puppe, L., 1999, *Catalysis and Zeolites: Fundamentals and Applications*, 2nd ed., Springer-Verlag Berlin Heidelberg.
- [21] Treacy, M.M.J., and Higgins, J.B., 2001, *Collection of Simulated XRD Powder Pattern for Zeolite*, 4th ed., Elsevier Science.
- [22] Wang, L., Yang, W., Xin, C., Ling, F., Sun, W., Fang, X., and Yang, R., 2012, Synthesis of nano-zeolite IM-5 by hydrothermal method with the aid of PEG and CTAB, *Mater. Lett.*, 69, 16–19.

Synthesis of Zeolite from Sugarcane Bagasse Ash Using Cetyltrimethylammonium Bromide as Structure Directing Agent

ORIGINALITY REPORT

18%

SIMILARITY INDEX

2%

INTERNET SOURCES

16%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|-------|--|----|
| 1 | Amit Bhatnagar, Kavindra Kumar Kesari, Narasinha Shurpali. "Multidisciplinary Approaches to Handling Wastes in Sugar Industries", Water, Air, & Soil Pollution, 2015
<small>Publication</small> | 2% |
| <hr/> | | |
| 2 | "Mesostructured Zeolites", Green Chemistry and Sustainable Technology, 2016.
<small>Publication</small> | 1% |
| <hr/> | | |
| 3 | Qianqian Yu, Zhigang Huang, Houxiang Sun, Lei Li, Xiaochun Zhu, Shenyong Ren, Baojian Shen. " Investigation on -Alkane Hydroisomerization, a Comparison of IM-5 to ZSM-5 Zeolites ", Industrial & Engineering Chemistry Research, 2018
<small>Publication</small> | 1% |
| <hr/> | | |
| 4 | ijesrt.com
<small>Internet Source</small> | 1% |
| <hr/> | | |
| 5 | Mohamed A. Moneim, Ezzat A. Ahmed. "Synthesis of Faujasite from Egyptian Clays: | 1% |

Characterizations and Removal of Heavy Metals", Geomaterials, 2015

Publication

6

Hongqiang Jin, Yang Liu, Chunyang Wang, Xunhui Lei, Min Guo, Fangqin Cheng, Mei Zhang. "Two-step modification towards enhancing the adsorption capacity of fly ash for both inorganic Cu(II) and organic methylene blue from aqueous solution", Environmental Science and Pollution Research, 2018

Publication

1%

7

Pham Thanh Huyen. "chapter 7 Clay Minerals Converted to Porous Materials and Their Application", IGI Global, 2017

Publication

1%

8

Sujata Subedi, Gabriel Arce, Marwa Hassan, Nitin Kumar, Michele Barbato, Maria Teresa Gutierrez-Wing. "Influence of Production Methodology on the Pozzolanic Activity of Sugarcane Bagasse Ash", MATEC Web of Conferences, 2019

Publication

1%

9

"Insights into the Chemistry of Organic Structure-Directing Agents in the Synthesis of Zeolitic Materials", Springer Nature, 2018

Publication

1%

10

Lijalem Ayele, Joaquín Pérez-Pariente, Yonas

Chebude, Isabel Díaz. "Conventional versus alkali fusion synthesis of zeolite A from low grade kaolin", Applied Clay Science, 2016

Publication

1 %

11

Yong Liu, Qiong Luo, Guodong Wang, Xianlong Li, Ping Na. "Synthesis and characterization of zeolite from coal fly ash", Materials Research Express, 2018

Publication

1 %

12

Eduardo J. M. de Paiva, Vinícius Kothe, Marcos Lúcio Corazza, Ângela Silva et al. "Chapter 12 General Assessment of the Currently Available Biodiesel Production Technologies", Springer Science and Business Media LLC, 2016

Publication

1 %

13

"Pollutants from Energy Sources", Springer Science and Business Media LLC, 2019

Publication

1 %

14

Nurudeen Salahudeen, Abdulkarim S. Ahmed. "Synthesis of hexagonal zeolite Y from Kankara kaolin using a split technique", Journal of Inclusion Phenomena and Macrocyclic Chemistry, 2016

Publication

1 %

15

Yong Liu, Guodong Wang, Qiong Luo, Xianlong Li, Zhao Wang. "The thermodynamics and kinetics for the removal of copper and nickel

1 %

ions by the zeolite Y synthesized from fly ash",
Materials Research Express, 2018

Publication

16

Hyun Jeong Lee, Su Hyun Kim, Joong Hyun Kim, Sung Jun Park, Sung June Cho. "Synthesis and characterization of zeolites MTT and MFI, with controlled morphologies using mixed structure directing agents", Microporous and Mesoporous Materials, 2014

Publication

<1 %

17

Changquan Zhang, Suqin Li. "Utilization of iron ore tailing for the synthesis of zeolite A by hydrothermal method", Journal of Material Cycles and Waste Management, 2018

Publication

<1 %

18

journal.uin-alauddin.ac.id

Internet Source

<1 %

19

Qing Xu, Tao Ji, San-Ji Gao, Zhengxian Yang, Nengsen Wu. "Characteristics and Applications of Sugar Cane Bagasse Ash Waste in Cementitious Materials", Materials, 2018

Publication

<1 %

20

Lee, Hyun Jeong, Su Hyun Kim, Joong Hyun Kim, Sung Jun Park, and Sung June Cho. "Synthesis and Characterization of Zeolites MTT and MFI, with Controlled Morphologies Using Mixed Structure Directing Agents", Microporous

<1 %

21

Min Bum Park, Sung June Cho, Suk Bong Hong. "Synthesis of Aluminosilicate and Gallosilicate Zeolites via a Charge Density Mismatch Approach and Their Characterization", Journal of the American Chemical Society, 2011

Publication

<1 %

22

Anderson Joel Schwanke, Rosana Balzer, Sibeles Pergher. "Chapter 43 Microporous and Mesoporous Materials from Natural and Inexpensive Sources", Springer Nature, 2019

Publication

<1 %

23

Bagci, E.. "Adsorption behaviour of xanthate and dithiophosphate from their mixtures on chalcopyrite", Minerals Engineering, 200708

Publication

<1 %

24

Asghar Zamani, Ahmad Poursattar Marjani, Nasim Abdollahpour. "Synthesis of high surface area boehmite and alumina by using walnut shell as template", International Journal of Nano and Biomaterials, 2019

Publication

<1 %

25

P S Utama, E Saputra, Khairat. "Effective utilizations of palm oil mill fly ash for synthetic amorphous silica and carbon zeolite composite

<1 %

26

Rongchapo, Wina, Onsulang Sophiphun,
Kamolwan Rintramee, Sanchai
Prayoonpokarach, and Jatuporn Wittayakun.
"Paraquat adsorption on porous materials
synthesized from rice husk silica", Water
Science & Technology, 2013.

Publication

27

S. Norsuraya, H. Fazlena, R. Norhasyimi.
"Sugarcane Bagasse as a Renewable Source of
Silica to Synthesize Santa Barbara Amorphous-
15 (SBA-15)", Procedia Engineering, 2016

Publication

28

Moisés, Murilo Pereira, Paula Pomaro de
Almeida, Cleiser Thiago Pereira da Silva,
Andrelson Wellington Rinaldi, Emerson Marcelo
Giroto, Joziane Gimenes Meneguín, Pedro
Augusto Arroyo, Ricardo Eugenio Bazan, Silvia
Luciana Fávaro, and Eduardo Radovanovic.
"Synthesis of zeolite from multilayer food
packing and sugar cane bagasse ash for CO₂
adsorption", RSC Advances, 2014.

Publication

29

Ana B. Pinar, Luis Gómez-Hortigüela, Lynne B.
McCusker, Joaquín Pérez-Pariente. "Controlling

<1 %

<1 %

<1 %

<1 %

the Aluminum Distribution in the Zeolite
Ferrierite via the Organic Structure Directing
Agent", Chemistry of Materials, 2013

Publication

30

Jatuporn Wittayakun. "Synthesis and
characterization of zeolite NaY from rice husk
silica", Korean Journal of Chemical Engineering,
07/2008

Publication

<1%

31

J. Theron, J. A. Walker, T. E. Cloete.
"Nanotechnology and Water Treatment:
Applications and Emerging Opportunities",
Critical Reviews in Microbiology, 2008

Publication

<1%

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off